

**Author's pre-print version, manuscript accepted at Brit J Pharmacol , Br J Pharmacol. 2016 Sep; 173(18):2671-701. doi: 10.1111/bph.13533. Epub 2016 Aug 10. Review. PMID: 27278329**

International union of basic and clinical pharmacology review

Acid-sensing ion channel and epithelial Na<sup>+</sup> channel nomenclature review: IUPHAR Review

Emilie Boscardin\*, Omar Alijevic\*, Edith Hummler, Simona Frateschi<sup>°</sup> and Stephan Kellenberger<sup>°</sup>

Department of Pharmacology and Toxicology, University of Lausanne, Lausanne, Switzerland

\*, these authors contributed equally to the manuscript

<sup>°</sup>, shared corresponding authorship

Running head: ASIC and ENaC nomenclature

Corresponding authors:

Stephan Kellenberger, [Stephan.Kellenberger@unil.ch](mailto:Stephan.Kellenberger@unil.ch), phone ++4121 692 5422 and Simona Frateschi, [Simona.Frateschi@unil.ch](mailto:Simona.Frateschi@unil.ch), phone ++41216925426, fax ++4121 692 5355, Department of Pharmacology and Toxicology, University of Lausanne, Rue du Bugnon 27, CH-1011 Lausanne, Switzerland

## Abstract

Acid-sensing ion channels (ASICs) and the Epithelial Na<sup>+</sup> channel (ENaC) are both members of the ENaC/DEG (degenerin) family of amiloride-sensitive Na<sup>+</sup> channels. ASICs act as proton sensors in the nervous system where they contribute, besides other roles, to fear behavior, learning and pain sensation. ENaC mediates Na<sup>+</sup> reabsorption across epithelia of the distal kidney and colon and of the airways. ENaC is a clinically used drug target in the context of hypertension and cystic fibrosis, while ASIC is an interesting potential target. Following a brief introduction we will review here selected aspects of ASIC and ENaC function. We discuss the origin and nature of pH changes in the brain and the involvement of ASICs in synaptic signaling. We expose how in the peripheral nervous system, ASICs cover together with other ion channels a wide pH range as proton sensors. We introduce the

mechanisms of aldosterone-dependent ENaC regulation and the evidence for an aldosterone-independent control of ENaC activity, such as regulation by dietary  $K^+$ . We then provide an overview of the regulation of ENaC by proteases, a topic with increasing interest over the last years. In spite of the profound differences in the physiological and pathological roles of ASICs and ENaC, these channels share many basic functional and structural properties. It is likely that further research will identify physiological contexts in which ASICs and ENaC have similar or overlapping roles.

## Abbreviations

AQP, aquaporin; ASDN, aldosterone-sensitive distal nephron; ASIC, acid-sensing ion channel; BASIC, bile acid-activated ion channel; BK, big calcium-activated  $K^+$  channel; CA, carbonic anhydrase; CAP-1, -2, or -3, channel activating proteases;  $Ca_v$ , voltage-gated  $Ca^{2+}$  channel; CCD, cortical collecting duct; CD, collecting duct; PHA-1, pseudohypoaldosteronism type 1; CFTR, cystic fibrosis transmembrane conductance regulator; CNT, connecting tubule; CF, cystic fibrosis; CLCN/Kb, voltage-sensitive chloride channel Kb; DCT, distal convoluted tubule; DRG, dorsal root ganglion; ENaC, amiloride-sensitive epithelial sodium channel; EPSP, excitatory post-synaptic potential; FaNaC, FMRFa-activated  $Na^+$  channel; FMRFa, Phe-Met-Arg-Phe-amide; GMQ, 2-guanidine-4-methylquinazoline; GPI, glycosylphosphatidyl-inositol; HAI, hepatocyte growth factor activator inhibitor; HCN, hyperpolarization-activated cyclic nucleotide-gated channel;  $I_A$ , A-type current of rapid inactivating  $K^+$  channels; iGluR, ionotropic glutamate receptor;  $I_K$ ,  $K^+$  current;  $I_{Na}$ ,  $Na^+$  current;  $I_h$ , current produced by HCN channels;  $I_{max}$ , maximal current amplitude;  $K_{ir}$ , inward rectifier  $K^+$  channel;  $K_v$ , voltage-gated  $K^+$  channel; MR, mineralocorticoid receptor;  $Na_v$ , voltage-gated  $Na^+$  channel; NBC,  $Na^+$ ,  $-HCO_3^-$  cotransporter; NCC,  $Na^+$ - $Cl^-$  cotransporter; NHE,  $Na^+$ - $H^+$  exchanger; OSR1/SPAK, Ste20-related protein kinases; P2X, purinergic receptor; PcTx1, Psalmotoxin 1;  $pH_{50}$ , pH of half-maximal effect; pHe, extracellular pH;  $pH_{50}Inh./Act.$ , pH of half-maximal inhibition/activation; PMCA, plasma membrane  $Ca^{2+}$ -ATPase; PNS, peripheral nervous system; PPK, pickpocket; ROMK, renal outer medullary potassium channel; SPLUNC1, the short palate, lung, and nasal epithelial clone 1; TASK, two-pore domain  $K^+$  channel; TRAAK, TWIK-related arachidonic acid-stimulated  $K^+$  channel; TREK, TWIK-related  $K^+$  channel; TRPM, transient receptor potential cation channel, subfamily M; TRPV, transient receptor potential cation channel subfamily V; TWIK, tandem of P-domain in a weak inwardly rectifying  $K^+$  channel; V-ATPase, vacuolar-type  $H^+$ -ATPase.

## Introduction

The ENaC/DEG superfamily of ion channels includes besides ENaC and ASICs the DEGs that are part of mechanotransduction complexes in *C. elegans* (Arnadottir, O'Hagan, Chen, Goodman & Chalfie, 2011), the peptide-gated channel FaNaC of snails (Lingueglia, Deval & Lazdunski, 2006), the mammalian “bile acid-sensitive ion channel” BASIC, (Wiemuth, Assmann & Grunder, 2014) as well as *Drosophila* ENaC/DEG channels such as pickpocket (PPK), ripped pocket and others (Adams, Anderson, Motto, Price, Johnson & Welsh, 1998) (Fig. 1A). The amino acid sequence identity between different ENaC/DEG subfamilies is 15-20%.

ASICs and FaNaC are expressed in the nervous system, DEGs are present in touch-sensitive neurons, BASIC shows highest expression in the brain, the liver and the intestine, and ENaC is found at highest levels in tight epithelia, while the *Drosophila* ENaC/DEG members are probably expressed in many different tissues. FaNaC is an excitatory ion channel of the nervous system of snails, DEGs are critical for *C. elegans* touch sensation, and *Drosophila* ENaC/DEG members may also be involved in touch sensation, among other roles. BASIC is activated by bile acids, its physiological role is however currently not known. ASICs are involved in fear behaviors, learning and memory functions, as well as pain sensation (Fig. 1B). They also contribute to neurodegeneration after ischemic stroke (rev. in (Kellenberger & Schild, 2015; Wemmie, Taugher & Kreple, 2013)). ENaC plays a well-established role in Na<sup>+</sup> reabsorption in the distal nephron and the distal colon and on airway epithelia, is involved in salt taste perception, epidermal differentiation and in skin barrier function (Fig. 1B).

Crystal structures of chicken ASIC1, which was in some studies co-crystallized with the ASIC toxins Psalmotoxin 1 (PcTx1) or Mit-Toxin, showed a channel made by three subunits (Bacongus, Bohlen, Goehring, Julius & Gouaux, 2014; Bacongus & Gouaux, 2012; Dawson et al., 2012; Gonzales, Kawate & Gouaux, 2009; Jasti, Furukawa, Gonzales & Gouaux, 2007). The shape of each subunit was compared to that of a hand holding a small ball, and accordingly the different extracellular domains were labeled palm, knuckle, finger, thumb and  $\beta$ -ball (Fig. 2A). The palm domain is the extracellular continuation of the transmembrane segments and forms a  $\beta$ -strand-rich scaffold of the extracellular channel part. The knuckle and  $\beta$ -ball are located on top and along the upper half of the palm, respectively (Fig. 2A-B). The finger and thumb are oriented towards the outside of the protein. Details of the crystal structures and their differences have been recently discussed (Grunder & Augustinowski,

2012; Kellenberger & Grutter, 2015; Kellenberger & Schild, 2015). Structure-function studies indicate that the ENaC and ASIC ectodomains play important roles in controlling the opening of the channel pore (rev. in (Kellenberger & Schild, 2015)). The sequence homology suggests that all ENaC/DEG members share the same subunit topology. Models of ENaC subunits have been constructed based on the ASIC crystal structures. The highest homology of the ectodomain between ASICs and ENaC is found in the palm and the  $\beta$ -ball (Kashlan et al., 2011; Kashlan & Kleyman, 2011). The predicted secondary structures of most other ENaC domains match the ASIC structure moderately well except for the finger that has the lowest homology, and contains a ~80 amino acid insertion in ENaC (Fig. 2C).

Stoichiometry predictions of ENaC that were based on functional and biochemical data strongly suggested that the functional ENaC at the plasma membrane is a tetramer (Anantharam & Palmer, 2007; Firsov, Gautschi, Merillat, Rossier & Schild, 1998; Kosari, Sheng, Li, Mak, Foskett & Kleyman, 1998). In contrast, all crystal structures describe ASIC as a trimer. In a recent study ASIC1a and ASIC2a containing fluorescently labeled subunits were expressed in *Xenopus* oocytes and the number of bleaching steps of plasma membrane-resident channels was counted to determine the subunit stoichiometry of these functional channels at the cell surface. This study concluded that the ASICs are trimers (Bartoi, Augustinowski, Polleichtner, Grunder & Ulbrich, 2014). A recent biochemical study shows however evidence for the existence of ASIC1a tetramers (van Bemmelen, Huser, Gautschi & Schild, 2015). The question of the ENaC/DEG subunit stoichiometry remains currently controversial.

## ASICs

### *Basic information on ASICs*

#### Physiological and pathological roles of ASICs

Of the ASIC subunits that participate in the formation of H<sup>+</sup>-activated channels, ASIC1a, -2a and -2b are widely expressed in the CNS; all ASIC subunits are found in the peripheral nervous system (PNS) (rev. in (Kellenberger & Schild, 2015; Wemmie, Taugher & Kreple, 2013). Since ASICs are Na<sup>+</sup>-selective ion channels, their activation is expected to induce a neuronal depolarization. Indeed, activation of ASICs in neurons of the CNS and PNS induces membrane depolarization and action potential induction (Fig. 3A) (Deval, Baron, Lingueglia, Mazarguil, Zajac & Lazdunski, 2003; Poirot, Berta, Decosterd & Kellenberger, 2006; Vukicevic & Kellenberger, 2004). ASIC1a shows in addition to its Na<sup>+</sup> permeability a small

permeability for  $\text{Ca}^{2+}$  that is likely important for some of its roles (Bassler, Ngo-Anh, Geisler, Ruppertsberg & Grunder, 2001; Boillat, Alijevic & Kellenberger, 2014; Waldmann, Champigny, Bassilana, Heurteaux & Lazdunski, 1997).

Synaptic signaling involves an acidification of the synaptic cleft, which can activate ASICs. Disruption of the ASIC1a-encoding gene in mice eliminated most of the ASIC currents of CNS neurons, impaired long-term potentiation in the hippocampus and induced a mild deficit in spatial learning (Wemmie et al., 2002). Recent studies have added important information on the role of ASICs in synaptic functions. ASIC1a is highly expressed in the amygdala, and there is strong evidence that ASIC1a of the amygdala contributes to fear behavior (Wemmie, Taugher & Kreple, 2013). Disruption of ASIC1a or inhibition of ASIC1a activity in the brain by PcTx1-containing venom reduced the infarct volume in an experimental stroke model by >50%, strongly suggesting that ASIC1a-mediated  $\text{Ca}^{2+}$  entry contributes to neurodegeneration in this situation (Xiong et al., 2004). Further studies showed that disruption of ASIC1a had a protective effect in several neurodegenerative diseases, including multiple sclerosis, and Huntington's and Parkinson's disease (rev. in (Wemmie, Taugher & Kreple, 2013)). The extracellular pH is lowered in inflammation and ischemia, which both involve pain. There is also strong evidence for a role of sensory neuron ASICs in pain sensation. ASICs belong to the same ion channel family as the *C. elegans* degenerins that form the channel parts of mechanotransduction complexes. A large number of studies with ASICs provide evidence for a regulatory role of ASICs in mechanosensation in several tissues and organs (rev. in (Omerbasic, Schuhmacher, Bernal Sierra, Smith & Lewin, 2015)).

### ASIC function and regulation

Exposure of ASICs to an acidic extracellular pH leads to rapid channel opening, followed by a slower entry into a non-conducting desensitized state. This results in a transient current (Fig. 3B). In some ASIC subtypes, such as ASIC3 and some heteromeric ASICs, desensitization is not complete and a small sustained current persists after the initial peak (Lingueglia et al., 1997; Waldmann, Champigny, Bassilana, Heurteaux & Lazdunski, 1997; Yagi, Wenk, Naves & McCleskey, 2006). Desensitization can also occur without apparent channel opening (termed steady-state desensitization in this case) during moderate lowering of the pH, and can limit the availability of ASICs for opening. The pH dependence of these two processes, channel activation and steady-state desensitization, is illustrated at the example of ASIC1a in Fig. 3C. The steady-state desensitization occurs at  $\text{pH} < 7.4$ , with a mid-point, termed pH of half-maximal desensitization of  $\sim 7.15$ . ASIC1a opening occurs at  $\text{pH} < 7$  and is characterized

by a pH of half-maximal activation ( $\text{pH}_{50}$ ) of  $\sim 6.5$ . These parameters determine the open probability of ASICs under given pH conditions. As shown in Table 2, ASIC1a and ASIC3 are the most sensitive ASICs and are activated by acidification to pH values only slightly below pH 7. In contrast, ASIC2a needs much more acidic pH ( $< 5.5$ ) for activation. ASIC1a is the ASIC isoform that is most sensitive to desensitization. It shows partial desensitization already at pH slightly below 7.4 (Fig. 3C and Table 2).

For several ASICs it has been shown with fast solution changes that the opening time constant at a pH that fully activates the channels is of the order of  $\sim 10$  ms (Bassler, Ngo-Anh, Geisler, Ruppertsberg & Grunder, 2001). The kinetics of current desensitization depend on the subunit composition, as shown in Table 2. Besides protons there are only very few ASIC activators known so far. The small synthetic molecule 2-guanidine-4-methylquinazoline (GMQ) activates ASIC3 at pH 7.4 and inhibits other ASIC subtypes by changing the pH dependence of activation and of steady-state desensitization (Alijevic & Kellenberger, 2012; Yu et al., 2010). Interestingly, endogenous arginine metabolites that contain a guanidinium group as does GMQ were found to have similar effects as GMQ (Li, Yu, Zhang, Cao & Xu, 2011).

ASIC activity is regulated by many different modulators, such as divalent and polyvalent cations, neuropeptides, arachidonic acid, protein kinases and proteases, as summarized in Table 3. Many of these regulatory mechanisms are likely active under physiological conditions. Divalent and polyvalent cations such as  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and spermine appear to compete with protons for binding sites, since an increase in their concentration shifts the ASIC pH dependence towards more acidic pH values. The peptide Phe-Met-Arg-Phe-amide (FMRFa) activates the ENaC/degenerin family member FaNaC (Lingueglia, Champigny, Lazdunski & Barbry, 1995). FMRFa and related mammalian neuropeptides slow the desensitization kinetics of ASIC1 and ASIC3 and induce a sustained current, with  $\text{EC}_{50}$  values that are for most peptides on the order of 10-50  $\mu\text{M}$  (Askwith, Cheng, Ikuma, Benson, Price & Welsh, 2000; Vick & Askwith, 2015). Arachidonic acid, whose tissue concentration is increased in ischemia and inflammation, potentiates ASIC currents in the CNS and the PNS by mechanisms that include an alkaline shift of the pH dependence of activation (Allen & Attwell, 2002; Deval et al., 2008; Smith, Cadiou & McNaughton, 2007). Trypsin was shown to shift the ASIC1a pH dependence of activation and steady-state desensitization to more acidic values by a cleavage in the ectodomain. This led to reduced acidification-induced ASIC currents and neuronal signaling at a physiological conditioning pH 7.4, and to an increased activity if the conditioning pH was slightly reduced (Vukicevic & Kellenberger, 2004; Vukicevic, Weder, Boillat, Boesch & Kellenberger, 2006). The serine protease tissue

kallikrein was also shown to cleave and regulate ASIC1a (Su, Tang, Liu, Zhou & Dong, 2011). Finally, ASIC function is modulated by interaction with other proteins, as discussed in (Wemmie, Price & Welsh, 2006) and updated in (Kellenberger & Schild, 2015).

### ASIC pharmacology

It needs to be noted that currently ASIC inhibition is not used clinically. The presented compounds except for amiloride, which is clinically used as ENaC inhibitor, have been characterized in cell systems and in part also in animal models. An interesting recent review of ASIC pharmacology is provided by (Baron & Lingueglia, 2015). Based on the known physiological and pathological roles of ASICs, drugs inhibiting them could have many uses, such as analgesia, neuroprotection after ischemic stroke, and treatment of anxiety.

Amiloride, the prototype ASIC small molecule blocker has a low potency ( $EC_{50}$  of 10-100  $\mu\text{M}$ ) and selectivity on ASIC peak currents and does not inhibit the sustained ASIC currents. Amiloride binds into the pore of ENaC and ASICs (Adams, Snyder & Welsh, 1999; Alijevic & Kellenberger, 2012; Schild, Schneeberger, Gautschi & Firsov, 1997). The site of action of other small molecule inhibitors on ASICs is not known. Amiloride derivatives modified at the 5-position of the pyrazine ring by hydrophobic groups increased the potency for ASIC3 inhibition by up to 100-fold (Kuduk et al., 2009). Nafamostat mesylate, an anti-inflammatory agent and protease inhibitor, contains a guanidinium moiety as do amiloride and GMQ, and was shown to inhibit ASIC currents, including the sustained current of ASIC3, with  $IC_{50}$  values of 2-70  $\mu\text{M}$  (Ugawa, Ishida, Ueda, Inoue, Nagao & Shimada, 2007). The chemically unrelated compound A-317567 inhibits peak and sustained currents of neuronal and recombinant ASICs with  $IC_{50}$  values between 2 and 30  $\mu\text{M}$  (Dube et al., 2005). Development of A-317567 derivatives yielded substances with higher affinity for ASICs, that lost however some of their selectivity ( $IC_{50}$  on ASIC3 of 400-500 nM, and for other neurotransmitter receptors of < 10  $\mu\text{M}$ ) (Kuduk et al., 2010).

Inflammation increases ASIC mRNA expression, and it was shown that several non-steroidal anti-inflammatory drugs at doses close to those used in clinics prevent or suppress the RNA overexpression (Voilley, de Weille, Mamet & Lazdunski, 2001). These drugs inhibit also ASIC currents, however with potencies that are orders of magnitude lower than that on the cyclooxygenases (Voilley, de Weille, Mamet & Lazdunski, 2001). Several antiprotozoal diarylamidines inhibit ASICs with  $IC_{50}$  of 0.3-38  $\mu\text{M}$  (Chen et al., 2010). A recent screening



of a fragment library followed by optimization led to ASIC3-inhibiting 2-aminopyridine derivatives with an IC<sub>50</sub> of ~3 μM (Wolkenberg et al., 2011).

Venom toxins acting on ASICs have been used to elucidate some of the physiological and pathological roles of ASICs (Wemmie, Taugher & Kreple, 2013). In addition, Complexes of ASIC with toxins were used to determine the crystal structure of ASICs in the presumable open conformation. The most important ASIC toxins are the gating modifiers PcTx1 of the spider *Psalmopoeus cambridgei* and the Mambalgins of the black mamba, the ASIC3 inhibitor APETx2 of the sea anemone *Anthopleura elegantissima* and the activating Mit-toxin of the Texas coral snake that generates a sustained ASIC opening at pH 7.4 (rev. in (Baron, Diochot, Salinas, Deval, Noel & Lingueglia, 2013)). Mit-toxin and Mambalgins target several ASIC subtypes, while PcTx1 is selective for ASIC1a homomers and ASIC1a/2b heteromers, and APETx2 for ASIC3-containing channels. These toxins have so far not been shown to target other channels besides ASICs, with the exception of APETx2 that also inhibits some voltage-gated Na<sup>+</sup> channel isoforms (Blanchard, Rash & Kellenberger, 2012; Peigneur, Beress, Moller, Mari, Forssmann & Tytgat, 2012). PcTx1 inhibits mammalian ASIC1a by an alkaline shift of the pH dependence of steady-state desensitization (leading to complete desensitization at pH 7.4), while Mambalgin inhibition is due to an acidic shift of the pH dependence of activation. The mechanisms of action of the other ASIC toxins are currently not known. Co-crystallization showed that PcTx1 binds to the acidic pocket of ASIC1, and that the much larger Mit-toxin binds to the wrist, palm and thumb domains, without however reaching into the acidic pocket (Baconguis, Bohlen, Goehring, Julius & Gouaux, 2014; Baconguis & Gouaux, 2012; Dawson et al., 2012). Site-directed mutagenesis indicated that Mambalgins also bind to the acidic pocket (Salinas et al., 2014; Schroeder et al., 2014).

### ***pH changes in the brain and role of ASICs in synaptic signaling in the CNS***

#### **Physiological extracellular pH changes during neuronal activity**

The resting extracellular pH (pHe) value in the interstitial fluid of brain tissues can vary between 7.15 and 7.33 depending on the brain area and is generally about 0.1-0.2 pH units more acidic than the blood pH (Mutch & Hansen, 1984; Syková & Svoboda, 1990). Since membrane depolarization, ion transport and metabolic activity affect the pHe, changes in pHe are commonly observed during neuronal activity and in changed metabolic states, such as e.g. ischemia (Chesler, 2003; Dmitriev & Mangel, 2001; Kraig, Ferreira-Filho & Nicholson, 1983; Krishtal, Osipchuk, Shelest & Smirnoff, 1987; Siesjö, von Hanwehr, Nergelius, Nevander & Ingvar, 1985). Active brain neurons generate ATP by oxidative phosphorylation, which leads



to production of CO<sub>2</sub>, while astroglial cells are primarily glycolytic, producing lactate (Deitmer, 2002). Information about the variations of extracellular pH in the CNS of mammals comes mainly from studies with anesthetized rodents and in vitro preparations in which the pHe was recorded using optical techniques or pH-sensitive microelectrodes (Chesler, 2003). Upon repetitive electrical stimulation the neuronal activity of the mammalian CNS produces a general pattern of pHe changes containing three phases (Chesler, 1990; Chesler, 2003; Makani & Chesler, 2010). An initial short (<200ms) transient alkaline shift of typically 0.01-0.2 pH units, occurring within tens of ms after the beginning of the stimulus, is followed by a long-lasting, slowly developing acidosis that reaches a maximal acidic shift of 0.1-0.25 pH units 20-60s after the beginning of the repetitive stimulation. In the third phase, the pH returns to the initial value. Since these measurements represent in most cases tissue pH changes it is possible that locally, as e.g. in synapses, the pH changes may have different amplitudes and kinetics. The brain pH is buffered by the bicarbonate-carbon dioxide buffer. The buffering capacity of the interstitial fluid depends therefore on the bicarbonate and CO<sub>2</sub> concentration and co-determines the magnitude of extracellular pH changes (Du et al., 2014; Highstein, Holstein, Mann & Rabbitt, 2014). The buffer capacity shows regional variations, and may be lower than originally thought (Chesler, 2003; Tong, Chen & Chesler, 2006).

The transient alkalinization mentioned above has been reported in hippocampus in several studies (Chen & Chesler, 2015; Chen & Chesler, 1992; Gottfried & Chesler, 1996; Krishtal, Osipchuk, Shelest & Smirnoff, 1987). Inhibition of the plasma membrane Ca<sup>2+</sup>-ATPase (PMCA) prevented the transient alkaline shift in hippocampal neurons effectively, suggesting that the activity of this transporter, that exchanges internal Ca<sup>2+</sup> for external H<sup>+</sup>, generates the alkaline shift (Kreitzer, Collis, Molina, Smith & Malchow, 2007; Makani & Chesler, 2010) (Makani *et al.*, 2010). Depending on the brain region, the initial alkaline shift can be more or less pronounced or even inexistent (Venton, Michael & Wightman, 2003; Yamamoto, Borgula & Steinberg, 1992). The regional differences in the magnitude of the initial alkaline shift may be due to different expression or activity of PMCA, ion channels and other acid-base transporters and to differences in buffering capacity. The alkalinization occurs in the same time frame as do excitatory post-synaptic currents and is therefore fast enough to modulate fast synaptic transmission (Gottfried & Chesler, 1996). Figure 4 illustrates the different mechanisms leading to pH changes in the synapse. The alkaline shift is also reduced by pharmacological blockade of different ion channels involved in neurotransmission, indicating that it is directly associated with neuronal activity (Smith & Chesler, 1999). This is not the case for the following acidification, which appears to be associated with another

process. Several observations suggest that the acidic change is mostly due to the activity of glial cells (Chesler, 2003; Jendelová & Syková, 1991). It was suggested that the  $\text{Na}^+\text{-HCO}_3^-$ -cotransporter NBC contributes to the lowering of the extracellular pH by the glia. Activation of NBC leads to transport of  $\text{Na}^+$  and  $\text{HCO}_3^-$  into the glial cell and thereby to extracellular acidification (Deitmer, 2002) (Fig 4).

#### Protons and ASICs, a neurotransmitter-receptor pair in the CNS?

In this section we present the experimental evidence for a role of protons as a neurotransmitter. We discuss then how ASICs contribute to synaptic signaling and long-term potentiation (LTP), thereby affecting learning and fear sensation. Finally, we review different observations suggesting that ASICs can strongly influence glutamate receptor function.

Indirect evidence for synaptic cleft acidification during electrical signaling has been provided by studies that measured the inhibition of presynaptic voltage-gated  $\text{Ca}^{2+}$  channel currents - that are inhibited by protons - and estimated the acidification in the synaptic cleft to 0.2-0.6 pH units (DeVries, 2001; Palmer, Hull, Vigh & von Gersdorff, 2003; Vessey et al., 2005). Recent studies with fluorescent pH indicators showed that presynaptic stimulation led to acidification of the synaptic cleft in synapses formed by vestibular hair cells and the calyx nerve terminal of the turtle *Trachemys scripta elegans* (Highstein, Holstein, Mann & Rabbitt, 2014) and in synapses of mouse lateral amygdala (Du *et al.*, 2014). These pH patterns differed from the alkalinization-acidification observed in earlier studies. The differences might be due to the different brain regions that were investigated or due to the configuration of the measurements. Several mechanisms may contribute to the acidification of the synaptic cleft. The synaptic vesicles containing neurotransmitters that are continuously replenished with protons by the  $\text{H}^+$ -ATPase pumps are an important source of protons in synapses (Beyenbach & Wiczorek, 2006; Vavassori & Mayer, 2014). There is evidence that the pH of cholinergic, glutamatergic and GABAergic synaptic vesicles can reach values as acidic as pH  $\sim 5.5$  (DeVries, 2001; Dietrich & Morad, 2010; Fuldner & Stadler, 1982; Michaelson & Angel, 1980; Miesenbock, De Angelis & Rothman, 1998; Palmer, Hull, Vigh & von Gersdorff, 2003). The synaptic vesicles fuse with the presynaptic neuronal plasma membrane during neurotransmission and release their acidic content together with the neurotransmitters into the synaptic cleft. The acidification of the synaptic cleft observed in the lateral amygdala had only a small amplitude ( $<0.1$  pH unit), and a time constant of 0.5-1 s, thus much slower than the release of synaptic vesicles (Du *et al.*, 2014). Highstein et al. also observed an acidification time course of  $\sim 0.5$  s (Highstein, Holstein, Mann & Rabbitt, 2014). The slow

kinetics may point to acid-base transporters as the source of this proton release into the synaptic cleft. Consistent with this hypothesis, it was demonstrated in cultured rat cerebellar granule cells that transient synaptic cleft acidification during GABAergic transmission was at least in part due to the activity of the  $\text{Na}^+/\text{H}^+$  exchanger (Dietrich & Morad, 2010).

The activity of many neuronal ion channels is pH-dependent (Table 4), suggesting that the pH changes during neuronal activity modulate ion channel function. In general, alkaline extracellular pH favors inward currents, thus enhancing excitability, while acidic extracellular pH depresses excitability in many circumstances (Chesler, 2003) and can be considered as negative feedback since it is caused by neuronal stimulation. The initial short alkalization is sufficient for relieving the proton block of a significant fraction of NMDA receptors and to activate different classes of voltage-gated calcium channels, thus increasing the glutamatergic excitation and boosting neurotransmitter release (Makani et al., 2012; Tombaugh & Somjen, 1996). ASICs are directly activated by extracellular protons, in contrast to most other excitatory ion channels, which are inhibited by acidification. Administration of specific ASIC1a antagonists or disruption of the ASIC1a gene eliminates the majority of the acid-induced currents in the CNS neurons (Wemmie, Taugher & Kreple, 2013; Wu et al., 2013). This demonstrates that the ASIC1a homomers and ASIC1a-containing heteromers are the principal sensors of rapid extracellular acidification in the brain. ASIC1a, -2a and -2b are widely expressed in the CNS (rev. in (Kellenberger & Schild, 2015; Wemmie, Taugher & Kreple, 2013)). Localization by immunohistochemistry studies, evidence for the interaction with the postsynaptic proteins PICK1 and AKAP150, as well as co-localization with PSD-95 in spines indicate together that ASIC1a has a somatodendritic distribution (Zha, 2013). The ASICs in the postsynaptic site are well situated to detect rapid synaptic pH changes.

A recent study established expression of a light-activated proton pump in mouse astrocytes to induce local extracellular acidification with light with high temporal precision to activate ASICs present on nearby neurons (Li, Yang & Canessa, 2014). Using this approach, efficient action potential induction by repetitive acidification was demonstrated.

In the lateral amygdala, presynaptic stimulation activated postsynaptic ASIC currents. Perfusion of glutamate receptor blockers inhibited 95% of the amplitude of the observed excitatory post-synaptic currents. The remaining 5% of the excitatory post-synaptic current amplitude were mediated by ASICs, since this current was absent in the presence of amiloride or if the ASIC1a gene was deleted (Du *et al.*, 2014). A similar situation with a contribution of ASICs to 5% of the excitatory post-synaptic current amplitude was also found in nucleus accumbens (Kreple et al., 2014). High-frequency stimulation at 100 Hz produced LTP of

excitatory post-synaptic potentials (EPSPs) in the hippocampus of wild type but not ASIC1a<sup>(-/-)</sup> mice (Wemmie et al., 2002). The ASIC1a<sup>(-/-)</sup> mice displayed mildly defective spatial learning and eyeblink conditioning, consistent with decreased LTP. There is currently a controversy regarding the involvement of ASIC1a in hippocampal LTP, since in a different model of ASIC1a knockout mice, in which ASIC1a was deleted at early embryonic stages – in contrast to the classical knockout used in the above study - normal LTP at CA3-CA1 synapses was observed (Wu et al., 2013). The origin of this discrepancy is not understood, but might involve roles of ASIC in synapse development. Another recent study found that pharmacological blockade of ASIC1a in hippocampal synapses impaired LTP, confirming thereby the initial observation by Wemmie et al. (Quintana et al., 2015).

The hippocampal CA1 area is particularly vulnerable to ischemia and undergoes delayed neuronal death in humans and several animal models after an ischemic attack (Kirino, 1982; Kuroiwa, Bonnekoh & Hossmann, 1990). The AMPA receptors show a high degree of post-ischemic plasticity to contribute to the excitotoxicity in the CA1 region by two mechanisms, anoxic LTP during the first hours, and an increased expression of Ca<sup>2+</sup>-permeable AMPA receptor types (lacking the GluA2 subunit) several hours later (Hsu & Huang, 1997; Pellegrini-Giampietro, Zukin, Bennett, Cho & Pulsinelli, 1992). After an oxygen-glucose deprivation, anoxic LTP was observed in organotypic hippocampal slice cultures of wt mice, but was absent in slice cultures of ASIC1a<sup>(-/-)</sup> mice or after pharmacological blockade of ASIC1a (Quintana et al., 2015). The presence of active ASIC1a was also required for the increased expression of Ca<sup>2+</sup>-permeable AMPARs after oxygen-glucose deprivation. Inhibition of ASIC1a or of Ca<sup>2+</sup>-permeable AMPARs was sufficient to protect neurons of the CA1 area, illustrating the important role of ASICs in neurodegeneration in this context.

ASIC1a<sup>(-/-)</sup> mice have reduced innate fear and show deficits in conditioned fear behavior (Wemmie, Taugher & Kreple, 2013). The fear-related behavior is in many cases correlated with CO<sub>2</sub> and acid sensing and was rescued in ASIC1a<sup>(-/-)</sup> mice by injection of a viral vector in the basolateral nuclei of the amygdala that restored ASIC1a expression locally (Coryell et al., 2009; Ziemann et al., 2009). Two recent studies showed that ASIC1a is critical for LTP at the synapses of the fear circuitry between the cortex and the basolateral nuclei of the amygdala (Chiang, Chien, Chen, Yanagawa & Lien, 2015; Du et al., 2014). The study by Chiang et al. investigated LTP at various synapses of amygdala neurons and found that the extent of LTP at different synapses correlated with the ASIC current density in postsynaptic neurons. Cell type-specific deletion of ASIC1a showed that ASIC-dependent LTP is required

at several amygdala synapses for fear learning. ASIC4 does not form functional channels but was shown to downregulate the expression of other ASIC subunits (Donier, Rugiero, Jacob & Wood, 2008). ASIC1a expression is therefore expected to be upregulated in ASIC4<sup>-/-</sup> mice, and it was indeed shown that ASIC4 knockout mice have an increased freezing response (Lin, Chien, Chiang, Liu, Lien & Chen, 2015). A recent study with rats suggests a species difference with regard to the role of ASICs in the fear circuitry. This study showed that ASIC1a activation reduces anxiety in rats, by enhancing inhibition in the basolateral amygdala (Pidoplichko et al., 2014). The reasons for this opposite role of ASICs in fear behavior of mice and rats is currently not understood, and the rat data rely so far on one single study. Given the complexity of the expression and role of ASIC1a in the mouse fear circuitry (Chiang *et al.*, 2015) it seems plausible that the ASIC expression pattern may be different between the fear circuitry of mice and rats. The role of ASICs in fear expressions of humans is currently not known.

Several studies have shown that ASICs interact functionally with glutamate receptors in synaptic signaling, and that ASIC function is required for LTP, as discussed above (Du et al., 2014; Kreple et al., 2014; Quintana et al., 2015; Wemmie et al., 2002). In the initial LTP study in hippocampus it was shown that activation of postsynaptic ASICs removes the Mg<sup>2+</sup> block of NMDA receptors, since LTP was only disrupted in ASIC1a<sup>(-/-)</sup> mice in physiological extracellular Mg<sup>2+</sup> concentrations, but was normal at low Mg<sup>2+</sup> concentrations (Wemmie et al., 2002). This does however not explain the more recent observations in the amygdala, the nucleus accumbens and in hippocampal cultures after oxygen-glucose deprivation. In the amygdala, the absence of ASIC1a decreased the EPSC amplitude only slightly, impaired however the LTP strongly (Du et al., 2014). Similarly, the presence or absence of ASIC1a strongly influenced glutamate receptor function in the nucleus accumbens (Kreple et al., 2014). The mechanism of this functional interaction of ASIC1a with glutamate receptors is not understood. There are however indications that the presence of ASICs can influence the density of dendritic spines and the glutamate receptor composition (Kreple et al., 2014; Zha, Wemmie, Green & Welsh, 2006). An earlier study had shown an interaction between NMDA receptors and ASICs in the context of ischemia by demonstrating that NMDA receptor activity leads to phosphorylation of ASIC1a by CaMKII that enhances ASIC currents and leads to ischemic cell death (Gao et al., 2005).

In summary, pH changes occur in the CNS during neuronal and metabolic activity. The synaptic cleft is acidified upon presynaptic stimulation, leading to the activation of

postsynaptic ASICs. In spite of their small contribution to the postsynaptic currents, ASICs play a critical role in synaptic signaling.

### *ASICs and other ion channels sense the extracellular pH in the PNS*

Nociceptive fibers conduct signals from the periphery to the CNS that are induced by a variety of potential tissue-damaging stimuli such as heat, pressure and chemicals. Many different substances that can modulate this signaling are released during tissue damage and inflammation. Although protons are the smallest modulators, they have dramatic effects on diverse properties of sensory neurons. Many different types of acid sensors are expressed by primary sensory neurons and especially by nociceptors, illustrating the vital importance of acid-base sensing and regulation. Tissue acidification occurs for example during ischemia, inflammation, cancer pain, and in muscle during exercise. It has been proposed that rapid localized pH changes may occur in the environment of peripheral nerves (Martin & Jain, 1994). In this section we review first the roles of ASICs in pain sensation. We then describe the ASIC currents in PNS neurons and provide evidence for species differences. We discuss the apparent paradox, that ASICs sense slowly developing and long-lasting pH changes although their activity is mostly transient. Finally, we present observations showing that ASICs are not the only pH sensors in the PNS and that they cover together with other ion channels proton sensing over a wide pH range.

### *Role of ASICs in pain sensation*

A large body of experimental data underlines the importance of sensory neuron ASICs in acid-induced nociception (Deval & Lingueglia, 2015; Sluka & Gregory, 2015). Studies with human volunteers showed that the pain due to iontophoresis or injection of acid solutions in the skin was inhibited by amiloride and that it followed the pH dependence of ASIC1a and ASIC3 (Jones, Slater, Cadiou, McNaughton & McMahon, 2004; Ugawa, Ueda, Ishida, Nishigaki, Shibata & Shimada, 2002). Deletion of ASIC3 in mice reduced several forms of inflammation- and chronic acidification-related pain (Sluka, Winter & Wemmie, 2009). Intrathecal administration of ASIC3-specific siRNA to rats diminished pain behaviors induced by inflammation or wounding (Deval et al., 2011; Deval et al., 2008). The role of ASIC3 in pain sensation was further confirmed when it was shown that local injection of the ASIC3 activator GMQ induced pain behaviors that depended on the presence of ASIC3 (Yu et al., 2010). The use of toxins showed that ASIC1 channels of the PNS are also involved in pain sensation. Injection of the ASIC-activating Mit-toxin of the Texas coral snake in the mouse paw induced pain that depended on the presence of ASIC1a (Bohlen et al., 2011).

Peripheral injection of Mambalgin, which inhibits several ASIC subtypes, exerted an analgesic action due to its inhibition of ASIC1b (Diochot et al., 2012). ASICs are also expressed in pain-processing areas of the CNS. Administration to the CNS of PcTx1 and of Mambalgin inhibited pain behaviors, indicating a role of central ASICs in pain sensation (Diochot et al., 2012; Mazzuca et al., 2007), in addition to the more obvious role of ASICs as pH sensors in the PNS. Two mouse models in which ASIC currents were suppressed – expression of a dominant negative ASIC3 mutant and a triple ASIC1a/ASIC2/ASIC3 knockout – showed increased pain behavior, indicating that some aspects of the role of ASICs in pain sensation are not understood yet (Kang et al., 2012; Mogil et al., 2005).

The neuropeptides Calcitonin gene-related peptide (CGRP) and substance P contribute to inflammation and are secreted from sensory nerve terminals in response to tissue acidification. CGRP induces vasodilation and substance P promotes increased vascular permeability leading to plasma extravasation during neurogenic inflammation (Hoyer & Bartfai, 2012; Walker, Conner, Poyner & Hay, 2010). Since the neuropeptide secretion is  $Ca^{2+}$ -dependent, TRPV1 and ASICs, which induce  $Ca^{2+}$  entry directly or via depolarization-induced activation of  $Ca_v$ s, may mediate acidification-induced neuropeptide secretion. Experiments using specific pharmacological inhibitors or gene knockout identified TRPV1 but not ASICs as the pH sensors for the acid-induced neuropeptide secretion from sensory neurons (Boillat, Alijevic & Kellenberger, 2014; Fischer, Reeh & Sauer, 2003; Pan, Gonzalez, Chang, Chacko, Wein & Malykhina, 2010; Weller, Reeh & Sauer, 2011). It is likely that the transient ASIC current is too short to induce neuropeptide release, and that a sustained current, as the one mediated by TRPV1 is required.

Peng et al. showed recently that disruption of ASIC3 in mice reduced itch and pain in response to local co-injection of acid and a pruritogen (Peng et al., 2015). A similar effect was obtained when GMQ was injected alone, suggesting that persistent ASIC3 activity may be sufficient to induce itch. Interestingly it has previously been shown that ASIC3 is potentiated by another pruritogen, serotonin (Wang et al., 2013). Whether the pruritogenic effect of serotonin is mediated by ASIC3 has however not been explored yet.

#### Expression pattern and cellular functions of ASICs in the PNS

Expression of the different ASIC subunits has been shown in primary afferent neurons innervating the skin, eye, ear, taste buds, heart, gut, skeletal muscle and joints (rev. in (Kellenberger & Schild, 2015; Wemmie, Taugher & Kreple, 2013)). ASIC function has been measured in small- and large-diameter neurons of the trigeminal, nodose and dorsal root



ganglia of mice and rats (Benson et al., 2002; Boillat, Alijevic & Kellenberger, 2014; Deval & Liguoglia, 2015; Poirot, Berta, Decosterd & Kellenberger, 2006). ASIC-like currents have also been measured from human DRG neurons (Baumann, Chaudhary & Martenson, 2004). Sensory neurons express homo- and heterotrimeric ASICs, producing current subtypes with different pH dependence. In rat DRG neurons,  $pH_{50}$  values ranging from 6.6 to  $<4.5$  were measured (Poirot, Berta, Decosterd & Kellenberger, 2006). The presence of ASIC2 in heteromeric ASICs tends to decrease the pH sensitivity of the channel (Table 4). Rat DRG neurons express heteromeric ASICs of different compositions or homomeric ASIC1a or ASIC3. The proportion of these current subtypes varied between studies (Deval et al., 2008; Poirot, Berta, Decosterd & Kellenberger, 2006). Rat DRG neurons express a high proportion of ASIC1 and ASIC3, with less ASIC2a and -2b (Boillat, Alijevic & Kellenberger, 2014; Deval et al., 2008; Poirot, Berta, Decosterd & Kellenberger, 2006). This is similar to the ASIC expression in human (Delaunay et al., 2012) but different from mouse DRGs that show predominant expression of ASIC2a and -2b (Drew et al., 2004; Hughes, Brierley, Young & Blackshaw, 2007).

Since ASIC currents are mostly transient due to rapid desensitization, they last only a few seconds, even if an acidic extracellular environment persists. In addition, at a pH slightly below the physiological 7.4, some ASIC subtypes desensitize without apparent opening and are not available for a subsequent activation in case of a further acidification. From these observed properties it is expected that ASICs are best adapted to sense rapid changes in pH, but will not report the persistence of a changed pH. If a pH change occurs slowly, as is the case in ischemia and inflammation, it may even desensitize ASICs rather than activating them. Two different mechanisms may allow the ASICs to sense pH in such conditions, 1) the sustained current of some ASICs, and 2) modulation of ASIC function by diverse endogenous mediators. Several studies detected a sustained fraction of ASIC currents in DRG neurons and in cells expressing ASIC3 (Deval et al., 2011; Poirot, Berta, Decosterd & Kellenberger, 2006; Yagi, Wenk, Naves & McCleskey, 2006). Native ASIC currents in DRG neurons are positively modulated by several inflammatory mediators. Serotonin, ATP, lactic acid, arachidonic acid and hypertonicity are able to enhance the proton-induced ASIC3 current, in several cases by changing its pH dependence (Baron & Liguoglia, 2015; Light, Huguen, Zhang, Rainier, Liu & Lee, 2008). Also, the synthetic molecule GMQ and related endogenous polyamines such as agmatine and arcaine have recently been shown to activate a sustained current in ASIC3 at physiological pHe (Li, Yu, Zhang, Cao & Xu, 2011; Yu et al., 2010).

### Extracellular pH modulation of ion channels in nociceptors

In the following paragraph we present the different pH sensors in PNS neurons, organized according to their pH sensitivity. Detailed information is provided in Table 4. The varied expression pattern of ASICs and other pH-sensitive ion channels suggests that pH changes may affect neuronal excitability to different extents in the various subpopulations of nociceptors.

*Fluctuations of the resting pH (pH 7.4-7.3):* members of the two-pore K<sup>+</sup> (K2P) channel family are constitutively open at resting membrane potentials. Their activity codetermines the hyperpolarized resting membrane potential of primary afferent neurons (Kang & Kim, 2006; Lesage & Lazdunski, 2000). Most K2P channels are inhibited by extracellular protons with a pH<sub>50</sub> of ~7.4. A slight lowering of the pH will therefore shift the resting membrane potential towards more positive values (Lesage & Barhanin, 2011)(Table 4). One specific K2P subtype, TREK2, is activated by extracellular protons (Sandoz, Douguet, Chatelain, Lazdunski & Lesage, 2009).

*Mild extracellular acidosis (acidification to pH 7.2-6.4)* inhibits different classes of K<sup>+</sup> channels. Studies with human and rat DRG neurons provided evidence for an acidification-induced depolarization that is mediated by K2P channels in this pH range (Baumann, Chaudhary & Martenson, 2004; Cooper, Johnson & Rau, 2004). For several K2P channels it was shown that their loss of function increased the sensation of pain (rev. in (Mathie & Veale, 2015)). Acidification activates several types of ASICs in DRG neurons (Baumann, Chaudhary & Martenson, 2004; Dirajlal, Pauers & Stucky, 2003; Poirot, Berta, Decosterd & Kellenberger, 2006). In this pH range the native ASIC currents are in many cases mediated by different ASIC heteromers containing ASIC1a and/or ASIC3 or by ASIC1a or ASIC3 homomers.

Mild extracellular acidosis inhibits efficiently glutamatergic (Traynelis & Cull-Candy, 1990; Traynelis & Cull-Candy, 1991) and GABAergic transmission (Valeyev, Hackman, Holohean, Wood, Katz & Davidoff, 1999; Zhai, Peoples & Li, 1998), neutralizing thus the excitatory and inhibitory tone on neurotransmission. Voltage-gated Ca<sup>2+</sup> channels are inhibited by extracellular acidification by mechanisms involving pore block and a shift in voltage dependence (Hille, 1992). Extracellular acidosis reduces in addition voltage-gated Na<sup>+</sup> channel currents (Daumas & Andersen, 1993; Tombaugh & Somjen, 1996).

*Severe extracellular acidosis (acidification to pH < 6.4)* activates the whole repertoire of ASIC subtypes, including heteromers with a strong ASIC2 contribution. ASIC3 and several ASIC heteromers develop a sustained current at severe acidosis below pH 4.5. Thus, severe acidosis provides persistent ASIC-mediated depolarizing inputs. Electrophysiology and Ca<sup>2+</sup> imaging experiments with rat DRG neurons showed that pH 6 induces an ASIC current in 50-70% of DRG neurons (Boillat, Alijevic & Kellenberger, 2014; Deval et al., 2008; Poirot, Berta, Decosterd & Kellenberger, 2006). 40-80 % of the putative nociceptors investigated in these studies showed TRPV1-like activity. TRPV1 is a polymodal ion channel activated by capsaicin and a number of other stimuli including noxious heat (>42°C) endogenous arachidonic acid derivatives, ethanol, camphor and protons (Caterina, Schumacher, Tominaga, Rosen, Levine & Julius, 1997; Mickle, Shepherd & Mohapatra, 2015). TRPV1-mediated sustained currents due to acidification are largely confined to sensory neurons with unmyelinated fibers and have a pH<sub>50</sub> of ~5.4. Therefore, TRPV1 activation by protons alone requires severely acidic conditions (pH≤6). TRPV1 is however also modulated by protons and has a substantially higher pH sensitivity for modulation (pH<sub>50</sub> ~7–6). Mild acidosis can therefore sensitize the TRPV1 to other stimuli such as capsaicin and temperature. As a consequence the temperature threshold of TRPV1 activation is lowered at acidic pH, leading to TRPV1 activation at normal body temperature (Tominaga et al., 1998). DRG neuron recordings of TRPV1 null mice showed a complete loss of capsaicin sensitivity and an impairment of thermal and pH sensitivity. Behavioral studies demonstrated an important role of TRPV1 in nociception associated with thermal/mechanical hypersensitivity, with models of inflammation and with different chronic pain conditions (Barton et al., 2006; Caterina et al., 2000; Honore et al., 2005; Keeble, Russell, Curtis, Starr, Pinter & Brain, 2005). Approximately 20 % of measured DRG neurons presented pH-sensitive inward currents that were not mediated by ASICs or TRPV1, indicating that they were mediated by other ion channels, as e.g. by the inhibition of K2P channels that would appear like an inward current. Upon acidification to pH 6, ASICs mediated larger currents and charge movements than TRPV1, due to their different pH sensitivity (Blanchard & Kellenberger, 2011; Poirot, Berta, Decosterd & Kellenberger, 2006).

P2X are trimeric cation channels that are activated by extracellular ATP. P2X2 and P2X3 are highly expressed in nociceptors (Chen, Akopian, Sivilotti, Colquhoun, Burnstock & Wood, 1995; Lewis, Neidhart, Holy, North, Buell & Surprenant, 1995). Disrupting P2X3 in mice decreased nociceptive behavior (Cockayne et al., 2000; Souslova et al., 2000). P2X receptor

activity is modulated by pH changes in sensory neurons. Extracellular acidification increases the P2X2 currents and inhibits P2X3 currents at low, and increases them at high ATP concentrations (Gerevich et al., 2007; King, Wildman, Ziganshina, Pintor & Burnstock, 1997). Simultaneous ATP release and extracellular lactic acidification occur in skeletal muscle under exercise. Interestingly, there is evidence for a physical interaction between P2X receptors and ASICs and it was shown that ATP-induced P2X2 activation potentiates the acid-induced ASIC3 current by twofold. It is therefore likely that ATP and acidosis can converge to increase nociceptor excitability (Birdsong et al., 2010).

## ENaC

### *Function and regulation of ENaC*

ENaC is composed of three different subunits, namely  $\alpha$ ,  $\beta$  and  $\gamma$ ENaC, encoded by three different genes (SCNN1A, SCNN1B and SCNN1G, respectively; see for review (Kellenberger & Schild, 2015)). In the kidney, ENaC is present in the aldosterone-sensitive distal nephron (ASDN), that is composed of the distal convoluted tubule (DCT), connecting tubule (CNT), and collecting duct (CD) segments (Fig. 6) (Rossier, Staub & Hummler, 2013), to allow fine-tuning of whole body  $\text{Na}^+$  homeostasis. Expression of ENaC is regulated transcriptionally but equally by post-translational modifications, which thereby determine its synthesis, trafficking and activity at the cell membrane (Rossier, 2014). This regulation occurs on the ENaC subunits themselves (Loffing et al., 2001), and/or on ENaC-regulating proteins such as the e.g. the serum- and glucocorticoid-regulated kinase 1 (Lang & Shumilina, 2013). Aldosterone modulates ENaC activity by transcription-dependent and -independent mechanisms (Thomas, McEneaney & Harvey, 2007). Due to space constraints in this review, we refer to the extensive literature on ENaC regulation (Kellenberger & Schild, 2015; Rossier, Baker & Studer, 2015; Verouti, Boscardin, Hummler & Frateschi, 2015).

### *ENaC function in different tissues*

#### *Role of ENaC in classical and non-classical tissues and organs*

The syndrome Pseudohypoaldosteronism-type-1 (PHA-1) was first described by Cheek and Perry in 1958 (Cheek & Perry, 1958). Patients suffering from PHA-1 present the following symptoms: renal salt wasting, hypovolemia and hypotension, hyperkalemia, metabolic acidosis, high plasma levels of renin and aldosterone, and respiratory illness. PHA-1 is caused by either mutations in mineralocorticoid receptors (autosomal dominant PHA-1) or by loss-of-function mutations in ENaC subunits (autosomal recessive PHA-1) (Furgeson & Linas,

2010). The genetic dissection of ENaC along the mouse nephron highly suggests that the early ASDN (DCT and/or CNT, Fig. 6) is crucial for Na<sup>+</sup> and K<sup>+</sup> balance. Whereas animals with ENaC deficiency in the CD survive well even under salt restriction or K<sup>+</sup> loading (Rubera et al., 2003), ENaC deficiency in the CNT and CD (Christensen et al., 2010), or reduced ENaC expression solely in CNT (Poulsen et al., 2015) alone is sufficient to induce a mild salt-losing phenotype. However, only the deletion of ENaC subunit expression in mice along the whole nephron mimics the severe human adult PHA-1 (Perrier et al., 2015). An additional relevant role of ENaC in the cortical collecting duct (CCD) can currently not be excluded. Table 5 provides an overview of studies with ENaC mouse models. The constitutive  $\alpha$ ENaC knockout mice develop respiratory distress syndrome and die soon after birth thus revealing an important role of the  $\alpha$ ENaC subunit in lung liquid clearance at birth (Hummler et al., 1996). The  $\gamma$ ENaC subunit seems to transiently facilitate neonatal lung liquid clearance at birth, suggesting equally important roles of the other ENaC subunits (Barker et al., 1998). Increased airway epithelial Na<sup>+</sup> absorption produces cystic fibrosis-like lung disease in transgenic mice (Mall, Grubb, Harkema, O'Neal & Boucher, 2004), unveiling an important role of ENaC in the regulation of the airway surface liquid volume (Mall et al., 2010). In colon,  $\alpha$ ENaC-deficiency causes Na<sup>+</sup> loss and aldosterone resistance, that is normally compensated by the renin-angiotensin-aldosterone system in the kidney (Malsure et al., 2014), thus these mice are able to maintain their Na<sup>+</sup> and K<sup>+</sup> homeostasis.

Apart from its *classical* role as a membrane constituent of many salt-reabsorbing epithelia that facilitates Na<sup>+</sup> movement across the tight epithelial in the distal nephron, distal colon, the ducts of salivary and sweat glands, and the lung, the expression of ENaC subunits is also reported in tissues and/or organs that seem not to be implicated in whole body Na<sup>+</sup> homeostasis, such as the skin, tongue, eye and blood vessels (rev. in (Kellenberger & Schild, 2015; Rossier, Staub & Hummler, 2013)). The analysis of genetically engineered mouse models started to reveal the role of ENaC in these tissues/organs, unveiling *non-classical* roles, namely in differentiation, barrier function, lipid synthesis and secretion (skin), but also migration of keratinocytes (Charles et al., 2008), and salt perception (tongue; (Chandrashekar et al., 2010), Table 5). In the human eye, all ENaC subunits ( $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ ) are expressed within the cornea, ciliary body, iris, and retina.  $\beta$  and  $\gamma$  ENaC subunits present distinct localizations (Krueger et al., 2012). The  $\beta$  subunit is present in basal regions of the limbal epithelium, while the  $\gamma$ -subunit is present throughout all layers of the corneal epithelium but not in the basal regions of the limbal epithelium. Measurements of electrical potential

difference confirmed functional ENaC-mediated  $\text{Na}^+$ -transport, which is likely implicated in maintaining hydration of the ocular surface (Yu et al., 2012). ENaC is also expressed in endothelial and vascular smooth muscle cells where its increased surface expression can lead to membrane stiffening and reduced release of nitric oxide (Jeggle et al., 2013). The contribution to hypertension of ENaC expressed in these cells remains however still to be determined (Kusche-Vihrog, Tarjus, Fels & Jaisser, 2014).

### *ENaC and hypertension*

According to the World Health Organization, hypertension or increased blood pressure strongly correlates with risk of cardiovascular events, stroke and kidney disease and thus represents the leading cause of death worldwide (Santulli, 2013). It was predicted that the number of hypertensive patients will reach 1.56 billion in 2025 (Kearney, Whelton, Reynolds, Muntner, Whelton & He, 2005) and will thus affect one third of adults in most developing and developed communities. Clinical practice guidelines have been written to provide a straightforward approach to manage hypertension (Weber & Anlauf, 2014). 95% of adults with high blood pressure have primary or essential hypertension with unknown cause, but it is widely accepted that genetic and environmental factors affect blood pressure. According to the American Society of Hypertension and the International Society of Hypertension, the treatment aims to control blood pressure and to deal with all risk factors for cardiovascular diseases, like e.g. overweight or stress (Weber & Anlauf, 2014). Lifestyle interventions have been identified to reduce blood pressure as e.g., weight loss, salt reduction, exercise and reduced alcohol consumption (Levenson, Skerrett & Gaziano, 2002). Lowering the salt intake decreases blood pressure (Bray et al., 2004; Vollmer et al., 2001) and prevents hypertension (Whelton, 1997). In addition, increased  $\text{K}^+$  intake improved blood pressure (Obel, 1989), although the combination of lower  $\text{Na}^+$  and higher  $\text{K}^+$  intake did not further decrease blood pressure (Chalmers et al., 1986). These findings highly suggest that food consumption may be primordial in the prevention and/or the treatment of hypertension, and the “Dietary Approaches to Stop Hypertension” reports that a diet composed of fruits, vegetables and low-fat dairy products rich in  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+$  exerts antihypertensive effects (Zemel, 1997). The diuretic hydrochlorothiazide, an inhibitor of the  $\text{Na}^+\text{Cl}^-$ -cotransporter NCC, is the oldest and most used antihypertensive drug. Hypertension treatment also includes angiotensin-converting enzyme inhibitors such as Captopril or, to avoid hypokalemia,  $\text{K}^+$ -sparing diuretics such as amiloride, an inhibitor of ENaC, which provides the final renal adjustment to  $\text{Na}^+$  balance in the distal kidney (see for review (Saha et al., 2005; Weber & Anlauf, 2014)).

Treatment with inhibitors of ENaC resulted in substantial improvement in blood pressure, highly suggesting that increase in  $\text{Na}^+$  transport by ENaC may be a common and requisite component of salt-dependent forms of hypertension (Pratt, 2005). Thus, the combination of two or more drugs that includes ENaC inhibitors was proposed (Pratt, 2005; Vidt & Borazanian, 2003). ENaC may play an even more central role in  $\text{Na}^+$  retention in the generation of hypertension than previously thought (Pratt, 2005). Thiazide diuretics are currently among the most prescribed anti-hypertension drugs. They are often combined with other anti-hypertensive drugs (Weber, 2014). Next generation diuretics may block synergistically NCC and ENaC, and possibly in addition the  $\text{Cl}^-/\text{HCO}_3^-$ -exchanger pendrin in patients with fluid overloads such as congestive heart failure, nephrotic syndrome, diuretic resistance or generalized edema. They may also block one or more pathways known to upregulate ENaC activity (rev. in (Rossier, 2014; Verouti, Boscardin, Hummler & Frateschi, 2015)).

#### ***ENaC and cystic fibrosis***

Cystic fibrosis (CF) is one of the most common hereditary life-threatening diseases. It is characterized by mutations in the gene coding for the cystic fibrosis transmembrane conductance regulator (CFTR) (Brennan & Schrijver, 2016). The most frequent mutation, the in-frame phenylalanine 508 deletion ( $\Delta\text{F508}$ ) is responsible for 70% of CF mutations worldwide, and exists since the Paleolithic period (Morral et al., 1994). The respiratory failure and finally the death of patients suffering from CF are due to airway obstruction, inflammation, and chronic bacterial infections. CFTR allows hydration and thus mucus clearance of the airway surface of the epithelia by secreting chloride to the lumen of the airway epithelia (Boucher, 2007). In the case of CFTR mutations, the mutated receptor cannot reach the plasma membrane due to a trafficking defect, and as a consequence the airway epithelium gets dehydrated because of the incapacity of CFTR to secrete chloride (Denning, Ostedgaard & Welsh, 1992; Kartner, Augustinas, Jensen, Naismith & Riordan, 1992). Although medical treatment strategies for CF have evolved rapidly over these last years, there is still today no cure. Among the new approaches, miRNAs allow the regulation of post-transcriptional gene expression. miRNAs can act either by repressing or by up-regulating genes of interest (Ramachandran et al., 2012). miRNA approaches have been recently proposed as promising therapeutic treatment of CF that could up-regulate the expression and the number of CFTR proteins that reach the cell surface, as demonstrated in primary human



airway epithelia (Ramachandran, Noorbakhsh, Defea & Hollenberg, 2012; Sonnevile et al., 2015).

The depth of surface liquid of airways is regulated by  $\text{Cl}^-$  secretion through CFTR and by  $\text{Na}^+$  absorption through ENaC. Reduced CFTR expression at the cell surface leads to ENaC upregulation, resulting in an increase of  $\text{Na}^+$  reabsorption and, consequently, to a dehydration of the airway surface (Clunes & Boucher, 2007). Interestingly, it was shown that a mouse model overexpressing the  $\beta$  subunit of the epithelial  $\text{Na}^+$  channel (ENaC) in the lower airways shows typical features of CF, namely reduction of the periciliary liquid weight in bronchi and tracheae, depletion of the airway surface volume, abnormal mucus transport, and reduction of the clearance of bacteria, with a mortality of  $\sim 50\%$  (Mall, Grubb, Harkema, O'Neal & Boucher, 2004). For 30 years inhalation of amiloride has been known to inhibit  $\text{Na}^+$  reabsorption in CF patients. ENaC inhibition improves mucociliary clearance and thereby retards the lung infection (Kohler, App, Schmitz-Schumann, Wurtemberger & Matthys, 1986). Amiloride therapy may reduce morbidity and mortality if it is given early in life before the onset of the lung disease (Zhou et al., 2008). As amiloride has a short duration of action on airway surfaces, second and third generation amiloride analogues, as benzamil and phenamil, were generated and tested to improve both the affinity to ENaC and decreasing the reversibility of the ENaC inhibition (Hirsh et al., 2006). Benzamil presents the highest potency and is rapidly absorbed from the mucosal surface to the cytosol. However, *in vivo* pharmacodynamic experiments in sheep have shown that amiloride and benzamil have the same efficiency (Hirsh et al., 2004). More recently, Parion compound N-(3,5-diamino-6-chloropyrazine-2-carbonyl)-N-4-[4-(2,3-dihydroxypropoxy)phenyl]butyl-guanine methanesulfonate (552-02) was tested in cystic fibrosis bronchial epithelial cells and in sheep for mucociliary clearance after aerosol dosing. Compared to amiloride, 552-02 induced greater airway surface liquid expansion in bronchial epithelial cells. In addition, 552-02 increased the mucociliary clearance in sheep at the time of administration and 4 to 6h later (Hirsh et al., 2008). In summary, the development of new ENaC blockers, as 552-02, with a higher potency, higher selectivity, and durability, is of clinical importance for the treatment of cystic fibrosis lung disease.

### ***Aldosterone-dependent and -independent regulation of ENaC***

#### **Aldosterone-dependent ENaC regulation**

Several hormones regulate ENaC activity (e.g., aldosterone, vasopressin, angiotensin, insulin, rev. by (Verouti, Boscardin, Hummler & Frateschi, 2015)). Among them, the

mineralocorticoid aldosterone acts on the principal cells in the ASDN (Fig. 6). Aldosterone is secreted by the adrenal gland in response to a small increase of  $K^+$  concentration in blood, or decrease of vascular volume, which activates the renin-angiotensin-aldosterone system, thereby maintaining the extracellular  $Na^+$  concentration and the blood pressure within physiological ranges (Bollag, 2014). Interestingly, independently of the renin-angiotensin-aldosterone system,  $K^+$  itself mediates aldosterone secretion from the adrenal gland (Williams, 2005). The genomic response to aldosterone has been extensively analyzed and occurs 4-24 hours following an aldosterone stimulus. In this phase, aldosterone is translocated into the nucleus, where it binds to the mineralocorticoid receptor (MR), and stimulates further synthesis of “aldosterone-induced proteins” like ion transporting proteins, such as the  $\alpha$  subunit of ENaC and the  $\alpha 1$  subunit of the  $Na^+ - K^+ - ATPase$  (Pearce, Soundararajan, Trimpert, Kashlan, Deen & Kohan, 2015). The rapid effects of aldosterone on target tissues are far less well understood and are still controversial, since they are coupled to MR or to a yet unidentified membrane-associated aldosterone receptor (Dooley, Harvey & Thomas, 2012). These non-genomic effects occur within minutes to 2h after aldosterone stimulus (Bollag, 2014). Thereby, the synthesis of ion transporter proteins is not required (Le Moellic et al., 2004). Aldosterone may activate the ERK1/2 MAPK cascade independently of MR (Rossol-Haseroth et al., 2004). Furthermore, activation of the PKC/protein kinase D signaling pathway through the c-Src-dependent trans-activation of the epidermal growth factor receptor may also contribute to early ENaC trafficking in response to aldosterone (Dooley, Harvey & Thomas, 2012). Thus, some of the early responses may potentiate the genomic effects of aldosterone through e.g., the phosphorylation of channels and/or transporters.

#### Aldosterone-independent ENaC regulation

ENaC activity is preserved in MR knockout mice (Berger et al., 1998), demonstrating that aldosterone is not the only modulator of ENaC. Furthermore, aldosterone synthase<sup>(-/-)</sup> mice survive even under low salt diet, demonstrating again an aldosterone-independent compensatory mechanism (Makhanova, Sequeira-Lopez, Gomez, Kim & Smithies, 2006). Interestingly, a study of Nesterov and colleagues clearly demonstrated by measuring ENaC activity in cut open tubules of aldosterone synthase<sup>(-/-)</sup> mice that aldosterone regulates ENaC only in the distal part of the nephron (CNT/CCD), whereas in the DCT2/CNT, ENaC regulation is aldosterone-independent (Nesterov, Dahlmann, Krueger, Bertog, Loffing & Korbmacher, 2012). Other hormones such as e.g. insulin, vasopressin, and angiotensin II

regulate ENaC activity as well (Verouti, Boscardin, Hummler & Frateschi, 2015). Indeed, in adrenalectomized mice, vasopressin is upregulated and ENaC function is not compromised (Mironova, Bugaj, Roos, Kohan & Stockand, 2012). Furthermore, more recently, angiotensin II has been proposed to upregulate  $\gamma$ ENaC in the DCT in AS (-/-) mice treated with high  $K^+$  diet (2%  $K^+$ ), thereby regulating  $Na^+$  and  $K^+$  homeostasis (Todkar et al., 2015) by increasing the electrochemical driving force for  $K^+$  excretion through renal  $K^+$  channels (van der Lubbe et al., 2013). This phenomenon is aldosterone-independent and involves a decrease of the upstream NCC activity (Sorensen et al., 2013; van der Lubbe et al., 2013). Several recent *in vivo* studies confirm that upon hyperkalemia, the distal  $Na^+$  delivery is increased, by decreasing the NCC activity, while upon hypovolemia, both NCC and ENaC are upregulated to improve  $Na^+$  absorption (Rengarajan, Lee, Oh, Delpire, Youn & McDonough, 2014; van der Lubbe, Zietse & Hoorn, 2013; Vitzthum, Seniuk, Schulte, Muller, Hetz & Ehmke, 2014). Thus, angiotensin II acts on the switch from  $K^+$  secretion upon hyperkalemia to  $Na^+$  reabsorption under hypovolemia, that is mediated via several kinases (Arroyo, Ronzaud, Lagnaz, Staub & Gamba, 2011; Mamenko et al., 2013). This so-called “aldosterone paradox” phenomenon (Arroyo & Gamba, 2012), determines NCC and ENaC as key players for  $Na^+$  and  $K^+$  balance within the DCT. As expected for an aldosterone-dependent ENaC regulation, treating WT mice with high  $Na^+$  and high  $K^+$  diet leads to an increase of the plasma aldosterone and an increase of the  $\beta$ ENaC subunit and serum- and glucocorticoid-regulated kinase 1 protein expression (Vitzthum, Seniuk, Schulte, Muller, Hetz & Ehmke, 2014). However, even in presence of the MR antagonist spironolactone or the ENaC blocker amiloride, blood pressure stays increased (Vitzthum, Seniuk, Schulte, Muller, Hetz & Ehmke, 2014). This highly suggests the existence of other, currently unknown regulatory mechanisms in a  $Na^+$ -repleted state. Several other aspects of ENaC regulation are still unresolved, as e.g. the molecular mechanisms underlying the aldosterone-dependent and independent regulation of ENaC, the differential distribution of ENaC channels along the nephron or the possibility of a cross-talk between NCC and ENaC within the DCT2.

#### Is ENaC regulated by potassium diet?

Up to now, evidence for ENaC regulation through  $K^+$  is poor. In case of hyperkalemia, angiotensin II is known to improve the delivery of  $Na^+$  to the distal part of the nephron to increase ENaC-mediated electrochemical secretion of  $K^+$  through renal  $K^+$  channels. As discussed above, this phenomenon is aldosterone-independent and leads to a decrease of the upstream NCC activity. To date, more is known about the regulation of NCC activity upon  $K^+$

loading than on ENaC regulation in this context. Low  $K^+$  diet is a strong stimulus to increase NCC activity (Sorensen et al., 2013; van der Lubbe et al., 2013) by an aldosterone-independent mechanism (Castaneda-Bueno et al., 2014; Vallon et al., 2009). Acute oral  $K^+$  loading resulted in a rapid NCC dephosphorylation and inactivation of NCC (Sorensen et al., 2013). In contrast, high  $Na^+$  diet mediates a decrease of NCC phosphorylation by oxidative-stress-responsive kinase 1/Ste20-related proline alanine rich kinase in an aldosterone-dependent manner, increasing thereby NCC activity (Chiga et al., 2008). ENaC activity within the DCT2/CNT segments is however independent of aldosterone (Nesterov, Dahlmann, Krueger, Bertog, Loffing & Korbmacher, 2012; Todkar et al., 2015) and the mechanism of ENaC regulation in the DCT2/CNT in this context is currently unknown. Furthermore, a regulation of ENaC through solely low  $K^+$  diet has not been investigated so far. The recent study by Perrier and colleagues (Perrier et al., 2015) clearly demonstrates that in nephron-specific  $\alpha$ ENaC knockout mice, hyperkalemia becomes the determining factor in regulating NCC activity, regardless of  $Na^+$  loss, and thus remains the predominant and life-threatening feature to be avoided.

#### **Regulation of ENaC by proteases**

In addition to the hormonal regulation through e.g. vasopressin (Ecelbarger, Kim, Wade & Knepper, 2001) and insulin (Marunaka, Hagiwara & Tohda, 1992), ENaC is regulated by changes in both extracellular and intracellular sodium. An increase in extracellular  $Na^+$  reduces ENaC activity (Fuchs, Larsen & Lindemann, 1977), and this  $Na^+$  self-inhibition can be abolished by treatment with extracellular proteases (Chraibi & Horisberger, 2002). The first evidence of ENaC regulation by proteases was reported in 1983 when Garty and Edelman applied high doses of trypsin (1 mg/ml) to toad urinary bladder and recorded a decrease in the amiloride-sensitive  $Na^+$  current probably as a result of proteolytic digestion of the ENaC channel (Garty & Edelman, 1983). When lower doses of trypsin or chymotrypsin (1-5  $\mu$ g/ml) were applied, ENaC activity expressed in the *Xenopus* oocytes was strongly stimulated by an increase in channel open probability (Chraibi, Vallet, Firsov, Hess & Horisberger, 1998). Since then, trypsin and chymotrypsin are commonly used as experimental tools to achieve maximal ENaC activation. In 1997, Vallet and colleagues screened a *Xenopus* A6 cell complementary DNA library to detect proteins involved in the control of ENaC activity, and isolated the first membrane-bound serine protease whose co-expression with ENaC induced a 3-fold increase in the ENaC  $Na^+$  current amplitude (Vallet, Chraibi,

Gaeggeler, Horisberger & Rossier, 1997). Consequently, this serine protease was termed channel-activating protease-1 (CAP1). It is encoded by the *Prss8* gene and is orthologous to human prostatic containing a glycosylphosphatidyl-inositol (GPI) domain as a plasma membrane anchor (Vallet, Chraïbi, Gaeggeler, Horisberger & Rossier, 1997). The mouse counterpart was cloned two years later from a CCD cell line derived from mouse kidney (Vuagniaux et al., 2000). Measurements of this cell line revealed that the ENaC-mediated Na<sup>+</sup> current was less sensitive (only ~50% of control) to the serine protease inhibitor aprotinin (Vuagniaux et al., 2000). This suggested that ENaC activation depends on more than one protease, and led to the discovery of two additional membrane-bound serine proteases that increased ENaC currents 6 to 10-fold. Accordingly, these type II-oriented membrane-bound serine proteases were called CAP2 (also known as Tmprss4) and CAP3 (also known as matriptase) (Vuagniaux et al., 2000). Interestingly, CAP3 has been shown to decrease an acid-activated ASIC current as measured by two-electrode voltage clamp in *Xenopus* oocytes (Clark, Jovov, Rooj, Fuller & Benos, 2010). Since then, several other proteases were added to the list of ENaC regulatory molecules, all increasing ENaC currents. TMPRSS3 which is mutated in deafness (Guipponi et al., 2002), furin (Hughey et al., 2004), neutrophil elastase (Caldwell, Boucher & Stutts, 2005), kallikrein (Patel, Chao & Palmer, 2012; Picard et al., 2008), plasmin (Buhl et al., 2014; Svenningsen et al., 2009), metalloprotease meprin  $\beta$  subunit (Garcia-Caballero et al., 2011), cathepsin-S (Haerteis et al., 2012), urokinase plasminogen activator (Chen et al., 2014; Ji, Zhao, Komissarov, Chang, Liu & Matthay, 2015), alkaline phosphatase from *Pseudomonas aeruginosa* (Butterworth, Zhang, Heidrich, Myerburg & Thibodeau, 2012; Butterworth, Zhang, Liu, Shanks & Thibodeau, 2014), and trypsin IV (Haerteis et al., 2014), are all proteases able to activate ENaC in *in vitro* settings. So far, only CAP1, kallikrein and the urokinase plasminogen activator have been confirmed as physiological regulators of ENaC activity by *in vivo* studies using genetically modified or spontaneous mutant mice (Chen et al., 2014; Frateschi et al., 2012; Malsure et al., 2014; Picard et al., 2008; Planes et al., 2009), and it was shown that the serine protease CAP2 is not implicated in ENaC regulation *in vivo* in kidney and colon (Keppner et al., 2015). The mechanism by which these proteases activate the channel is still largely debated. Several researchers claim and provided evidence for a direct activation mediated by proteolytic cleavage of ENaC that would lead to the removal of inhibitory domains and/or changes in the three-dimensional structure of the channel resulting in its activation (Shi, Carattino, Hughey & Kleyman, 2013). Indeed, this hypothesis is quite tempting, as most of these enzymes are membrane-bound proteins facing the extracellular side of the plasma membrane or soluble

molecules and thus potentially located in close proximity with ENaC. Although cleavage products of the different ENaC subunits have been observed under various conditions, functional evidence is still missing. In this regard, CAP1 is peculiar and its action on ENaC still controversial. A CAP1 cleavage site (RKRK) was identified in the  $\gamma$  subunit of ENaC downstream the proposed consensus cleavage site for furin. On the one hand, the mutation of either the CAP1 cleavage site or the furin cleavage site abolished the activation of ENaC by CAP1 in *Xenopus* oocytes, and prevented the appearance of a smaller fragment of the  $\gamma$  subunit (Bruns et al., 2007; Carattino, Hughey & Kleyman, 2008). However, mutation of the CAP1 catalytic triad that should render the protease catalytically-inactive, did not prevent full ENaC activation and moreover induced  $\gamma$  subunit cleavage as well as its wild-type counterpart (Andreasen, Vuagniaux, Fowler-Jaeger, Hummler & Rossier, 2006; Bruns et al., 2007; Carattino et al., 2014). Other studies indicated that there is no consistent correlation between  $\gamma$  subunit cleavage and ENaC activation (Fejes-Toth, Frindt, Naray-Fejes-Toth & Palmer, 2008; Harris, Garcia-Caballero, Stutts, Firsov & Rossier, 2008). In addition, although CAP1 is indeed a positive regulator of ENaC *in vivo*, since CAP1 loss-of-function or absence significantly decreases ENaC basal activity (Frateschi et al., 2012; Malsure et al., 2014; Planes et al., 2009), the cleaved fragment originating from  $\gamma$ ENaC was still observed in the colon or in the lung of tissue-specific CAP1 knockouts (Malsure et al., 2014; Planes et al., 2009). Interestingly, both the wild-type and the catalytically inactive mutant CAP1 are able to elicit skin pathology when overexpressed in mouse models of ichthyotic skin disorders (Crisante et al., 2014; Frateschi et al., 2011). This indicates that as for ENaC activation, the catalytic triad of CAP1 is dispensable for provoking pathophysiological effects. The presence of cleaved fragments originating from ENaC subunits may nevertheless have physiological meanings, even if they most likely do not origin from direct cleavage by CAP1. Very recently, Zachar et al. reported that, in the human kidney,  $\gamma$ ENaC is subject to proteolytic processing, yielding fragments compatible with furin cleavage, and that proteinuria is accompanied with cleavage at the putative CAP1 site of  $\gamma$ ENaC (Zachar et al., 2015). Table 6 highlights the proteases involved in the regulation of ENaC with particular emphasis on their physiological implication.

Protease inhibitors achieve another level of regulation of ENaC activity. Using a proteomic approach, Garcia-Caballero et al. identified SPLUNC1 (the short palate, lung, and nasal epithelial clone 1) as an ENaC inhibitor highly expressed in the airways, colon and kidney (Garcia-Caballero et al., 2009), with the inhibitory domain located at the N terminus (Hobbs et al., 2013).  $\alpha$ 1-antitrypsin, an anti-serine protease found in human plasma and lung

epithelial fluid (Lazrak et al., 2009), inhibits amiloride-sensitive Na<sup>+</sup> transport in *Xenopus* oocytes and alveolar fluid clearance in mice (Lazrak et al., 2009). A role of SerpinB1, a highly efficient inhibitor of neutrophil serine proteases, was described in the cystic fibrosis airways suggesting a possible link to ENaC regulation (Cooley, Sontag, Accurso & Remold-O'Donnell, 2011). Although a putative impact of the pulmonary protease inhibitors secretory leukocyte protease inhibitor I and trappin-2 on ENaC and Na<sup>+</sup> transport in the airways remains unknown, their antibacterial and anti-fungal properties towards common microorganisms in cystic fibrosis patients have been reported (Sallenave, 2010; Zani et al., 2011). Similarly, loss of either of the two Kunitz-type transmembrane serine protease inhibitors, hepatocyte growth factor activator inhibitor (HAI)-1 or -2, expressed by tubular epithelium in kidney (Yamauchi, Kataoka, Itoh, Seguchi, Hasui & Osada, 2004), is associated with embryonic lethality in mice (Mitchell et al., 2001; Tanaka et al., 2005). All developmental defects in HAI-1- and HAI-2-deficient embryos are rescued by simultaneous CAP1 deficiency, but not by absence of ENaC, suggesting that ENaC activity is not involved (Szabo et al., 2012). The protease activity of CAP1 can be inhibited by the serine protease inhibitor protease nexin-1 (also known as SerpinE2) (Chen, Zhang & Chai, 2004; Crisante et al., 2014). Nexin-1 is known to be an endogenous inhibitor for  $\alpha$ -thrombin, plasmin, and plasminogen activators, and Wakida et al. showed an inhibitory effect of nexin-1 on CAP1-induced ENaC currents and proposed that nexin-1 could exert a natriuretic role by inhibiting CAP1 activity (Wakida et al., 2006). We also recently reported an inhibitory effect of nexin-1 on both wild-type and catalytically inactive mutant CAP1 *in vitro* and in an *in vivo* model, indicating that the catalytic site of CAP1 is dispensable for nexin-1 inhibition (Crisante et al., 2014), and demonstrating a novel inhibitory interaction between CAP1 and nexin-1. This opens the search for specific CAP1 antagonists that are independent of its catalytic activity.

Altogether, proteases (in particular serine proteases) and their inhibitors are important regulators of ENaC function. Although progress has been made on the mechanism of such regulations, the precise signaling pathways remain to be disclosed, which necessitates the usage of *in vivo* models to define their physiological significance.

## Outlook

As discussed above, all ENaC/DEG family members share the same subunit topology and form Na<sup>+</sup> channels that are inhibited by amiloride (Kellenberger & Schild, 2015). More recently it was found that ENaC and ASICs are both regulated by proteases, and that also ENaC is pH-dependent (Table 4). The tissue expression and physiological roles are however



very different between ASICs and ENaC. Whereas ENaC is well known for its role in mediating Na<sup>+</sup> entry in epithelial cells, especially in the kidney, lung and colon, ASICs are predominantly found in the CNS and the PNS. The question arises whether in tissues expressing both ASIC and ENaC channels, overlapping functions can be identified. No function has so far been found for ENaC in the ear and in DRG neurons, in contrast to ASICs (Raouf et al., 2012; Rusch & Hummler, 1999). There is currently evidence for roles of both ENaC and ASICs in mechanosensation in arterial baroreceptors (Drummond, Welsh & Abboud, 2001; Lu et al., 2009). ENaC  $\beta$  and  $\gamma$  subunits are also expressed in vascular smooth muscle cells where they contribute to pressure-induced constriction. Similarly, ASICs in vascular smooth muscle cells likely regulate vascular tone and chemotactic migration (Drummond, Jernigan & Grifoni, 2008). Other overlapping functions of ENaC and ASICs may still be discovered. Interestingly, two recent studies indicate that overexpression of ASICs together with ENaC may lead to the formation of hybrid ENaC/ASIC channels (Jeggle, Smith, Stewart, Haerteis, Korbmacher & Edwardson, 2015; Kapoor et al., 2011).

What will be the next challenges and progresses in ENaC and ASIC research? For ASICs, it will be important to identify and develop specific and potent small molecule inhibitors, and this with two aims, first to confirm the physiological and pathological roles of ASICs, and second to take advantage of ASIC pharmacology to treat human diseases. The available structural information will together with complementary approaches lead to a better understanding of the molecular mechanisms of ASIC function, allowing a rational drug development. For the ENaC field, ENaC distribution and function have still to be unveiled in classical and non-classical tissues and organs and may reveal so far unknown functions of ENaC as e.g. a defect in lipid synthesis and processing in skin (Charles et al., 2008). Conditional and inducible gene targeting of single ENaC subunits in adult mice may help to shed some light on these important issues.

### **Acknowledgements**

Work in the author's laboratories was funded by Swiss national science foundation grants 31003A\_163347 (to E.H.) and 31003A\_153419 (to S.K.).

### **Conflict of interest**

The authors declare that they don't have any conflicts of interest.

## References

- Abdrakhmanova G, Cleemann L, Lindstrom J, & Morad M (2004). Differential modulation of beta2 and beta4 subunits of human neuronal nicotinic acetylcholine receptors by acidification. *Mol Pharmacol* 66: 347-355.
- Abdrakhmanova G, Dorfman J, Xiao Y, & Morad M (2002). Protons enhance the gating kinetics of the alpha3/beta4 neuronal nicotinic acetylcholine receptor by increasing its apparent affinity to agonists. *Mol Pharmacol* 61: 369-378.
- Adams CM, Anderson MG, Motto DG, Price MP, Johnson WA, & Welsh MJ (1998). Ripped Pocket and Pickpocket, Novel Drosophila Deg/Enac Subunits Expressed in Early Development and in Mechanosensory Neurons. *J Cell Biol* 140: 143-152.
- Adams CM, Snyder PM, & Welsh MJ (1999). Paradoxical stimulation of a DEG/ENaC channel by amiloride. *J Biol Chem* 274: 15500-15504.
- Alijevic O, & Kellenberger S (2012). Subtype-specific modulation of acid-sensing ion channel (ASIC) function by 2-guanidine-4-methylquinazoline. *J Biol Chem* 287: 36059-36070.
- Allen NJ, & Attwell D (2002). Modulation of ASIC channels in rat cerebellar Purkinje neurons by ischaemia-related signals. *J Physiol-London* 543: 521-529.
- Anantharam A, & Palmer LG (2007). Determination of epithelial Na<sup>+</sup> channel subunit stoichiometry from single-channel conductances. *J Gen Physiol* 130: 55-70.
- Andreasen D, Vuagniaux G, Fowler-Jaeger N, Hummler E, & Rossier BC (2006). Activation of Epithelial Sodium Channels by Mouse Channel Activating Proteases (mCAP) Expressed in Xenopus Oocytes Requires Catalytic Activity of mCAP3 and mCAP2 but not mCAP1. *J Am Soc Nephrol* 17: 968-976.
- Arnadottir J, O'Hagan R, Chen Y, Goodman MB, & Chalfie M (2011). The DEG/ENaC protein MEC-10 regulates the transduction channel complex in *Caenorhabditis elegans* touch receptor neurons. *J Neurosci* 31: 12695-12704.
- Arroyo JP, & Gamba G (2012). Advances in WNK signaling of salt and potassium metabolism: clinical implications. *Am J Nephrol* 35: 379-386.
- Arroyo JP, Ronzaud C, Lagnaz D, Staub O, & Gamba G (2011). Aldosterone paradox: differential regulation of ion transport in distal nephron. *Physiology* 26: 115-123.
- Askwith CC, Cheng C, Ikuma M, Benson C, Price MP, & Welsh MJ (2000). Neuropeptide FF and FMRFamide potentiate acid-evoked currents from sensory neurons and proton-gated DEG/ENaC channels. *Neuron* 26: 133-141.
- Babini E, Paukert M, Geisler HS, & Grunder S (2002). Alternative splicing and interaction with di- and polyvalent cations control the dynamic range of acid-sensing ion channel 1 (ASIC1). *J Biol Chem* 277: 41597-41603.

Baconguis I, Bohlen CJ, Goehring A, Julius D, & Gouaux E (2014). X-ray structure of Acid-sensing ion channel 1-snake toxin complex reveals open state of a Na<sup>+</sup>-selective channel. *Cell* 156: 717-729.

Baconguis I, & Gouaux E (2012). Structural plasticity and dynamic selectivity of acid-sensing ion channel-spider toxin complexes. *Nature* 489: 400-405.

Barker PM, Nguyen MS, Gatzky JT, Grubb B, Norman H, Hummler E, *et al.* (1998). Role of gammaENaC subunit in lung liquid clearance and electrolyte balance in newborn mice. Insights into perinatal adaptation and pseudohypoaldosteronism. *J Clin Invest* 102: 1634-1640.

Baron A, Diochot S, Salinas M, Deval E, Noel J, & Lingueglia E (2013). Venom toxins in the exploration of molecular, physiological and pathophysiological functions of acid-sensing ion channels. *Toxicon* 75: 187-204.

Baron A, & Lingueglia E (2015). Pharmacology of acid-sensing ion channels - Physiological and therapeutical perspectives. *Neuropharmacology* 94: 19-35.

Bartoi T, Augustinowski K, Polleichtner G, Grunder S, & Ulbrich MH (2014). Acid-sensing ion channel (ASIC) 1a/2a heteromers have a flexible 2:1/1:2 stoichiometry. *Proc Natl Acad Sci U S A* 111: 8281-8286.

Barton NJ, McQueen DS, Thomson D, Gauldie SD, Wilson AW, Salter DM, *et al.* (2006). Attenuation of experimental arthritis in TRPV1R knockout mice. *Exp Mol Pathol* 81: 166-170.

Bassler EL, Ngo-Anh TJ, Geisler HS, Ruppertsberg JP, & Grunder S (2001). Molecular and functional characterization of acid-sensing ion channel (ASIC) 1b. *J Biol Chem* 276: 33782-33787.

Baumann TK, Chaudhary P, & Martenson ME (2004). Background potassium channel block and TRPV1 activation contribute to proton depolarization of sensory neurons from humans with neuropathic pain. *Eur J Neurosci* 19: 1343-1351.

Benson CJ, Xie JH, Wemmie JA, Price MP, Henss JM, Welsh MJ, *et al.* (2002). Heteromultimers of DEG/ENaC subunits form H<sup>+</sup>-gated channels in mouse sensory neurons. *Proc Natl Acad Sci USA* 99: 2338-2343.

Berger S, Bleich M, Schmid W, Cole TJ, Peters J, Watanabe H, *et al.* (1998). Mineralocorticoid receptor knockout mice: pathophysiology of Na<sup>+</sup> metabolism. *Proc Natl Acad Sci U S A* 95: 9424-9429.

Bertog M, Cuffe JE, Pradervand S, Hummler E, Hartner A, Porst M, *et al.* (2008). Aldosterone responsiveness of the epithelial sodium channel (ENaC) in colon is increased in a mouse model for Liddle's syndrome. *J Physiol* 586: 459-475.

Beyenbach KW, & Wieczorek H (2006). The V-type H<sup>+</sup> ATPase: molecular structure and function, physiological roles and regulation. *J Exp Biol* 209: 577-589.

- Birdsong WT, Fierro L, Williams FG, Spelta V, Naves LA, Knowles M, *et al.* (2010). Sensing muscle ischemia: coincident detection of acid and ATP via interplay of two ion channels. *Neuron* 68: 739-749.
- Blanchard MG, & Kellenberger S (2011). Effect of a temperature increase in the non-noxious range on proton-evoked ASIC and TRPV1 activity. *Pflugers Arch* 461: 123-139.
- Blanchard MG, Rash LD, & Kellenberger S (2012). Inhibition of voltage-gated Na<sup>+</sup> currents in sensory neurones by the sea anemone toxin APETx2. *Br J Pharmacol* 165: 2167-2177.
- Blot-Chabaud M, Djelidi S, Courtois-Coutry N, Fay M, Cluzeaud F, Hummler E, *et al.* (2001). Coordinate control of Na,K-atpase mRNA expression by aldosterone, vasopressin and cell sodium delivery in the cortical collecting duct. *Cellular and molecular biology* 47: 247-253.
- Bohlen CJ, Chesler AT, Sharif-Naeini R, Medzihradzsky KF, Zhou S, King D, *et al.* (2011). A heteromeric Texas coral snake toxin targets acid-sensing ion channels to produce pain. *Nature* 479: 410-414.
- Boillat A, Alijevic O, & Kellenberger S (2014). Calcium entry via TRPV1 but not ASICs induces neuropeptide release from sensory neurons. *Molecular and cellular neurosciences* 61: 13-22.
- Bollag WB (2014). Regulation of aldosterone synthesis and secretion. *Compr Physiol* 4: 1017-1055.
- Bonora M, Riffault L, Marie S, Mall M, Clement A, & Tabary O (2011). Morphological analysis of the trachea and pattern of breathing in betaENaC-Tg mice. *Respiratory physiology & neurobiology* 178: 346-348.
- Boucher RC (2007). Airway surface dehydration in cystic fibrosis: pathogenesis and therapy. *Annu Rev Med* 58: 157-170.
- Bray GA, Vollmer WM, Sacks FM, Obarzanek E, Svetkey LP, Appel LJ, *et al.* (2004). A further subgroup analysis of the effects of the DASH diet and three dietary sodium levels on blood pressure: results of the DASH-Sodium Trial. *Am J Cardiol* 94: 222-227.
- Brennan ML, & Schrijver I (2016). Cystic Fibrosis: A Review of Associated Phenotypes, Use of Molecular Diagnostic Approaches, Genetic Characteristics, Progress, and Dilemmas. *J Mol Diagn* 18: 3-14.
- Bruns JB, Carattino MD, Sheng S, Maarouf AB, Weisz OA, Pilewski JM, *et al.* (2007). Epithelial Na<sup>+</sup> channels are fully activated by furin- and prostasin-dependent release of an inhibitory peptide from the gamma-subunit. *J Biol Chem* 282: 6153-6160.
- Buhl KB, Oxlund CS, Friis UG, Svenningsen P, Bistrup C, Jacobsen IA, *et al.* (2014). Plasmin in urine from patients with type 2 diabetes and treatment-resistant hypertension activates ENaC in vitro. *J Hypertens* 32: 1672-1677.

- Butterworth MB, Zhang L, Heidrich EM, Myerburg MM, & Thibodeau PH (2012). Activation of the epithelial sodium channel (ENaC) by the alkaline protease from *Pseudomonas aeruginosa*. *J Biol Chem* 287: 32556-32565.
- Butterworth MB, Zhang L, Liu X, Shanks RM, & Thibodeau PH (2014). Modulation of the epithelial sodium channel (ENaC) by bacterial metalloproteases and protease inhibitors. *PLoS one* 9: e100313.
- Cadiou H, Studer M, Jones NG, Smith ES, Ballard A, McMahon SB, *et al.* (2007). Modulation of acid-sensing ion channel activity by nitric oxide. *J Neurosci* 27: 13251-13260.
- Caldwell RA, Boucher RC, & Stutts MJ (2005). Neutrophil elastase activates near-silent epithelial Na<sup>+</sup> channels and increases airway epithelial Na<sup>+</sup> transport. *Am J Physiol Lung Cell Mol Physiol* 288: L813-819.
- Carattino MD, Hughey RP, & Kleyman TR (2008). Proteolytic Processing of the Epithelial Sodium Channel {gamma} Subunit Has a Dominant Role in Channel Activation. *J Biol Chem* 283: 25290-25295.
- Carattino MD, Mueller GM, Palmer LG, Frindt G, Rued AC, Hughey RP, *et al.* (2014). Proximin interacts with the epithelial Na<sup>+</sup> channel and facilitates cleavage of the gamma-subunit by a second protease. *Am J Physiol Renal Physiol* 307: F1080-1087.
- Castaneda-Bueno M, Cervantes-Perez LG, Rojas-Vega L, Arroyo-Garza I, Vazquez N, Moreno E, *et al.* (2014). Modulation of NCC activity by low and high K(+) intake: insights into the signaling pathways involved. *Am J Physiol Renal Physiol* 306: F1507-1519.
- Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, *et al.* (2000). Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288: 306-313.
- Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, & Julius D (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389: 816-824.
- Chalmers J, Morgan T, Doyle A, Dickson B, Hopper J, Mathews J, *et al.* (1986). Australian National Health and Medical Research Council dietary salt study in mild hypertension. *J Hypertens Suppl* 4: S629-637.
- Chandrashekar J, Kuhn C, Oka Y, Yarmolinsky DA, Hummler E, Ryba NJ, *et al.* (2010). The cells and peripheral representation of sodium taste in mice. *Nature* 464: 297-301.
- Chang CT, Bens M, Hummler E, Boulkroun S, Schild L, Teulon J, *et al.* (2005). Vasopressin-stimulated CFTR Cl<sup>-</sup> currents are increased in the renal collecting duct cells of a mouse model of Liddle's syndrome. *J Physiol* 562: 271-284.
- Charles RP, Guitard M, Leyvraz C, Breiden B, Haftek M, Haftek-Terreau Z, *et al.* (2008). Postnatal requirement of the epithelial sodium channel for maintenance of epidermal barrier function. *J Biol Chem* 283: 2622-2630.

- Cheek DB, & Perry JW (1958). A salt wasting syndrome in infancy. *Arch Dis Child* 33: 252-256.
- Chen CC, Akopian AN, Sivilotti L, Colquhoun D, Burnstock G, & Wood JN (1995). A P2X purinoceptor expressed by a subset of sensory neurons. *Nature* 377: 428-431.
- Chen H-Y, & Chesler M (2015). Autocrine boost of NMDAR current in hippocampal CA1 pyramidal neurons by a PMCA-dependent, perisynaptic, extracellular pH shift. *J Neurosci* 35: 873-877.
- Chen JC, & Chesler M (1992). Modulation of extracellular pH by glutamate and GABA in rat hippocampal slices. *J Neurophysiol* 67: 29-36.
- Chen LM, Zhang X, & Chai KX (2004). Regulation of prostasin expression and function in the prostate. *The Prostate* 59: 1-12.
- Chen X, Qiu L, Li M, Durrnagel S, Orser BA, Xiong ZG, *et al.* (2010). Diarylamidines: high potency inhibitors of acid-sensing ion channels. *Neuropharmacology* 58: 1045-1053.
- Chen Z, Dillon GH, & Huang R (2004). Molecular determinants of proton modulation of glycine receptors. *J Biol Chem* 279: 876-883.
- Chen Z, Zhao R, Zhao M, Liang X, Bhattarai D, Dhiman R, *et al.* (2014). Regulation of epithelial sodium channels in urokinase plasminogen activator deficiency. *Am J Physiol Lung Cell Mol Physiol* 307: L609-617.
- Chesler M (1990). The regulation and modulation of pH in the nervous system. *Progr Neurobiol* 34: 401-427.
- Chesler M (2003). Regulation and modulation of pH in the brain. *Physiol Rev* 83: 1183-1221.
- Chiang PH, Chien TC, Chen CC, Yanagawa Y, & Lien CC (2015). ASIC-dependent LTP at multiple glutamatergic synapses in amygdala network is required for fear memory. *Sci Rep* 5: 10143.
- Chiga M, Rai T, Yang SS, Ohta A, Takizawa T, Sasaki S, *et al.* (2008). Dietary salt regulates the phosphorylation of OSR1/SPAK kinases and the sodium chloride cotransporter through aldosterone. *Kidney Int* 74: 1403-1409.
- Chraibi A, & Horisberger JD (2002). Na self inhibition of human epithelial Na channel: temperature dependence and effect of extracellular proteases. *J Gen Physiol* 120: 133-145.
- Chraibi A, Vallet V, Firsov D, Hess SK, & Horisberger JD (1998). Protease modulation of the activity of the epithelial sodium channel expressed in *Xenopus* oocytes. *J Gen Physiol* 111: 127-138.
- Christensen BM, Perrier R, Wang Q, Zuber AM, Maillard M, Mordasini D, *et al.* (2010). Sodium and potassium balance depends on alphaENaC expression in connecting tubule. *J Am Soc Nephrol* 21: 1942-1951.

Christensen BM, Zuber AM, Loffing J, Stehle JC, Deen PM, Rossier BC, *et al.* (2011).  $\alpha$ ENaC-mediated lithium absorption promotes nephrogenic diabetes insipidus. *J Am Soc Nephrol* 22: 253-261.

Chu X-P, Zhu X-M, Wei W-L, Li G-H, Simon RP, MacDonald JF, *et al.* (2003). Acidosis decreases low  $\text{Ca}^{2+}$ -induced neuronal excitation by inhibiting the activity of calcium-sensing cation channels in cultured mouse hippocampal neurons. *J Physiol (Lond)* 550: 385-399.

Cichy A, Ackels T, Tsitoura C, Kahan A, Gronloh N, Sochtig M, *et al.* (2015). Extracellular pH regulates excitability of vomeronasal sensory neurons. *J Neurosci* 35: 4025-4039.

Clark EB, Jovov B, Rooj AK, Fuller CM, & Benos DJ (2010). Proteolytic cleavage of human acid-sensing ion channel 1 by the serine protease matriptase. *J Biol Chem* 285: 27130-27143.

Claydon TW, Boyett MR, Sivaprasadarao A, Ishii K, Owen JM, O'Beirne HA, *et al.* (2000). Inhibition of the  $\text{K}^+$  channel kv1.4 by acidosis: protonation of an extracellular histidine slows the recovery from N-type inactivation. *J Physiol (Lond)* 526 Pt 2: 253-264.

Clunes MT, & Boucher RC (2007). Cystic Fibrosis: The Mechanisms of Pathogenesis of an Inherited Lung Disorder. *Drug Discov Today Dis Mech* 4: 63-72.

Cockayne DA, Hamilton SG, Zhu QM, Dunn PM, Zhong Y, Novakovic S, *et al.* (2000). Urinary bladder hyporeflexia and reduced pain-related behaviour in P2X(3)-deficient mice. *Nature* 407: 1011-1015.

Cooley J, Sontag MK, Accurso FJ, & Remold-O'Donnell E (2011). SerpinB1 in cystic fibrosis airway fluids: quantity, molecular form and mechanism of elastase inhibition. *Eur Resp J* 37: 1083-1090.

Cooper BY, Johnson RD, & Rau KK (2004). Characterization and function of TWIK-related acid sensing  $\text{K}^+$  channels in a rat nociceptive cell. *Neurosci* 129: 209-224.

Coryell MW, Wunsch AM, Haenfler JM, Allen JE, Schnizler M, Ziemann AE, *et al.* (2009). Acid-sensing ion channel-1a in the amygdala, a novel therapeutic target in depression-related behavior. *J Neurosci* 29: 5381-5388.

Crisante G, Battista L, Iwaszkiewicz J, Nesca V, Merillat AM, Sergi C, *et al.* (2014). The CAP1/Prss8 catalytic triad is not involved in PAR2 activation and protease nexin-1 (PN-1) inhibition. *Faseb J* 28: 4792-4805.

Dahlmann A, Pradervand S, Hummler E, Rossier BC, Frindt G, & Palmer LG (2003). Mineralocorticoid regulation of epithelial  $\text{Na}^+$  channels is maintained in a mouse model of Liddle's syndrome. *Am J Physiol Renal Physiol* 285: F310-318.

Daumas P, & Andersen OS (1993). Proton block of rat brain sodium channels. Evidence for two proton binding sites and multiple occupancy. *J Gen Physiol* 101: 27-43.

- Dawson RJ, Benz J, Stohler P, Tetaz T, Joseph C, Huber S, *et al.* (2012). Structure of the Acid-sensing ion channel 1 in complex with the gating modifier Psalmotoxin 1. *Nature communications* 3: 936.
- Deitmer JW (2002). A role for CO<sub>2</sub> and bicarbonate transporters in metabolic exchanges in the brain. *J Neurochem* 80: 721-726.
- Delaunay A, Gasull X, Salinas M, Noel J, Friend V, Lingueglia E, *et al.* (2012). Human ASIC3 channel dynamically adapts its activity to sense the extracellular pH in both acidic and alkaline directions. *Proc Natl Acad Sci U S A* 109: 13124-13129.
- Denning GM, Ostedgaard LS, & Welsh MJ (1992). Abnormal localization of cystic fibrosis transmembrane conductance regulator in primary cultures of cystic fibrosis airway epithelia. *J Cell Biol* 118: 551-559.
- Deval E, Baron A, Lingueglia E, Mazarguil H, Zajac JM, & Lazdunski M (2003). Effects of neuropeptide SF and related peptides on acid sensing ion channel 3 and sensory neuron excitability. *Neuropharmacol* 44: 662-671.
- Deval E, & Lingueglia E (2015). Acid-Sensing Ion Channels and nociception in the peripheral and central nervous systems. *Neuropharmacol* 94: 49-57.
- Deval E, Noel J, Gasull X, Delaunay A, Alloui A, Friend V, *et al.* (2011). Acid-sensing ion channels in postoperative pain. *J Neurosci* 31: 6059-6066.
- Deval E, Noel J, Lay N, Alloui A, Diochot S, Friend V, *et al.* (2008). ASIC3, a sensor of acidic and primary inflammatory pain. *EMBO J* 27: 3047-3055.
- DeVries SH (2001). Exocytosed protons feedback to suppress the Ca<sup>2+</sup> current in mammalian cone photoreceptors. *Neuron* 32: 1107-1117.
- Dietrich CJ, & Morad M (2010). Synaptic acidification enhances GABA<sub>A</sub> signaling. *J Neurosci* 30: 16044-16052.
- Diochot S, Baron A, Salinas M, Douguet D, Scarzello S, Dabert-Gay AS, *et al.* (2012). Black mamba venom peptides target acid-sensing ion channels to abolish pain. *Nature* 490: 552-555.
- Dirajlal S, Pauers LE, & Stucky CL (2003). Differential response properties of IB4-positive and -negative unmyelinated sensory neurons to protons and capsaicin. *J Neurophysiol* 89: 513-524.
- Dmitriev AV, & Mangel SC (2001). Circadian clock regulation of pH in the rabbit retina. *J Neurosci* 21: 2897-2902.
- Donier E, Rugiero F, Jacob C, & Wood JN (2008). Regulation of ASIC activity by ASIC4-- new insights into ASIC channel function revealed by a yeast two-hybrid assay. *Eur J Neurosci* 28: 74-86.



- Dooley R, Harvey BJ, & Thomas W (2012). Non-genomic actions of aldosterone: from receptors and signals to membrane targets. *Mol Cell Endocrinol* 350: 223-234.
- Drew LJ, Rohrer DK, Price MP, Blaver KE, Cockayne DA, Cesare P, *et al.* (2004). Acid-sensing ion channels ASIC2 and ASIC3 do not contribute to mechanically activated currents in mammalian sensory neurones. *J Physiol (Lond)* 556: 691-710.
- Drummond HA, Jernigan NL, & Grifoni SC (2008). Sensing tension: epithelial sodium channel/acid-sensing ion channel proteins in cardiovascular homeostasis. *Hypertension* 51: 1265-1271.
- Drummond HA, Welsh MJ, & Abboud FM (2001). ENaC subunits are molecular components of the arterial baroreceptor complex. *Ann New York Acad Sci* 940: 42-47.
- Du J, Reznikov LR, Price MP, Zha XM, Lu Y, Moninger TO, *et al.* (2014). Protons are a neurotransmitter that regulates synaptic plasticity in the lateral amygdala. *Proc Natl Acad Sci U S A* 111: 8961-8966.
- Dube GR, Lehto SG, Breese NM, Baker SJ, Wang X, Matulenko MA, *et al.* (2005). Electrophysiological and in vivo characterization of A-317567, a novel blocker of acid sensing ion channels. *Pain* 117: 88-96.
- Ecelbarger CA, Kim GH, Wade JB, & Knepper MA (2001). Regulation of the abundance of renal sodium transporters and channels by vasopressin. *Exp Neurol* 171: 227-234.
- Egli M, Duplain H, Lepori M, Cook S, Nicod P, Hummler E, *et al.* (2004). Defective respiratory amiloride-sensitive sodium transport predisposes to pulmonary oedema and delays its resolution in mice. *J Physiol* 560: 857-865.
- Ehling P, Cerina M, Budde T, Meuth SG, & Bittner S (2015). The CNS under pathophysiologic attack--examining the role of K<sub>2p</sub> channels. *Pflugers Arch* 467: 959-972.
- Fejes-Toth G, Frindt G, Naray-Fejes-Toth A, & Palmer LG (2008). Epithelial Na<sup>+</sup> channel activation and processing in mice lacking SGK1. *Am J Physiol Renal Physiol* 294: F1298-1305.
- Firsov D, Gautschi I, Merillat AM, Rossier BC, & Schild L (1998). The heterotetrameric architecture of the epithelial sodium channel (ENaC). *EMBO J* 17: 344-352.
- Fischer MJ, Reeh PW, & Sauer SK (2003). Proton-induced calcitonin gene-related peptide release from rat sciatic nerve axons, in vitro, involving TRPV1. *Eur J Neurosci* 18: 803-810.
- Frateschi S, Camerer E, Crisante G, Rieser S, Membrez M, Charles RP, *et al.* (2011). PAR2 absence completely rescues inflammation and ichthyosis caused by altered CAP1/Prss8 expression in mouse skin. *Nature communications* 2: 161.
- Frateschi S, Keppner A, Malsure S, Iwaszkiewicz J, Sergi C, Merillat AM, *et al.* (2012). Mutations of the serine protease CAP1/Prss8 lead to reduced embryonic viability, skin defects, and decreased ENaC activity. *Am J Pathol* 181: 605-615.

- Fu Y, Gerasimova M, Batz F, Kuczkowski A, Alam Y, Sanders PW, *et al.* (2015). PPARgamma agonist-induced fluid retention depends on alphaENaC expression in connecting tubules. *Nephron* 129: 68-74.
- Fuchs W, Larsen EH, & Lindemann B (1977). Current-voltage curve of sodium channels and concentration dependence of sodium permeability in frog skin. *J Physiol* 267: 137-166.
- Fuldner HH, & Stadler H (1982). <sup>31</sup>P-NMR analysis of synaptic vesicles. Status of ATP and internal pH. *Eur J Biochem / FEBS* 121: 519-524.
- Furgeson SB, & Linas S (2010). Mechanisms of type I and type II pseudohypoaldosteronism. *J Am Soc Nephrol* 21: 1842-1845.
- Gao J, Duan B, Wang DG, Deng XH, Zhang GY, Xu L, *et al.* (2005). Coupling between NMDA receptor and acid-sensing ion channel contributes to ischemic neuronal death. *Neuron* 48: 635-646.
- Garcia-Caballero A, Dang Y, He H, & Stutts MJ (2008). ENaC proteolytic regulation by channel-activating protease 2. *J Gen Physiol* 132: 521-535.
- Garcia-Caballero A, Ishmael SS, Dang Y, Gillie D, Bond JS, Milgram SL, *et al.* (2011). Activation of the epithelial sodium channel by the metalloprotease meprin beta subunit. *Channels (Austin)* 5: 14-22.
- Garcia-Caballero A, Rasmussen JE, Gaillard E, Watson MJ, Olsen JC, Donaldson SH, *et al.* (2009). SPLUNC1 regulates airway surface liquid volume by protecting ENaC from proteolytic cleavage. *Proc Natl Acad Sci U S A* 106: 11412-11417.
- Garty H, & Edelman IS (1983). Amiloride-sensitive trypsinization of apical sodium channels. Analysis of hormonal regulation of sodium transport in toad bladder. *J Gen Physiol* 81: 785-803.
- Gerevich Z, Zadori ZS, Koles L, Kopp L, Milius D, Wirkner K, *et al.* (2007). Dual effect of acid pH on purinergic P2X3 receptors depends on the histidine 206 residue. *J Biol Chem* 282: 33949-33957.
- Gille T, Randrianarison-Pellan N, Goolaerts A, Dard N, Uzunhan Y, Ferrary E, *et al.* (2014). Hypoxia-induced inhibition of epithelial Na(+) channels in the lung. Role of Nedd4-2 and the ubiquitin-proteasome pathway. *Am J Respir Cell Mol Biol* 50: 526-537.
- Gonzales EB, Kawate T, & Gouaux E (2009). Pore architecture and ion sites in acid-sensing ion channels and P2X receptors. *Nature* 460: 599-604.
- Gottfried JA, & Chesler M (1996). Temporal resolution of activity-dependent pH shifts in rat hippocampal slices. *J Neurophysiol* 76: 2804-2807.
- Grunder S, & Augustinowski K (2012). Toxin binding reveals two open state structures for one acid-sensing ion channel. *Channels (Austin)* 6: 409-413.

- Guipponi M, Vuagniaux G, Wattenhofer M, Shibuya K, Vazquez M, Dougherty L, *et al.* (2002). The transmembrane serine protease (TMPRSS3) mutated in deafness DFNB8/10 activates the epithelial sodium channel (ENaC) in vitro. *Hum Mol Genet* 11: 2829-2836.
- Haerteis S, Krappitz A, Krappitz M, Murphy JE, Bertog M, Krueger B, *et al.* (2014). Proteolytic activation of the human epithelial sodium channel by trypsin IV and trypsin I involves distinct cleavage sites. *J Biol Chem* 289: 19067-19078.
- Haerteis S, Krappitz M, Bertog M, Krappitz A, Baraznenok V, Henderson I, *et al.* (2012). Proteolytic activation of the epithelial sodium channel (ENaC) by the cysteine protease cathepsin-S. *Pflugers Arch* 464: 353-365.
- Harris M, Firsov D, Vuagniaux G, Stutts MJ, & Rossier BC (2007). A novel neutrophil elastase inhibitor prevents elastase activation and surface cleavage of the epithelial sodium channel expressed in *Xenopus laevis* oocytes. *J Biol Chem* 282: 58-64.
- Harris M, Garcia-Caballero A, Stutts MJ, Firsov D, & Rossier BC (2008). Preferential assembly of epithelial sodium channel (ENaC) subunits in *Xenopus* oocytes: role of furin-mediated endogenous proteolysis. *J Biol Chem* 283: 7455-7463.
- Hesselager M, Timmermann DB, & Ahring PK (2004). pH dependency and desensitization kinetics of heterologously expressed combinations of acid-sensing ion channel subunits. *J Biol Chem* 279: 11006-11015.
- Highstein SM, Holstein GR, Mann MA, & Rabbitt RD (2014). Evidence that protons act as neurotransmitters at vestibular hair cell-calyx afferent synapses. *Proc Natl Acad Sci USA* 111: 5421-5426.
- Hille B (1992) *Ionic Channels of Excitable Membranes*. 2nd edn. Sinauer Associates, Inc.: Sunderland, MA.
- Hirsh AJ, Molino BF, Zhang J, Astakhova N, Geiss WB, Sargent BJ, *et al.* (2006). Design, synthesis, and structure-activity relationships of novel 2-substituted pyrazinoylguanidine epithelial sodium channel blockers: drugs for cystic fibrosis and chronic bronchitis. *J Med Chem* 49: 4098-4115.
- Hirsh AJ, Sabater JR, Zamurs A, Smith RT, Paradiso AM, Hopkins S, *et al.* (2004). Evaluation of second generation amiloride analogs as therapy for cystic fibrosis lung disease. *J Pharmacol Exp Ther* 311: 929-938.
- Hirsh AJ, Zhang J, Zamurs A, Fleegle J, Thelin WR, Caldwell RA, *et al.* (2008). Pharmacological properties of N-(3,5-diamino-6-chloropyrazine-2-carbonyl)-N'-4-[4-(2,3-dihydroxypropoxy)phenyl] butyl-guanidine methanesulfonate (552-02), a novel epithelial sodium channel blocker with potential clinical efficacy for cystic fibrosis lung disease. *J Pharmacol Exp Ther* 325: 77-88.
- Hobbs CA, Blanchard MG, Alijevic O, Tan CD, Kellenberger S, Bencharit S, *et al.* (2013). Identification of the SPLUNC1 ENaC-inhibitory domain yields novel strategies to treat sodium hyperabsorption in cystic fibrosis airway epithelial cultures. *Am J Physiol Lung Cell Mol Physiol* 305: L990-L1001.

Honore P, Wismer CT, Mikusa J, Zhu CZ, Zhong C, Gauvin DM, *et al.* (2005). A-425619 [1-isoquinolin-5-yl-3-(4-trifluoromethyl-benzyl)-urea], a novel transient receptor potential type V1 receptor antagonist, relieves pathophysiological pain associated with inflammation and tissue injury in rats. *J Pharmacol Exp Ther* 314: 410-421.

Hoyer D, & Bartfai T (2012). Neuropeptides and neuropeptide receptors: drug targets, and peptide and non-peptide ligands: a tribute to Prof. Dieter Seebach. *Chemistry & biodiversity* 9: 2367-2387.

Hsu KS, & Huang CC (1997). Characterization of the anoxia-induced long-term synaptic potentiation in area CA1 of the rat hippocampus. *Br J Pharmacol* 122: 671-681.

Hughes PA, Brierley SM, Young RL, & Blackshaw LA (2007). Localization and comparative analysis of acid-sensing ion channel (ASIC1, 2, and 3) mRNA expression in mouse colonic sensory neurons within thoracolumbar dorsal root ganglia. *J Comp Neurol* 500: 863-875.

Hughey RP, Bruns JB, Kinlough CL, Harkleroad KL, Tong Q, Carattino MD, *et al.* (2004). Epithelial sodium channels are activated by furin-dependent proteolysis. *J Biol Chem* 279: 18111-18114.

Hughey RP, Mueller GM, Bruns JB, Kinlough CL, Poland PA, Harkleroad KL, *et al.* (2003). Maturation of the epithelial Na<sup>+</sup> channel involves proteolytic processing of the alpha- and gamma-subunits. *J Biol Chem* 278: 37073-37082.

Hummler E, Barker P, Gatzky J, Beermann F, Verdumo C, Schmidt A, *et al.* (1996). Early death due to defective neonatal lung liquid clearance in alpha-ENaC-deficient mice. *Nat Genet* 12: 325-328.

Hummler E, Barker P, Talbot C, Wang Q, Verdumo C, Grubb B, *et al.* (1997). A mouse model for the renal salt-wasting syndrome pseudohypoaldosteronism. *Proc Natl Acad Sci U S A* 94: 11710-11715.

Immke DC, & McCleskey EW (2003). Protons open acid-sensing ion channels by catalyzing relief of Ca<sup>2+</sup> blockade. *Neuron* 37: 75-84.

Jasti J, Furukawa H, Gonzales EB, & Gouaux E (2007). Structure of acid-sensing ion channel 1 at 1.9 Å resolution and low pH. *Nature* 449: 316-323.

Jeggle P, Callies C, Tarjus A, Fassot C, Fels J, Oberleithner H, *et al.* (2013). Epithelial sodium channel stiffens the vascular endothelium in vitro and in Liddle mice. *Hypertension* 61: 1053-1059.

Jeggle P, Smith ES, Stewart AP, Haerteis S, Korbmacher C, & Edwardson JM (2015). Atomic force microscopy imaging reveals the formation of ASIC/ENaC cross-clade ion channels. *Biochem Biophys Res Commun* 464: 38-44.

Jendelová P, & Syková E (1991). Role of glia in K<sup>+</sup> and pH homeostasis in the neonatal rat spinal cord. *Glia* 4: 56-63.

- Ji HL, & Benos DJ (2004). Degenerin sites mediate proton activation of dbg-epithelial sodium channel. *J Biol Chem* 279: 26939-26947.
- Ji HL, Zhao R, Komissarov AA, Chang Y, Liu Y, & Matthay MA (2015). Proteolytic regulation of epithelial sodium channels by urokinase plasminogen activator: cutting edge and cleavage sites. *J Biol Chem* 290: 5241-5255.
- Jones NG, Slater R, Cadiou H, McNaughton P, & McMahon SB (2004). Acid-Induced Pain and Its Modulation in Humans. *J Neurosci* 24: 10974-10979.
- Kang D, & Kim D (2006). TREK-2 (K2P10.1) and TRESK (K2P18.1) are major background K<sup>+</sup> channels in dorsal root ganglion neurons. *Am J Physiol Cell Physiol* 291: C138-146.
- Kang S, Jang JH, Price MP, Gautam M, Benson CJ, Gong H, *et al.* (2012). Simultaneous disruption of mouse ASIC1a, ASIC2 and ASIC3 genes enhances cutaneous mechanosensitivity. *PloS one* 7: e35225.
- Kapoor N, Lee W, Clark E, Bartoszewski R, McNicholas CM, Latham CB, *et al.* (2011). Interaction of ASIC1 and ENaC subunits in human glioma cells and rat astrocytes. *Am J Physiol Cell Physiol* 300: C1246-1259.
- Kartner N, Augustinas O, Jensen TJ, Naismith AL, & Riordan JR (1992). Mislocalization of delta F508 CFTR in cystic fibrosis sweat gland. *Nat Genet* 1: 321-327.
- Kashlan OB, Adelman JL, Okumura S, Blobner BM, Zuzek Z, Hughey RP, *et al.* (2011). Constraint-based, homology model of the extracellular domain of the epithelial Na<sup>+</sup> channel alpha subunit reveals a mechanism of channel activation by proteases. *J Biol Chem* 286: 649-660.
- Kashlan OB, & Kleyman TR (2011). ENaC structure and function in the wake of a resolved structure of a family member. *Am J Physiol Renal Physiol* 301: F684-696.
- Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, & He J (2005). Global burden of hypertension: analysis of worldwide data. *Lancet* 365: 217-223.
- Keeble J, Russell F, Curtis B, Starr A, Pinter E, & Brain SD (2005). Involvement of transient receptor potential vanilloid 1 in the vascular and hyperalgesic components of joint inflammation. *Arthritis and rheumatism* 52: 3248-3256.
- Kehl SJ, Eduljee C, Kwan DCH, Zhang S, & Fedida D (2002). Molecular determinants of the inhibition of human Kv1.5 potassium currents by external protons and Zn(2+). *J Physiol (Lond)* 541: 9-24.
- Kellenberger S, & Grutter T (2015). Architectural and functional similarities between trimeric ATP-gated P2X receptors and acid-sensing ion channels. *J Mol Biol* 427: 54-66.
- Kellenberger S, & Schild L (2015). International Union of Basic and Clinical Pharmacology. XCI. Structure, Function, and Pharmacology of Acid-Sensing Ion Channels and the Epithelial Na<sup>+</sup> Channel. *Pharmacol Rev* 67: 1-35.

- Keppner A, Andreassen D, Merillat AM, Bapst J, Ansermet C, Wang Q, *et al.* (2015). Epithelial Sodium Channel-Mediated Sodium Transport Is Not Dependent on the Membrane-Bound Serine Protease CAP2/Tmprss4. *PLoS one* 10: e0135224.
- King BF, Wildman SS, Ziganshina LE, Pintor J, & Burnstock G (1997). Effects of extracellular pH on agonism and antagonism at a recombinant P2X2 receptor. *Br J Pharmacol* 121: 1445-1453.
- Kirino T (1982). Delayed neuronal death in the gerbil hippocampus following ischemia. *Brain Res* 239: 57-69.
- Kohler D, App E, Schmitz-Schumann M, Wurtemberger G, & Matthys H (1986). Inhalation of amiloride improves the mucociliary and the cough clearance in patients with cystic fibrosis. *Eur J Respir Dis Suppl* 146: 319-326.
- Kosari F, Sheng S, Li J, Mak DO, Foskett JK, & Kleyman TR (1998). Subunit stoichiometry of the epithelial sodium channel. *J Biol Chem* 273: 13469-13474.
- Kota P, Garcia-Caballero A, Dang H, Gentsch M, Stutts MJ, & Dokholyan NV (2012). Energetic and structural basis for activation of the epithelial sodium channel by matriptase. *Biochemistry* 51: 3460-3469.
- Kovacikova J, Winter C, Loffing-Cueni D, Loffing J, Finberg KE, Lifton RP, *et al.* (2006). The connecting tubule is the main site of the furosemide-induced urinary acidification by the vacuolar H<sup>+</sup>-ATPase. *Kidney Int* 70: 1706-1716.
- Kraig RP, Ferreira-Filho CR, & Nicholson C (1983). Alkaline and acid transients in cerebellar microenvironment. *J Neurophysiol* 49: 831-850.
- Kreitzer MA, Collis LP, Molina AJA, Smith PJS, & Malchow RP (2007). Modulation of extracellular proton fluxes from retinal horizontal cells of the catfish by depolarization and glutamate. *J Gen Physiol* 130: 169-182.
- Kreple CJ, Lu Y, Taugher RJ, Schwager-Gutman AL, Du J, Stump M, *et al.* (2014). Acid-sensing ion channels contribute to synaptic transmission and inhibit cocaine-evoked plasticity. *Nat Neurosci* 17: 1083-1091.
- Krishtal OA, Osipchuk YV, Shelest TN, & Smirnoff SV (1987). Rapid extracellular pH transients related to synaptic transmission in rat hippocampal slices. *Brain Res* 436: 352-356.
- Krueger B, Schlotzer-Schrehardt U, Haerteis S, Zenkel M, Chankiewicz VE, Amann KU, *et al.* (2012). Four subunits (alpha-beta-gamma-delta) of the epithelial sodium channel (ENaC) are expressed in the human eye in various locations. *Invest Ophthalmol Vis Sci* 53: 596-604.
- Kuduk SD, Di Marco CN, Bodmer-Narkevitch V, Cook SP, Cato MJ, Jovanovska A, *et al.* (2010). Synthesis, structure-activity relationship, and pharmacological profile of analogs of the ASIC-3 inhibitor A-317567. *ACS Chem Neurosci* 1: 19-24.

- Kuduk SD, Di Marco CN, Chang RK, Dipardo RM, Cook SP, Cato MJ, *et al.* (2009). Amiloride derived inhibitors of acid-sensing ion channel-3 (ASIC3). *Bioorg Med Chem Lett* 19: 2514-2518.
- Kuroiwa T, Bonnekoh P, & Hossmann KA (1990). Therapeutic window of CA1 neuronal damage defined by an ultrashort-acting barbiturate after brain ischemia in gerbils. *Stroke* 21: 1489-1493.
- Kusche-Vihrog K, Tarjus A, Fels J, & Jaisser F (2014). The epithelial Na<sup>+</sup> channel: a new player in the vasculature. *Curr Opin Nephrol Hypertens* 23: 143-148.
- Lang F, & Shumilina E (2013). Regulation of ion channels by the serum- and glucocorticoid-inducible kinase SGK1. *Faseb J* 27: 3-12.
- Lazrak A, Nita I, Subramaniam D, Wei S, Song W, Ji HL, *et al.* (2009). Alpha(1)-antitrypsin inhibits epithelial Na<sup>+</sup> transport in vitro and in vivo. *Am J Respir Cell Mol Biol* 41: 261-270.
- Le Moellic C, Ouvrard-Pascaud A, Capurro C, Cluzeaud F, Fay M, Jaisser F, *et al.* (2004). Early nongenomic events in aldosterone action in renal collecting duct cells: PKC $\alpha$  activation, mineralocorticoid receptor phosphorylation, and cross-talk with the genomic response. *J Am Soc Nephrol* 15: 1145-1160.
- Lei S, Orser BA, Thatcher GR, Reynolds JN, & MacDonald JF (2001). Positive allosteric modulators of AMPA receptors reduce proton-induced receptor desensitization in rat hippocampal neurons. *J Neurophysiol* 85: 2030-2038.
- Lesage F, & Barhanin J (2011). Molecular Physiology of pH-Sensitive Background K2P Channels. *Physiology* 26: 424-437.
- Lesage F, & Lazdunski M (2000). Molecular and functional properties of two-pore-domain potassium channels. *Am J Physiol Renal Physiol* 279: F793-801.
- Levenson JW, Skerrett PJ, & Gaziano JM (2002). Reducing the global burden of cardiovascular disease: the role of risk factors. *Prev Cardiol* 5: 188-199.
- Leviel F, Hubner CA, Houillier P, Morla L, El Moghrabi S, Brideau G, *et al.* (2010). The Na<sup>+</sup>-dependent chloride-bicarbonate exchanger SLC4A8 mediates an electroneutral Na<sup>+</sup> reabsorption process in the renal cortical collecting ducts of mice. *J Clin Invest* 120: 1627-1635.
- Lewis C, Neidhart S, Holy C, North RA, Buell G, & Surprenant A (1995). Coexpression of P2X2 and P2X3 receptor subunits can account for ATP-gated currents in sensory neurons. *Nature* 377: 432-435.
- Li T, Yang Y, & Canessa CM (2014). A method for activation of endogenous acid-sensing ion channel 1a (ASIC1a) in the nervous system with high spatial and temporal precision. *J Biol Chem* 289: 15441-15448.
- Li WG, Yu Y, Zhang ZD, Cao H, & Xu TL (2011). ASIC3 channels integrate agmatine and multiple inflammatory signals through the nonproton ligand sensing domain. *Mol Pain* 6: 88.

- Light AR, Hughen RW, Zhang J, Rainier J, Liu Z, & Lee J (2008). Dorsal root ganglion neurons innervating skeletal muscle respond to physiological combinations of protons, ATP, and lactate mediated by ASIC, P2X, and TRPV1. *J Neurophysiol* 100: 1184-1201.
- Lin S-H, Chien Y-C, Chiang W-W, Liu Y-Z, Lien C-C, & Chen C-C (2015). Genetic mapping of ASIC4 and contrasting phenotype to ASIC1a in modulating innate fear and anxiety. *Eur J Neurosci* 41: 1553-1568.
- Lingueglia E, Champigny G, Lazdunski M, & Barbry P (1995). Cloning of the amiloride-sensitive FMRFamide peptide-gated sodium channel. *Nature* 378: 730-733.
- Lingueglia E, de Weille JR, Bassilana F, Heurteaux C, Sakai H, Waldmann R, *et al.* (1997). A modulatory subunit of acid sensing ion channels in brain and dorsal root ganglion cells. *J Biol Chem* 272: 29778-29783.
- Lingueglia E, Deval E, & Lazdunski M (2006). FMRFamide-gated sodium channel and ASIC channels: a new class of ionotropic receptors for FMRFamide and related peptides. *Peptides* 27: 1138-1152.
- Loffing J, Zecevic M, Feraille E, Kaissling B, Asher C, Rossier BC, *et al.* (2001). Aldosterone induces rapid apical translocation of ENaC in early portion of renal collecting system: possible role of SGK. *Am J Physiol Renal Physiol* 280: F675-682.
- Lu Y, Ma X, Sabharwal R, Snitsarev V, Morgan D, Rahmouni K, *et al.* (2009). The ion channel ASIC2 is required for baroreceptor and autonomic control of the circulation. *Neuron* 64: 885-897.
- Makani S, Chen HY, Esquenazi S, Shah GN, Waheed A, Sly WS, *et al.* (2012). NMDA receptor-dependent afterdepolarizations are curtailed by carbonic anhydrase 14: regulation of a short-term postsynaptic potentiation. *J Neurosci* 32: 16754-16762.
- Makani S, & Chesler M (2010). Rapid rise of extracellular pH evoked by neural activity is generated by the plasma membrane calcium ATPase. *J Neurophysiol* 103: 667-676.
- Makhanova N, Sequeira-Lopez ML, Gomez RA, Kim HS, & Smithies O (2006). Disturbed homeostasis in sodium-restricted mice heterozygous and homozygous for aldosterone synthase gene disruption. *Hypertension* 48: 1151-1159.
- Mall M, Grubb BR, Harkema JR, O'Neal WK, & Boucher RC (2004). Increased airway epithelial Na<sup>+</sup> absorption produces cystic fibrosis-like lung disease in mice. *Nat Med* 10: 487-493.
- Mall MA, Button B, Johannesson B, Zhou Z, Livraghi A, Caldwell RA, *et al.* (2010). Airway surface liquid volume regulation determines different airway phenotypes in liddle compared with betaENaC-overexpressing mice. *J Biol Chem* 285: 26945-26955.
- Malsure S, Wang Q, Charles RP, Sergi C, Perrier R, Christensen BM, *et al.* (2014). Colon-specific deletion of epithelial sodium channel causes sodium loss and aldosterone resistance. *J Am Soc Nephrol* 25: 1453-1464.



- Mamenko M, Zaika O, Prieto MC, Jensen VB, Doris PA, Navar LG, *et al.* (2013). Chronic angiotensin II infusion drives extensive aldosterone-independent epithelial Na<sup>+</sup> channel activation. *Hypertension* 62: 1111-1122.
- Martin GR, & Jain RK (1994). Noninvasive measurement of interstitial pH profiles in normal and neoplastic tissue using fluorescence ratio imaging microscopy. *Cancer research* 54: 5670-5674.
- Marunaka Y, Hagiwara N, & Tohda H (1992). Insulin activates single amiloride-blockable Na channels in a distal nephron cell line (A6). *The American journal of physiology* 263: F392-400.
- Mathie A, & Veale EL (2015). Two-pore domain potassium channels: potential therapeutic targets for the treatment of pain. *Pflugers Arch* 467: 931-943.
- Mauro T, Guitard M, Behne M, Oda Y, Crumrine D, Komuves L, *et al.* (2002). The ENaC channel is required for normal epidermal differentiation. *J Invest Dermatol* 118: 589-594.
- Mazucca M, Heurteaux C, Alloui A, Diochot S, Baron A, Voilley N, *et al.* (2007). A tarantula peptide against pain via ASIC1a channels and opioid mechanisms. *Nat Neurosci* 10: 943-945.
- McDonald FJ, Yang B, Hrstka RF, Drummond HA, Tarr DE, McCray PB, Jr., *et al.* (1999). Disruption of the beta subunit of the epithelial Na<sup>+</sup> channel in mice: hyperkalemia and neonatal death associated with a pseudohypoaldosteronism phenotype. *Proc Natl Acad Sci U S A* 96: 1727-1731.
- Michaelson DM, & Angel I (1980). Determination of delta pH in cholinergic synaptic vesicles: its effect on storage and release of acetylcholine. *Life Sci* 27: 39-44.
- Mickle AD, Shepherd AJ, & Mohapatra DP (2015). Sensory TRP channels: the key transducers of nociception and pain. *Prog Mol Biol Transl Sci* 131: 73-118.
- Miesenbock G, De Angelis DA, & Rothman JE (1998). Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* 394: 192-195.
- Mironova E, Bugaj V, Roos KP, Kohan DE, & Stockand JD (2012). Aldosterone-independent regulation of the epithelial Na<sup>+</sup> channel (ENaC) by vasopressin in adrenalectomized mice. *Proc Natl Acad Sci U S A* 109: 10095-10100.
- Mitchell KJ, Pinson KI, Kelly OG, Brennan J, Zupicich J, Scherz P, *et al.* (2001). Functional analysis of secreted and transmembrane proteins critical to mouse development. *Nat Genet* 28: 241-249.
- Mogil JS, Breese NM, Witty M-F, Ritchie J, Rainville M-L, Ase A, *et al.* (2005). Transgenic Expression of a Dominant-Negative ASIC3 Subunit Leads to Increased Sensitivity to Mechanical and Inflammatory Stimuli. *J Neurosci* 25: 9893-9901.

- Mordasini D, Loffing-Cueni D, Loffing J, Beatrice R, Maillard MP, Hummler E, *et al.* (2015). ENaC activity in collecting ducts modulates NCC in cirrhotic mice. *Pflugers Arch* 467: 2529-2539.
- Morral N, Bertranpetit J, Estivill X, Nunes V, Casals T, Gimenez J, *et al.* (1994). The origin of the major cystic fibrosis mutation (delta F508) in European populations. *Nat Genet* 7: 169-175.
- Mott DD, Washburn MS, Zhang S, & Dingledine RJ (2003). Subunit-dependent modulation of kainate receptors by extracellular protons and polyamines. *J Neurosci* 23: 1179-1188.
- Mutch WA, & Hansen AJ (1984). Extracellular pH changes during spreading depression and cerebral ischemia: mechanisms of brain pH regulation. *J Cereb Blood Flow Metab* 4: 17-27.
- Nesterov V, Dahlmann A, Krueger B, Bertog M, Loffing J, & Korbmayer C (2012). Aldosterone-dependent and -independent regulation of the epithelial sodium channel (ENaC) in mouse distal nephron. *Am J Physiol Renal Physiol* 303: F1289-1299.
- Nichols DP, Jiang D, Happoldt C, Berman R, & Chu HW (2015). Therapeutic Effects of alpha1-Antitrypsin on *Pseudomonas aeruginosa* Infection in ENaC Transgenic Mice. *PloS one* 10: e0141232.
- Obel AO (1989). Placebo-controlled trial of potassium supplements in black patients with mild essential hypertension. *J Cardiovasc Pharmacol* 14: 294-296.
- Olivier R, Scherrer U, Horisberger JD, Rossier BC, & Hummler E (2002). Selected contribution: limiting Na(+) transport rate in airway epithelia from alpha-ENaC transgenic mice: a model for pulmonary edema. *J Appl Physiol* 93: 1881-1887.
- Omerbasic D, Schuhmacher LN, Bernal Sierra YA, Smith ES, & Lewin GR (2015). ASICs and mammalian mechanoreceptor function. *Neuropharmacol* 94: 80-86.
- Palmer MJ, Hull C, Vigh J, & von Gersdorff H (2003). Synaptic cleft acidification and modulation of short-term depression by exocytosed protons in retinal bipolar cells. *J Neurosci* 23: 11332-11341.
- Pan XQ, Gonzalez JA, Chang S, Chacko S, Wein AJ, & Malykhina AP (2010). Experimental colitis triggers the release of substance P and calcitonin gene-related peptide in the urinary bladder via TRPV1 signaling pathways. *Exp Neurol* 225: 262-273.
- Passero CJ, Mueller GM, Myerburg MM, Carattino MD, Hughey RP, & Kleyman TR (2012). TMPRSS4-dependent activation of the epithelial sodium channel requires cleavage of the gamma-subunit distal to the furin cleavage site. *Am J Physiol Renal Physiol* 302: F1-8.
- Passero CJ, Mueller GM, Rondon-Berrios H, Tofovic SP, Hughey RP, & Kleyman TR (2008). Plasmin activates epithelial Na<sup>+</sup> channels by cleaving the gamma subunit. *J Biol Chem* 283: 36586-36591.
- Patel AB, Chao J, & Palmer LG (2012). Tissue kallikrein activation of the epithelial Na channel. *Am J Physiol Renal Physiol* 303: F540-550.

- Paukert M, Babini E, Pusch M, & Grunder S (2004). Identification of the Ca<sup>2+</sup> blocking site of acid-sensing ion channel (ASIC) 1: implications for channel gating. *J Gen Physiol* 124: 383-394.
- Pearce D, Soundararajan R, Trimpert C, Kashlan OB, Deen PMT, & Kohan DE (2015). Collecting Duct Principal Cell Transport Processes and Their Regulation. *Clin J Am Soc Nephro* 10: 135-146.
- Pech V, Thumova M, Dikalov SI, Hummler E, Rossier BC, Harrison DG, *et al.* (2013). Nitric oxide reduces Cl<sup>-</sup> absorption in the mouse cortical collecting duct through an ENaC-dependent mechanism. *Am J Physiol Renal Physiol* 304: F1390-1397.
- Pech V, Thumova M, Kim YH, Agazatian D, Hummler E, Rossier BC, *et al.* (2012). ENaC inhibition stimulates Cl<sup>-</sup> secretion in the mouse cortical collecting duct through an NKCC1-dependent mechanism. *Am J Physiol Renal Physiol* 303: F45-55.
- Peigneur S, Beress L, Moller C, Mari F, Forssmann WG, & Tytgat J (2012). A natural point mutation changes both target selectivity and mechanism of action of sea anemone toxins. *FASEB J* 26: 5141-5151.
- Pellegrini-Giampietro DE, Zukin RS, Bennett MV, Cho S, & Pulsinelli WA (1992). Switch in glutamate receptor subunit gene expression in CA1 subfield of hippocampus following global ischemia in rats. *Proc Natl Acad Sci U S A* 89: 10499-10503.
- Peng Z, Li WG, Huang C, Jiang YM, Wang X, Zhu MX, *et al.* (2015). ASIC3 Mediates Itch Sensation in Response to Coincident Stimulation by Acid and Nonproton Ligand. *Cell reports* 13: 387-398.
- Perrier R, Boscardin E, Malsure S, Sergi C, Maillard MP, Loffing J, *et al.* (2015). Severe Salt-Losing Syndrome and Hyperkalemia Induced by Adult Nephron-Specific Knockout of the Epithelial Sodium Channel alpha-Subunit. *J Am Soc Nephrol* in press: 10.1681/ASN.2015020154.
- Perrier R, Richard S, Sainte-Marie Y, Rossier BC, Jaisser F, Hummler E, *et al.* (2005). A direct relationship between plasma aldosterone and cardiac L-type Ca<sup>2+</sup> current in mice. *J Physiol* 569: 153-162.
- Petroff EY, Price MP, Snitsarev V, Gong H, Korovkina V, Abboud FM, *et al.* (2008). Acid-sensing ion channels interact with and inhibit BK K<sup>+</sup> channels. *Proc Natl Acad Sci U S A* 105: 3140-3144.
- Picard N, Eladari D, El Moghrabi S, Planes C, Bourgeois S, Houillier P, *et al.* (2008). Defective ENaC processing and function in tissue kallikrein-deficient mice. *J Biol Chem* 283: 4602-4611.
- Pidoplichko VI, Aroniadou-Anderjaska V, Prager EM, Figueiredo TH, Almeida-Suhett CP, Miller SL, *et al.* (2014). ASIC1a Activation Enhances Inhibition in the Basolateral Amygdala and Reduces Anxiety. *J Neurosci* 34: 3130-3141.

- Planes C, Randrianarison NH, Charles RP, Frateschi S, Cluzeaud F, Vuagniaux G, *et al.* (2009). ENaC-mediated alveolar fluid clearance and lung fluid balance depend on the channel-activating protease 1. *EMBO molecular medicine* 2: 26-37.
- Plant LD (2012). A Role for K2P Channels in the Operation of Somatosensory Nociceptors. *Front Mol Neurosci* 5: 21.
- Poirot O, Berta T, Decosterd I, & Kellenberger S (2006). Distinct ASIC currents are expressed in rat putative nociceptors and are modulated by nerve injury. *J Physiol* 576: 215-234.
- Poirot O, Vukicevic M, Boesch A, & Kellenberger S (2004). Selective regulation of acid-sensing ion channel 1 by serine proteases. *J Biol Chem* 279: 38448-38457.
- Poulsen SB, Praetorius J, Damkier HH, Miller L, Nelson RD, Hummler E, *et al.* (2015). Reducing alphaENaC expression in kidney connecting tubule induces pseudohypoaldosteronism type 1 symptoms during K<sup>+</sup> loading. *Am J Physiol Renal Physiol*: ajrenal.00258.02015.
- Pradervand S, Barker PM, Wang Q, Ernst SA, Beermann F, Grubb BR, *et al.* (1999a). Salt restriction induces pseudohypoaldosteronism type 1 in mice expressing low levels of the beta-subunit of the amiloride-sensitive epithelial sodium channel. *Proc Natl Acad Sci U S A* 96: 1732-1737.
- Pradervand S, Vandewalle A, Bens M, Gautschi I, Loffing J, Hummler E, *et al.* (2003). Dysfunction of the epithelial sodium channel expressed in the kidney of a mouse model for Liddle syndrome. *J Am Soc Nephrol* 14: 2219-2228.
- Pradervand S, Wang Q, Burnier M, Beermann F, Horisberger JD, Hummler E, *et al.* (1999b). A mouse model for Liddle's syndrome. *J Am Soc Nephrol* 10: 2527-2533.
- Pratt JH (2005). Central role for ENaC in development of hypertension. *J Am Soc Nephrol* 16: 3154-3159.
- Prole DL, Lima PA, & Marrion NV (2003). Mechanisms underlying modulation of neuronal KCNQ2/KCNQ3 potassium channels by extracellular protons. *J Gen Physiol* 122: 775-793.
- Quintana P, Soto D, Poirot O, Zonouzi M, Kellenberger S, Muller D, *et al.* (2015). Acid-sensing ion channel 1a drives AMPA receptor plasticity following ischaemia and acidosis in hippocampal CA1 neurons. *J Physiol* 593: 4373-4386.
- Ramachandran R, Noorbakhsh F, Defea K, & Hollenberg MD (2012). Targeting proteinase-activated receptors: therapeutic potential and challenges. *Nat Rev Drug Discov* 11: 69-86.
- Ramachandran S, Karp PH, Jiang P, Ostedgaard LS, Walz AE, Fisher JT, *et al.* (2012). A microRNA network regulates expression and biosynthesis of wild-type and DeltaF508 mutant cystic fibrosis transmembrane conductance regulator. *Proc Natl Acad Sci U S A* 109: 13362-13367.

- Randrianarison N, Clerici C, Ferreira C, Fontayne A, Pradervand S, Fowler-Jaeger N, *et al.* (2008). Low expression of the beta-ENaC subunit impairs lung fluid clearance in the mouse. *Am J Physiol Lung Cell Mol Physiol* 294: L409-416.
- Randrianarison N, Escoubet B, Ferreira C, Fontayne A, Fowler-Jaeger N, Clerici C, *et al.* (2007). beta-Liddle mutation of the epithelial sodium channel increases alveolar fluid clearance and reduces the severity of hydrostatic pulmonary oedema in mice. *J Physiol* 582: 777-788.
- Raouf R, Rugiero F, Kieseewetter H, Hatch R, Hummler E, Nassar MA, *et al.* (2012). Sodium channels and mammalian sensory mechanotransduction. *Mol Pain* 8: 21.
- Rengarajan S, Lee DH, Oh YT, Delpire E, Youn JH, & McDonough AA (2014). Increasing plasma [K<sup>+</sup>] by intravenous potassium infusion reduces NCC phosphorylation and drives kaliuresis and natriuresis. *Am J Physiol Renal Physiol* 306: F1059-1068.
- Rossier BC (2014). Epithelial sodium channel (ENaC) and the control of blood pressure. *Current opinion in pharmacology* 15C: 33-46.
- Rossier BC, Baker ME, & Studer RA (2015). Epithelial sodium transport and its control by aldosterone: the story of our internal environment revisited. *Physiol Rev* 95: 297-340.
- Rossier BC, Staub O, & Hummler E (2013). Genetic dissection of sodium and potassium transport along the aldosterone-sensitive distal nephron: Importance in the control of blood pressure and hypertension. *FEBS Lett* 587: 1929-1941.
- Rossol-Haseroth K, Zhou Q, Braun S, Boldyreff B, Falkenstein E, Wehling M, *et al.* (2004). Mineralocorticoid receptor antagonists do not block rapid ERK activation by aldosterone. *Biochem Biophys Res Commun* 318: 281-288.
- Rubera I, Loffing J, Palmer LG, Frindt G, Fowler-Jaeger N, Sauter D, *et al.* (2003). Collecting duct-specific gene inactivation of alphaENaC in the mouse kidney does not impair sodium and potassium balance. *J Clin Invest* 112: 554-565.
- Rusch A, & Hummler E (1999). Mechano-electrical transduction in mice lacking the alpha-subunit of the epithelial sodium channel. *Hearing research* 131: 170-176.
- Saha C, Eckert GJ, Ambrosius WT, Chun TY, Wagner MA, Zhao Q, *et al.* (2005). Improvement in blood pressure with inhibition of the epithelial sodium channel in blacks with hypertension. *Hypertension* 46: 481-487.
- Salinas M, Besson T, Delettre Q, Diochot S, Boulakirba S, Douguet D, *et al.* (2014). Binding site and inhibitory mechanism of the mambalgin-2 pain-relieving peptide on acid-sensing ion channel 1a. *J Biol Chem* 289: 13363-13373.
- Sallenave JM (2010). Secretory leukocyte protease inhibitor and elafin/trappin-2: versatile mucosal antimicrobials and regulators of immunity. *Am J Respir Cell Mol Biol* 42: 635-643.

Sandoz G, Douguet D, Chatelain F, Lazdunski M, & Lesage F (2009). Extracellular acidification exerts opposite actions on TREK1 and TREK2 potassium channels via a single conserved histidine residue. *Proc Natl Acad Sci U S A* 106: 14628-14633.

Santulli G (2013). Epidemiology of cardiovascular disease in the 21st century: updated numbers and updated facts. *J Cardiovasc Disease* 1: 1-2.

Schild L, Schneeberger E, Gautschi I, & Firsov D (1997). Identification of amino acid residues in the alpha, beta, and gamma subunits of the epithelial sodium channel (ENaC) involved in amiloride block and ion permeation. *J Gen Physiol* 109: 15-26.

Schroeder CI, Rash LD, Vila-Farres X, Rosengren KJ, Mobli M, King GF, *et al.* (2014). Chemical synthesis, 3D structure, and ASIC binding site of the toxin mambalgin-2. *Angew Chem Int Ed Engl* 53: 1017-1020.

Sheng S, Carattino MD, Bruns JB, Hughey RP, & Kleyman TR (2006). Furin cleavage activates the epithelial Na<sup>+</sup> channel by relieving Na<sup>+</sup> self-inhibition. *Am J Physiol Renal Physiol* 290: F1488-1496.

Sherwood TW, Lee KG, Gormley MG, & Askwith CC (2011). Heteromeric acid-sensing ion channels (ASICs) composed of ASIC2b and ASIC1a display novel channel properties and contribute to acidosis-induced neuronal death. *J Neurosci* 31: 9723-9734.

Shi S, Carattino MD, Hughey RP, & Kleyman TR (2013). ENaC regulation by proteases and shear stress. *Curr Mol Pharmacol* 6: 28-34.

Siesjö BK, von Hanwehr R, Nergelius G, Nevander G, & Ingvar M (1985). Extra- and intracellular pH in the brain during seizures and in the recovery period following the arrest of seizure activity. *J Cereb Blood Flow Metab* 5: 47-57.

Sluka KA, & Gregory NS (2015). The dichotomized role for acid sensing ion channels in musculoskeletal pain and inflammation. *Neuropharmacology* 94: 58-63.

Sluka KA, Winter OC, & Wemmie JA (2009). Acid-sensing ion channels: A new target for pain and CNS diseases. *Curr Opin Drug Discov Devel* 12: 693-704.

Smith ES, Cadiou H, & McNaughton PA (2007). Arachidonic acid potentiates acid-sensing ion channels in rat sensory neurons by a direct action. *Neuroscience* 145: 686-698.

Smith SE, & Chesler M (1999). Effect of divalent cations on AMPA-evoked extracellular alkaline shifts in rat hippocampal slices. *J Neurophysiol* 82: 1902-1908.

Somodi S, Varga Z, Hajdu P, Starkus JG, Levy DI, Gaspar R, *et al.* (2004). pH-dependent modulation of Kv1.3 inactivation: role of His399. *Am J Physiol Cell Physiol* 287: C1067-1076.

Sonneville F, Ruffin M, Guillot L, Rousselet N, Le Rouzic P, Corvol H, *et al.* (2015). New insights about miRNAs in cystic fibrosis. *Am J Pathol* 185: 897-908.

- Sorensen MV, Grossmann S, Roesinger M, Gresko N, Todkar AP, Barmettler G, *et al.* (2013). Rapid dephosphorylation of the renal sodium chloride cotransporter in response to oral potassium intake in mice. *Kidney Int* 83: 811-824.
- Souslova V, Cesare P, Ding Y, Akopian AN, Stanfa L, Suzuki R, *et al.* (2000). Warm-coding deficits and aberrant inflammatory pain in mice lacking P2X3 receptors. *Nature* 407: 1015-1017.
- Starkus JG, Fleig A, & Penner R (2010). The calcium-permeable non-selective cation channel TRPM2 is modulated by cellular acidification. *J Physiol (Lond)* 588: 1227-1240.
- Su J, Tang Y, Liu L, Zhou H, & Dong Q (2011). Regulation of acid-sensing ion channel 1a function by tissue kallikrein may be through channel cleavage. *Neurosci Lett* 490: 46-51.
- Suzuki M, Mizuno A, Kodaira K, & Imai M (2003). Impaired pressure sensation in mice lacking TRPV4. *J Biol Chem* 278: 22664-22668.
- Svenningsen P, Bistrup C, Friis UG, Bertog M, Haerteis S, Krueger B, *et al.* (2009). Plasmin in nephrotic urine activates the epithelial sodium channel. *J Am Soc Nephrol* 20: 299-310.
- Syková E, & Svoboda J (1990). Extracellular alkaline-acid-alkaline transients in the rat spinal cord evoked by peripheral stimulation. *Brain Res* 512: 181-189.
- Szabo R, Uzzun Sales K, Kosa P, Shylo NA, Godiksen S, Hansen KK, *et al.* (2012). Reduced prostatic (CAP1/PRSS8) activity eliminates HAI-1 and HAI-2 deficiency-associated developmental defects by preventing matriptase activation. *PLoS genetics* 8: e1002937.
- Tanaka H, Nagaike K, Takeda N, Itoh H, Kohama K, Fukushima T, *et al.* (2005). Hepatocyte growth factor activator inhibitor type 1 (HAI-1) is required for branching morphogenesis in the chorioallantoic placenta. *Mol Cell Biol* 25: 5687-5698.
- Thomas W, McEneaney V, & Harvey BJ (2007). Rapid responses to steroid hormones in the kidney. *Nephron Physiology* 107: p1-9.
- Todkar A, Picard N, Loffing-Cueni D, Sorensen MV, Mihailova M, Nesterov V, *et al.* (2015). Mechanisms of renal control of potassium homeostasis in complete aldosterone deficiency. *J Am Soc Nephrol* 26: 425-438.
- Tombaugh GC, & Somjen GG (1996). Effects of extracellular pH on voltage-gated Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>2+</sup> currents in isolated rat CA1 neurons. *J Physiol (Lond)* 493 ( Pt 3): 719-732.
- Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K, *et al.* (1998). The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21: 531-543.
- Tong C-K, Chen K, & Chesler M (2006). Kinetics of activity-evoked pH transients and extracellular pH buffering in rat hippocampal slices. *J Neurophysiol* 95: 3686-3697.
- Traynelis SF, & Cull-Candy SG (1990). Proton inhibition of N-methyl-D-aspartate receptors in cerebellar neurons. *Nature* 345: 347-350.

- Traynelis SF, & Cull-Candy SG (1991). Pharmacological properties and H<sup>+</sup> sensitivity of excitatory amino acid receptor channels in rat cerebellar granule neurones. *J Physiol (Lond)* 433: 727-763.
- Ugawa S, Ishida Y, Ueda T, Inoue K, Nagao M, & Shimada S (2007). Nafamostat mesilate reversibly blocks acid-sensing ion channel currents. *Biochem Biophys Res Commun* 363: 203-208.
- Ugawa S, Ueda T, Ishida Y, Nishigaki M, Shibata Y, & Shimada S (2002). Amiloride-blockable acid-sensing ion channels are leading acid sensors expressed in human nociceptors. *J Clin Invest* 110: 1185-1190.
- Ugawa S, Yamamoto T, Ueda T, Ishida Y, Inagaki A, Nishigaki M, *et al.* (2003). Amiloride-insensitive currents of the acid-sensing ion channel- 2a (ASIC2a)/ASIC2b heteromeric sour-taste receptor channel. *J Neurosci* 23: 3616-3622.
- Valeyev AY, Hackman JC, Holohean AM, Wood PM, Katz JL, & Davidoff RA (1999). GABA-Induced Cl<sup>-</sup> current in cultured embryonic human dorsal root ganglion neurons. *J Neurophysiol* 82: 1-9.
- Vallet V, Chraïbi A, Gaeggeler HP, Horisberger JD, & Rossier BC (1997). An epithelial serine protease activates the amiloride-sensitive sodium channel. *Nature* 389: 607-610.
- Vallon V, Hummler E, Rieg T, Pochynyuk O, Bugaj V, Schroth J, *et al.* (2009). Thiazolidinedione-induced fluid retention is independent of collecting duct alphaENaC activity. *J Am Soc Nephrol* 20: 721-729.
- van Bemmelen MX, Huser D, Gautschi I, & Schild L (2015). The Human Acid-Sensing Ion Channel ASIC1a: Evidence for a Homotetrameric Assembly State at the Cell Surface. *PLoS one* 10: e0135191.
- van der Lubbe N, Moes AD, Rosenbaek LL, Schoep S, Meima ME, Danser AH, *et al.* (2013). K<sup>+</sup>-induced natriuresis is preserved during Na<sup>+</sup> depletion and accompanied by inhibition of the Na<sup>+</sup>-Cl<sup>-</sup> cotransporter. *Am J Physiol Renal Physiol* 305: F1177-1188.
- van der Lubbe N, Zietse R, & Hoorn EJ (2013). Effects of angiotensin II on kinase-mediated sodium and potassium transport in the distal nephron. *Curr Opin Nephrol Hypertens* 22: 120-126.
- Vavassori S, & Mayer A (2014). A new life for an old pump: V-ATPase and neurotransmitter release. *J Cell Biol* 205: 7-9.
- Venton BJ, Michael DJ, & Wightman RM (2003). Correlation of local changes in extracellular oxygen and pH that accompany dopaminergic terminal activity in the rat caudate-putamen. *J Neurochem* 84: 373-381.
- Verouti SN, Boscardin E, Hummler E, & Frateschi S (2015). Regulation of blood pressure and renal function by NCC and ENaC: lessons from genetically engineered mice. *Curr Opin Pharmacol* 21: 60-72.



- Vessey JP, Stratis AK, Daniels BA, Da Silva N, Jonz MG, Lalonde MR, *et al.* (2005). Proton-mediated feedback inhibition of presynaptic calcium channels at the cone photoreceptor synapse. *J Neurosci* 25: 4108-4117.
- Vick JS, & Askwith CC (2015). ASICs and neuropeptides. *Neuropharmacol* 94: 36-41.
- Vidt DG, & Borazanian RA (2003). Treat high blood pressure sooner: tougher, simpler JNC 7 guidelines. *Cleve Clin J Med* 70: 721-728.
- Vilin YY, Peters CH, & Ruben PC (2012). Acidosis differentially modulates inactivation in na(v)1.2, na(v)1.4, and na(v)1.5 channels. *Front Pharmacol* 3: 109.
- Vitzthum H, Seniuk A, Schulte LH, Muller ML, Hetz H, & Ehmke H (2014). Functional coupling of renal K<sup>+</sup> and Na<sup>+</sup> handling causes high blood pressure in Na<sup>+</sup> replete mice. *J Physiol* 592: 1139-1157.
- Voilley N, de Weille J, Mamet J, & Lazdunski M (2001). Nonsteroid anti-inflammatory drugs inhibit both the activity and the inflammation-induced expression of acid-sensing ion channels in nociceptors. *J Neurosci* 21: 8026-8033.
- Vollmer WM, Sacks FM, Ard J, Appel LJ, Bray GA, Simons-Morton DG, *et al.* (2001). Effects of diet and sodium intake on blood pressure: subgroup analysis of the DASH-sodium trial. *Ann Intern Med* 135: 1019-1028.
- Vuagniaux G, Vallet V, Jaeger NF, Pfister C, Bens M, Farman N, *et al.* (2000). Activation of the amiloride-sensitive epithelial sodium channel by the serine protease mCAP1 expressed in a mouse cortical collecting duct cell line. *J Am Soc Nephrol* 11: 828-834.
- Vukicevic M, & Kellenberger S (2004). Modulatory effects of acid-sensing ion channels on action potential generation in hippocampal neurons. *Am J Physiol Cell Physiol* 287: C682-690.
- Vukicevic M, Weder G, Boillat A, Boesch A, & Kellenberger S (2006). Trypsin Cleaves Acid-sensing Ion Channel 1a in a Domain That Is Critical for Channel Gating. *J Biol Chem* 281: 714-722.
- Wakida N, Kitamura K, Tuyen DG, Maekawa A, Miyoshi T, Adachi M, *et al.* (2006). Inhibition of prostasin-induced ENaC activities by PN-1 and regulation of PN-1 expression by TGF-beta1 and aldosterone. *Kidney Int* 70: 1432-1438.
- Waldmann R, Bassilana F, Deweille J, Champigny G, Heurteaux C, & Lazdunski M (1997). Molecular cloning of a non-inactivating proton-gated Na<sup>+</sup> channel specific for sensory neurons. *J Biol Chem* 272: 20975-20978.
- Waldmann R, Champigny G, Bassilana F, Heurteaux C, & Lazdunski M (1997). A proton-gated cation channel involved in acid-sensing. *Nature* 386: 173-177.
- Walker CS, Conner AC, Poyner DR, & Hay DL (2010). Regulation of signal transduction by calcitonin gene-related peptide receptors. *Trends Pharmacol Sci* 31: 476-483.

Wang Q, Clement S, Gabbiani G, Horisberger JD, Burnier M, Rossier BC, *et al.* (2004). Chronic hyperaldosteronism in a transgenic mouse model fails to induce cardiac remodeling and fibrosis under a normal-salt diet. *Am J Physiol Renal Physiol* 286: F1178-1184.

Wang Q, Hummler E, Maillard M, Nussberger J, Rossier BC, Brunner HR, *et al.* (2001). Compensatory up-regulation of angiotensin II subtype 1 receptors in alpha ENaC knockout heterozygous mice. *Kidney Int* 59: 2216-2221.

Wang X, Li WG, Yu Y, Xiao X, Cheng J, Zeng WZ, *et al.* (2013). Serotonin facilitates peripheral pain sensitivity in a manner that depends on the nonproton ligand sensing domain of ASIC3 channel. *J Neurosci* 33: 4265-4279.

Weber F, & Anlauf M (2014). Treatment resistant hypertension--investigation and conservative management. *Dtsch Arztebl Int* 111: 425-431.

Weber MA (2014). The evolving clinical management of hypertension. *J Clin Hypertens (Greenwich)* 16: 917-924.

Weller K, Reeh PW, & Sauer SK (2011). TRPV1, TRPA1, and CB1 in the isolated vagus nerve-axonal chemosensitivity and control of neuropeptide release. *Neuropeptides* 45: 391-400.

Wemmie JA, Chen J, Askwith CC, Hruska-Hageman AM, Price MP, Nolan BC, *et al.* (2002). The acid-activated ion channel ASIC contributes to synaptic plasticity, learning, and memory. *Neuron* 34: 463-477.

Wemmie JA, Price MP, & Welsh MJ (2006). Acid-sensing ion channels: advances, questions and therapeutic opportunities. *Trends Neurosci* 29: 578-586.

Wemmie JA, Taugher RJ, & Kreple CJ (2013). Acid-sensing ion channels in pain and disease. *Nature Rev Neurosci* 14: 461-471.

Whelton PK *et al.* (1997). Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Arch Intern Med* 157: 657-667.

Wiemuth D, Assmann M, & Grunder S (2014). The bile acid-sensitive ion channel (BASIC), the ignored cousin of ASICs and ENaC. *Channels (Austin)* 8: 29-34.

Wilkins ME, Hosie AM, & Smart TG (2005). Proton modulation of recombinant GABA(A) receptors: influence of GABA concentration and the beta subunit TM2-TM3 domain. *J Physiol* 567: 365-377.

Williams GH (2005). Aldosterone biosynthesis, regulation, and classical mechanism of action. *Heart Fail Rev* 10: 7-13.

- Wolkenberg SE, Zhao Z, Mulhearn JJ, Harrison ST, Sanders JM, Cato MJ, *et al.* (2011). High concentration electrophysiology-based fragment screen: discovery of novel acid-sensing ion channel 3 (ASIC3) inhibitors. *Bioorg Med Chem Lett* 21: 2646-2649.
- Wu P-Y, Huang Y-Y, Chen C-C, Hsu T-T, Lin Y-C, Weng J-Y, *et al.* (2013). Acid-sensing ion channel-1a is not required for normal hippocampal LTP and spatial memory. *J Neurosci* 33: 1828-1832.
- Xiong ZG, Zhu XM, Chu XP, Minami M, Hey J, Wei WL, *et al.* (2004). Neuroprotection in ischemia: Blocking calcium-permeable acid-sensing ion channels. *Cell* 118: 687-698.
- Yagi J, Wenk HN, Naves LA, & McCleskey EW (2006). Sustained currents through ASIC3 ion channels at the modest pH changes that occur during myocardial ischemia. *Circulation research* 99: 501-509.
- Yamamoto F, Borgula GA, & Steinberg RH (1992). Effects of light and darkness on pH outside rod photoreceptors in the cat retina. *Exp Eye Res* 54: 685-697.
- Yamamura H, Ugawa S, Ueda T, Nagao M, & Shimada S (2004). Protons activate the delta-subunit of the epithelial Na<sup>+</sup> channel in humans. *J Biol Chem* 279: 12529-12534.
- Yamauchi M, Kataoka H, Itoh H, Seguchi T, Hasui Y, & Osada Y (2004). Hepatocyte growth factor activator inhibitor types 1 and 2 are expressed by tubular epithelium in kidney and down-regulated in renal cell carcinoma. *J Urol* 171: 890-896.
- Yang HY, Charles RP, Hummler E, Baines DL, & Isseroff RR (2013). The epithelial sodium channel mediates the directionality of galvanotaxis in human keratinocytes. *J Cell Sci* 126: 1942-1951.
- Yu D, Thelin WR, Rogers TD, Stutts MJ, Randell SH, Grubb BR, *et al.* (2012). Regional differences in rat conjunctival ion transport activities. *Am J Physiol Cell Physiol* 303: C767-780.
- Yu Y, Chen Z, Li WG, Cao H, Feng EG, Yu F, *et al.* (2010). A nonproton ligand sensor in the acid-sensing ion channel. *Neuron* 68: 61-72.
- Zachar RM, Skjodt K, Marcussen N, Walter S, Toft A, Nielsen MR, *et al.* (2015). The epithelial sodium channel gamma-subunit is processed proteolytically in human kidney. *J Am Soc Nephrol* 26: 95-106.
- Zani ML, Tanga A, Saidi A, Serrano H, Dallet-Choisy S, Baranger K, *et al.* (2011). SLPI and trappin-2 as therapeutic agents to target airway serine proteases in inflammatory lung diseases: current and future directions. *Biochem Soc Trans* 39: 1441-1446.
- Zemel MB (1997). Dietary pattern and hypertension: the DASH study. *Dietary Approaches to Stop Hypertension. Nutr Rev* 55: 303-305.
- Zha XM (2013). Acid-sensing ion channels: trafficking and synaptic function. *Mol Brain* 6: 1.

Zha XM, Wemmie JA, Green SH, & Welsh MJ (2006). Acid-sensing ion channel 1a is a postsynaptic proton receptor that affects the density of dendritic spines. *Proc Natl Acad Sci U S A* 103: 16556-16561.

Zhai J, Peoples RW, & Li C (1998). Proton inhibition of GABA-activated current in rat primary sensory neurons. *Pflugers Arch* 435: 539-545.

Zhou Z, Duerr J, Johannesson B, Schubert SC, Treis D, Harm M, *et al.* (2011). The ENaC-overexpressing mouse as a model of cystic fibrosis lung disease. *J Cystic Fibrosis* 10 Suppl 2: S172-182.

Zhou Z, Treis D, Schubert SC, Harm M, Schatterny J, Hirtz S, *et al.* (2008). Preventive but not late amiloride therapy reduces morbidity and mortality of lung disease in betaENaC-overexpressing mice. *Am J Respir Crit Care Med* 178: 1245-1256.

Zhu G, Chanchevalap S, Cui N, & Jiang C (1999). Effects of intra- and extracellular acidifications on single channel Kir2.3 currents. *J Physiol (Lond)* 516 ( Pt 3): 699-710.

Ziemann AE, Allen JE, Dahdaleh NS, Drebot, II, Coryell MW, Wunsch AM, *et al.* (2009). The amygdala is a chemosensor that detects carbon dioxide and acidosis to elicit fear behavior. *Cell* 139: 1012-1021.

## Figure legends

### Figure 1. Relation and roles of ENaC and ASICs.

(A) Phylogenetic tree of the ENaC/degenerin family, showing besides ASIC and ENaC the subfamilies pickpocket (PPK), degenerin (DEG), the FMRFa-activated channel FaNaC and the bile acid-sensing ion channel BASIC (also known as hINaC or BLINaC). (B) Illustration of the different physiological and pathological roles of ASICs and ENaC.

### Figure 2. ASIC and ENaC subunit organization.

(A) Schematic view of one subunit in the context of the trimeric ASIC, highlighting the different domains finger (purple), knuckle (turquoise),  $\beta$ -ball (orange), palm (yellow), thumb (blue) and transmembrane domains (red). (B) Structure of an ASIC1 subunit, based on the crystal structure obtained from chicken ASIC1 binding MitTx (Bacongus, Bohlen, Goehring, Julius & Gouaux, 2014). The domains are colored as in (A). (C) Model of the extracellular part of  $\alpha$ ENaC (Kashlan & Kleyman, 2011). Coloring as in (A); the transmembrane part was not modeled.

### Figure 3. Functional properties of ASIC.

(A) Action potential induction by extracellular acidification to pH 6.4, mediated by ASICs, measured by whole-cell current-clamp from a mouse hippocampal neuron. (B) pH5-induced

current recorded in whole-cell voltage-clamp to -60 mV from a Chinese hamster ovary cell stably transfected with ASIC1a. (C) pH dependence of steady state desensitization (open circles) and of activation (black squares) of ASIC1a. In steady-state desensitization experiments, cells were exposed for 55 s to the indicated conditioning pH, and a stimulation pH 6 solution was applied for 5 s to open the not yet desensitized channels. The normalized current response at pH 6 is plotted as a function of the conditioning pH. For ASIC1a activation, the cells were perfused by a pH 7.4 solution and once per minute this solution was changed to one of acidic pH to open the channels. The normalized current is plotted as a function of the stimulation pH.

#### **Figure 4. Extracellular pH changes during neurotransmission in CNS.**

Illustration based on experiments performed in hippocampal, photoreceptor, amygdala and calyx synapses, showing a synapse between a pre- and postsynaptic neuronal terminal and an astrocyte. The numbers 1-3 (colored in blue) represent potential mechanisms for the initial alkaline shift recorded in hippocampal synapses: neurotransmission stimulates calcium signaling in the perisynaptic cells. As a consequence the mechanisms of calcium buffering system are stimulated. One of these mechanisms involves the PMCA, which pumps the accumulated intracellular calcium ions out of the cell in exchange for external protons, leading to extracellular alkalinization. The numbers 4-6 (colored in red) represent potential mechanisms underlying the acidic shift. The glial  $\text{Na}^+\text{-HCO}_3^-$ -cotransporter (NBC) causes a gradual extracellular acidification (4). Intense neurotransmission increases astrocytic energy demand, resulting in lactate and  $\text{CO}_2$  production (5). For clarity only the lactate into presynaptic terminals is shown. Lactate is transported outside the astrocyte by the  $\text{H}^+$ -coupled monocarboxylate transporter (MCT) and  $\text{CO}_2$  can freely diffuse, also leading to extracellular acidification at the synaptic cleft. Fast acidifications at the synaptic cleft may also occur as a result of synaptic vesicle release and also by non-vesicular release involving an undefined transporter (6). For simplification, only the transport of protons and of  $\text{HCO}_3^-$  is shown in the figure (grey arrows; protons are red dots). The black arrows indicate the activation of synaptic vesicle release by calcium entry.

#### **Figure 5. Extracellular pH sensitive ion channels in the PNS.**

Illustration based on experiments performed in small-diameter DRG neurons innervating peripheral organs. The scheme indicates the pathway of a signal from the sensory nerve

terminals in an organ (in the example the heart) to the spinal cord and the brain and focuses on two specific locations, the peripheral nerve terminal (left) and the signaling along the sensory nerve axon (right). Only the ion channels that are modulated by pHe on the nerve terminal and along the axon are shown, indicated by the symbols “+” (stimulation by lowering of pHe) and “-” (inhibition by lowering of pHe). Red symbols indicate that this modulation increases excitability, and blue symbols indicate that it decreases excitability. The number of symbols correlates with the pH dependence of the regulation (Table 4) as indicated at the bottom of the figure.

**Figure 6. ENaC expression along the aldosterone-sensitive distal nephron**

Schematic view of a nephron, with detailed views of the ion transport mechanisms in cells of the DCT, CNT and CCD. CLCN/Kb, voltage-sensitive chloride channel Kb; ROMK, renal outer medullary potassium channel.

## Tables

**Table 1. Links to online information in the IUPHAR/BPS Guide to PHARMACOLOGY**

### Targets

Acid-sensing (proton-gated) ion channels (ASICs)	118
Cystic fibrosis transmembrane conductance regulator	129
Ionotropic glutamate receptors	75
Epithelial sodium channels (ENaC)	122
Na-Cl symporter	970
P2X receptors	77
Prostasin	2400
Renin	2413
Sodium-dependent HCO <sub>3</sub> <sup>-</sup> transporter	171
Transient Receptor Potential channels	78
Two-P potassium channels	79
Voltage-gated Na <sup>+</sup> channels	82
WNK lysine deficient protein kinase	643

### Ligands

amiloride	2421
A-317567	4116
APETx2	4135
agmatine	4127
aldosterone	2872
arcaine	4137
benzamil	4145
Calcitonin gene-related peptide	681
GMQ	4203
nafamostat	4262
P552-02	4280
phenamil	4281
Psalmotoxin 1	4292
triamterene	4329
vasopressin	2168

(The numbers on the right column represent the target or ligand identification number in the database. They will be necessary for making the hyperlinks, should however not appear in the final table.)

**Table 2 Biophysical properties of ASICs**

	pH <sub>50</sub> activation	pHD <sub>50</sub> Steady-state desensitization	Desensitization time constant (s)	Most important sites of expression
ASIC1a	6.2-6.6	~7.2	~0.4	CNS, PNS
ASIC1b	5.9-6.3	~6.7	~0.9	PNS
ASIC2a	4.0-4.9	~5.6	~1.4	CNS, PNS
ASIC3	6.4-6.7	~7.1	~0.3	PNS

The desensitization time constant is indicated for stimulation pH close to the pH<sub>50</sub>. These data are from different articles and reviews (Alijevic & Kellenberger, 2012; Blanchard & Kellenberger, 2011; Hesselager, Timmermann & Ahring, 2004; Poirot, Vukicevic, Boesch & Kellenberger, 2004; Wemmie, Taugher & Kreple, 2013). ASIC4, not indicated in the table, has not been shown so far to form functional channels and is mainly expressed in the CNS (Lin, Chien, Chiang, Liu, Lien & Chen, 2015).



**Table 3 ASIC modulators**

<b>Modulator class</b>	<b>Important examples</b>	<b>Effect</b>	<b>Site of action</b>	<b>References</b>
Divalent cations	Ca <sup>2+</sup> , Mg <sup>2+</sup> , Ba <sup>2+</sup>	Acidic shift of pH dependence	Not known	(Babini, Paukert, Geisler & Grunder, 2002; Immke & McCleskey, 2003)
		Pore block	Acidic residues in pore entry of ASIC1a	(Paukert, Babini, Pusch & Grunder, 2004)
Neuropeptides	Dynorphin, FMRFa	Acidic shift of pH dependence of steady-state desensitization, slowing of desensitization and induction of sustained current	Not known	(Askwith, Cheng, Ikuma, Benson, Price & Welsh, 2000; Vick & Askwith, 2015)
Proteases	Tissue kallikrein, trypsin	Acidic shift of pH dependence	Cleavage in finger domain	(Poirot, Vukicevic, Boesch & Kellenberger, 2004; Su, Tang, Liu, Zhou & Dong, 2011; Vukicevic, Weder, Boillat, Boesch & Kellenberger, 2006)
Protein kinases	PKA, PKC	Changes expression and function	Intracellular	Rev. in (Kellenberger & Schild, 2015; Wemmie, Taugher & Kreple, 2013)
Other	Arachidonic acid	Increase of peak current amplitude	Not known	(Allen & Attwell, 2002; Smith, Cadiou & McNaughton, 2007)
	Nitric oxide	Increase of peak current amplitude	Extracellular	(Cadiou et al., 2007)

**Table 4: extracellular pH-sensitive ion channels expressed in CNS and PNS neurons**

<b>Inhibition by pHe (pH<sub>50</sub>Inh.)</b>	<b>Ion channels</b>	<b>Mechanisms / Comments</b>
pH 8.6 (PNS)	(1) TASK2	Decrease of K <sup>+</sup> pore occupancy and open probability
pH 7.4 - pH 7.3 (CNS, PNS)	(2) TREK1, (3) Kv1.4, (4) TASK1, (5) TRAAK, (6) KCNQ2/3	(2) C-type inactivation enhanced, (4) open probability reduced, (6) I max reduced, depolarizing shift of the activation
pH 7.3 - pH 7 (CNS, PNS)	(7) NMDAR, (8) GABAAR, (9) voltage-gated calcium channels (Ca <sub>v</sub> ), (10) glycine receptors	(7) Reduced burst duration, decreased opening frequency, (8), (10) subunit- and agonist-dependent, (9) I max reduced, depolarizing shift of the activation
Mixed pH effects (CNS, PNS)	(11) TWIK1	From pH 7.5 to pH 6.5: current enhanced. Below pH 6.5: current inhibited.
pH 6.7 - pH 6.5 (CNS, PNS)	(12) Kir2.3, (13) TASK3, (14) TRPM2,	(12), (13) conductance and open probability reduced
pH 6.3 - pH 6.2 (CNS, PNS)	(15) AMPAR, (16) Ca <sup>2+</sup> -sensing non-selective cation channels, (17) Kv1.5	(15) enhancing desensitization, (17) facilitation of a non-conducting state
pH 6 - pH 4.6 (CNS, PNS)	(18) Kainate receptors, (19) Kv1.3, (20) I <sub>A</sub> , I <sub>K</sub> , (21) I <sub>Na2</sub> , (22) Nav1.2	(18) Subunit dependent, (20) depolarizing shift of the inactivation, (19), (21) I max reduced, (21), (22) depolarizing shift of the activation
<b>Stimulation by pHe (pH<sub>50</sub>Act.)</b>	<b>Ion channels</b>	<b>Mechanisms / Comments</b>
pH 7.3 (CNS, PNS)	(23) TREK2, (24) P2X2	(24) Sensitization to activation by ATP
pH 6.5 - pH6 (CNS, PNS)	(25) ASIC1a, (26) ASIC1a/2b, (27) ASIC3, (28) δβγ-ENaC	
pH 5.8 - pH 3 (CNS, PNS)	(29) ASIC1a/2a, (30) ASIC2a, (31) ASIC2a/2b	
Mixed pH effects (CNS, PNS)	(32) BK and (33) Kv1.3 associated with ASICs	~50% inhibition of BK and Kv1.3 currents at pH 7.4
Mixed pH effects (CNS, PNS)	(34) P2X3	The ATP concentration dependence of the current is shifted to higher concentrations and its maximal current amplitude is increased
(37), (38) ~ pH 5.4 for proton activation (37) pH 7 - pH 6 for	(35) TRPV1, (36) TRPV4	(35), (36) Direct activation by strong acidosis (< pH 6). (35) Sensitization to other stimuli such as capsaicin and temperature by mild acidosis (pH 7-6)

---

proton  
sensitization  
(CNS, PNS)

---

pH 6.5-pH

4 (CNS, PNS) (37) nAChRs  $\alpha 3/\beta 4$  (38) HCN,  $I_h$

(37) Increased of the agonist-induced current and changed kinetics. Agonist- and subunit- dependent.  
(38) Activation of  $I_h$  current

---

*The ion channels are classified based on their pH sensitivity. Table citations: (1), (2), (4), (5), (13), (23):* (Ehling, Cerina, Budde, Meuth & Bittner, 2015; Lesage & Barhanin, 2011) (3): (Claydon et al., 2000). (4), (13): (Plant, 2012). (6): (Prole, Lima & Marrion, 2003). (7), (15), (18): (Traynelis & Cull-Candy, 1990; Traynelis & Cull-Candy, 1991). (8) (Wilkins, Hosie & Smart, 2005; Zhai, Peoples & Li, 1998). (9), (20), (21) (Tombaugh & Somjen, 1996). (10) (Chen, Dillon & Huang, 2004). (11) (12) (Zhu, Chanchevalap, Cui & Jiang, 1999). (14) (Starkus, Fleig & Penner, 2010). (15) (Lei, Orser, Thatcher, Reynolds & MacDonald, 2001). (16) (Chu et al., 2003). (17) (Kehl, Eduljee, Kwan, Zhang & Fedida, 2002). (18) (Mott, Washburn, Zhang & Dingleline, 2003). (19) (Petroff et al., 2008; Somodi et al., 2004). (21) (Daumas & Andersen, 1993). (22) (Vilin, Peters & Ruben, 2012). (24) (King, Wildman, Ziganshina, Pintor & Burnstock, 1997). (25), (30) (Alijevic & Kellenberger, 2012) (26) (Sherwood, Lee, Gormley & Askwith, 2011). (27) (Waldmann, Bassilana, Deweille, Champigny, Heurteaux & Lazdunski, 1997). (28) (Ji & Benos, 2004; Yamamura, Ugawa, Ueda, Nagao & Shimada, 2004). (29) (Bartoi, Augustinowski, Polleichtner, Grunder & Ulbrich, 2014). (31) (Ugawa et al., 2003). (32), (33) (Petroff et al., 2008). (34) (Gerevich et al., 2007) (Tominaga et al., 1998). (36), (Suzuki, Mizuno, Kodaira & Imai, 2003). (37) (Abdrakhmanova, Cleemann, Lindstrom & Morad, 2004; Abdrakhmanova, Dorfman, Xiao & Morad, 2002). (38) (Cichy et al., 2015). (41) (Birdsong et al., 2010). **Table abbreviations:** BK, big calcium-activated  $K^+$  channel; HCN, hyperpolarization-activated cyclic nucleotide-gated channel;  $I_A$ , A-type current of rapid inactivating  $K^+$  channels;  $I_h$ , current produced by HCN channels;  $I_K$ , delayed rectifier  $K^+$  current;  $I_{max}$ , maximal current amplitude;  $I_{Na}$ ,  $Na^+$  current;  $K_{ir}$ , inward rectifier  $K^+$  channel;  $K_v$ , voltage-gated  $K^+$  channel; nAChR, nicotinic acetylcholine receptor;  $Na_v$ , voltage-gated  $Na^+$  channel; P2X, Purinergic receptor;  $pH_{50Inh./Act.}$ , pH of half-maximal inhibition/activation; TASK, two-pore domain  $K^+$  channel; TRAAK, TWIK-related arachidonic acid-stimulated  $K^+$  channel; TREK, TWIK-related  $K^+$  channel; TRPM, transient receptor potential cation channel, subfamily M; TRPV, transient receptor potential cation channel subfamily V; TWIK, tandem of P-domain in a weak inwardly rectifying  $K^+$  channel.

**Table 5. ENaC animal models**

Mouse model	tissue / organ	Phenotype	Reference
$\alpha$ ENaC KO (constitutive)	kidney and lung	neonatal death, lung fluid clearance failure, hyperkalemia and sodium loss	(Hummler et al., 1996)
		increased number of angiotensin II subtype 1 receptors in renal tissues and lowered blood pressure during the angiotensin II receptor blockade	(Wang et al., 2001)
	skin	decreased transcription of the $\alpha 1$ -subunit of $\text{Na}^+, \text{K}^+$ -ATPase	(Blot-Chabaud et al., 2001)
		increased transepidermal water loss, higher skin surface pH, disturbed stratum corneum lipid composition and lamellar body secretion	(Charles et al., 2008)
		impaired directionality of galvanotaxis in keratinocytes	(Yang, Charles, Hummler, Baines & Isseroff, 2013)
		altered epidermal differentiation	(Mauro et al., 2002)
	embryo	inability to rescue the lethal embryonic phenotype in double $\alpha$ ENaC and HAI-1 (hepatocyte growth factor activator inhibitor-1) KOs	(Szabo et al., 2012)
ear	normal mechano-electrical transducer apparatus	(Rusch & Hummler, 1999)	
$\beta$ ENaC KO (constitutive)	kidney	hyperkalemia, neonatal death and type 1 pseudohypoaldosteronism	(McDonald et al., 1999)
$\gamma$ ENaC KO (constitutive)	kidney	hyperkalemia, neonatal death and type 1 pseudohypoaldosteronism	(Barker et al., 1998)
CNT-specific $\alpha$ ENaC KO (Hoxb7, constitutive)	kidney	normal sodium and potassium balance	(Rubera et al., 2003)
		normal ascites development	(Mordasini et al., 2015)
		abolished benzamil-sensitive component of $\text{Cl}^-$ absorption	(Pech et al., 2012)
		protection from lithium-induced nephrogenic diabetes insipidus	(Christensen et al., 2011)
		presence of electroneutral, amiloride-resistant, thiazide-sensitive, transepithelial $\text{NaCl}$ absorption in CNT	(Leviel et al., 2010)
		rosiglitazone-induced fluid retention	(Vallon et al., 2009)
		normal urinary acidification following furosemide alone and in combination with hydrochlorothiazide treatment	(Kovacikova et al., 2006)
CNT-specific $\alpha$ ENaC KO (AQP2, constitutive)	kidney	higher urinary sodium excretion and hyperkalemia under $\text{Na}^+$ -deficient diet	(Christensen et al., 2010)
		attenuation of body weight and water increase following RGZ (rosiglitazone) treatment	(Fu et al., 2015)
nephron-specific $\alpha$ ENaC KO (inducible)		hyperkalemia and body weight loss under regular salt diet	(Perrier et al., 2015)
CNT-specific $\alpha$ ENaC KO (constitutive)		type 1 pseudohypoaldosteronism symptoms during high dietary $\text{K}^+$ loading	(Poulsen et al., 2015)
$\beta$ and $\gamma$ ENaC KOs (constitutive)	sensory neurons	normal mechanosensory behavior	(Raouf et al., 2012)
airway-specific overexpression $\beta$ ENaC-transgene (constitutive)	lung	increased airway $\text{Na}^+$ absorption, airway mucus obstruction and chronic airway inflammation	(Zhou et al., 2011)
		unaltered development of airway cartilage	(Bonora, Riffault, Marie, Mall, Clement & Tabary, 2011)
		therapeutic effects of $\alpha 1$ -antitrypsin on Pseudomonas aeruginosa infection	(Nichols, Jiang, Happoldt, Berman & Chu, 2015)
		defective regulation of airway surface liquid volume and ENaC-mediated $\text{Na}^+$ absorption	(Mall et al., 2010)
$\alpha$ ENaC KO (constitutive)	tongue	complete loss of salt attraction and sodium taste response	(Chandrashekar et al., 2010)
$\beta$ ENaC -	kidney	high blood pressure, metabolic alkalosis, hypokalemia with cardiac and renal	(Pradervand et al., 1999b)

Liddle knock-in (constitutive)		hypertrophy under high salt intake		
		constitutive hyperactivity of ENaC in cortical connecting ducts	(Pradervand et al., 2003)	
		maintained mineralocorticoid regulation of ENaC	(Dahlmann, Pradervand, Hummler, Rossier, Frindt & Palmer, 2003)	
		increased vasopressin-stimulated CFTR Cl <sup>-</sup> currents in CCD cells	(Chang et al., 2005)	
	lung		no effect of hypoxia on amiloride-sensitive alveolar fluid clearance	(Gille et al., 2014)
			increased alveolar fluid clearance and reduced severity of hydrostatic pulmonary edema	(Randrianarison et al., 2007)
			pendrin gene ablation in Liddle homozygous does not eliminate nitric oxide-sensitive net Cl <sup>-</sup> flux and transepithelial potential difference	(Pech et al., 2013)
			intact regulation of airway surface liquid volume and ENaC-mediated Na <sup>+</sup> absorption	(Mall et al., 2010)
	intestine	increased aldosterone responsiveness of ENaC in colon	(Bertog et al., 2008)	
	heart	decreased action potential duration, intracellular Ca <sup>2+</sup> transient amplitude and contraction	(Perrier et al., 2005)	
βENaC hypomorphic mice (constitutive)	kidney	weight loss, hyperkalemia and decreased blood pressure on low-salt diets	(Pradervand et al., 1999a)	
	lung	impaired lung fluid clearance	(Randrianarison et al., 2008)	
αENaC(-/-)Tg (αENaC CMV) (constitutive)	lung, kidney and intestine	rescued perinatal lethal pulmonary phenotype and partially restored Na <sup>+</sup> transport in renal, colonic, and pulmonary epithelia	(Hummler et al., 1997)	
	lung	reduced Na <sup>+</sup> transport rate probably insufficient for airway fluid clearance and favoring pulmonary edema	(Olivier, Scherrer, Horisberger, Rossier & Hummler, 2002)	
		predisposition to pulmonary edema and delayed resolution	(Egli et al., 2004)	
	heart	absence of cardiac remodeling and fibrosis under a normal-salt diet	(Wang et al., 2004)	
increased action potential duration, intracellular Ca <sup>2+</sup> transient amplitude and contraction		(Perrier et al., 2005)		
colon-specific αENaC KO (constitutive)	intestine	sodium loss and aldosterone resistance	(Malsure et al., 2014)	

**Table 6 Proteases modulating ENaC in vitro and in vivo activity**

protease	type	ENaC regulation <i>in vitro</i>	ENaC cleavage	ENaC regulation <i>in vivo</i>
CAP1/Prss8, also known as prostaticin	Membrane-bound serine protease, GPI-anchored	Catalytically-independent and increased ENaC activity in <i>X. oocytes</i> and mCCD cells (Vallet, Chraibi, Gaeggeler, Horisberger & Rossier, 1997; Vuagniaux et al., 2000) (Andreasen, Vuagniaux, Fowler-Jaeger, Hummler & Rossier, 2006).	Cleavage of $\gamma$ ENaC in transfected MDCK cells through catalytically active and inactive CAP1 (Bruns et al., 2007). Probable involvement of a second protease (Carattino et al., 2014).	40% decrease in ENaC-mediated $\text{Na}^+$ currents in alveolar epithelial cells; unaltered ENaC cleavage in lung-specific CAP1 KO mice (Planes et al., 2009). Reduced $\Delta$ PDamil but normal ENaC processing in intestine-specific CAP1 KO (Malsure et al., 2014).
CAP2/Tmprss4	Membrane-bound serine protease, type II	Catalytically-dependent and increased ENaC current in <i>X. oocytes</i> (Vuagniaux et al., 2000).	Cleavage of all three ENaC subunits in <i>X. oocytes</i> (Garcia-Caballero, Dang, He & Stutts, 2008). Cleavage only in $\gamma$ ENaC (Passero, Mueller, Myerburg, Carattino, Hughey & Kleyman, 2012).	Normal $\Delta$ PDamil and ENaC processing in constitutive CAP2 KO (Keppner et al., 2015).
CAP3/matriptase	Membrane-bound serine protease, type II	Catalytically-dependent and increased ENaC current in <i>X. oocytes</i> (Vuagniaux et al., 2000).	Mediates cleavage of ENaC at basic residues near the $\gamma$ -ENaC furin site (Kota, Garcia-Caballero, Dang, Gentzsch, Stutts & Dokholyan, 2012).	Not determined.
TMPRSS3	Membrane-bound serine protease, type II	Increased ENaC current in <i>X. oocytes</i> . TMPRSS3 mutants fail to activate ENaC (Guipponi et al., 2002).	Not determined.	Not determined.
Furin	Intracellular (subtilisin-like proprotein convertase family)	Activated ENaC through suppression $\text{Na}^+$ self-inhibition (Sheng, Carattino, Bruns, Hughey & Kleyman, 2006).	Cleaved $\alpha$ and $\gamma$ ENaC in <i>X. oocytes</i> (Hughey et al., 2004; Hughey et al., 2003).	Not determined.
neutrophil elastase	soluble serine protease	increased ENaC activity in cultured human bronchial epithelia. Increased open probability of near-silent channels in NIH-3T3 cells (Caldwell, Boucher & Stutts, 2005).	neutrophil elastase-dependent activation of $\gamma$ ENaC (Caldwell, Boucher & Stutts, 2005; Harris, Firsov, Vuagniaux, Stutts & Rossier, 2007).	not determined.
kallikrein	soluble serine proteinase	activated ENaC in <i>X. oocytes</i> (Patel, Chao & Palmer, 2012).	ENaC activation and cleavage abolished in mouse ENaC mutants (Patel, Chao & Palmer, 2012).	blunted urinary $\text{Na}^+$ excretion after amiloride injection and decreased colonic $\Delta$ PDamil (Picard et al., 2008), and absence of the cleaved (70-kDa) $\gamma$ ENaC in kallikrein KO mice (Picard et al., 2008).
plasmin	soluble serine proteinase	activated ENaC in M-1 cells and <i>X. oocytes</i> (Svenningsen et al., 2009).	released $\gamma$ ENaC peptide (Passero, Mueller, Rondon-Berrios, Tofovic, Hughey & Kleyman, 2008; Svenningsen et al., 2009).	not determined.
metalloprotease meprin $\beta$ subunit	cell surface or secreted meprin metalloproteinase	increased ENaC in <i>X. oocytes</i> and epithelial cells (Garcia-Caballero et al., 2011).	cleavage of $\alpha$ and $\gamma$ -ENaC subunits (Garcia-Caballero et al., 2011).	not determined.

cathepsin-S	lysosomal cysteine protease	activated ENaC in <i>X.</i> oocytes and M-1 cells (Haerteis et al., 2012).	$\gamma$ ENaC cleavage (Haerteis et al., 2012).	not determined.
urokinase plasminogen activator	membrane-bound serine protease, GPI-anchored	catalytically-dependent activation of ENaC in <i>X.</i> oocytes (Ji, Zhao, Komissarov, Chang, Liu & Matthay, 2015).	$\gamma$ but not $\alpha$ ENaC cleavage in <i>X.</i> oocytes (Ji, Zhao, Komissarov, Chang, Liu & Matthay, 2015).	decreased ENaC activity in primary tracheal epithelial cells from urokinase plasminogen activator KO (Chen et al., 2014).
alkaline phosphatase from <i>Pseudomonas aeruginosa</i>	phosphomonoesterase (periplasmic space in Gram-negative bacteria)	activated ENaC in immortalized and primary human bronchial epithelial cells from both cystic fibrosis and non-cystic fibrosis patients (Butterworth, Zhang, Heidrich, Myerburg & Thibodeau, 2012; Butterworth, Zhang, Liu, Shanks & Thibodeau, 2014).	cleaved $\gamma$ ENaC (Butterworth, Zhang, Heidrich, Myerburg & Thibodeau, 2012).	not determined.
trypsin IV and I		Increased ENaC activity by trypsin IV in human airway epithelial cells (Haerteis et al., 2014).	ENaC cleavage and activation by trypsin IV (but not by trypsin I) (Haerteis et al., 2014).	not determined.




Figure 1, Boscardin et al.






Figure 2, Boscardin et al.




Figure 3, Boscardin et al.




Figure 4, Boscardin et al.




Figure 5, Boscardin et al.




Figure 6, Boscardin et al.