

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: The salivary microbiome for differentiating individuals: proof of principle.

Authors: Leake SL, Pagni M, Falquet L, Taroni F, Greub G

Journal: Microbes and infection / Institut Pasteur

Year: 2016 Jun

Volume: 18

Issue: 6

Pages: 399-405

DOI: 10.1016/j.micinf.2016.03.011

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

1 The salivary microbiome for differentiating individuals: proof of principle

2

3 Sarah L. Leake^a, Marco Pagni^b, Laurent Falquet^{b,c}, Franco Taroni^{aΔ}, Gilbert Greub^{dΔ*}

4

5 a: School of Criminal Justice, University of Lausanne, Lausanne, Switzerland

6

7 b: Swiss Institute of Bioinformatics, Vital-IT group, Lausanne, Switzerland

8

9 c: Department of Biology, University of Fribourg, Fribourg, Switzerland

10

11 d: Institute of Microbiology, Lausanne, Switzerland

12

13

14 * Corresponding author:

15

16 Prof Gilbert Greub

17 Institute of Microbiology

18 University of Lausanne

19 Bugnon 48

20 1011 Lausanne

21 041 21 314 49 79

22 gilbert.greub@chuv.ch

23

24 Δ: Both authors contributed equally to this work

25

26

27

28

29 Keywords: saliva microbiota; forensic science; Illumina sequencing; human identification

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

Abstract

Human identification has played a prominent role in forensic science for the past two decades. Identification based on unique genetic traits is driving the field. However, this may have limitations, for instance, for twins. Moreover, high-throughput sequencing techniques are now available and may provide a high amount of data likely useful in forensic science.

This study investigates the potential for bacteria found in the salivary microbiome to be used to differentiate individuals. Two different targets (16S rRNA and *rpoB*) were chosen to maximise coverage of the salivary microbiome and when combined, they increase the power of differentiation (identification). Paired-end Illumina high-throughput sequencing was used to analyse the bacterial composition of saliva from two different people at four different time points (t=0 and t=28 days and then one year later at t=0 and t=28 days). Five major phyla dominate the samples: Firmicutes, Proteobacteria, Actinobacteria, Bacteroidetes and Fusobacteria. *Streptococcus*, a Firmicutes, is one of the most abundant aerobic genera found in saliva and targeting *Streptococcus rpoB* has enabled a deeper characterisation of the different streptococci species, which cannot be differentiated using 16S rRNA alone. We have observed that samples from the same person group together regardless of time of sampling. The results indicate that it is possible to distinguish two people using the bacterial microbiota present in their saliva.

53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102

1. Introduction

Current methods of human identification in forensic science rely heavily upon the analysis of human DNA. However, there are limitations to the use of human DNA namely its degradation and low quantity. For example, in sexual assault cases, the DNA from the perpetrator is often masked by the DNA of the victim making identification difficult. In such cases saliva is commonly found due to it being transferred through, amongst others, biting, kissing and licking. To overcome the current unsatisfactory situation, the potential of other targets, for example bacteria, needs to be investigated. Why is bacterial DNA interesting in this context? Firstly, bacterial DNA is better protected than human DNA and more resistant to degradation. Therefore, bacterial DNA will persist better once deposited on a surface. Secondly, it may be possible to distinguish twins using bacterial DNA [1], a feat impossible with current human DNA based methods.

It has been estimated that 99% of bacteria found in the environment cannot be cultured [2]. However, with the arrival of next generation sequencing (NGS) the analysis of bacterial community composition has reached depths previously unachievable. There is now potential to exploit bacteria for forensic purposes. Fierer et al. demonstrated that the analysis of the skin microbiome could be used to link an individual to an object they touched and that the bacterial community found on the object was more similar to the community on the owners hand than to 270 other hands, indicating the potential of this technique for forensic identification [3]. This study extends the idea presented by Fierer et al. by demonstrating the potential of NGS analysis of the salivary microbiota for forensic identification.

A number of studies showing saliva bacterial community composition using NGS have been published [1,4-9]. To date the main gene targeted is 16S rRNA because it is ubiquitous and essential for bacterial life [10,11]. However, there are limitations to targeting 16S rRNA namely, intra-genomic heterogeneity, mosaicism and the lack of a universal threshold sequence identity value [12]. Therefore, in order to have a more complete picture of a microbiome, analysing a second (single-copy) target is essential. In this study the second gene targeted was *rpoB* which, encodes the beta-subunit of RNA polymerase, a very important enzyme that is highly conserved throughout bacteria. It has been shown that like the 16S rRNA gene the *rpoB* gene contains alternating variable and conserved regions [13]. The hypervariable regions of *rpoB* have shown promise for bacterial identification down to the species and subspecies levels [14-16]. Specifically studies have shown that humans have many different strains of the same *Streptococcus* species, the most prevalent genus in saliva, with many strains being unique to individuals [17,18]. Using 16S rRNA alone these strains would not be detected and therefore an important part of the salivary microbiome would be missed out. By combining *rpoB* with 16S rRNA a deeper level of identification is possible.

Saliva unlike sperm and blood, the other main biological fluids found in criminal cases, is not sterile. Indeed, saliva contains, as many as 500 million bacterial cells per millilitre (ml) and at least 700 different bacterial species [19]. The average composition of the salivary microbiome being known [1,8], we wondered whether there is enough variation to differentiate salivary microbiomes of two different people. To date, studies have shown that differences in salivary microbial communities between individuals are present [5,20], however whether these differences are great enough to differentiate individuals has yet to be explored. Additionally,

103 the salivary microbiome has been shown to be stable over a couple of months [5,8] but no
104 longer, however studies on gut microbiota show stability over a few years [21,22], further
105 work is required to see if this pattern is observed in saliva microbiota. Thus, this study
106 investigates the intra and inter-individual variation of the salivary microbiome of two healthy
107 subjects to investigate the potential of saliva microbiota in forensic science.

108

109 2. Materials and Methods

110

111 2.1. Sampling and DNA extraction

112 This study was approved by the Ethics Committee of the Canton of Vaud, Switzerland
113 (protocol 357/11). Saliva samples were obtained from two healthy adult individuals at four
114 time points; t=0 and t=30 days and one year later at t=0 and t=30, with informed consent.
115 Volunteers were asked to brush their teeth in the morning and not eat or drink one hour before
116 sampling. The saliva was collected by spitting into a sterile tube and then stored at -20°C until
117 processing. DNA extraction was performed using the automated MagNA Pure 96 DNA and
118 Viral Nucleic Acid small volume kit (Roche) following the Pathogen Universal 200 v2.0
119 protocol [23]. Samples were then stored at -20°C.

120

121 2.2. PCR and sequencing

122 In order to maximise coverage of the salivary microbiome, two different targets were chosen;
123 16S rRNA and *rpoB*. Practically two different pairs of primers targeting *rpoB* were used to
124 investigate the biodiversity of streptococci (*rpoB1*) and other bacteria (*rpoB2*). For 16S
125 rRNA, primers were designed to amplify the V5 region and for *rpoB*, two sets of primers
126 covered the V1 region. Primers were designed using general target species then checked
127 against species known to be found in saliva (see table Table 1 for final primer sequences).
128 Each target was amplified separately in a reaction containing 5 µl of DNA extract, 0.5 µM of
129 both forward and reverse primer, 1x Phusion® HF buffer, 200 µM each dNTP, 0.02U/µl
130 Phusion® Hot Start II DNA polymerase, 3% DMSO and 1mM MgCl₂ in a total volume of 50
131 µl. The following thermal cycling parameters were used: initial denaturation at 98°C for 30
132 seconds, 35 cycles of denaturation at 98°C for 5 seconds, primer dependant annealing
133 temperature (see Table 1 for annealing temperatures) for 15 seconds and extension at 72°C for
134 10 seconds with a final extension of 5 minutes at 72°C.

135

136 Table 1

137

138 All amplified targets from the same sample were pooled together and the pooled sample
139 barcoded. To pool samples equal molar amounts of each sample are necessary, in this case
140 approximately ten picomoles of each were used. The samples were then purified using
141 Agencourt AMPure XP PCR purification (Beckman Coulter). The purified products were
142 then separated on an agarose gel and the band corresponding to the target size (120bp)
143 excised. Finally, the sequencing libraries were prepared using the TruSeq DNA sample
144 preparation kit (Illumina) [24]. Then, 100 cycles of paired-end sequencing were performed on
145 a HiSeq 2000 (Illumina).

146

147 2.3. Sequence analysis

148 Base-calling was performed by HCS 2.0.12/RTA 1.17.21.3 and quality control by the
149 CASAVA 1.8.2 pipeline using standard parameters. Specifically FastQC was used for quality
150 control, by running FastQC in Casava mode the sequences which did not pass the quality
151 threshold were removed [25]. FLASH was used to overlap the paired reads [26]. As each
152 sample contained the sequences for three targets, each target was separated out using barcode

153 splitter (from the FASTX-tool kit [27]) with exact matching for the primer sequence
154 (sequences available in the European Nucleotide Archive under accession number
155 PRJEB6052). This step also removes chimeric sequences.

156
157 Sequences were clustered into operational taxonomic units (OTUs) using CD-HIT-EST 4.5.4
158 [28]. For 16S rRNA 97% identity was used and for *rpoB* 95%. Any clusters containing less
159 than twenty sequences were removed helping to reduce the number of OTUs resulting from
160 sequencing errors and contamination. Then a representative sequence for each cluster was
161 inputted into BLAST and compared against the entire nucleotide database using the best-hit
162 algorithm to give the ‘top’ hit. The same process was carried out for both targets to enable
163 direct comparison of results.

164
165 In order to compare the taxa abundances between the two experiments the data was
166 normalised using DESeq [29], despite it being designed for RNAseq data, it can also be
167 applied to microbiome data [30]. To minimise the effect of highly abundant taxa the data was
168 then transformed by taking the $\log_{10}(x+1)$ of each count (x). To compare the taxa abundances,
169 the samples from each individual were combined and the mean calculated, producing a mean
170 abundance for each individual per taxon, per target gene. Two statistical inferential
171 approaches have been performed. On one side, from a frequentist perspective, a 2-tailed
172 unpaired t-test was used to compare the means (θ_1 for individual 1 and θ_2 for individual 2,
173 respectively) and then the taxa were ranked by p-values. On the other hand, a Bayesian
174 perspective was adopted by calculating Bayes factors (BF) to test the hypothesis $H_0: \theta_1 -$
175 $\theta_2 = 0$ versus $H_1: \theta_1 - \theta_2 \neq 0$. Due to the small sample size hierarchical clustering
176 using the Ward method was used to group the data and a dendrogram used to visualise the
177 grouping. The R packages hclust and as.dendrogram were used to carry out the clustering
178 analyses. To combine data from different targets taxa considered as significant from each
179 target were inputted into a table and hierarchical analysis performed.

180

181

182 **3. Results**

183 *3.1. Illumina sequencing results*

184 The saliva microbiome composition of 2 individuals was explored at 4 different time points.
185 The samples were split into two sequencing runs with samples taken one month apart being
186 sequenced together. Therefore, each run contained two samples per individual making 4
187 samples in total, per run. Run one was performed one year before run two. In total, run one
188 produced 193,221,302 reads. After quality control, pairing and filtering 59,971,947 reads
189 were used for analysis with the following target breakdown: 16S rRNA - 21,534,203, *rpoB1* -
190 29,693,058 and *rpoB2* - 8,744,686. In total, run two produced 201,692,619 reads. After
191 quality filtering and pairing 56,762,234 reads were used for analysis with the following target
192 breakdown: 16S rRNA - 30,604,336, *rpoB1* - 17,007,924 and *rpoB2* - 9,149,974. A
193 breakdown of the number of different OTUs found per sample, per target can be found in
194 Table 2.

195

196 *3.2. Microbiome composition*

197 The use of three targets enables the microbiome composition to be analysed to a greater
198 depth. Fig.1 shows the proportion of the top five phyla per individual and per target. For both
199 *rpoB1* and 16S rRNA, Firmicutes is the most common phyla constituting over 90% and 70%
200 of the population respectively. For *rpoB2* the population is composed of over 90%
201 Actinobacteria. The large difference in taxa found by each *rpoB* primer pair is expected as

202 they were designed to amplify different taxa, demonstrating the benefit of targeting more than
203 one region of the same target gene.

204

205 Fig.1

206

207 The addition of *rpoB* enables certain genera to be analysed down to the species and even
208 strain level. Specifically, with 16S rRNA *Streptococcus* can be detected at the genus level and
209 occasionally the species level (9 different OTUs); however, with *rpoB* it can be detected to
210 the species/strain level (53 different OTUs) enabling a deeper characterisation of this part of
211 the saliva microbiome. This is important as *Streptococcus* makes up about 80% of Firmicutes,
212 the most abundant phylum.

213

214 3.3. Minimum sequences required

215 This study used the HiSeq2000 to analyse the samples, a machine which can produce over
216 one billion reads, as at the outset of this study the number of sequences required to separate
217 two individuals was unknown. To calculate the minimum number of sequences necessary the
218 data were randomly sub-sampled at different levels: 1000, 10000, 50000, 100000, 500000 and
219 1000000 sequences. The analysis was performed to the end and the relative distances
220 calculated between the samples at all levels are shown in Fig.2 For *rpoB2* that provides the
221 smallest separation, at least 50000 sequences were required to adequately discriminate the
222 two investigated individuals. 16S rRNA provides the best separation when looking at the
223 targets individually. However, when 16S rRNA and *rpoB1* are combined the separation is
224 improved. Combining all three targets produces the best separation, however the addition of
225 *rpoB2* does not greatly improve the separation except at 50000 sequences where the
226 separation is significantly improved.

227

228 Fig.2

229

230 3.4. Clustering threshold

231 Unlike previous studies the main aim of this study was to investigate whether the bacteria
232 found in saliva could be used to separate samples from different individuals and not just
233 characterise the microbiome. Different clustering thresholds were tested to see which one
234 gave the best separation taking into account analysis time i.e. the total time required to
235 analyse the data after sequencing. Fig.3 shows that as the percent identity, generally, increases
236 so does the relative distance between the two individuals. The results for both *rpoB* targets are
237 shown in Fig.3A where the dashed line indicates the chosen threshold of 95%. In Fig.3B the
238 dashed line highlights the chosen threshold for 16S rRNA of 97%. These percentages
239 correspond to previously published studies for species level characterisation for *rpoB* and 16S
240 rRNA, respectively [10,31]. For both targets 100% identity provides the best separation
241 however the analysis time, for 16S rRNA especially, is very long and therefore it is not the
242 most efficient solution.

243

244 Fig.3

245

246 3.5. Hierarchical clustering

247 Firstly the normalised logged data was filtered by performing a 2-tailed unpaired t-test and
248 ranking the taxa by p-value and only the taxa with a p-value < 0.1 (and a BF <1) were kept for
249 analysis. The data was further filtered by removing any taxa that did not appear in both
250 experiments. Hierarchical clustering was performed by first calculating the Euclidean distance
251 and then using the Ward method to produce relative distances between each sample. Fig.4

252 shows the dendrograms representing the relative distances between the samples, for each
253 target, (A-C) and then for all targets combined (D). For all targets, samples from different
254 individuals are separated, due to a significant inter-individual variation. Concerning the intra-
255 individual variation samples sequenced in the same run are expected to be more similar and
256 therefore logically grouped together as seen in Fig.4B and D. Conversely, the intra-individual
257 separation for *rpoB1* (Fig.4A) and 16S rRNA (Fig.4C) is not ideal. However, when all three
258 targets are combined good inter and intra-individual separation could also be achieved,
259 demonstrating the benefit of analysing more than one target gene.

260

261 Fig.4

262

263 4. Discussion

264

265 This paper presented the first study into the use of the salivary microbiome for human
266 identification. It has shown that the salivary microbiome exhibits a significant biodiversity
267 and by using a PCR-based metagenomic approach the discrimination of two unrelated
268 individuals was possible. The biodiversity revealed in all samples was similar to that found by
269 previous studies, showing that the designed primers are robust. However, the abundances do
270 differ but this has been observed previously [1].

271

272 Previous studies [1,6,8] have shown that the most common phyla found in saliva are:
273 Firmicutes, Proteobacteria, Actinobacteria, Bacteroidetes and Fusobacteria and this study
274 concurs with these findings; however the abundances differ slightly. Stahringer et al. analysed
275 264 saliva samples and showed that bacterial abundances varied greatly, this study falls
276 within the observed variation. In the same study they defined a genus-level core microbiome
277 containing eight genera [1]. By combining three targets in this study a genus-level core
278 microbiome of 58 genera was observed. This high number of genera covers about 95% of the
279 population of each individual implying that most differences come from the species/strain
280 level. However, this study is limited by a small sample size and more samples may reveal the
281 core microbiome to be similar to previous studies. Such a small sample size was chosen, as
282 the depth of sequencing required to differentiate two individuals was unknown. Therefore,
283 this was one of the major goals of the research. Had too many samples been analysed in one
284 run, the minimum number of sequences required may not have been achieved, so we
285 remained conservative with regards to sample size.

286

287 The results showed that the minimum number of sequences for this type of analysis is
288 100,000 as this provided a good separation between individuals with all targets. However, the
289 addition of *rpoB2* did not significantly increase the discrimination. One of the main
290 advantages of *rpoB* is that it identifies a fair number of species/strains and both primer pairs
291 identify different species. However, *rpoB2* identifies much less than *rpoB1*. Even though the
292 best separation is achieved with sequences of all three target genes, very good separation is
293 still achieved when combining only 16S rRNA and *rpoB1*. Therefore, the choice of target
294 combination would depend on how many samples were to be sequenced in one run. By only
295 using two target genes, more samples could be sequenced, making the technique more
296 economical whilst achieving rather similar results. By choosing a clustering threshold, which
297 enables identification down to the species/strain level whilst remaining time efficient, the
298 whole analysis could be carried out in about one week, depending on which high-throughput
299 sequencer is used.

300

301 To perform the hierarchical clustering the data was filtered to only use the taxa found to be
302 significant by a 2-tailed unpaired t-test (and a $BF < 1$ meaning a support, generally with
303 values that very strongly support the hypothesis H_1). Due to the number of OTUs found
304 obviously not all of them are useful for separating samples from different individuals. To
305 reduce analysis complexity, only OTUs found in both sequencing runs were kept as they
306 could be more accurately attributed to an individual and techniques used in forensic science
307 are required to be as robust as possible. Inevitably there is some natural variation in saliva
308 microbiota due to it being a dynamic fluid and certain bacteria will not always be detected,
309 being either absent or in too few numbers. To ensure that no sequencing errors were included,
310 any clusters containing less than twenty sequences were removed prior to analysis. Even with
311 this highly conservative algorithm, samples from one individual can be successfully separated
312 from those of a second individual (Fig.4) whilst minimising the intra-individual variation.
313 Altogether, our technique proved to be highly robust and is innovative not only for its
314 putative application in forensic science, but also by using a combination of a highly
315 discriminative gene (*rpoB*) with the 16S rRNA target generally used for PCR-based
316 metagenomics. However, the present work only represents a first proof of principle and we
317 need to study twins in order to confirm that saliva microbiota may indeed differentiate twins.
318 A recent study by Stahringer et al. showed that for twins aged between 12-24 years their
319 salivary microbiome was not statistically more similar than for any other pair [1]. This
320 indicates that overall there is very little or no genetic influence on salivary microbiome
321 composition and that the differences observed between twins mainly come from
322 environmental factors. Indeed a number of environmental factors such as diet, oral hygiene,
323 smoking, alcohol and drug consumption may influence the salivary microbiome [1].
324 Therefore a person's microbiome could be used as intelligence to inform about their lifestyle.

325
326 One major environmental factor is antibiotics. Lazarevic et al. described the effects of
327 amoxicillin treatment on the salivary microbiota in children with acute otitis media. They
328 showed that directly after treatment there was a change in the microbiota in terms of both
329 species richness and diversity [32]. However, three weeks after the end of treatment the
330 microbiota had mainly recovered back to pre-antibiotic diversity. This, would only impact
331 cases where the saliva was deposited on a crime scene whilst the perpetrator was taking
332 antibiotics. In such cases, presence of antibiotics in the sample might be determined and an
333 additional sample might then be obtained upon treatment with the same antimicrobial
334 substance. In the case where the perpetrator is taking antibiotics when apprehended a
335 reference sample could be taken at a later date once the salivary microbiome had recovered.

336
337 Another important point to consider with regards to forensic traces is how resistant the traces
338 (i.e. here the bacterial DNA) are to external factors. Indeed, UV light, heat and humidity can
339 degrade human DNA, environmental conditions which are often found at crime scenes. One
340 advantage of microbiota based forensic investigation is that bacterial DNA is better protected
341 from degradation than human DNA as bacterial DNA is circular often highly condensed as
342 "nucleoid" and therefore harder to be degraded by enzymes. Moreover, prokaryotic cells have
343 a cell wall, which is chemically complex with a peptidoglycan matrix that better protects the
344 contents of the cell compared to the cell membrane of eukaryotic cells. Therefore bacterial
345 DNA should be more resistant than eukaryotic DNA to external factors taking longer to be
346 degraded.

347
348 The goal of this technique is not to replace current methods used for human identification but
349 to be complementary. When these methods do not produce satisfactory results there is no
350 other option from a biological identification standpoint. By analysing the salivary

351 microbiome, new options become available that previously were not possible. There are two
352 main applications of this technique in forensic science: human identification and intelligence.
353 The first will only be possible if a reference sample is available. The second application uses
354 the same data but looks at the presence of specific bacteria, which could indicate a certain
355 lifestyle. This information might be used to help guide an investigation. If an identification is
356 not possible then the data acquired could still provide valuable information to a case.
357 However, much more work is needed to relate given species to given lifestyle habits.

358
359 In conclusion, Illumina high-throughput sequencing of the salivary microbiome can be used
360 to identify saliva samples from two different individuals. This technique shows promise for
361 human identification, specifically for twins and other cases where standard DNA typing does
362 not provide satisfactory results due to degradation of human DNA. The results could also be
363 used for intelligence purposes by providing information concerning a person's lifestyle.
364 Further work is required to investigate the benefit and limitations of this technique.

366 **Acknowledgments**

367
368
369 The authors would like to thank Sebastien Aeby (Institute of Microbiology) for his technical
370 support as well as the Lausanne Genomic Technologies Facility for carrying out the
371 sequencing and Vital-IT (Swiss Institute for Bioinformatics) for computing power.

372
373

374 **References**

- 375
- 376 [1] Stahringer SS, Clemente JC, Corley RP, Hewitt J, Knights D, Walters WA, et al.
377 Nurture trumps nature in a longitudinal survey of salivary bacterial communities in
378 twins from early adolescence to early adulthood. *Genome Research* 2012;22:2146–
379 52.
- 380 [2] Handelsman J. Metagenomics: Application of Genomics to Uncultured
381 Microorganisms. *Microbiol Mol Biol Rev* 2004;68:669–85.
- 382 [3] Fierer N, Lauber CL, Zhou N, McDonald D, Costello EK, Knight R. Forensic
383 identification using skin bacterial communities. *P Natl Acad Sci Usa* 2010;107:6477–
384 81.
- 385 [4] Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the Normal Bacterial
386 Flora of the Oral Cavity. *Journal of Clinical Microbiology* 2005;43:5721–32.
- 387 [5] Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, Knight R. Bacterial
388 Community Variation in Human Body Habitats Across Space and Time. *Science*
389 2009;326:1694–7.
- 390 [6] Lazarevic V, Whiteson K, Huse S, Hernandez D, Farinelli L, Østerås M, et al.
391 Metagenomic study of the oral microbiota by Illumina high-throughput sequencing.
392 *Journal of Microbiological Methods* 2009;79:266–71.
- 393 [7] Zaura E, Keijser BJF, Huse SM, Crielaard W. Defining the healthy core microbiome
394 of oral microbial communities. *BMC Microbiology* 2009;9.
- 395 [8] Lazarevic V, Whiteson K, Hernandez D, Francois P, Schrenzel J. Study of inter- and
396 intra-individual variations in the salivary microbiota. *BMC Genomics* 2010;11:523.
- 397 [9] Caporaso JG, Lauber C, Costello E, Berg-Lyons D, Gonzalez A, Stombaugh J, et al.
398 Moving pictures of the human microbiome. *Genome Biology* 2011;12:R50.
- 399 [10] Case RJ, Boucher Y, Dahllorf I, Holmstrom C, Doolittle WF, Kjelleberg S. Use of
400 16S rRNA and rpoB Genes as Molecular Markers for Microbial Ecology Studies.
401 *Applied and Environmental Microbiology* 2007;73:278–88.
- 402 [11] Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification
403 for phylogenetic study. *Journal of Bacteriology* 1991;173:697–703.
- 404 [12] Rajendhran J, Gunasekaran P. Microbial phylogeny and diversity: Small subunit
405 ribosomal RNA sequence analysis and beyond. *Microbiological Research*
406 2011;166:99–110.
- 407 [13] Boor KJ, Duncan ML, Price CW. Genetic and Transcriptional Organization of the
408 Region Encoding the Subunit of *Bacillus subtilis* RNA Polymerase. *Journal of*
409 *Biological Chemistry* 1995;270:20329–36.
- 410 [14] Mollet C, Drancourt M, Raoult D. rpoB sequence analysis as a novel basis for
411 bacterial identification. *Molecular Microbiology* 1997;26:1005–11.
- 412 [15] Adekambi T, Colson P, Drancourt M. rpoB-Based Identification of Nonpigmented
413 and Late-Pigmenting Rapidly Growing Mycobacteria. *Journal of Clinical*
414 *Microbiology* 2003;41:5699–708.
- 415 [16] Scola BL, Bui LTM, Baranton G, Khamis A, Raoult D. Partial rpoB gene sequencing
416 for identification of *Leptospira* species. *FEMS Microbiology Letters* 2006;263:142–
417 7.
- 418 [17] Rudney JD, Larson CJ. Use of restriction fragment polymorphism analysis of rRNA
419 genes to assign species to unknown clinical isolates of oral viridans streptococci.
420 *Journal of Clinical Microbiology* 1994;32:437–43.
- 421 [18] Wisplinghoff H, Reinert RR, Cornely O, Seifert H. Molecular Relationships and
422 Antimicrobial Susceptibilities of Viridans Group Streptococci Isolated from Blood of
423 Neutropenic Cancer Patients. *Journal of Clinical Microbiology* 1999;37:1876–80.

- 424 [19] Paster BJ, Olsen I, Aas JA, Dewhirst FE. The breadth of bacterial diversity in the
425 human periodontal pocket and other oral sites. *Periodontology* 2000 2006;42:80–7.
- 426 [20] Consortium THMP. Structure, function and diversity of the healthy human
427 microbiome. *Nature* 2012;486:207–14.
- 428 [21] Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, et
429 al. The Long-Term Stability of the Human Gut Microbiota. *Science* 2013;341.
- 430 [22] Schloissnig S, Arumugam M, Sunagawa S, Mitreva M, Tap J, Zhu A, et al. Genomic
431 variation landscape of the human gut microbiome. *Nature* 2013;493:45–50.
- 432 [23] Roche Diagnostics GmbH. MagNA Pure 96 DNA and Viral NA Small Volume Kit.
433 Version 08 2012.
- 434 [24] Illumina, Inc. TruSeq DNA Sample Prep Kits. Illumina; 2012.
- 435 [25] <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
436 BioinformaticsBabrahamAcUk n.d.
- 437 [26] Magofç T, Salzberg SL. FLASH: Fast Length Adjustment of Short Reads to Improve
438 Genome Assemblies. *Bioinformatics* 2011.
- 439 [27] http://hannonlab.cshl.edu/fastx_toolkit/index.html. HannonlabCshlEdu n.d.
- 440 [28] Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of
441 protein or nucleotide sequences. *Bioinformatics* 2006;22:1658–9.
- 442 [29] Anders S, Huber W. Differential expression analysis for sequence count data.
443 *Genome Biology* 2010;11:R106.
- 444 [30] McMurdie PJ, Holmes S. Waste Not, Want Not: Why Rarefying Microbiome Data Is
445 Inadmissible. *PLoS Comput Biol* 2014;10:e1003531.
- 446 [31] Drancourt M, Raoult D. rpoB Gene Sequence-Based Identification of *Staphylococcus*
447 Species. *Journal of Clinical Microbiology* 2002;40:1333–8.
- 448 [32] Lazarevic V, Manzano S, Gaia N, Girard M, Whiteson K, Hibbs J, et al. Effects of
449 amoxicillin treatment on the salivary microbiota in children with acute otitis media.
450 *Clinical Microbiology and Infection* 2013;19:E335–42.
- 451
452
453
454
455
456
457
458
459
460
461
462

463 **Figure legends**

464

465 **Fig.1. Relative abundance of the top five phyla, per individual, per target gene.** A and B
466 are different individuals and the target genes are shown in brackets.

467

468 **Fig.2. Number of sequences required for sample separation.** The relative distance
469 corresponds to the distance between two individuals calculated using the Euclidean distance
470 and the Ward method of hierarchical clustering, on the normalised and logged species
471 abundance. Only species with a p-value < 0.1 from a t-test between the samples from each
472 individual or a BF < 1 were used.

473

474 **Fig.3. Comparison of clustering thresholds for the separation of individuals.** The percent
475 identity is that used for clustering the sequences into OTUs with CD-HIT. The relative
476 distance corresponds to the distance between two individuals calculated using the Euclidean
477 distance and the Ward method of hierarchical clustering, on the normalised and logged
478 species abundance. Only species with a p-value < 0.1 from a t-test between the samples from
479 each individual or a BF < 1 were used. A = both rpoB targets and B = 16S rRNA. The dashed
480 line highlights the chosen threshold.

481

482 **Fig.4. Hierarchical clustering of all eight samples for each target.** The relative distance
483 corresponds to the distance between two individuals (A and B) calculated using the Euclidean
484 distance and the Ward method of hierarchical clustering, on the normalised and logged
485 species abundance. Only species occurring in both experiments and with a p-value < 0.1 from
486 a t-test between the samples from each individual or a BF < 1 were used.

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511 **Table 1. Primers designed for each gene target.** Primer name for 16S rRNA and *rpoB2*
 512 corresponds to the *Escherichia coli* positions and for *rpoB1* to the *Streptococcus bovis*
 513 positions.
 514

Gene	Primer name	Primer sequence (5'-3')	Tm (°C)
16S rRNA	792 F	AGGATTAGATACCCTGGTAG	56
	891R	CGTACTCCCCAGGCGG	
<i>rpoB1</i>	130F	GGACCTGGTGGTTTGAC	64
	220R	CGATGTTAGGTCCTTCAGG	
<i>rpoB2</i>	340F	GGACCAGAACAACCCG	60
	434R	GGGTGTCCGTCTCGAAC	

515
 516
 517
 518

Table 2. Species-level OTUs for all samples, per target.

Sample	No. OTUs 16S rRNA	No. OTUs <i>rpoB1</i>	No. OTUs <i>rpoB2</i>
Experiment 1			
A1	810	145	20
A2	793	147	23
B1	839	149	25
B2	828	144	29
Experiment 2			
A3	1273	182	46
A4	1267	185	44
B3	1291	169	44
B4	1283	171	48

519

Figure 1




Figure 2





Figure 3

A



B




Figure 4

