

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Estimation of malnutrition prevalence using administrative data: Not as simple as it seems.

Authors: Khalatbari-Soltani S, Waeber G, Marques-Vidal P

Journal: Clinical nutrition (Edinburgh, Scotland)

Year: 2015 Dec

Volume: 34

Issue: 6

Pages: 1276-7

DOI: 10.1016/j.clnu.2015.09.001

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

1 **Estimation of malnutrition prevalence using administrative data: Not as simple as it seems.**

2 Saman Khalatbari-Soltani, PhD student ^{a,b}, Gerard Waeber, MD ^b and Pedro Marques-Vidal,

3 MD, PhD ^b

4 ^a Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, Biopole 2,

5 Route de la Corniche 10, CH-1010 Lausanne, Switzerland and ^b Department of Internal

6 Medicine, Internal Medicine, Lausanne University Hospital (CHUV), rue du Bugnon 46, 1011

7 Lausanne, Switzerland

8 **Authors' emails:**

9 Saman Khalatbari-Soltani: Saman.KhalatbariSoltani@unil.ch

10 Gerard Waeber Gerard.Waeber@chuv.ch

11 Pedro Marques-Vidal: Pedro-Manuel.Marques-Vidal@chuv.ch

12 **Address for correspondence and reprints**

13 Pedro Marques-Vidal

14 Department of Internal Medicine, Internal Medicine

15 Room BH10-642

16 Lausanne University Hospital (CHUV)

17 Rue du Bugnon 46

18 1011 Lausanne

19 **Switzerland**

20 Phone: +41 21 314 09 34

21 Email: Pedro-Manuel.Marques-Vidal@chuv.ch

22 Conflict of interest: none.

23 Dear Editor,

24 Malnutrition is a common finding among hospitalized patients, leading to increased
25 morbidity and higher length of stay and costs (1). Several studies have assessed the prevalence of
26 malnutrition using hospital discharge data, i.e. using international classification of diseases (ICD)
27 codes. Nowadays, most hospital data is available in electronic format, including prealbumin, a
28 marker of malnutrition (2). Whether malnutrition defined by low prealbumin levels is actively
29 reported in discharge data has seldom been assessed. Thus, we assessed the prevalence of
30 malnutrition according to four definitions: 1) any ICD-10 code E40 to E46; 2) ICD-10 code R63
31 or R64; 3) prealbumin levels <15 mg/dL (measurements performed in 996 patients devoid of
32 liver disease or infection) and 4) any one of the previous three. We used data from 2002 to 2013
33 from the Department of Internal Medicine of the Lausanne University Hospital, Switzerland
34 (32,850 patients aged ≥ 18 years).

35 Prevalence (95% confidence interval – CI) of malnutrition using the entire sample was
36 2.7% (2.5-2.9); 1.7% (1.6-1.9); 2.0% (1.8-2.2) and 6.1% (5.9-6.4) for definitions 1, 2, 3 and 4,
37 respectively. These findings are in agreement with other studies conducted in the United States
38 and Spain, which reported 3.2% and 1.4% malnutrition diagnosis at discharge according to ICD-
39 9 codes (3, 4). No differences in prevalence were found between genders or age groups using
40 definitions 1 and 2, while higher prevalence were found in men using definition 3 and in younger
41 patients using definitions 3 and 4 (Table). Among the 996 patients for whom prealbumin was
42 measured, 656 [65.9% (62.8-68.8)] had prealbumin levels <15 mg/dL, but of these 656 patients
43 with possible malnutrition, only 56 (8.5%) were reported as malnourished in discharge data
44 (definition 1) and only 17 (2.6%) were considered as malnourished according to definition 2.
45 These findings are in accordance with the literature (4, 5), showing that results from objective

46 nutritional assessments are rarely translated into ICD codes at discharge. This underreporting of
47 malnutrition has considerable consequences for health planning, as health statistics and most
48 public health decisions are based solely on hospital discharge data.

49 We conclude that the prevalence and determinants of hospital malnutrition vary
50 significantly according to the definition applied. Results from objectively assessed malnutrition
51 are not actively coded, leading to a considerable underestimation of malnutrition prevalence in
52 hospital discharge data. Professionals filling the discharge letter should be more sensitized
53 towards malnutrition.

54 **Table:** Malnutrition prevalence according to gender and age groups. Results are expressed as
 55 number of hospitalizations and (%).

	Malnutrition Definition			
	ICD-10 code	ICD-10 code	Prealbumin	All
	E40 to E46	R63 or R64	<15 mg/dL	
Gender				
Women	450 (2.8)	298 (1.9)	286 (1.8)	993 (6.2)
Men	432 (2.6)	269 (1.6)	370 (2.2)	1027 (6.2)
* <i>P</i> -value	0.265	0.104	0.004	0.99
Age group				
30-59	226 (2.9)	142 (1.8)	174 (2.2)	509 (6.5)
60-69	166 (3.0)	94 (1.7)	151 (2.7)	393 (7.1)
70-79	201 (2.6)	109 (1.5)	177 (2.3)	473 (6.2)
80-89	221 (2.4)	172 (1.9)	127 (1.4)	503 (5.5)
90+	68 (2.5)	50 (1.8)	27 (1.0)	142 (5.3)
* <i>P</i> -value	0.131	0.219	< 0.001	< 0.001

56 ICD, International Classification of Diseases. **P*-value for between group comparisons using

57 Chi-square test

58 **References:**

- 59 1. Norman K, Pichard C, Lochs H, Pirlich M. Prognostic impact of disease-related
60 malnutrition. *Clin Nutr.* 2008;27(1):5-15.
- 61 2. Robinson MK, Trujillo EB, Mogensen KM, Rounds J, McManus K, Jacobs DO.
62 Improving nutritional screening of hospitalized patients: the role of prealbumin. *JPEN Journal of*
63 *parenteral and enteral nutrition.* 2003;27(6):389-95; quiz 439.
- 64 3. Corkins MR, Guenter P, DiMaria-Ghalili RA, Jensen GL, Malone A, Miller S, et al.
65 Malnutrition Diagnoses in Hospitalized Patients: United States, 2010. *JPEN.* 2014;38(2):186-95.
- 66 4. Marco J, Barba R, Zapatero A, Matia P, Plaza S, Losa JE, et al. Prevalence of the
67 notification of malnutrition in the departments of internal medicine and its prognostic
68 implications. *Clin Nutr.* 2011;30(4):450-4.
- 69 5. Polednak AP. Indicators of nutritional screening in hospital records of newly diagnosed
70 Hispanic and Asian-American adult cancer patients in Connecticut. *Nutrition.* 2008;24(10):1053-
71 6.
72