

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Genital Chlamydia trachomatis: understanding the roles of innate and adaptive immunity in vaccine research.

Authors: Vasilevsky S, Greub G, Nardelli-Haefliger D, Baud D

Journal: Clinical microbiology reviews

Year: 2014 Apr

Volume: 27

Issue: 2

Pages: 346-70

DOI: 10.1128/CMR.00105-13

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Genital Chlamydia trachomatis: understanding the roles of innate and adaptive immunity in vaccine research.

3

4

5 **Sam Vasilevsky^{1,2}, Gilbert Greub², Denise Nardelli-Haefliger³ and David Baud^{1,2#}**

6

7 ¹ Materno-fetal & Obstetrics Research Unit, Department of Obstetrics and Gynecology,
8 University hospital, 1011 Lausanne, Switzerland

9 ² Center for Research on Intracellular Bacteria, Institute of Microbiology, Faculty of Biology
10 and Medicine, University of Lausanne, 1011 Lausanne, Switzerland

11 ³ Dpt. of Urology, Centre Hospitalier Universitaire Vaudois and University of Lausanne, CH-
12 1011 Lausanne, Switzerland

13

14 Running Head: Immunity and *Chlamydia* Vaccine

15 Keywords: *Chlamydia*, vaccines, immunology, macrophages, dendritic cells, T cells

16

17 **#Corresponding author:** David Baud, MD PhD

18 Department of Obstetrics and Gynecology

19 University hospital

20 Centre Hospitalier Universitaire Vaudois (CHUV)

21 1011 Lausanne

22 SWITZERLAND

23 Phone: (00) 41 79 556 13 51

24 Email: david.baud@chuv.ch

25		
26	SUMMARY	3
27	INTRODUCTION	3
28	CHLAMYDIA BIOLOGY	4
29	IMMUNITY TO CHLAMYDIA	5
30	T cells	5
31	Dendritic Cells	7
32	Macrophages	8
33	B cells/Antibodies	9
34	VACCINES	10
35	Intact attenuated organisms	11
36	Subunit antigenic determinants	13
37	Recombinant proteins	14
38	Plasmid DNA	16
39	OTHER CHLAMYDIAL VACCINES AND DELIVERY SYSTEMS	17
40	Bacterial ghosts	17
41	Biodegradable polymers	18
42	Gas vesicles	19
43	ADJUVANTS	19
44	VACCINATION ROUTES	20
45	CONCLUSIONS	21
46	REFERENCES	23
47		
48		
49		
50		
51		
52		
53		

54 SUMMARY

55 Despite significant advances in the understanding of the host response to chlamydial
56 infection and over 30 years of vaccine research, *Chlamydia trachomatis* remains the leading
57 cause of bacterial sexually transmitted disease worldwide. This gram-negative obligate
58 intracellular bacterium, that often remains asymptomatic, may cause pelvic inflammatory
59 disease (PID), ectopic pregnancies, scarring of the fallopian tubes, miscarriage and infertility
60 when left untreated. In the genital tract, *Chlamydia trachomatis* primarily infects epithelium
61 cells and requires Th1 immunity for optimal clearance. This review first focuses on the
62 immune cells important in a chlamydial infection. Secondly, we will summarize the research
63 and challenges associated with developing a chlamydial vaccine that elicits a protective Th1-
64 mediated immune response without inducing adverse immunopathologies.

65

66 INTRODUCTION

67 *Chlamydia trachomatis* is the leading cause of bacterial sexually transmitted diseases
68 in humans. According to the WHO in 2008, there was 105 million new cases of STDs each
69 year due to *C. trachomatis* worldwide and the infection rate has been steadily increasing (9,
70 103). When symptomatic, *C. trachomatis* can lead to mucopurulent endocervical discharge,
71 hypertrophic cervix, and post coital bleeding. In 20-40% of untreated women *C. trachomatis*
72 may reach the fallopian tubes via the endometrial epithelium and cause pelvic inflammatory
73 disease (PID). However, *C. trachomatis* genital tract infections are often asymptomatic (75-
74 90%) and therefore remain undiagnosed and untreated. This can lead to tubal factor infertility
75 or ectopic pregnancies (68, 69), which is a life threatening condition. *C. trachomatis* can be
76 easily treated with antibiotics such as erythromycin, azithromycin or doxycycline. However,

77 several studies have documented that within a year after treatment for a *C. trachomatis*
78 infection, 13-26% of individuals showed evidence of persistent or recurrent infections (38,
79 51). Therefore, due to the high rate of asymptomatic infections, recurrent infections and the
80 severity of pathologies induced by *Chlamydia*, the development of a vaccine is paramount.
81 This review focuses on *C. trachomatis* and *C. muridarum* (a model organism that naturally
82 infects rodents and largely used for animal experiments) immunity and the challenges
83 associated with generating a vaccine against this bacterium. Table 1 summarizes recent
84 developments in chlamydial research including *Chlamydia* strain or antigen used, cell type
85 affected and immune response elicited.

86

87 **CHLAMYDIA BIOLOGY**

88 The genus *Chlamydia* includes species that infect humans (*C. trachomatis*, *C.*
89 *pneumoniae*), and animals (*C. muridarum*, *C. suis*, *C. abortus*) (21). Presently, there have
90 been 18 identified serovars of *C. trachomatis* based on the reactivity of patient sera to the
91 major outer membrane protein (MOMP) (135). Some serovars are associated with ocular
92 tissue infections (A-C) while others primarily infect genital tissues (D-K) (7). *C. trachomatis*
93 is a gram-negative obligate intracellular bacterium that in the genital tissues normally infects
94 the epithelium layer of the cervix of women and the urethra of men (12).

95 *Chlamydia* exists in two developmental forms, the infectious extracellular non-
96 replicating elementary body (EB) and the non-infectious intracellular replicating reticulate
97 body (RB). The EB displays no metabolic activity, is resistant to both chemical and physical
98 factors, and is adapted for prolonged extracellular survival. Infection begins when the small
99 (~0.2-0.3 μ m) EB attaches to the host cell and is internalized inside an entry vacuole which
100 avoids fusion with the lysosome. After 8-10 hours the vesicle bound EB (termed an inclusion)

101 replicates by binary fission into the larger (~0.8 μm) RB (138). Following several rounds of
102 division, the RB's reorganize and revert back to the EB (131). Inside host cells, *C.*
103 *trachomatis* circumvents endogenous stress mechanisms, prevents lysosomal fusion and
104 escapes intracellular destruction by replicating in an inclusion outside of the endocytic
105 pathway (138). *C. trachomatis*-infected cells have increased inducible oxide synthase (iNOS)
106 and increased pro-inflammatory molecules such as activins, which may be involved in
107 scarring (1, 115).

108

109 **IMMUNITY TO *CHLAMYDIA***

110 **T cells**

111 A critical role for T cells in immunity to *Chlamydia* was demonstrated almost 30 years
112 ago when Rank *et al.* observed that athymic nude mice established chronic infection with *C.*
113 *muridarum* after intravaginal inoculation whereas wild-type (wt) mice resolved the infection
114 in 20 days (114). In human and mouse models, both CD4+ and CD8+ T cells can be detected
115 at the site of *C. trachomatis* infection (65, 71, 101, 129). T cells are unable to recognize
116 pathogens or antigens without the help of antigen presenting cells (APC) such as dendritic
117 cells (DC), macrophages, or B cells. APC are able to phagocytose chlamydial EBs in the
118 extracellular space or engulf infected cells harboring RBs. After phagocytosis, APC degrade
119 chlamydial components and present the peptides via MHC class II-antigen complex to CD4 +
120 T cells or MHC class I-antigen complex to CD8 + T cells. In fact, numerous *C. trachomatis*
121 antigens have been identified which can be recognized by human CD4+ and CD8+ T cells
122 including the cysteine-rich outer membrane protein 2 (Omp2) (40), polymorphic outer
123 membrane protein D (POMP-D) (41), MOMP (50, 72, 104), heat shock protein 60 (hsp 60)

124 (25, 50), chlamydial protease activating factor (CPAF) (75), PmpG, PmpF, and RpIF (66,
125 101). High-throughput proteomic screening has identified even more potential
126 immunodominant *C. trachomatis* antigens including 36 that have been shown to react with
127 sera from three strains of mice immunized with live *Chlamydia* and two protein antigens that
128 were able to induce a polyfunctional Th1 CD4⁺ T cell response and high Th1 antibody titers
129 (112, 126). Although *Chlamydia* is able to induce a Th2 response characterized by IL-4 and
130 Th2-associated antibodies such as IgG1, a Th1 response predominates characterized by the
131 production of IL-12 by APC (17) and the subsequent activation of IFN- γ producing T cells
132 and plasma B cells that secrete Th1-associated antibodies such as IgG2a and IgG3 (97, 100).
133 However, a recent study demonstrated that CD4⁺ T cells from women with genital tract *C.*
134 *trachomatis* infection that were restimulated *ex vivo* with inactivated (γ -irradiated) EB secrete
135 significantly more IL-4 than TNF- α and IFN- γ . This study suggests that the type of immune
136 response (Th1 vs Th2) to *C. trachomatis* may be tissue specific (132).

137 While there is ample evidence that CD4⁺ T cells play an integral part in the resolution
138 of *C. trachomatis* and *C. muridarum* infections (34, 36, 39, 60), the role for CD8⁺ T cell has
139 been controversial even though CD8⁺ T cells are induced following infection and *Chlamydia*-
140 specific human and mouse CD8⁺ T cells are cytotoxic for *Chlamydia*-infected target cells
141 (137). A recent study by Murthy *et al.* demonstrated that wt and CD8⁺ T cell-deficient mice
142 displayed similar clearance of *C. muridarum* following vaginal chlamydial challenge (92).
143 These data support previous studies which demonstrated that CD8⁺ T cells are not critical for
144 *C. trachomatis* clearance (87, 88, 121). Furthermore, the CD8⁺ T cell-deficient mice
145 demonstrated reduced oviduct pathology (hydrosalpinx) compared to wt, suggesting a role of
146 CD8⁺ T cells in chlamydial pathogenesis (92). An interesting study demonstrated that the
147 majority of CD8⁺ T cells in the cervix before and after a *C. trachomatis* infection do not
148 express perforin (53). Perforin is a cytolytic protein found in the granules of CD8⁺ T cells

149 which forms a pore by inserting itself into the cells plasma membrane resulting in lysis of the
150 target cell. Therefore, the lack of perforin in endocervix CD8+ T cells may explain why
151 CD8+ T cells do not play a critical role in the elimination of genital chlamydial infection.
152 Although CD8+ T cells appear not to be critical in resolving a chlamydial infection and may
153 even contribute to chlamydial sequelae, they nonetheless may play a contributory albeit
154 secondary role by regulating other cells and by their own production of IFN- γ (137).

155 **Dendritic Cells**

156 Dendritic cells (DC) are known to be the quintessential antigen presenting cells.
157 Immature DC are highly phagocytic and after internalization of pathogens they degrade the
158 components and present the peptides to T cells via MHC receptors activating the T cells and
159 initiating a cell- mediated and / or humoral immune response. The capacity of DCs to present
160 chlamydial antigens to T cells, secrete Th1 cytokines such as IL-12 and TNF- α and the
161 importance of MHC class molecules in chlamydial infection has been demonstrated both *in*
162 *vitro* and *in vivo* (64, 84, 87, 99, 122). An early study conducted by Lu and Zhong showed
163 that bone marrow derived dendritic cells (BMDC) pulsed with heat-killed *C. trachomatis* and
164 adoptively transferred into a naive mouse was protective against a subsequent challenge with
165 live *C. trachomatis* in a mouse lung infection model (82). This protection was mediated by a
166 Th1 response further demonstrating a correlation between Th1 skewed immunity and
167 protection against chlamydial infection. In contrast, DCs that were pulsed with recombinant
168 MOMP and adoptively transferred into mice elicited primarily the Th2-associated antibody
169 IgG1 (119). Furthermore, IL-10 (Th2-associated cytokine) KO DC pulsed with UV-
170 inactivated *C. trachomatis* and adoptively transferred activated a high frequency of Th1 cells
171 (47). These data have direct relevance to vaccine development because it indicates that the
172 type of cytokines produced and antigens processed by the DC and presented to CD4+ T cells

173 is essential in the Th1/Th2 balance of the immune response to *Chlamydia*. There is also
174 evidence that live *Chlamydia* is required for an optimal and protective immune response. Rey-
175 Ladino and colleagues demonstrated that the level of protection induced by DC pulsed with
176 UV-inactivated *C. trachomatis* EB and adoptively transferred into mice was significantly less
177 than in mice that were challenged with live EB pulsed DC (116). A more recent study
178 discovered that murine DCs pulsed with live *C. muridarum* EBs presented 45 MHC class II
179 peptides mapping 13 proteins whereas dead EBs presented only six MHC class II peptides
180 mapping to three proteins (143). However, *C. trachomatis* has developed strategies to limit
181 the presentation of these antigens to T cells by downregulating MHC expression on APC (54).
182 *C. trachomatis* has been shown to inhibit MHC molecules within infected cells through the
183 degradation of the MHC class I transcription factor RFX-5 and the MHC class II transcription
184 factor USF-1 by secreting the chlamydial protease CPAF into the cytosol (29, 111, 145, 146).
185 DC are important to vaccine research because they are the critical links between innate and
186 adaptive immunity. Two recent studies using DC transfected with a recombinant adenovirus
187 carrying *C. trachomatis* MOMP antigen (81) and DC pulsed *ex vivo* with recombinant *C.*
188 *trachomatis* protease-like factor (rCPAF) (75) illustrate the ability of DC to induce protective
189 immunity against genital *C. trachomatis* and *C. muridarum* challenge respectively.

190 **Macrophages**

191 Studies using both *C. trachomatis* and *C. muridarum* have shown that macrophages
192 are recruited to sites of infection (88) and are capable of phagocytosing *Chlamydia* (8).
193 Macrophages are also a source of both proinflammatory cytokines such as IL-8, IL-6 and
194 TNF- α (6, 141). However, unlike epithelial cells, macrophages are not a hospitable niche for
195 chlamydial intracellular replication illustrated by the fact that compared to epithelial cells
196 only a small fraction of chlamydial RBs are detected in macrophages (124). *C. trachomatis*

197 destruction inside the macrophage has been associated with host cell autophagy, a process by
198 which cells degrade cytoplasmic proteins and organelles (2, 124, 140), and studies have
199 demonstrated that macrophage autophagy can enhance antigen presentation to T cells (22).
200 Furthermore, INF- γ has been shown to enhance both autophagy and upregulation of MHC
201 class molecules in macrophages (2, 15). This is relevant because, in addition to activating
202 primed T cells, there is evidence suggesting that macrophages are able to initiate a humoral
203 response in naive mice (130). Therefore, enhanced upregulation of MHC molecules
204 containing chlamydial antigens may induce T cells to initiate both a cell-mediated and
205 antibody immune response against *Chlamydia*. However, Jendro *et al.* demonstrated that
206 human macrophages infected with *C. trachomatis* can induce T cell apoptosis (61, 62). In
207 addition to efficiently eliminating *Chlamydia* and presenting the peptides to T cells,
208 macrophages may also have an effect on chlamydial infection by inducing T cell death and
209 perpetuating a persistent infection.

210 **B cells/Antibodies**

211 Previous studies demonstrated that anti-*Chlamydia* antibodies correlated with
212 protective immunity against *C. trachomatis* in humans (4, 59) and numerous *C. trachomatis*
213 proteins have been shown to induce antigen specific antibodies (36). However, even though
214 anti-*Chlamydia* antibodies are able to neutralize infection *in vitro* (5, 13) growing evidence
215 show that B cells may not play a critical role in controlling a primary chlamydial infection but
216 are important for a secondary memory response (89, 90). Several mechanisms have been
217 proposed on how B cells contribute to immunity during re-infection. These mechanisms
218 include antibody-mediated neutralization and opsonization (5), antibody-dependent cellular
219 cytotoxicity (ADCC) (86) (a mechanism of cell-mediated immune defense whereby cells that
220 have antibodies attached to their surface are targeted for lysis), and the formation of antigen-

221 antibody complexes that bind Fc receptors on the APC which then enhances phagocytosis and
222 antigen presentation to the CD4+ T cell (57).

223 Heat shock proteins (hsp), which are found in both eukaryotic and prokaryotic
224 organisms, are stress-proteins that are involved in the correct folding of intracellular proteins.
225 *C. trachomatis* is known to secrete hsp's during an infection and antigenic epitopes from the
226 bacterial hsp's have proven to be strong inducers of cellular and humoral immunity.
227 Chlamydial hps60 exhibits over 70% sequence homology and 100% amino acid homology of
228 four defined epitopes with human hsp60 (3) and several studies have suggested that
229 autoimmunity to human hsp60 is a result of cross reactivity after a chlamydial infection (27,
230 136). However, a recent study demonstrated an association with tubal factor infertility (TFI)
231 and antibodies to MOMP and hsp60 from *C. trachomatis* but no connection between TFI and
232 antibodies to human hsp60 (49) pointing to an infectious rather than an autoimmune response
233 as the cause of TFI.

234 In conclusion cell-mediated immunity that activates macrophages, neutrophils and
235 mediators such as IL-12, IFN- γ and TNF- α is required for initial clearance. However, for
236 protective immunity both cell-mediated and humoral immunity are needed including antigen-
237 specific T cells and antibodies that enhance uptake, processing and presentation of chlamydial
238 antigens by DC for a rapid Th1-mediated chlamydial clearance.

239

240 **VACCINES**

241 Due to increasing rates of mainly asymptomatic *C. trachomatis* infections worldwide
242 and the adverse long term consequences resulting from these infections (ectopic pregnancy,
243 infertility, preterm birth) developing an anti-chlamydial vaccine is paramount. However, a
244 human vaccine that elicits both T cell and B cell immunity has been elusive. Although the two

245 murine models using *C. trachomatis* and *C. muridarum* are the most common models used for
246 chlamydial vaccine research non-human primates, pigs and guinea pigs have also been
247 utilized (24). Our poor understanding of the immune response in the female genital tract,
248 which is highly regulated by sex hormones during the menstrual cycle (52), the lack of
249 adjuvants that not only optimize the immune response to *Chlamydia* antigens but can target
250 the vaccine-specific-immune responses to the site of infection and limited understanding of
251 what type of chlamydial antigens induce a protective immune response hinder the
252 development of a human *C. trachomatis* vaccine. *C. trachomatis* vaccine has to induce both
253 mucosal and systemic immune responses, but autoimmune cross reactions with human
254 antigens and unregulated inflammation that causes pathology has to be avoided. Table 2
255 summarizes recent chlamydial antigens, delivery systems, routes of vaccination and infection
256 and the subsequent immune responses elicited.

257 **Intact attenuated organisms**

258 Successful vaccines against ovine enzootic abortions have been available for many
259 years (32). These vaccines consisted of live attenuated or inactivated *C. abortus* strains and
260 provided proof of principle that a successful vaccine against *Chlamydia* was possible in
261 mammals. However, these vaccines did not prevent infectivity and lacked the rigorous
262 immunization schedules, efficacy, safety and toxicity standards required for human vaccine
263 (56, 85). Nonetheless, because of the success of these vaccines, live attenuated *C. trachomatis*
264 bacteria were used as the first human *Chlamydia* vaccines (43). Attenuation was induced by
265 either mutagenesis or by growing the organisms in culture. In the latter approach, after several
266 passages, one or more mutations arise, which may result in a nonvirulent attenuated strain.
267 Vaccines with live organisms are generally considered optimal because they contain virtually
268 all of their antigenic determinants in the correct three dimensional conformation. Moreover,

269 they replicate similarly to the target pathogen thus promoting the processing and presentation
270 of antigens similar to natural infection and eliciting humoral and cell-mediated immunity.
271 However, using live attenuated organisms for vaccines has drawbacks because large scale
272 production of pure *Chlamydia* is extremely complex and these vaccines need cold storage.
273 Even more importantly, they can also revert to virulent wild-type strains resulting in disease
274 or persistent infection (26).

275 Initial vaccine human trials using live attenuated *C. trachomatis* led to partial short-
276 lived protection, however some individuals who were re-exposed to *Chlamydia* developed a
277 more severe pathological delayed-type hypersensitivity (DTH) response than those that did
278 not receive the vaccine (43). Because of the safety issues of live vaccines, research switched
279 to organisms that were heat or chemically-inactivated. The major disadvantage of these types
280 of vaccines is the absence of replication and poor induction of cell-mediated immunity, which
281 is critical for the clearance of *Chlamydia*, necessitating the need for re-vaccination and
282 adjuvants. Heat or chemical bacterial inactivation may also release unwanted and detrimental
283 components which can have deleterious effects or degrade protein antigenic determinants
284 thereby reducing the degree of protection. Recently, plasmid-deficient *Chlamydia* strains
285 have been used in vaccine research with conflicting results. O'Connell *et al.* demonstrated
286 that a plasmid-deficient strain of *C. muridarum* (Nigg) that is defective in its ability to
287 accumulate glycogen did not cause inflammatory pathology in mice. Furthermore, the
288 plasmid-deficient bacterium protected mice against a secondary infection with plasmid-
289 competent virulent *C. muridarum* (97). However, a different group demonstrated that mice
290 vaccinated with an attenuated plasmidless *C. trachomatis* (L2R) were not protected from
291 colonization and inflammatory pathology after a secondary challenge with wild-type *C.*
292 *trachomatis* (serovar D) although there was a reduction in infectious burden at early time
293 points (100).

294 **Subunit antigenic determinants**

295 Another vaccine strategy utilized is the administration of purified antigenic
296 determinants known to elicit an immune response. Subunit vaccines are safer than attenuated
297 or heat/chemically inactivated organisms because they cannot revert to a virulent form and
298 undesirable antigens that might induce immunopathology can be avoided. One of the most
299 studied vaccine candidate for *C. trachomatis* is the structurally and immunologically
300 dominant protein in the chlamydial outer membrane MOMP. This membrane protein contains
301 several conserved CD4+, CD8+, and B cell epitopes (96) . An early study conducted by Pal
302 and colleagues demonstrated that *C. muridarum* COMP (chlamydial outer membrane
303 complex), a chlamydial outer membrane with a cysteine cross-linked protein shell,
304 significantly protected mice against genital challenge whereas MOMP did not (108). Several
305 years later the same group administered purified and refolded preparation of *C. muridarum*
306 MOMP along with Freund's adjuvant. The refolded MOMP-Freunds adjuvant conferred a
307 significant level of protection in the vaccinated mice against a genital infection demonstrating
308 the importance of adjuvants and a correct MOMP configuration in eliciting a protective
309 immune response (109). Tiffrea *et al.* discovered that a polymer that keeps membrane proteins
310 soluble (amphipol) in aqueous solution was able to stabilize MOMP and enhance its
311 protective ability as a vaccine (128). Another group immunized mice with a *C. trachomatis*
312 MOMP-ISCOM vaccine. ISCOM (immune stimulating complex) which are mainly composed
313 of cholesterol, phospholipids and saponin, are known to induce both a cell-mediated and
314 humoral response when used as vaccine adjuvants. Inoculation with MOMP-ISCOM was able
315 elicit a Th1 antigen-specific response and vaginal infection was cleared within one week (58).
316 A *C. muridarum* MOMP native preparation combined with an adjuvant consisting of
317 nontoxic subunit B cholera toxin conjugated to CpG (CTB-CpG) elicited a significant cell

318 mediated and antigen-specific antibody response against a pulmonary challenge with *C.*
319 *muridarum* (18). A non-human primate model was used to demonstrate the efficacy of a
320 vaccine formulated with native MOMP. Rhesus macaques that were immunized
321 intramuscularly and subcutaneously along with the adjuvants CpG-2395 and Montanide ISA
322 720 produced high levels of Th1 cytokines (INF- γ , TNF- α) and *C. trachomatis*-specific IgG
323 and IgA (19). Drawbacks of subunits vaccines include the fact that extracting, refolding and
324 purifying protein complexes such as MOMP is very expensive and purifications are not
325 standardized so difference in extraction methods may influence the conformation of the
326 protein epitopes and vaccine efficacy.

327 **Recombinant proteins**

328 The advent of recombinant DNA technology has made it possible to produce large
329 quantities of bacterial proteins. Thus, different attempts were made to use rMOMP in *C.*
330 *trachomatis* vaccine. Unfortunately, producing rMOMP with its native conformational
331 epitopes intact on a large scale is challenging and in some expression systems full-length
332 rMOMP is toxic (78, 147). In 2009, a comparison of vaccines using native or recombinant
333 MOMP demonstrated that the degree of protection obtained with recombinant MOMP was
334 not as robust as that achieved with native MOMP preparation (123). However, other studies
335 using rMOMP with and without adjuvants demonstrated protection against *Chlamydia* (98,
336 127). In 2011, Kalbina and colleagues designed a chimeric gene construct containing two
337 antigenic regions of MOMP and introduced the construct into a bacteria (*Escherichia coli*)
338 and two plants (*Arabidopsis thaliana*, *Daucus carota*). The construct was successfully
339 expressed in *E. coli*, and stable integration of the transgene was demonstrated in *A. thaliana*
340 and *D. carota* over several generations. The rMOMP purified from *E. coli* was used to
341 produce antibodies in rabbits and these antibodies recognized the proteins in both *E. coli*, *A.*

342 *thaliana*, *D. carota* as well as in inactivated *C. trachomatis* elementary bodies. The stability
343 of the construct in the offspring plants suggests that this system may be useful for large scale
344 production of rMOMP and the authors plan to use the transgenic plants as edible vaccine
345 vectors for laboratory animal experiments (67).

346 Other recombinant proteins besides MOMP have also been shown to be potential
347 vaccine candidates. In 2007, Murphy *et al.* investigated the potential of recombinant CPAF to
348 elicit an immune response that would resolve chlamydial infection. Mice immunized
349 intranasally with rCPAF and IL-12 (Th1 cytokine) demonstrated increased IFN- γ , and
350 minimal IL-4 (Th2 cytokine) production, elevated IgG2a (Th1) and IgA (mucosal) antibody
351 levels, displayed a markedly reduced bacterial burden upon *C. muridarum* genital inoculation
352 and were protected against pathological consequences of *Chlamydia* infection compared with
353 mock immunized mice (91). The same group demonstrated that rCPAF intranasal vaccination
354 may prevent infertility from repeated genital *C. muridarum* infections in mice (93). Mice
355 immunized with a recombinant chlamydial glycogen phosphorylase (GlgP) and intravaginally
356 challenged with live *C. muridarum* elicited a Th1-dominant T cell response that included anti-
357 chlamydial antibodies and reduced hydrosalpinx severity. Additionally, the GlgP-immunized
358 mice exhibited a significant reduction of vaginal shedding on day 14 post-infection (76).
359 Olsen *et al.* utilized two recombinant proteins in a subunit chlamydial vaccine. The fusion
360 protein CTH1 consisted of CT443 (omcB) which is known to elicit both a humoral and cell-
361 mediated response and CT521 (rl 16) a known target for cells during natural infection in
362 humans. Immunization with CTH1 along with the strong Th1 inducing adjuvant CAF01
363 elicited TNF- α , IL-2 and INF- γ production from T cells and high titers of both Th1 (IgG2a)
364 and Th2 (IgG1) CTH1-specific antibodies. The vaccine significantly reduced bacterial
365 shedding after a vaginal challenge with live *C. trachomatis* and *C. muridarum* and protection
366 was demonstrated to be solely CD4⁺ T cell-mediated in the *C. muridarum* model (102). Lu

367 and colleagues screened 5 recombinant chlamydial antigens (ABC transporter [ArtJ], outer
368 membrane complex protein B [OmcB], macrophage infectivity potentiator [Mip], inclusion
369 membrane protein [Inc (crpA:TC0726)], and an hypothetical protein), that were previously
370 found to react with sera from intravaginally *C. muridarum* infected mice for their ability to
371 induce protection against chlamydial infection. Only Mip induced pronounced protection
372 which was characterized by Th1-dominant T cell response and anti-Mip antibodies (80).

373 **Plasmid DNA**

374 DNA vaccines work by injecting a plasmid that encodes a specific gene of interest
375 within the host. The product of the gene can then be expressed inducing an immune response.
376 DNA vaccines have several advantages compared with other vaccination strategies. DNA can
377 be easily and inexpensively purified and plasmid vectors can be rapidly constructed and easily
378 tested (79). Additionally, DNA vaccines can encode for multiple epitopes that are in the
379 native three dimensional configuration and avoid the problem associated with attenuated
380 organisms which are able to revert back to virulent forms. However, as with other vaccine
381 strategies, DNA vaccines have some disadvantages. In autoimmune diseases such as lupus
382 anti-DNA antibodies are produced and there is the possibility that bacterial DNA injection
383 could elicit a humoral response that cross-reacts with the host DNA. Also, because DNA
384 encodes for proteins, DNA vaccines cannot be utilized for non-protein based antigens such as
385 polysaccharides or lipids (73) and although likely rare, there is the risk that DNA could
386 integrate into the host chromosome (35). In 1999 Pal and colleagues immunized mice with a
387 DNA vaccine that encoded for the MOMP gene of *C. trachomatis*. When the mice were
388 vaginally challenged with *C. trachomatis* the immune response was modest and the mice were
389 not protected against infection (106). The following year Dong-Ji *et al.* demonstrated that
390 immunization with DNA-MOMP and boosting with MOMP-ISCOM conferred higher

391 protection against *C. trachomatis* when compared with mice that were only immunized with
392 MOMP-ISCOM (28). More recently two studies using a pig model assessed the efficacy of
393 DNA chlamydial vaccines. Schautteet *et al.* combined aerosol-vaginal delivery of DNA
394 vaccine encoding for MOMP co-administered with DNA encoding for three different
395 adjuvants (GM-CSF, *E. coli* enterotoxin subunit A and B). Vaccination induced significant
396 protection against genital *C. trachomatis* challenge although the infection was not completely
397 resolved (117). Ou and colleagues demonstrated that an OmpA-based DNA vaccine against
398 *Chlamydia abortus* in piglets elicited higher antigen-specific IgG antibodies and T cell
399 proliferative response compared with controls (105). Mammalian cells transfected with a
400 plasmid encoding for MOMP epitopes inserted in a human papillomavirus (HPV) major
401 capsid protein L1 was used in a murine model of *C. trachomatis* genital infection.
402 Intramuscular administration elicited a Th1 response characterized by low IL-4 production
403 and antibodies against MOMP (139). All of these recent studies demonstrate the feasibility of
404 DNA-based vaccine and this approach thus deserves further study.

405

406 **OTHER CHLAMYDIAL VACCINES AND DELIVERY SYSTEMS**

407 **Bacterial ghosts**

408 Bacterial ghosts (BGs) are cell envelopes derived from Gram-negative bacteria. BGs
409 are devoid of all cytoplasmic content but have a preserved cellular morphology including all
410 cell surface structures. BGs are non living but retain all of the antigenic components of their
411 living counterparts and the inside of the BG envelope can be loaded with peptides, drugs, or
412 DNA (74). In 2007, a vaccine system in which a DNA plasmid that encoded for *C.*
413 *trachomatis* MOMP and the porin protein (PorB) was inserted into a recombinant *Vibrio*

414 *cholerae* ghost (rVCG) was used. Animals that were immunized intramuscularly with the
415 DNA-bacterial ghost vaccine completely resolved a *C. trachomatis* genital infection after two
416 weeks post-infection. The inflammatory response was Th1, characterized by high levels of
417 IgA and IgG2a (55). More recently, Eko and colleagues used the rVCG that contained PorB
418 and chlamydial polymorphic membrane protein-D (PmpD) proteins to evaluate its ability to
419 induce chlamydial immunity. Intramuscular immunization elicited high levels of Th1-
420 associated antibody IgG2a, mucosal-associated antibody IgA, IFN- γ (Th1) and low levels of
421 IL-5 (Th2) in response to an intravaginal *C. muridarum* infection (31).

422 **Biodegradable polymers**

423 PLGA (poly D, L-lactide-co-glycolide) is an FDA approved polysaccharide that can
424 encapsulate peptide, proteins or DNA. PLGA's are efficiently phagocytosed by DC and
425 macrophages (83, 134) and PLGA antigens are able to be presented on both MHC class I and
426 II molecules thus activating CD4+ and CD8+ T cells (45, 133). Chitosan is a linear
427 polysaccharide derived from the deacetylation of chitin. The glucosamine units of chitosan
428 have a density of amine groups which permits strong electrostatic interactions with proteins
429 and genes. Additionally chitosan is mucoadhesive and has enhanced penetration capacity
430 across mucosal barriers.(10). Both of these nanoparticles are biodegradable, relatively non-
431 toxic and have been used as delivery systems for chlamydial vaccines. Two recent studies
432 using recombinant MOMP encapsulated in PLGA demonstrated enhanced capacity of the
433 peptide to induce Th1 cytokine, cellular and antibody immune response (33, 125). Cambridge
434 *et al.* demonstrated that MOMP was expressed in the muscle tissues and spleens of mice that
435 were intramuscularly injected with chitosan nanoparticles containing recombinant MOMP
436 DNA (14).

437 **Gas vesicles**

438 Gas vesicles are gas containing structures found in some bacteria and Archaea. These protein
439 structures are hollow, rigid, lipid-free, allow diffusion of gases across its membrane, and are
440 able to express peptides from various genes. Studies have shown that in the absence of
441 adjuvants, *Halobacteria* gas vesicles that displayed viral peptides elicited robust long-lived
442 immune response characterized by immunological memory in mice (120). *Halobacteria*-
443 derived gas vesicles that were loaded with gene fragments coding for MOMP, OmcB (outer
444 membrane complex B), and PompB (polymorphic outer membrane B) and expressed on the
445 surface were able to elicit a Th1 cytokine profile in human foreskin fibroblasts *in vitro*. In
446 addition the presence of the recombinant proteins were confirmed by anti-*Chlamydia*
447 antibodies and from *Chlamydia*-positive patient serum suggesting this could be an effective
448 antigen delivery system for a *Chlamydia* vaccine (20).

449

450 **ADJUVANTS**

451 Adjuvants enhance immunity and one of the main challenges in developing an
452 effective chlamydial vaccine is identifying antigen/adjuvant combinations that elicit a
453 protective immune response *in vivo*. Various adjuvant such as the ones mentioned in this
454 review (e.g. Freund's adjuvant, ISCOM's, CTB-CpG, CpG, bacterial ghosts) have been used
455 in chlamydial vaccine research with varying results. Recent research has added other new
456 antigen/adjuvant candidates with encouraging results. A study by Yu and colleagues
457 evaluated the chlamydial protein PmpG and five adjuvants, including three cationic liposome
458 formulations, Montanide ISA720-CpG-ODN1826 and alum. The results demonstrated that the
459 cationic liposomal adjuvants DDA-MPL and DDa-TDB elicited the best protective immune

460 responses against *C. muridarum*. Additionally, using DDA-MPL as an adjuvant along with 7
461 different T cell antigens (PmpG, PmpE, Aasf, Rp1F, TC0420, TC0825) conferred equal or
462 better protection than the vaccine antigen MOMP alone (142). This highlights the various
463 opportunities to further improve vaccine candidates by identifying the optimal
464 epitope/adjuvant combination

465

466 **VACCINATION ROUTES**

467 Vaccine efficacy is not only defined by the type of antigen and adjuvant used but also
468 by the administration route, since lymphocytes primed by antigens in vivo are endowed with
469 specialized homing programs guiding their migration to specific mucosal sites (95). Once
470 naive T cells are primed in a lymph node, a global switch of their homing program occurs
471 which enables them, while trafficking through the blood circulation, to detect chemokines and
472 adhesion molecules which direct them to their tissue destination. Lymphocytes, activated by
473 antigen presentation occurring in lymph nodes draining a mucosal site, acquire specialized
474 homing programs leading them to preferentially migrate to the same or other specific mucosal
475 sites. Of note, T cell homing to the genital mucosa involves either $\alpha 1\beta 1$, $\alpha 4\beta 1$ (110) or
476 $\alpha 4\beta 7/E$ selectin (70) in *Chlamydia* infected mice. Both systemic and mucosal immunization
477 routes have been shown to be able to induce both humoral and cell-mediated immune
478 responses in the genital tract with intranasal immunization being often more effective (11,
479 94). Overall, mucosal immunization routes were more effective at preventing genital
480 challenges with a variety of pathogens (37, 44, 63, 77, 144).

481 Numerous immunization routes have been used for chlamydial vaccinations including
482 oral (48), intranasal (i.n.) (46), intravaginal (i.vag) (118), subcutaneous (107), intramuscular
483 (30), perivaginal (107), perisacral (107), sublingual v(113) and colonic (16). A study using

484 purified MOMP with a *Borrelia* surface protein as an adjuvant demonstrated that in two
485 different mouse strains "intramuscular + subcutaneous" and "perivaginal + perisacral"
486 immunization elicited high systemic (IgG) and mucosal (IgA) serum antibodies. In contrast,
487 the mice that received the MOMP-adjuvant i.n. had low IgG and IgA serum antibodies (107).
488 However, a recent study showed that i.n. immunization with rMOMP resulted in MOMP-
489 specific IgA and IgG antibodies in genital tract secretions demonstrating i.n. administration
490 may target immunity to the reproductive tract (23). Several studies comparing the protective
491 ability of various vaccination routes demonstrated that a combined mucosal and systemic
492 inoculation may be optimal. Using rMOMP with the adjuvants CpG and Montanide for
493 systemic route (intramuscular and subcutaneous) and rMOMP with cholera toxin for the
494 mucosal routes (sublingual and colonic), the authors demonstrated that following i.n. *C.*
495 *trachomatis* challenge the sublingual + intramuscular + subcutaneous group showed the best
496 protection (113). Another group demonstrated that mice immunized by combined mucosal
497 and systemic routes with *C. muridarum* recombinant MOMP plus the adjuvants CpG and
498 Montanide not only elicited the strongest chlamydia-specific humoral and cell-mediated
499 response after vaginal challenge with *C. muridarum* but also protected against infertility (16).
500

501 CONCLUSIONS

502 Chlamydial infection is a public health concern worldwide and a vaccine that
503 stimulates multiple arms of the adaptive immune system and avoids immunopathological
504 consequences would be the best solution for the control of this sexually transmitted disease.
505 Unfortunately, a partial or fully protective vaccine has yet to be developed highlighting the
506 complex nature of the immunobiology mounted against this intracellular parasitic bacterium.
507 The immune response to chlamydial infection is dynamic and involves cells and mediators

508 from both arms of the host's immune system. Clearance of a chlamydial infection requires a
509 coordinated immune response between innate immune cells such as macrophages, DC and
510 cells important in both cell-mediated and humoral adaptive responses such as CD4+T cells,
511 CD8+ T cells and B cells. Activation and clonal expansion of T cells occurs through cognate
512 interactions with DC that present chlamydial antigens on their MHC molecules and B cells
513 produce anti-chlamydial antibodies through interaction with these clonal T cells. However,
514 persistent infection seems to induce chronic inflammation and tissue damage. A shift from
515 Th1 to Th2 also appears to induce scarring and immune pathology. It is therefore essential to
516 understand these immunological dynamics in order to develop a vaccine that is both effective,
517 long-lasting and does not have the deleterious effects associated with unregulated
518 inflammation. Further research is needed to identify novel adjuvants that enhance the immune
519 response and antigens that induce a protective T cell response and anti-chlamydial antibodies.

520 A mathematical model developed by Gray and colleagues demonstrated that a fully
521 protective vaccine, administered to adolescents before they are sexually active, would be able
522 to eradicate *Chlamydia* infection in 20 years. In addition, the model predicted that vaccinating
523 100% of women would have a greater epidemiological impact than vaccinating both sexes
524 (42). Unfortunately there are risks and ethical questions associated with vaccination programs
525 as demonstrated by the first *Chlamydia* vaccine using live attenuated bacterium (43). Thus,
526 research is needed to develop an efficient and safe chlamydial vaccine.

527

528

529 **REFERENCES**

- 530 1. **Agrawal, T., A. R. Bhengraj, V. Vats, S. Salhan, and A. Mittal.** 2011. Expression of TLR 2, TLR
531 4 and iNOS in cervical monocytes of Chlamydia trachomatis-infected women and their role in
532 host immune response. *Am J Reprod Immunol* **66**:534-43.
- 533 2. **Al-Zeer, M. A., H. M. Al-Younes, D. Lauster, M. Abu Lubad, and T. F. Meyer.** 2013.
534 Autophagy restricts Chlamydia trachomatis growth in human macrophages via IFNG-
535 inducible guanylate binding proteins. *Autophagy* **9**:50-62.
- 536 3. **Bachmaier, K., N. Neu, L. M. de la Maza, S. Pal, A. Hessel, and J. M. Penninger.** 1999.
537 Chlamydia infections and heart disease linked through antigenic mimicry. *Science* **283**:1335-
538 9.
- 539 4. **Barenfanger, J., and A. B. MacDonald.** 1974. The role of immunoglobulin in the
540 neutralization of trachoma infectivity. *J Immunol* **113**:1607-17.
- 541 5. **Bartolini, E., E. Ianni, E. Frigimelica, R. Petracca, G. Galli, F. B. Scorza, N. Norais, D. Laera, F.**
542 **Giusti, A. Pierleoni, M. Donati, R. Cevenini, O. Finco, G. Grandi, and R. Grifantini.** 2013.
543 Recombinant outer membrane vesicles carrying Chlamydia muridarum HtrA induce
544 antibodies that neutralize chlamydial infection in vitro. *Journal of Extracellular vesicles* **2**.
- 545 6. **Bas, S., L. Neff, M. Vuillet, U. Spenato, T. Seya, M. Matsumoto, and C. Gabay.** 2008. The
546 proinflammatory cytokine response to Chlamydia trachomatis elementary bodies in human
547 macrophages is partly mediated by a lipoprotein, the macrophage infectivity potentiator,
548 through TLR2/TLR1/TLR6 and CD14. *J Immunol* **180**:1158-68.
- 549 7. **Baud, D., L. Regan, and G. Greub.** 2008. Emerging role of Chlamydia and Chlamydia-like
550 organisms in adverse pregnancy outcomes. *Curr Opin Infect Dis* **21**:70-6.
- 551 8. **Beagley, K. W., W. M. Huston, P. M. Hansbro, and P. Timms.** 2009. Chlamydial infection of
552 immune cells: altered function and implications for disease. *Crit Rev Immunol* **29**:275-305.

- 553 9. **Bebear, C., and B. de Barbeyrac.** 2009. Genital Chlamydia trachomatis infections. Clin
554 Microbiol Infect **15**:4-10.
- 555 10. **Boyoglu, S., K. Vig, S. Pillai, V. Rangari, V. A. Dennis, F. Khazi, and S. R. Singh.** 2009.
556 Enhanced delivery and expression of a nanoencapsulated DNA vaccine vector for respiratory
557 syncytial virus. Nanomedicine **5**:463-72.
- 558 11. **Brandtzaeg, P.** 2009. Mucosal immunity: induction, dissemination, and effector functions.
559 Scand J Immunol **70**:505-15.
- 560 12. **Brunham, R. C., and J. Rey-Ladino.** 2005. Immunology of Chlamydia infection: implications
561 for a Chlamydia trachomatis vaccine. Nat Rev Immunol **5**:149-61.
- 562 13. **Byrne, G. I., R. S. Stephens, G. Ada, H. D. Caldwell, H. Su, R. P. Morrison, B. Van der Pol, P.
563 Bavoil, L. Bobo, S. Everson, and et al.** 1993. Workshop on in vitro neutralization of Chlamydia
564 trachomatis: summary of proceedings. J Infect Dis **168**:415-20.
- 565 14. **Cambridge, C. D., S. R. Singh, A. B. Waffo, S. J. Fairley, and V. A. Dennis.** 2013. Formulation,
566 characterization, and expression of a recombinant MOMP Chlamydia trachomatis DNA
567 vaccine encapsulated in chitosan nanoparticles. Int J Nanomedicine **8**:1759-71.
- 568 15. **Cao, H., R. G. Wolff, M. S. Meltzer, and R. M. Crawford.** 1989. Differential regulation of class
569 II MHC determinants on macrophages by IFN-gamma and IL-4. J Immunol **143**:3524-31.
- 570 16. **Carmichael, J. R., S. Pal, D. Tifrea, and L. M. de la Maza.** 2011. Induction of protection
571 against vaginal shedding and infertility by a recombinant Chlamydia vaccine. Vaccine
572 **29**:5276-83.
- 573 17. **Chen, L., L. Lei, Z. Zhou, J. He, S. Xu, C. Lu, J. Chen, Z. Yang, G. Wu, I. T. Yeh, G. Zhong, and Y.
574 Wu.** 2013. Contribution of interleukin-12 p35 (IL-12p35) and IL-12p40 to protective immunity
575 and pathology in mice infected with Chlamydia muridarum. Infect Immun **81**:2962-71.
- 576 18. **Cheng, C., I. Bettahi, M. I. Cruz-Fisher, S. Pal, P. Jain, Z. Jia, J. Holmgren, A. M. Harandi, and
577 L. M. de la Maza.** 2009. Induction of protective immunity by vaccination against Chlamydia

- 578 trachomatis using the major outer membrane protein adjuvanted with CpG
579 oligodeoxynucleotide coupled to the nontoxic B subunit of cholera toxin. *Vaccine* **27**:6239-
580 46.
- 581 19. **Cheng, C., S. Pal, I. Bettahi, K. L. Oxford, P. A. Barry, and L. M. de la Maza.** 2011.
582 Immunogenicity of a vaccine formulated with the *Chlamydia trachomatis* serovar F, native
583 major outer membrane protein in a nonhuman primate model. *Vaccine* **29**:3456-64.
- 584 20. **Childs, T. S., and W. C. Webley.** 2012. In vitro assessment of halobacterial gas vesicles as a
585 *Chlamydia* vaccine display and delivery system. *Vaccine* **30**:5942-8.
- 586 21. **Corsaro, D., and G. Greub.** 2006. Pathogenic potential of novel *Chlamydiae* and diagnostic
587 approaches to infections due to these obligate intracellular bacteria. *Clin Microbiol Rev*
588 **19**:283-97.
- 589 22. **Crotzer, V. L., and J. S. Blum.** 2009. Autophagy and its role in MHC-mediated antigen
590 presentation. *J Immunol* **182**:3335-41.
- 591 23. **Cunningham, K. A., A. J. Carey, L. Hafner, P. Timms, and K. W. Beagley.** 2011. *Chlamydia*
592 *muridarum* major outer membrane protein-specific antibodies inhibit in vitro infection but
593 enhance pathology in vivo. *Am J Reprod Immunol* **65**:118-26.
- 594 24. **de Clercq, E., I. Kalmar, and D. Vanrompay.** 2013. Animal models for studying *Chlamydia*
595 *trachomatis* female genital tract infection. *Infect. Immun.*
- 596 25. **Deane, K. H., R. M. Jecock, J. H. Pearce, and J. S. Gaston.** 1997. Identification and
597 characterization of a DR4-restricted T cell epitope within *chlamydia* heat shock protein 60.
598 *Clin Exp Immunol* **109**:439-45.
- 599 26. **Detmer, A., and J. Glenting.** 2006. Live bacterial vaccines--a review and identification of
600 potential hazards. *Microb Cell Fact* **5**:23.

- 601 27. **Domeika, M., K. Domeika, J. Paavonen, P. A. Mardh, and S. S. Witkin.** 1998. Humoral
602 immune response to conserved epitopes of *Chlamydia trachomatis* and human 60-kDa heat-
603 shock protein in women with pelvic inflammatory disease. *J Infect Dis* **177**:714-9.
- 604 28. **Dong-Ji, Z., X. Yang, C. Shen, H. Lu, A. Murdin, and R. C. Brunham.** 2000. Priming with
605 *Chlamydia trachomatis* major outer membrane protein (MOMP) DNA followed by MOMP
606 ISCOM boosting enhances protection and is associated with increased immunoglobulin A and
607 Th1 cellular immune responses. *Infect Immun* **68**:3074-8.
- 608 29. **Dong, F., M. Pirbhai, Y. Zhong, and G. Zhong.** 2004. Cleavage-dependent activation of a
609 chlamydia-secreted protease. *Mol Microbiol* **52**:1487-94.
- 610 30. **Eko, F. O., E. Ekong, Q. He, C. M. Black, and J. U. Igietseme.** 2011. Induction of immune
611 memory by a multisubunit chlamydial vaccine. *Vaccine* **29**:1472-80.
- 612 31. **Eko, F. O., D. N. Okenu, U. P. Singh, Q. He, C. Black, and J. U. Igietseme.** 2011. Evaluation of
613 a broadly protective *Chlamydia*-cholera combination vaccine candidate. *Vaccine* **29**:3802-10.
- 614 32. **Entrican, G., N. Wheelhouse, S. R. Wattegedera, and D. Longbottom.** 2012. New challenges
615 for vaccination to prevent chlamydial abortion in sheep. *Comp Immunol Microbiol Infect Dis*
616 **35**:271-6.
- 617 33. **Fairley, S. J., S. R. Singh, A. N. Yilma, A. B. Waffo, P. Subbarayan, S. Dixit, M. A. Taha, C. D.**
618 **Cambridge, and V. A. Dennis.** 2013. *Chlamydia trachomatis* recombinant MOMP
619 encapsulated in PLGA nanoparticles triggers primarily T helper 1 cellular and antibody
620 immune responses in mice: a desirable candidate nanovaccine. *Int J Nanomedicine* **8**:2085-
621 99.
- 622 34. **Farris, C. M., S. G. Morrison, and R. P. Morrison.** 2010. CD4+ T cells and antibody are
623 required for optimal major outer membrane protein vaccine-induced immunity to *Chlamydia*
624 *muridarum* genital infection. *Infect Immun* **78**:4374-83.

- 625 35. **Faurez, F., D. Dory, V. Le Moigne, R. Gravier, and A. Jestin.** 2010. Biosafety of DNA vaccines:
626 new generation of DNA vectors and current knowledge on the fate of plasmids after
627 injection. *Vaccine* **28**.
- 628 36. **Finco, O., E. Frigimelica, F. Buricchi, R. Petracca, G. Galli, E. Faenzi, E. Meoni, A. Bonci, M.**
629 **Agnusdei, F. Nardelli, E. Bartolini, M. Scarselli, E. Caproni, D. Laera, L. Zedda, D. Skibinski, S.**
630 **Giovinazzi, R. Bastone, E. Ianni, R. Cevenini, G. Grandi, and R. Grifantini.** 2011. Approach to
631 discover T- and B-cell antigens of intracellular pathogens applied to the design of *Chlamydia*
632 *trachomatis* vaccines. *Proc Natl Acad Sci U S A* **108**:9969-74.
- 633 37. **Gallichan, W. S., and K. L. Rosenthal.** 1998. Long-term immunity and protection against
634 herpes simplex virus type 2 in the murine female genital tract after mucosal but not systemic
635 immunization. *J Infect Dis* **177**:1155-61.
- 636 38. **Gaydos, C. A., C. Wright, B. J. Wood, G. Waterfield, S. Hobson, and T. C. Quinn.** 2008.
637 *Chlamydia trachomatis* reinfection rates among female adolescents seeking rescreening in
638 school-based health centers. *Sex Transm Dis* **35**:233-7.
- 639 39. **Gondek, D. C., A. J. Olive, G. Stary, and M. N. Starnbach.** 2012. CD4+ T cells are necessary
640 and sufficient to confer protection against *Chlamydia trachomatis* infection in the murine
641 upper genital tract. *J Immunol* **189**:2441-9.
- 642 40. **Goodall, J. C., H. Beacock-Sharp, K. H. Deane, and J. S. Gaston.** 2001. Recognition of the 60
643 kilodalton cysteine-rich outer membrane protein OMP2 by CD4(+) T cells from humans
644 infected with *Chlamydia trachomatis*. *Clin Exp Immunol* **126**:488-93.
- 645 41. **Goodall, J. C., G. Yeo, M. Huang, R. Raggiaschi, and J. S. Gaston.** 2001. Identification of
646 *Chlamydia trachomatis* antigens recognized by human CD4+ T lymphocytes by screening an
647 expression library. *Eur J Immunol* **31**:1513-22.

- 648 42. **Gray, R. T., K. W. Beagley, P. Timms, and D. P. Wilson.** 2009. Modeling the impact of
649 potential vaccines on epidemics of sexually transmitted *Chlamydia trachomatis* infection. *J*
650 *Infect Dis* **199**:1680-8.
- 651 43. **Grayston, J. T., and S. P. Wang.** 1978. The potential for vaccine against infection of the
652 genital tract with *Chlamydia trachomatis*. *Sex Transm Dis* **5**:73-7.
- 653 44. **Gupta, S., R. Janani, Q. Bin, P. Luciw, C. Greer, S. Perri, H. Legg, J. Donnelly, S. Barnett, D.**
654 **O'Hagan, J. M. Polo, and M. Vajdy.** 2005. Characterization of human immunodeficiency virus
655 Gag-specific gamma interferon-expressing cells following protective mucosal immunization
656 with alphavirus replicon particles. *J Virol* **79**:7135-45.
- 657 45. **Hamdy, S., O. Molavi, Z. Ma, A. Haddadi, A. Alshamsan, Z. Gobti, S. Elhasi, J. Samuel, and A.**
658 **Lavasanifar.** 2008. Co-delivery of cancer-associated antigen and Toll-like receptor 4 ligand in
659 PLGA nanoparticles induces potent CD8+ T cell-mediated anti-tumor immunity. *Vaccine*
660 **26**:5046-57.
- 661 46. **He, Q., L. Martinez-Sobrido, F. O. Eko, P. Palese, A. Garcia-Sastre, D. Lyn, D. Okenu, C.**
662 **Bandea, G. A. Ananaba, C. M. Black, and J. U. Igietseme.** 2007. Live-attenuated influenza
663 viruses as delivery vectors for *Chlamydia* vaccines. *Immunology* **122**:28-37.
- 664 47. **He, Q., T. T. Moore, F. O. Eko, D. Lyn, G. A. Ananaba, A. Martin, S. Singh, L. James, J. Stiles,**
665 **C. M. Black, and J. U. Igietseme.** 2005. Molecular basis for the potency of IL-10-deficient
666 dendritic cells as a highly efficient APC system for activating Th1 response. *J Immunol*
667 **174**:4860-9.
- 668 48. **Hickey, D. K., F. E. Aldwell, and K. W. Beagley.** 2010. Oral immunization with a novel lipid-
669 based adjuvant protects against genital *Chlamydia* infection. *Vaccine* **28**:1668-72.
- 670 49. **Hjelholt, A., G. Christiansan, T. G. Johansson, H. J. Ingerslev, and S. Birkelund.** 2011. Tubal
671 factor infertility is associated with antibodies against *Chlamydia trachomatis* heat shock
672 protein 60 (HSP60) but not human HSP60. *Human Reproduction* **0**.

- 673 50. **Holland, M. J., D. J. Conway, T. J. Blanchard, O. M. Mahdi, R. L. Bailey, H. C. Whittle, and D.**
674 **C. Mabey.** 1997. Synthetic peptides based on *Chlamydia trachomatis* antigens identify
675 cytotoxic T lymphocyte responses in subjects from a trachoma-endemic population. *Clin Exp*
676 *Immunol* **107**:44-9.
- 677 51. **Hosenfeld, C. B., K. A. Workowski, S. Berman, A. Zaidi, J. Dyson, D. Mosure, G. Bolan, and**
678 **H. M. Bauer.** 2009. Repeat infection with *Chlamydia* and gonorrhea among females: a
679 systematic review of the literature. *Sex Transm Dis* **36**:478-89.
- 680 52. **Huston, W. M., M. Harvie, A. Mittal, P. Timms, and K. W. Beagley.** 2012. Vaccination to
681 protect against infection of the female reproductive tract. *Expert Rev Clin Immunol* **8**:81-94.
- 682 53. **Ibana, J. A., L. Myers, C. Porretta, M. Lewis, S. N. Taylor, D. H. Martin, and A. J. Quayle.**
683 2012. The major CD8 T cell effector memory subset in the normal and *Chlamydia*
684 *trachomatis*-infected human endocervix is low in perforin. *BMC Immunol* **13**:66.
- 685 54. **Ibana, J. A., D. J. Schust, J. Sugimoto, T. Nagamatsu, S. J. Greene, and A. J. Quayle.** 2011.
686 *Chlamydia trachomatis* immune evasion via downregulation of MHC class I surface
687 expression involves direct and indirect mechanisms. *Infect Dis Obstet Gynecol* **2011**:420905.
- 688 55. **Ifere, G. O., Q. He, J. U. Igietseme, G. A. Ananaba, D. Lyn, W. Lubitz, K. L. Kellar, C. M. Black,**
689 **and F. O. Eko.** 2007. Immunogenicity and protection against genital *Chlamydia* infection and
690 its complications by a multisubunit candidate vaccine. *J Microbiol Immunol Infect* **40**:188-
691 200.
- 692 56. **Igietseme, J. U., and C. M. Black.** 2013. *Chlamydial Infection: A Clinical and Public Health*
693 *Perspective*, vol. 7.
- 694 57. **Igietseme, J. U., F. O. Eko, Q. He, and C. M. Black.** 2004. Antibody regulation of Tcell
695 immunity: implications for vaccine strategies against intracellular pathogens. *Expert Rev*
696 *Vaccines* **3**:23-34.

- 697 58. **Igietseme, J. U., and A. Murdin.** 2000. Induction of protective immunity against Chlamydia
698 trachomatis genital infection by a vaccine based on major outer membrane protein-lipophilic
699 immune response-stimulating complexes. *Infect Immun* **68**:6798-806.
- 700 59. **Jawetz, E., L. Rose, L. Hanna, and P. Thygeson.** 1965. Experimental inclusion conjunctivitis in
701 man: measurements of infectivity and resistance. *JAMA* **194**:620-32.
- 702 60. **Jayarapu, K., M. Kerr, S. Ofner, and R. M. Johnson.** 2010. Chlamydia-specific CD4 T cell
703 clones control Chlamydia muridarum replication in epithelial cells by nitric oxide-dependent
704 and -independent mechanisms. *J Immunol* **185**:6911-20.
- 705 61. **Jendro, M. C., T. Deutsch, B. Korber, L. Kohler, J. G. Kuipers, B. Krausse-Opatz, J.**
706 **Westermann, E. Raum, and H. Zeidler.** 2000. Infection of human monocyte-derived
707 macrophages with Chlamydia trachomatis induces apoptosis of T cells: a potential
708 mechanism for persistent infection. *Infect Immun* **68**:6704-11.
- 709 62. **Jendro, M. C., F. Fingerle, T. Deutsch, A. Liese, L. Kohler, J. G. Kuipers, E. Raum, M. Martin,**
710 **and H. Zeidler.** 2004. Chlamydia trachomatis-infected macrophages induce apoptosis of
711 activated T cells by secretion of tumor necrosis factor-alpha in vitro. *Med Microbiol Immunol*
712 **193**:45-52.
- 713 63. **Jiang, J. Q., A. Patrick, R. B. Moss, and K. L. Rosenthal.** 2005. CD8+ T-cell-mediated cross-
714 clade protection in the genital tract following intranasal immunization with inactivated
715 human immunodeficiency virus antigen plus CpG oligodeoxynucleotides. *J Virol* **79**:393-400.
- 716 64. **Jiang, X., C. Shen, J. Rey-Ladino, H. Yu, and R. C. Brunham.** 2008. Characterization of murine
717 dendritic cell line JAWS II and primary bone marrow-derived dendritic cells in Chlamydia
718 muridarum antigen presentation and induction of protective immunity. *Infect Immun*
719 **76**:2392-401.

- 720 65. **Johansson, M., and N. Lycke.** 2001. Immunological memory in B-cell-deficient mice conveys
721 long-lasting protection against genital tract infection with *Chlamydia trachomatis* by rapid
722 recruitment of T cells. *Immunology* **102**:199-208.
- 723 66. **Johnson, R. M., H. Yu, M. S. Kerr, J. E. Slaven, K. P. Karunakaran, and R. C. Brunham.** 2012.
724 PmpG303-311, a protective vaccine epitope that elicits persistent cellular immune responses
725 in *Chlamydia muridarum*-immune mice. *Infect Immun* **80**:2204-11.
- 726 67. **Kalbina, I., A. Wallin, I. Lindh, P. Engstrom, S. Andersson, and K. Strid.** 2011. A novel
727 chimeric MOMP antigen expressed in *Escherichia coli*, *Arabidopsis thaliana*, and *Daucus*
728 *carota* as a potential *Chlamydia trachomatis* vaccine candidate. *Protein Expr Purif* **80**:194-
729 202.
- 730 68. **Karaer, A., I. Mert, S. Cavkaytar, and S. Batioglu.** 2013. Serological investigation of the role
731 of selected sexually transmitted infections in the aetiology of ectopic pregnancy. *Eur J*
732 *Contracept Reprod Health Care* **18**:68-74.
- 733 69. **Kavanagh, K., L. A. Wallace, C. Robertson, P. Wilson, and A. Scoular.** 2013. Estimation of the
734 risk of tubal factor infertility associated with genital chlamydial infection in women: a
735 statistical modelling study. *Int J Epidemiol* **42**:493-503.
- 736 70. **Kelly, K. A., A. M. Chan, A. Butch, and T. Darville.** 2009. Two different homing pathways
737 involving integrin beta7 and E-selectin significantly influence trafficking of CD4 cells to the
738 genital tract following *Chlamydia muridarum* infection. *Am J Reprod Immunol* **61**:438-45.
- 739 71. **Kelly, K. A., J. C. Walker, S. H. Jameel, H. L. Gray, and R. G. Rank.** 2000. Differential
740 regulation of CD4 lymphocyte recruitment between the upper and lower regions of the
741 genital tract during *Chlamydia trachomatis* infection. *Infect Immun* **68**:1519-28.
- 742 72. **Kim, S. K., L. Devine, M. Angevine, R. DeMars, and P. B. Kavathas.** 2000. Direct detection
743 and magnetic isolation of *Chlamydia trachomatis* major outer membrane protein-specific
744 CD8+ CTLs with HLA class I tetramers. *J Immunol* **165**:7285-92.

- 745 73. **kumar, U., S. Kumar, S. Varghese, R. Chamoli, and P. Barthwal.** 2013. DNA Vaccine: A
746 modern biotechnological approach towards human welfare and clinical trials. *International*
747 *Journal of Research in Biomedicine and Biotechnology* **3**:17-20.
- 748 74. **Langemann, T., V. J. Koller, A. Muhammad, P. Kudela, U. B. Mayr, and W. Lubitz.** 2010. The
749 Bacterial Ghost platform system: production and applications. *Bioeng Bugs* **1**:326-36.
- 750 75. **Li, W., A. K. Murthy, B. K. Chaganty, M. N. Guentzel, J. Seshu, J. P. Chambers, G. Zhong, and**
751 **B. P. Arulanandam.** 2011. Immunization with dendritic cells pulsed ex vivo with recombinant
752 chlamydial protease-like activity factor induces protective immunity against genital
753 chlamydia muridarum challenge. *Front Immunol* **2**:73.
- 754 76. **Li, Z., C. Lu, B. Peng, H. Zeng, Z. Zhou, Y. Wu, and G. Zhong.** 2012. Induction of protective
755 immunity against *Chlamydia muridarum* intravaginal infection with a chlamydial glycogen
756 phosphorylase. *PLoS One* **7**:e32997.
- 757 77. **Lindqvist, M., J. Persson, K. Thorn, and A. M. Harandi.** 2009. The mucosal adjuvant effect of
758 alpha-galactosylceramide for induction of protective immunity to sexually transmitted viral
759 infection. *J Immunol* **182**:6435-43.
- 760 78. **Longbottom, D.** 2003. Chlamydial vaccine development. *J Med Microbiol* **52**:537-40.
- 761 79. **Longbottom, D., and M. Livingstone.** 2006. Vaccination against chlamydial infections of man
762 and animals. *Vet J* **171**:263-75.
- 763 80. **Lu, C., B. Peng, Z. Li, L. Lei, Z. Li, L. Chen, Q. He, G. Zhong, and Y. Wu.** 2013. Induction of
764 protective immunity against *Chlamydia muridarum* intravaginal infection with the chlamydial
765 immunodominant antigen macrophage infectivity potentiator. *Microbes Infect* **15**:329-38.
- 766 81. **Lu, H., H. Wang, H. M. Zhao, L. Zhao, Q. Chen, M. Qi, J. Liu, H. Yu, X. P. Yu, X. Yang, and W.**
767 **M. Zhao.** 2010. Dendritic cells (DCs) transfected with a recombinant adenovirus carrying
768 chlamydial major outer membrane protein antigen elicit protective immune responses
769 against genital tract challenge infection. *Biochem Cell Biol* **88**:757-65.

- 770 82. **Lu, H., and G. Zhong.** 1999. Interleukin-12 production is required for chlamydial antigen-
771 pulsed dendritic cells to induce protection against live *Chlamydia trachomatis* infection.
772 *Infect Immun* **67**:1763-9.
- 773 83. **Luzardo-Alvarez, A., N. Blarer, K. Peter, J. F. Romero, C. Reymond, G. Corradin, and B.**
774 **Gander.** 2005. Biodegradable microspheres alone do not stimulate murine macrophages in
775 vitro, but prolong antigen presentation by macrophages in vitro and stimulate a solid
776 immune response in mice. *J Control Release* **109**:62-76.
- 777 84. **Matyszak, M. K., J. L. Young, and J. S. Gaston.** 2002. Uptake and processing of *Chlamydia*
778 *trachomatis* by human dendritic cells. *Eur J Immunol* **32**:742-51.
- 779 85. **Meeusen, E. N., J. Walker, A. Peters, P. P. Pastoret, and G. Jungersen.** 2007. Current status
780 of veterinary vaccines. *Clin Microbiol Rev* **20**:489-510, table of contents.
- 781 86. **Moore, T., G. A. Ananaba, J. Bolier, S. Bowers, T. Belay, F. O. Eko, and J. U. Igietseme.** 2002.
782 Fc receptor regulation of protective immunity against *Chlamydia trachomatis*. *Immunology*
783 **105**:213-21.
- 784 87. **Morrison, R. P., K. Feilzer, and D. B. Tumas.** 1995. Gene knockout mice establish a primary
785 protective role for major histocompatibility complex class II-restricted responses in
786 *Chlamydia trachomatis* genital tract infection. *Infect Immun* **63**:4661-8.
- 787 88. **Morrison, S. G., and R. P. Morrison.** 2000. In situ analysis of the evolution of the primary
788 immune response in murine *Chlamydia trachomatis* genital tract infection. *Infect Immun*
789 **68**:2870-9.
- 790 89. **Morrison, S. G., and R. P. Morrison.** 2005. A predominant role for antibody in acquired
791 immunity to chlamydial genital tract reinfection. *J Immunol* **175**:7536-42.
- 792 90. **Murthy, A. K., B. K. Chaganty, W. Li, M. N. Guentzel, J. P. Chambers, J. Seshu, G. Zhong, and**
793 **B. P. Arulanandam.** 2009. A limited role for antibody in protective immunity induced by

- 794 rCPAF and CpG vaccination against primary genital *Chlamydia muridarum* challenge. *FEMS*
795 *Immunol Med Microbiol* **55**:271-9.
- 796 91. **Murthy, A. K., J. P. Chambers, P. A. Meier, G. Zhong, and B. P. Arulanandam.** 2007.
797 Intranasal vaccination with a secreted chlamydial protein enhances resolution of genital
798 *Chlamydia muridarum* infection, protects against oviduct pathology, and is highly dependent
799 upon endogenous gamma interferon production. *Infect Immun* **75**:666-76.
- 800 92. **Murthy, A. K., W. Li, B. K. Chaganty, S. Kamalakaran, M. N. Guentzel, J. Seshu, T. G.**
801 **Forsthuber, G. Zhong, and B. P. Arulanandam.** 2011. Tumor necrosis factor alpha production
802 from CD8+ T cells mediates oviduct pathological sequelae following primary genital
803 *Chlamydia muridarum* infection. *Infect Immun* **79**:2928-35.
- 804 93. **Murthy, A. K., W. Li, M. N. Guentzel, G. Zhong, and B. P. Arulanandam.** 2011. Vaccination
805 with the defined chlamydial secreted protein CPAF induces robust protection against female
806 infertility following repeated genital chlamydial challenge. *Vaccine* **29**:2519-22.
- 807 94. **Naz, R. K.** 2012. Female genital tract immunity: distinct immunological challenges for vaccine
808 development. *J Reprod Immunol* **93**:1-8.
- 809 95. **Nolz, J. C., G. R. Starbeck-Miller, and J. T. Harty.** 2011. Naive, effector and memory CD8 T-
810 cell trafficking: parallels and distinctions. *Immunotherapy* **3**:1223-33.
- 811 96. **Nunes, A., P. J. Nogueira, M. J. Borrego, and J. P. Gomes.** 2010. Adaptive evolution of the
812 *Chlamydia trachomatis* dominant antigen reveals distinct evolutionary scenarios for B- and T-
813 cell epitopes: worldwide survey. *PLoS One* **5**.
- 814 97. **O'Connell, C. M., R. R. Ingalls, C. W. Andrews, Jr., A. M. Scurlock, and T. Darville.** 2007.
815 Plasmid-deficient *Chlamydia muridarum* fail to induce immune pathology and protect against
816 oviduct disease. *J Immunol* **179**:4027-34.
- 817 98. **O'Meara, C. P., C. W. Armitage, M. C. Harvie, P. Timms, N. Y. Lycke, and K. W. Beagley.**
818 2013. Immunization with a MOMP-based vaccine protects mice against a pulmonary

- 819 Chlamydia challenge and identifies a disconnection between infection and pathology. PLoS
820 One **8**:e61962.
- 821 99. **Ojcius, D. M., Y. Bravo de Alba, J. M. Kanellopoulos, R. A. Hawkins, K. A. Kelly, R. G. Rank,**
822 **and A. Dautry-Varsat.** 1998. Internalization of Chlamydia by dendritic cells and stimulation of
823 Chlamydia-specific T cells. J Immunol **160**:1297-303.
- 824 100. **Olivares-Zavaleta, N., W. Whitmire, D. Gardner, and H. D. Caldwell.** 2010. Immunization
825 with the attenuated plasmidless Chlamydia trachomatis L2(25667R) strain provides partial
826 protection in a murine model of female genitourinary tract infection. Vaccine **28**:1454-62.
- 827 101. **Olive, A. J., D. C. Gondek, and M. N. Starnbach.** 2011. CXCR3 and CCR5 are both required for
828 T cell-mediated protection against C. trachomatis infection in the murine genital mucosa.
829 Mucosal Immunol **4**:208-16.
- 830 102. **Olsen, A. W., M. Theisen, D. Christensen, F. Follmann, and P. Andersen.** 2010. Protection
831 against Chlamydia promoted by a subunit vaccine (CTH1) compared with a primary intranasal
832 infection in a mouse genital challenge model. PLoS One **5**:e10768.
- 833 103. **Organization, W. H.** 2008. Global incidence and prevalence of selected curable sexually
834 transmitted infections. World Health Organization, Geneva, Switzerland.
- 835 104. **Ortiz, L., M. Angevine, S. K. Kim, D. Watkins, and R. DeMars.** 2000. T-cell epitopes in
836 variable segments of Chlamydia trachomatis major outer membrane protein elicit serovar-
837 specific immune responses in infected humans. Infect Immun **68**:1719-23.
- 838 105. **Ou, C., D. Tian, Y. Ling, Q. Pan, Q. He, F. O. Eko, and C. He.** 2013. Evaluation of an ompA-
839 based phage-mediated DNA vaccine against Chlamydia abortus in piglets. Int
840 Immunopharmacol **16**:505-10.
- 841 106. **Pal, S., K. M. Barnhart, Q. Wei, A. M. Abai, E. M. Peterson, and L. M. de la Maza.** 1999.
842 Vaccination of mice with DNA plasmids coding for the Chlamydia trachomatis major outer

843 membrane protein elicits an immune response but fails to protect against a genital
844 challenge. *Vaccine* **17**:459-65.

845 107. **Pal, S., C. J. Luke, A. G. Barbour, E. M. Peterson, and L. M. de la Maza.** 2003. Immunization
846 with the *Chlamydia trachomatis* major outer membrane protein, using the outer surface
847 protein A of *Borrelia burgdorferi* as an adjuvant, can induce protection against a chlamydial
848 genital challenge. *Vaccine* **21**:1455-65.

849 108. **Pal, S., I. Theodor, E. M. Peterson, and L. M. de la Maza.** 1997. Immunization with an
850 acellular vaccine consisting of the outer membrane complex of *Chlamydia trachomatis*
851 induces protection against a genital challenge. *Infect Immun* **65**:3361-9.

852 109. **Pal, S., I. Theodor, E. M. Peterson, and L. M. de la Maza.** 2001. Immunization with the
853 *Chlamydia trachomatis* mouse pneumonitis major outer membrane protein can elicit a
854 protective immune response against a genital challenge. *Infect Immun* **69**:6240-7.

855 110. **Perry, L. L., K. Feilzer, J. L. Portis, and H. D. Caldwell.** 1998. Distinct homing pathways direct
856 T lymphocytes to the genital and intestinal mucosae in *Chlamydia*-infected mice. *J Immunol*
857 **160**:2905-14.

858 111. **Peschel, G., L. Kernschmidt, C. Cirl, N. Wantia, T. Ertl, S. Durr, H. Wagner, T. Miethke, and**
859 **N. Rodriguez.** 2010. *Chlamydomydia pneumoniae* downregulates MHC-class II expression by
860 two cell type-specific mechanisms. *Mol Microbiol* **76**:648-61.

861 112. **Picard, M. D., K. P. Cohane, T. M. Gierahn, D. E. Higgins, and J. B. Flechtner.** 2012. High-
862 throughput proteomic screening identifies *Chlamydia trachomatis* antigens that are capable
863 of eliciting T cell and antibody responses that provide protection against vaginal challenge.
864 *Vaccine* **30**:4387-93.

865 113. **Ralli-Jain, P., D. Tifrea, C. Cheng, S. Pal, and L. M. de la Maza.** 2010. Enhancement of the
866 protective efficacy of a *Chlamydia trachomatis* recombinant vaccine by combining systemic
867 and mucosal routes for immunization. *Vaccine* **28**:7659-66.

- 868 114. **Rank, R. G., L. S. Soderberg, and A. L. Barron.** 1985. Chronic chlamydial genital infection in
869 congenitally athymic nude mice. *Infect Immun* **48**:847-9.
- 870 115. **Refaat, B., M. Al-Azemi, I. Geary, A. Eley, and W. Ledger.** 2009. Role of activins and inducible
871 nitric oxide in the pathogenesis of ectopic pregnancy in patients with or without *Chlamydia*
872 *trachomatis* infection. *Clin Vaccine Immunol* **16**:1493-503.
- 873 116. **Rey-Ladino, J., K. M. Koochesfahani, M. L. Zaharik, C. Shen, and R. C. Brunham.** 2005. A live
874 and inactivated *Chlamydia trachomatis* mouse pneumonitis strain induces the maturation of
875 dendritic cells that are phenotypically and immunologically distinct. *Infect Immun* **73**:1568-
876 77.
- 877 117. **Schautteet, K., E. De Clercq, Y. Jonsson, S. Lagae, K. Chiers, E. Cox, and D. Vanrompay.** 2012.
878 Protection of pigs against genital *Chlamydia trachomatis* challenge by parenteral or mucosal
879 DNA immunization. *Vaccine* **30**:2869-81.
- 880 118. **Schautteet, K., E. Stuyven, D. S. Beeckman, S. Van Acker, M. Carlon, K. Chiers, E. Cox, and D.**
881 **Vanrompay.** 2011. Protection of pigs against *Chlamydia trachomatis* challenge by
882 administration of a MOMP-based DNA vaccine in the vaginal mucosa. *Vaccine* **29**:1399-407.
- 883 119. **Shaw, J., V. Grund, L. Durling, D. Crane, and H. D. Caldwell.** 2002. Dendritic cells pulsed with
884 a recombinant chlamydial major outer membrane protein antigen elicit a CD4(+) type 2
885 rather than type 1 immune response that is not protective. *Infect Immun* **70**:1097-105.
- 886 120. **Sremac, M., and E. S. Stuart.** 2008. Recombinant gas vesicles from *Halobacterium* sp.
887 displaying SIV peptides demonstrate biotechnology potential as a pathogen peptide delivery
888 vehicle. *BMC Biotechnol* **8**:9.
- 889 121. **Su, H., and H. D. Caldwell.** 1995. CD4+ T cells play a significant role in adoptive immunity to
890 *Chlamydia trachomatis* infection of the mouse genital tract. *Infect Immun* **63**:3302-8.

- 891 122. **Su, H., R. Messer, W. Whitmire, E. Fischer, J. C. Portis, and H. D. Caldwell.** 1998. Vaccination
892 against chlamydial genital tract infection after immunization with dendritic cells pulsed ex
893 vivo with nonviable Chlamydiae. *J Exp Med* **188**:809-18.
- 894 123. **Sun, G., S. Pal, J. Weiland, E. M. Peterson, and L. M. de la Maza.** 2009. Protection against an
895 intranasal challenge by vaccines formulated with native and recombinant preparations of the
896 Chlamydia trachomatis major outer membrane protein. *Vaccine* **27**:5020-5.
- 897 124. **Sun, H. S., E. W. Eng, S. Jeganathan, A. T. Sin, P. C. Patel, E. Gracey, R. D. Inman, M. R.**
898 **Terebiznik, and R. E. Harrison.** 2012. Chlamydia trachomatis vacuole maturation in infected
899 macrophages. *J Leukoc Biol* **92**:815-27.
- 900 125. **Taha, M. A., S. R. Singh, and V. A. Dennis.** 2012. Biodegradable PLGA85/15 nanoparticles as
901 a delivery vehicle for Chlamydia trachomatis recombinant MOMP-187 peptide.
902 *Nanotechnology* **23**:325101.
- 903 126. **Teng, A., M. I. Cruz-Fisher, C. Cheng, S. Pal, G. Sun, P. Ralli-Jain, D. M. Molina, P. L. Felgner,**
904 **X. Liang, and L. M. de la Maza.** 2012. Proteomic identification of immunodominant
905 chlamydial antigens in a mouse model. *J Proteomics* **77**:176-86.
- 906 127. **Tifrea, D. F., P. Ralli-Jain, S. Pal, and L. M. de la Maza.** 2013. Vaccination with the
907 recombinant major outer membrane protein elicits antibodies to the constant domains and
908 induces cross-serovar protection against intranasal challenge with Chlamydia trachomatis.
909 *Infect Immun* **81**:1741-50.
- 910 128. **Tifrea, D. F., G. Sun, S. Pal, G. Zardeneta, M. J. Cocco, J. L. Popot, and L. M. de la Maza.**
911 2011. Amphipols stabilize the Chlamydia major outer membrane protein and enhance its
912 protective ability as a vaccine. *Vaccine* **29**:4623-31.
- 913 129. **Van Voorhis, W. C., L. K. Barrett, Y. T. Sweeney, C. C. Kuo, and D. L. Patton.** 1996. Analysis of
914 lymphocyte phenotype and cytokine activity in the inflammatory infiltrates of the upper

- 915 genital tract of female macaques infected with *Chlamydia trachomatis*. *J Infect Dis* **174**:647-
916 50.
- 917 130. **Vasilevsky, S., J. Colino, R. Puliaev, D. H. Canaday, and C. M. Snapper.** 2008. Macrophages
918 pulsed with *Streptococcus pneumoniae* elicit a T cell-dependent antibody response upon
919 transfer into naive mice. *J Immunol* **181**:1787-97.
- 920 131. **Vergara, M. R. C., A. J. Buendia Marin, L. del Rio Alonzo, F. C. Gijon, N. O. Hernandez, M. C.**
921 **G. Ruiz, and J. S. Lorente.** 2005. *Chlamydia trachomatis* genital infection: Immunity and
922 prospects for vaccine development. *Immunologia* **24**:298-312.
- 923 132. **Vicetti Miguel, R. D., S. A. Harvey, W. A. LaFramboise, S. D. Reighard, D. B. Matthews, and**
924 **T. L. Cherpes.** 2013. Human female genital tract infection by the obligate intracellular
925 bacterium *Chlamydia trachomatis* elicits robust Type 2 immunity. *PLoS One* **8**:e58565.
- 926 133. **Waeckerle-Men, Y., B. Gander, and M. Groettrup.** 2005. Delivery of tumor antigens to
927 dendritic cells using biodegradable microspheres. *Methods Mol Med* **109**:35-46.
- 928 134. **Waeckerle-Men, Y., and M. Groettrup.** 2005. PLGA microspheres for improved antigen
929 delivery to dendritic cells as cellular vaccines. *Adv Drug Deliv Rev* **57**:475-82.
- 930 135. **Wang, S. P., and J. T. Grayston.** 1991. Three new serovars of *Chlamydia trachomatis*: Da, Ia,
931 and L2a. *J Infect Dis* **163**:403-5.
- 932 136. **Witkin, S. S.** 2002. Immunological aspects of genital chlamydia infections. *Best Pract Res Clin*
933 *Obstet Gynaecol* **16**:865-74.
- 934 137. **Wizel, B., J. Nystrom-Asklin, C. Cortes, and A. Tvinnereim.** 2008. Role of CD8(+)T cells in the
935 host response to *Chlamydia*. *Microbes Infect* **10**:1420-30.
- 936 138. **Wyrick, P. B.** 2010. *Chlamydia trachomatis* persistence in vitro: an overview. *J Infect Dis* **201**
937 **Suppl 2**:S88-95.

- 938 139. **Xu, W., J. Liu, W. Gong, J. Chen, S. Zhu, and L. Zhang.** 2011. Protective immunity against
939 Chlamydia trachomatis genital infection induced by a vaccine based on the major outer
940 membrane multi-epitope human papillomavirus major capsid protein L1. *Vaccine* **29**:2672-8.
- 941 140. **Yasir, M., N. D. Pachikara, X. Bao, Z. Pan, and H. Fan.** 2011. Regulation of chlamydial
942 infection by host autophagy and vacuolar ATPase-bearing organelles. *Infect Immun* **79**:4019-
943 28.
- 944 141. **Yilma, A. N., S. R. Singh, S. J. Fairley, M. A. Taha, and V. A. Dennis.** 2012. The anti-
945 inflammatory cytokine, interleukin-10, inhibits inflammatory mediators in human epithelial
946 cells and mouse macrophages exposed to live and UV-inactivated Chlamydia trachomatis.
947 *Mediators Inflamm* **2012**:520174.
- 948 142. **Yu, H., K. P. Karunakaran, X. Jiang, C. Shen, P. Andersen, and R. C. Brunham.** 2012.
949 Chlamydia muridarum T cell antigens and adjuvants that induce protective immunity in mice.
950 *Infect Immun* **80**:1510-8.
- 951 143. **Yu, H., K. P. Karunakaran, I. Kelly, C. Shen, X. Jiang, L. J. Foster, and R. C. Brunham.** 2011.
952 Immunization with live and dead Chlamydia muridarum induces different levels of protective
953 immunity in a murine genital tract model: correlation with MHC class II peptide presentation
954 and multifunctional Th1 cells. *J Immunol* **186**:3615-21.
- 955 144. **Zhang, X., A. A. Chentoufi, G. Dasgupta, A. B. Nesburn, M. Wu, X. Zhu, D. Carpenter, S. L.**
956 **Wechsler, S. You, and L. BenMohamed.** 2009. A genital tract peptide epitope vaccine
957 targeting TLR-2 efficiently induces local and systemic CD8+ T cells and protects against
958 herpes simplex virus type 2 challenge. *Mucosal Immunol* **2**:129-43.
- 959 145. **Zhang, X. L., I. S. Tsui, C. M. Yip, A. W. Fung, D. K. Wong, X. Dai, Y. Yang, J. Hackett, and C.**
960 **Morris.** 2000. Salmonella enterica serovar typhi uses type IVB pili to enter human intestinal
961 epithelial cells. *Infect Immun* **68**:3067-73.

962 146. **Zhong, G., P. Fan, H. Ji, F. Dong, and Y. Huang.** 2001. Identification of a chlamydial protease-
963 like activity factor responsible for the degradation of host transcription factors. *J Exp Med*
964 **193**:935-42.

965 147. **Zhu, S., J. Chen, M. Zheng, W. Gong, X. Xue, W. Li, and L. Zhang.** 2010. Identification of
966 immunodominant linear B-cell epitopes within the major outer membrane protein of
967 *Chlamydia trachomatis*. *Acta Biochim Biophys Sin (Shanghai)* **42**:771-8.

968

969

Table 1. Summary of recent developments in chlamydial research including chlamydial strain/antigen utilized, cell type infected and immune response elicited

Cell type	Chlamydia/Antigen	Immune Response	Ref.
Mouse macrophage cell line (J774) and human macrophages	<i>C. trachomatis</i>	<ul style="list-style-type: none"> Live and inactivated <i>Chlamydia</i> induced elevated IL-8, IL-1β, TNF-α, IL-6. 	(3, 24)
Mouse (RAW) and human (THP-1) macrophage cell line	<i>C. trachomatis</i>	<ul style="list-style-type: none"> Live <i>Chlamydia</i> induced autophagy. 	(1, 22)
Human macrophages	<i>C. trachomatis</i>	<ul style="list-style-type: none"> Live <i>Chlamydia</i> infected macrophages induced T cell apoptosis. 	(10, 11)
Mouse BMDC	<i>C. muridarum</i>	<ul style="list-style-type: none"> DC pulsed with UV-inactivated <i>Chlamydia</i> in vitro secreted elevated levels of IL-12. DC pulsed with UV-inactivated <i>Chlamydia</i> and adoptively transferred into naive mice induced strong protection against live chlamydial lung infection. IL-12^{-/-} DC failed to induce Th-1 dominant response and did not induce strong protection against chlamydial infection. 	(14)
Mouse BMDC	rMOMP	<ul style="list-style-type: none"> DC pulsed with rMOMP secreted IL-12 and induced infection-sensitized CD4+T cells to secrete IFN-γ. DC pulsed with rMOMP and adoptively transferred into naive mice generated a Th2 anti-MOMP immune response. 	(21)
Mouse BMDC	<i>C. trachomatis</i>	<ul style="list-style-type: none"> IL-10^{-/-} DC pulsed with UV-inactivated <i>Chlamydia</i> caused early DC maturation, activation, increased ability to process and present antigens and enhanced the rate of Th1 activation. 	(7)
Mouse BMDC	<i>C. muridarum</i>	<ul style="list-style-type: none"> DC incubated with UV-inactivated <i>Chlamydia</i> expressed low levels of CD40 and CD80, secreted low levels of proinflammatory cytokines and exhibited reduced recognition by <i>Chlamydia</i>-specific CD4+ T cells. Adoptive transfer of live EB-pulsed DC was more effective than UV <i>Chlamydia</i> at protecting mice against a live intranasal chlamydial challenge. 	(20)
Mouse BMDC	<i>C. muridarum</i>	<ul style="list-style-type: none"> DC pulsed with live EBs presented 45 MHC class II <i>C. muridarum</i> peptides mapping to 13 proteins. In contrast DC pulsed with heat or UV-inactivated <i>Chlamydia</i> presented only six MHC class II chlamydial peptides mapping to 3 proteins. Only two epitopes were shared in common between live and inactivated <i>C. muridarum</i>. 	(25)
Mouse BMDC	Recombinant adenovirus carrying <i>C. trachomatis</i> MOMP	<ul style="list-style-type: none"> DC exhibited increased CD80 and MHC class II, IL-12 and were able to stimulate CD4+ T cell proliferation and IFN-γ. Adoptively transferred MOMP transfected DC generated Th1-biased cytokine production, mucosal IgA and protected mice against chlamydial genital tract infection. 	(13)
Mouse BMDC	UV <i>C. muridarum</i> + CpG or rCPAF + CpG	<ul style="list-style-type: none"> DC pulsed with rCPAF + CpG exhibited increased CD86, CD80, CD40, MHC class II, IL-12 but not IL-10 and IL-4. Mice adoptively immunized with rCPAF + CpG or UV <i>C. muridarum</i> + CpG pulsed DC produced elevated IFN-γ, IG1, IgG2a and exhibited reduced <i>Chlamydia</i> shedding and reduced oviduct pathology compared to infected mock-immunized mice. 	(12)

Mouse T cells	<i>C. muridarum</i>	<ul style="list-style-type: none"> Athymic nude mice established chronic genital tract infection whereas wild-type mice resolved infection in 20 days. 	(19)
Mouse T cells	<i>C. trachomatis</i> T cell antigens + AbISCO-100	<ul style="list-style-type: none"> Potent CD8+ T response, polyfunctional Th1-polarized CD4+ T cell responses (INF-γ, TNF-α, IL-2) and high protein specific Th1-skewed antibody response (IgG2c). Adoptive transfer of CD4+ T cells and CD8+ T cells to naive non-immunized mice protected against <i>C. trachomatis</i> vaginal challenge whereas passive transfer of immune sera did not. 	(18)
Mouse T cells	<i>C. muridarum</i> MOMP + CpG and Montanide ISA	<ul style="list-style-type: none"> Vaccinated mice were depleted of CD4+ and CD8+ T cells and challenged vaginally with live <i>C. muridarum</i>. Depletion of CD4+ T cells, but not CD8+ T cells diminished vaccine-induced protection. 	(4)
Mouse CD4+ T cells	<i>C. trachomatis</i>	<ul style="list-style-type: none"> Genital tract <i>C. trachomatis</i> infection stimulated the activation and memory development of <i>C. trachomatis</i>-specific CD4+ T cells. CD4+ T cells are necessary to confer protection against <i>C. trachomatis</i> infection. 	(6)
Mouse CD4 + T cells	<i>C. muridarum</i>	<ul style="list-style-type: none"> CD4 T cell clone-induced epithelial NO production was critical for controlling replication. Most potent CD4+ T cell clones were dependent on T cell degranulation for chlamydial replication control. 	(9)
Human CD4+ T cells	<i>C. trachomatis</i>	<ul style="list-style-type: none"> CD4+ T cells from women with genital tract infection that were pulsed ex vivo with EB secreted significantly more IL-4 than TNF-α and INF-γ. 	(23)
Mouse CD8+ T cells	<i>C. muridarum</i>	<ul style="list-style-type: none"> TNF-α from CD8+ T cells contributed significantly to oviduct pathological sequelae, but not bacterial clearance, following genital chlamydial challenge. 	(17)
Human CD8+ T cells	<i>C. trachomatis</i>	<ul style="list-style-type: none"> Endocervix effector memory CD8+ T cells from <i>C. trachomatis</i> infected women expressed low perforin levels. 	(8)
Human B cells/Antibodies		<ul style="list-style-type: none"> Identified 21 antibody inducing antigens from <i>C. trachomatis</i>-infected patients sera. 	(5)
Mouse B cells/Antibodies	Recombinant outer membrane vesicles carrying <i>C. muridarum</i> HtrA	<ul style="list-style-type: none"> Mice immunized with outer membrane vesicles carrying HtrA induced anti-HtrA-specific antibodies that neutralized <i>C. muridarum</i> infectivity in vitro. 	(2)
Mouse B cells/Antibodies	<i>C. muridarum</i> or MOMP monoclonal antibody (mAb)	<ul style="list-style-type: none"> Passive immunization with serum from <i>C. muridarum</i> infected mice conferred a marked level of protection from <i>C. muridarum</i> genital reinfection and shortened the time of infection. MOMP mAbs conferred significant level of immunity to reinfection and reduced shedding. 	(15)
Mouse B cells/Antibodies	rCPAF + CpG	<ul style="list-style-type: none"> Both wild-type and B cell deficient (μmT) mice vaccinated intranasally with rCPAF + CpG and challenged with live <i>C. muridarum</i> vaginally demonstrated comparable clearance and similar reductions in pathology. 	(16)

--	--	--	--

1. **Al-Zeer, M. A., H. M. Al-Younes, D. Lauster, M. Abu Lubad, and T. F. Meyer.** 2013. Autophagy restricts Chlamydia trachomatis growth in human macrophages via IFNG-inducible guanylate binding proteins. *Autophagy* **9**:50-62.
2. **Bartolini, E., E. Ianni, E. Frigimelica, R. Petracca, G. Galli, F. B. Scorza, N. Norais, D. Laera, F. Giusti, A. Pierleoni, M. Donati, R. Cevenini, O. Finco, G. Grandi, and R. Grifantini.** 2013. Recombinant outer membrane vesicles carrying Chlamydia muridarum HtrA induce antibodies that neutralize chlamydial infection in vitro. *Journal of Extracellular vesicles* **2**.
3. **Bas, S., L. Neff, M. Vuillet, U. Spenato, T. Seya, M. Matsumoto, and C. Gabay.** 2008. The proinflammatory cytokine response to Chlamydia trachomatis elementary bodies in human macrophages is partly mediated by a lipoprotein, the macrophage infectivity potentiator, through TLR2/TLR1/TLR6 and CD14. *J Immunol* **180**:1158-68.
4. **Farris, C. M., S. G. Morrison, and R. P. Morrison.** 2010. CD4+ T cells and antibody are required for optimal major outer membrane protein vaccine-induced immunity to Chlamydia muridarum genital infection. *Infect Immun* **78**:4374-83.
5. **Finco, O., E. Frigimelica, F. Buricchi, R. Petracca, G. Galli, E. Faenzi, E. Meoni, A. Bonci, M. Agnusdei, F. Nardelli, E. Bartolini, M. Scarselli, E. Caproni, D. Laera, L. Zedda, D. Skibinski, S. Giovinazzi, R. Bastone, E. Ianni, R. Cevenini, G. Grandi, and R. Grifantini.** 2011. Approach to discover T- and B-cell antigens of intracellular pathogens applied to the design of Chlamydia trachomatis vaccines. *Proc Natl Acad Sci U S A* **108**:9969-74.
6. **Gondek, D. C., A. J. Olive, G. Stary, and M. N. Starnbach.** 2012. CD4+ T cells are necessary and sufficient to confer protection against Chlamydia trachomatis infection in the murine upper genital tract. *J Immunol* **189**:2441-9.
7. **He, Q., T. T. Moore, F. O. Eko, D. Lyn, G. A. Ananaba, A. Martin, S. Singh, L. James, J. Stiles, C. M. Black, and J. U. Igietseme.** 2005. Molecular basis for the potency of IL-10-deficient dendritic cells as a highly efficient APC system for activating Th1 response. *J Immunol* **174**:4860-9.
8. **Ibana, J. A., L. Myers, C. Porretta, M. Lewis, S. N. Taylor, D. H. Martin, and A. J. Quayle.** 2012. The major CD8 T cell effector memory subset in the normal and Chlamydia trachomatis-infected human endocervix is low in perforin. *BMC Immunol* **13**:66.
9. **Jayarapu, K., M. Kerr, S. Ofner, and R. M. Johnson.** 2010. Chlamydia-specific CD4 T cell clones control Chlamydia muridarum replication in epithelial cells by nitric oxide-dependent and -independent mechanisms. *J Immunol* **185**:6911-20.
10. **Jendro, M. C., T. Deutsch, B. Korber, L. Kohler, J. G. Kuipers, B. Krausse-Opatz, J. Westermann, E. Raum, and H. Zeidler.** 2000. Infection of human monocyte-derived macrophages with Chlamydia trachomatis induces apoptosis of T cells: a potential mechanism for persistent infection. *Infect Immun* **68**:6704-11.
11. **Jendro, M. C., F. Fingerle, T. Deutsch, A. Liese, L. Kohler, J. G. Kuipers, E. Raum, M. Martin, and H. Zeidler.** 2004. Chlamydia trachomatis-infected macrophages induce apoptosis of activated T cells by secretion of tumor necrosis factor-alpha in vitro. *Med Microbiol Immunol* **193**:45-52.

12. **Li, W., A. K. Murthy, B. K. Chaganty, M. N. Guentzel, J. Seshu, J. P. Chambers, G. Zhong, and B. P. Arulanandam.** 2011. Immunization with dendritic cells pulsed ex vivo with recombinant chlamydial protease-like activity factor induces protective immunity against genital chlamydia muridarum challenge. *Front Immunol* **2**:73.
13. **Lu, H., H. Wang, H. M. Zhao, L. Zhao, Q. Chen, M. Qi, J. Liu, H. Yu, X. P. Yu, X. Yang, and W. M. Zhao.** 2010. Dendritic cells (DCs) transfected with a recombinant adenovirus carrying chlamydial major outer membrane protein antigen elicit protective immune responses against genital tract challenge infection. *Biochem Cell Biol* **88**:757-65.
14. **Lu, H., and G. Zhong.** 1999. Interleukin-12 production is required for chlamydial antigen-pulsed dendritic cells to induce protection against live *Chlamydia trachomatis* infection. *Infect Immun* **67**:1763-9.
15. **Morrison, S. G., and R. P. Morrison.** 2005. A predominant role for antibody in acquired immunity to chlamydial genital tract reinfection. *J Immunol* **175**:7536-42.
16. **Murthy, A. K., B. K. Chaganty, W. Li, M. N. Guentzel, J. P. Chambers, J. Seshu, G. Zhong, and B. P. Arulanandam.** 2009. A limited role for antibody in protective immunity induced by rCPAF and CpG vaccination against primary genital *Chlamydia muridarum* challenge. *FEMS Immunol Med Microbiol* **55**:271-9.
17. **Murthy, A. K., W. Li, B. K. Chaganty, S. Kamalakaran, M. N. Guentzel, J. Seshu, T. G. Forsthuber, G. Zhong, and B. P. Arulanandam.** 2011. Tumor necrosis factor alpha production from CD8+ T cells mediates oviduct pathological sequelae following primary genital *Chlamydia muridarum* infection. *Infect Immun* **79**:2928-35.
18. **Picard, M. D., K. P. Cohane, T. M. Gierahn, D. E. Higgins, and J. B. Flechtner.** 2012. High-throughput proteomic screening identifies *Chlamydia trachomatis* antigens that are capable of eliciting T cell and antibody responses that provide protection against vaginal challenge. *Vaccine* **30**:4387-93.
19. **Rank, R. G., L. S. Soderberg, and A. L. Barron.** 1985. Chronic chlamydial genital infection in congenitally athymic nude mice. *Infect Immun* **48**:847-9.
20. **Rey-Ladino, J., K. M. Koochesfahani, M. L. Zaharik, C. Shen, and R. C. Brunham.** 2005. A live and inactivated *Chlamydia trachomatis* mouse pneumonitis strain induces the maturation of dendritic cells that are phenotypically and immunologically distinct. *Infect Immun* **73**:1568-77.
21. **Shaw, J., V. Grund, L. Durling, D. Crane, and H. D. Caldwell.** 2002. Dendritic cells pulsed with a recombinant chlamydial major outer membrane protein antigen elicit a CD4(+) type 2 rather than type 1 immune response that is not protective. *Infect Immun* **70**:1097-105.
22. **Sun, H. S., E. W. Eng, S. Jeganathan, A. T. Sin, P. C. Patel, E. Gracey, R. D. Inman, M. R. Terebiznik, and R. E. Harrison.** 2012. *Chlamydia trachomatis* vacuole maturation in infected macrophages. *J Leukoc Biol* **92**:815-27.
23. **Vicetti Miguel, R. D., S. A. Harvey, W. A. LaFramboise, S. D. Reighard, D. B. Matthews, and T. L. Cherpes.** 2013. Human female genital tract infection by the obligate intracellular bacterium *Chlamydia trachomatis* elicits robust Type 2 immunity. *PLoS One* **8**:e58565.
24. **Yilma, A. N., S. R. Singh, S. J. Fairley, M. A. Taha, and V. A. Dennis.** 2012. The anti-inflammatory cytokine, interleukin-10, inhibits inflammatory mediators in human epithelial cells and mouse macrophages exposed to live and UV-inactivated *Chlamydia trachomatis*. *Mediators Inflamm* **2012**:520174.

25. **Yu, H., K. P. Karunakaran, I. Kelly, C. Shen, X. Jiang, L. J. Foster, and R. C. Brunham.** 2011. Immunization with live and dead *Chlamydia muridarum* induces different levels of protective immunity in a murine genital tract model: correlation with MHC class II peptide presentation and multifunctional Th1 cells. *J Immunol* **186**:3615-21.

Table 2. Summary of recent developments in chlamydial vaccine research

<u>Vaccines</u>	<u>Advantages</u>	<u>Disadvantages</u>	<u>Ag/Adjuvants</u>	<u>Ag Immunization Route</u>	<u>Model/ Chlamydia Infection Route</u>	<u>Immune Response</u>	<u>Ref.</u>
Intact <i>Chlamydia</i>	<ul style="list-style-type: none"> • Intact Ag • Native configuration • Replication • Humoral/Cellular immunity 	<ul style="list-style-type: none"> • Requires refrigeration • Potential reverting to virulent strains • Large scale production difficult • Possible transmission to unvaccinated individuals 	Plasmid-deficient <i>Chlamydia</i> (CM972, CM3.1)		Mouse / i.v.	<ul style="list-style-type: none"> • Elevated IgG2a (Th1); low levels of IgG1 (Th2) • Mutants do not stimulate TLR2-dependent cytokine production • Infected mice with mutant <i>Chlamydia</i> and challenged with wt <i>Chlamydia</i> are protected against oviduct disease 	(17)
			Plasmid-Deficient <i>Chlamydia</i> (L2)		Mouse / i.v	<ul style="list-style-type: none"> • Elevated IgG2a; low IgG1; no IgA (mucosal) • No pathology in the urogenital tract induced by L2 • Mice vaccinated with plasmid-deficient bacterium were not protected from infection/inflammation with secondary wt chlamydial infection 	(19)
Purified Subunits	<ul style="list-style-type: none"> • Do not revert to virulent strains • Avoids undesirable antigens 	<ul style="list-style-type: none"> • Expensive to produce • Purification not standardized • Difficult to maintain native conformation of antigen complex 	MOMP + subunit B cholera toxin conjugated to CpG	i.m. + s.c.	Mouse / i.n.	<ul style="list-style-type: none"> • Elevated IgG2a, IgG3 (Th1); lower IgG1 • Elevated INF-γ (Th1) 	(2)
			MOMP-ISCOM	i.n. or i.m.	Mouse / i.n.	<ul style="list-style-type: none"> • i.m. induced highest INF-γ and IL-4 (Th2) 	(10)
			MOMP + Freund's adjuvant	i.m + s.c	Mouse / i.v.	<ul style="list-style-type: none"> • Vortexed MOMP elicited higher IgG2a vs IgG1 • Sonicated MOMP elicited higher IgG1 vs IgG2a 	(23)
			MOMP + IC31	i.m + s.c	Mouse / i.n.	<ul style="list-style-type: none"> • Higher IgG1 than IgG2a 	(3)
			MOMP + CpG/Montanide	i.m + s.c	Rhesus macaque	<ul style="list-style-type: none"> • Elevated IgG ,IgA, INF-γ and TNF-α 	(4)
Recombinant Proteins	<ul style="list-style-type: none"> • High yields • Inexpensive 	<ul style="list-style-type: none"> • Some proteins require post-translational modification • If produced in <i>E. coli</i> possibility of endotoxin contamination 	rMOMP + Cholera toxin/CpG or CTA1	s.l. or t.c. or i.n.	Mouse / i.n.	<ul style="list-style-type: none"> • Elevated IFN-γ and TNF-α • i.n. immunization with MOMP + either adjuvant protected mice from infection but not pathology • t.c. immunization with MOMP and CTA1-DD protected mice from pathology but <i>Chlamydia</i> burden was same as control mice 	(18)
			rMOMP + CpG /Montonide	i.m + s.c.	Mouse / i.n.	<ul style="list-style-type: none"> • Vaccination protected against fibrotic scarring in lungs • Elevated IgG2a and lower levels of IgG1 	(27)
			rCPAF + IL-12	i.n.	Mouse / i.v.	<ul style="list-style-type: none"> • Increased IFN-γ; minimal IL-4 • Elevated IgG2a and IgA 	(15)
			rCPAF + CpG	i.n.	Mouse / i.v.	<ul style="list-style-type: none"> • Vaccination significantly prevented infertility 	(16)
			rCTH1 + CAF01	s.c.	Mouse / i.v.	<ul style="list-style-type: none"> • T cell production of TNF-α/IL-2/IFN-γ • anti-CTH1 IgG2a, IgG1 • Protection was solely CD4+T cell-mediated 	(20)
			rGlgP + CpG	i.m.	Mouse / i.v.	<ul style="list-style-type: none"> • Th1-dominant T cell response • Reduced hydrosalpinx severity 	(12)
			rMIP	i.m.	Mouse / i.v.	<ul style="list-style-type: none"> • More IgG2a vs IgG1 	(13)

			rCT043	i.m.	Mouse / i.n.	<ul style="list-style-type: none"> Elevated IFN-γ and no IL-4 Reduced hydrosalpinx severity rCT043 reduces bacterial load in a mouse model of i.n. infection 	(14)
			rCT823 + ISCOM and CT144 + ISCOM	s.c.	Mouse / i.v.	<ul style="list-style-type: none"> Elevated INF-γ, TNF-α, IL-2 No detectable IL-4 and IL-10 Elevated IgG2c (Th1) but not IgG1 	(24)
DNA Vaccines	<ul style="list-style-type: none"> Cheap Easy to produce Can encode for multiple epitopes Native conformation of antigenic determinants 	<ul style="list-style-type: none"> Safety Possible genome integration Anti-DNA antibodies Not possible for non-proteins 	DNA MOMP	i.m.	Mouse / i.v.	<ul style="list-style-type: none"> Elevated levels of IgG2a and IgG1 	(22)
			Priming with MOMP and secondary boost with DNA MOMP-ISCOM	i.m.	Mouse / i.n.	<ul style="list-style-type: none"> Elevated levels of IgG2a, IgA and IFN-γ 	(6)
			DNA MOMP + GM-CSF, enterotoxin (<i>E. coli</i>) A & B	i.n. + i.v.	Pig / i.v.	<ul style="list-style-type: none"> Vaccination induced significant protection against genital challenge Protection correlated with efficient T cell priming and elevated IgA anti-MOMP antibodies and low IL-4 production 	(25)
			ompA	i.m.	Pig / i.m.		(21)
Bacterial Ghosts	<ul style="list-style-type: none"> Inactivation not required therefore relevant antigenic determinants are not denatured Easy to produce Require no refrigeration Carriage of different antigens, DNA and drugs simultaneously Recognition and phagocytosis by APC 	<ul style="list-style-type: none"> Presence of LPS 	MOMP & PorB DNA plasmid	i.m.	Mouse / i.v.	<ul style="list-style-type: none"> High levels of IgG2a and IgA 	(9)
			PmpD & PorB DNA plasmid	i.m.	Mouse / i.v.	<ul style="list-style-type: none"> High levels of IgG2a and IgA, IFN-γ and low levels of IL-5 (Th2) 	(7)
Biodegradable Polymers	<ul style="list-style-type: none"> Biodegradable Non-toxic High encapsulation capacity PLGA's are efficiently phagocytosed by DC and macrophages Chitosan has mucosal adhesiveness properties and enhanced penetration across mucosal barrier 		rMOMP encapsulated in PLGA	s.c.	Mouse	<ul style="list-style-type: none"> Elevated CD4+ and CD8+ T cells Elevated INF-γ, IL-12; reduced IL-4, IL-10 Elevated IgG2a; reduced IgG1 	(8, 26)
			Chitosan containing rMOMP DNA	i.m.			(1)

Vaccines from Transgenic Plants	<ul style="list-style-type: none"> • Low cost production • Ease of use 	<ul style="list-style-type: none"> • Requirement for strong adjuvant 	MOMP introduced into <i>A. thaliana</i> and <i>D. carota</i>				(11)
Gas Vesicles	<ul style="list-style-type: none"> • Able to express peptides from various genes 		Gen fragments coding for MOMP, OmcB, Pomp loaded into <i>Halobacterium</i> -derived gas vesicles			<ul style="list-style-type: none"> • Elicited Th-1 cytokines in human foreskin fibroblasts 	(5)

* i.m. (intramuscular), s.c. (subcutaneous), i.n. (intranasal), s.l. (sublingual), t.c. (truncutaneous), i.v. (intravaginal)

1. Cambridge, C. D., S. R. Singh, A. B. Waffo, S. J. Fairley, and V. A. Dennis. 2013. Formulation, characterization, and expression of a recombinant MOMP *Chlamydia trachomatis* DNA vaccine encapsulated in chitosan nanoparticles. *Int J Nanomedicine* 8:1759-71.
2. Cheng, C., I. Bettahi, M. I. Cruz-Fisher, S. Pal, P. Jain, Z. Jia, J. Holmgren, A. M. Harandi, and L. M. de la Maza. 2009. Induction of protective immunity by vaccination against *Chlamydia trachomatis* using the major outer membrane protein adjuvanted with CpG oligodeoxynucleotide coupled to the nontoxic B subunit of cholera toxin. *Vaccine* 27:6239-46.
3. Cheng, C., M. I. Cruz-Fisher, D. Tifrea, S. Pal, B. Wizel, and L. M. de la Maza. 2011. Induction of protection in mice against a respiratory challenge by a vaccine formulated with the *Chlamydia* major outer membrane protein adjuvanted with IC31(R). *Vaccine* 29:2437-43.
4. Cheng, C., S. Pal, I. Bettahi, K. L. Oxford, P. A. Barry, and L. M. de la Maza. 2011. Immunogenicity of a vaccine formulated with the *Chlamydia trachomatis* serovar F, native major outer membrane protein in a nonhuman primate model. *Vaccine* 29:3456-64.
5. Clifton, D. R., K. A. Fields, S. S. Grieshaber, C. A. Dooley, E. R. Fischer, D. J. Mead, R. A. Carabeo, and T. Hackstadt. 2004. A chlamydial type III translocated protein is tyrosine-phosphorylated at the site of entry and associated with recruitment of actin. *Proc Natl Acad Sci U S A* 101:10166-71.
6. Dong-Ji, Z., X. Yang, C. Shen, H. Lu, A. Murdin, and R. C. Brunham. 2000. Priming with *Chlamydia trachomatis* major outer membrane protein (MOMP) DNA followed by MOMP ISCOM boosting enhances protection and is associated with increased immunoglobulin A and Th1 cellular immune responses. *Infect Immun* 68:3074-8.
7. Eko, F. O., D. N. Okenu, U. P. Singh, Q. He, C. Black, and J. U. Igietseme. 2011. Evaluation of a broadly protective *Chlamydia*-cholera combination vaccine candidate. *Vaccine* 29:3802-10.
8. Fairley, S. J., S. R. Singh, A. N. Yilma, A. B. Waffo, P. Subbarayan, S. Dixit, M. A. Taha, C. D. Cambridge, and V. A. Dennis. 2013. *Chlamydia trachomatis* recombinant MOMP encapsulated in PLGA nanoparticles triggers primarily T helper 1 cellular and antibody immune responses in mice: a desirable candidate nanovaccine. *Int J Nanomedicine* 8:2085-99.
9. Ifere, G. O., Q. He, J. U. Igietseme, G. A. Ananaba, D. Lyn, W. Lubitz, K. L. Kellar, C. M. Black, and F. O. Eko. 2007. Immunogenicity and protection against genital *Chlamydia* infection and its complications by a multisubunit candidate vaccine. *J Microbiol Immunol Infect* 40:188-200.
10. Igietseme, J. U., and A. Murdin. 2000. Induction of protective immunity against *Chlamydia trachomatis* genital infection by a vaccine based on major outer membrane protein-lipophilic immune response-stimulating complexes. *Infect Immun* 68:6798-806.
11. Kalbina, I., A. Wallin, I. Lindh, P. Engstrom, S. Andersson, and K. Strid. 2011. A novel chimeric MOMP antigen expressed in *Escherichia coli*, *Arabidopsis thaliana*, and *Daucus carota* as a potential *Chlamydia trachomatis* vaccine candidate. *Protein Expr Purif* 80:194-202.
12. Li, Z., C. Lu, B. Peng, H. Zeng, Z. Zhou, Y. Wu, and G. Zhong. 2012. Induction of protective immunity against *Chlamydia muridarum* intravaginal infection with a chlamydial glycogen phosphorylase. *PLoS One* 7:e32997.
13. Lu, C., B. Peng, Z. Li, L. Lei, Z. Li, L. Chen, Q. He, G. Zhong, and Y. Wu. 2013. Induction of protective immunity against *Chlamydia muridarum* intravaginal infection with the chlamydial immunodominant antigen macrophage infectivity potentiator. *Microbes Infect* 15:329-38.

14. Meoni, E., E. Faenzi, E. Frigimelica, L. Zedda, D. Skibinski, S. Giovinazzi, A. Bonci, R. Petracca, E. Bartolini, G. Galli, M. Agnusdei, F. Nardelli, F. Buricchi, N. Norais, I. Ferlenghi, M. Donati, R. Cevenini, O. Finco, G. Grandi, and R. Grifantini. 2009. CT043, a protective antigen that induces a CD4+ Th1 response during *Chlamydia trachomatis* infection in mice and humans. *Infect Immun* 77:4168-76.
15. Murthy, A. K., J. P. Chambers, P. A. Meier, G. Zhong, and B. P. Arulanandam. 2007. Intranasal vaccination with a secreted chlamydial protein enhances resolution of genital *Chlamydia muridarum* infection, protects against oviduct pathology, and is highly dependent upon endogenous gamma interferon production. *Infect Immun* 75:666-76.
16. Murthy, A. K., W. Li, M. N. Guentzel, G. Zhong, and B. P. Arulanandam. 2011. Vaccination with the defined chlamydial secreted protein CPAF induces robust protection against female infertility following repeated genital chlamydial challenge. *Vaccine* 29:2519-22.
17. O'Connell, C. M., R. R. Ingalls, C. W. Andrews, Jr., A. M. Scurlock, and T. Darville. 2007. Plasmid-deficient *Chlamydia muridarum* fail to induce immune pathology and protect against oviduct disease. *J Immunol* 179:4027-34.
18. O'Meara, C. P., C. W. Armitage, M. C. Harvie, P. Timms, N. Y. Lycke, and K. W. Beagley. 2013. Immunization with a MOMP-based vaccine protects mice against a pulmonary *Chlamydia* challenge and identifies a disconnection between infection and pathology. *PLoS One* 8:e61962.
19. Olivares-Zavaleta, N., W. Whitmire, D. Gardner, and H. D. Caldwell. 2010. Immunization with the attenuated plasmidless *Chlamydia trachomatis* L2(25667R) strain provides partial protection in a murine model of female genitourinary tract infection. *Vaccine* 28:1454-62.
20. Olsen, A. W., M. Theisen, D. Christensen, F. Follmann, and P. Andersen. 2010. Protection against *Chlamydia* promoted by a subunit vaccine (CTH1) compared with a primary intranasal infection in a mouse genital challenge model. *PLoS One* 5:e10768.
21. Ou, C., D. Tian, Y. Ling, Q. Pan, Q. He, F. O. Eko, and C. He. 2013. Evaluation of an ompA-based phage-mediated DNA vaccine against *Chlamydia abortus* in piglets. *Int Immunopharmacol* 16:505-10.
22. Pal, S., K. M. Barnhart, Q. Wei, A. M. Abai, E. M. Peterson, and L. M. de la Maza. 1999. Vaccination of mice with DNA plasmids coding for the *Chlamydia trachomatis* major outer membrane protein elicits an immune response but fails to protect against a genital challenge. *Vaccine* 17:459-65.
23. Pal, S., I. Theodor, E. M. Peterson, and L. M. de la Maza. 2001. Immunization with the *Chlamydia trachomatis* mouse pneumonitis major outer membrane protein can elicit a protective immune response against a genital challenge. *Infect Immun* 69:6240-7.
24. Picard, M. D., K. P. Cohane, T. M. Gierahn, D. E. Higgins, and J. B. Flechtner. 2012. High-throughput proteomic screening identifies *Chlamydia trachomatis* antigens that are capable of eliciting T cell and antibody responses that provide protection against vaginal challenge. *Vaccine* 30:4387-93.
25. Schautteet, K., E. De Clercq, Y. Jonsson, S. Lagae, K. Chiers, E. Cox, and D. Vanrompay. 2012. Protection of pigs against genital *Chlamydia trachomatis* challenge by parenteral or mucosal DNA immunization. *Vaccine* 30:2869-81.
26. Taha, M. A., S. R. Singh, and V. A. Dennis. 2012. Biodegradable PLGA85/15 nanoparticles as a delivery vehicle for *Chlamydia trachomatis* recombinant MOMP-187 peptide. *Nanotechnology* 23:325101.
27. Tifrea, D. F., P. Ralli-Jain, S. Pal, and L. M. de la Maza. 2013. Vaccination with the recombinant major outer membrane protein elicits antibodies to the constant domains and induces cross-serovar protection against intranasal challenge with *Chlamydia trachomatis*. *Infect Immun* 81:1741-50.