

Gerster, F.M . ; Vernez, D. ; Wild, P.P. ; Hopf, N.B. **Hazardous substances in frequently used professional cleaning products.** International Journal of Occupational and Environmental Health, 20(1):46-60, 2014.

Postprint version	Final draft post-refereeing
Journal website	<a href="http://www.maneyonline.com/loi/oeht">http://www.maneyonline.com/loi/oeht</a>
Pubmed link	<a href="http://www.ncbi.nlm.nih.gov/pubmed/24804339">http://www.ncbi.nlm.nih.gov/pubmed/24804339</a>
DOI	<a href="https://doi.org/10.1179/2049396713Y.0000000052">DOI:10.1179/2049396713Y.0000000052</a>

## **Abstract**

A growing number of studies identified cleaners as a group at risk for adverse health effects of the skin and the respiratory tract. Chemical substances present in cleaning products could be responsible for these effects. Currently, only limited information is available about irritant and health hazardous chemical substances found in cleaning products. We hypothesized that chemical substances present in cleaning products are known health hazardous substances that might be involved in adverse health effects of the skin and the respiratory tract. A systematic review of cleaning products used in the Swiss cleaning sector and selected after discussing with Swiss professional cleaning companies (n=1476) to identify their most used products (n=105), was performed. Safety Data Sheets were reviewed and hazardous substances present in cleaning products were tabulated with current European and Global Harmonized System hazard labels. Professional cleaning products are mixtures of substances (arithmetic mean  $3.5 \pm 2.8$ ), and more than 130 different chemical substances were identified in 105 products. The main groups of chemicals were fragrances, glycol ethers, surfactants, solvents; and to a lesser extent phosphates, salts, detergents, pH-stabilizers, acids, and bases. Up to 75% of products contained irritant (Xi), 64% harmful (Xn) and 28% corrosive (C) labeled substances. Hazards for eyes (59%), skin (50%) and by ingestion (60%) were the most reported. Cleaning products give potentially rise to simultaneous exposures to different chemical substances. As professional cleaners represent a large workforce, and cleaning products are ubiquitously used, it is a major public health issue to better understand these exposures. The list of substances provided in this study contains important information for future occupational exposure assessment studies.

Keywords: respiratory irritant, skin irritant, cleaning, exposure hazard, fragrance, glycol ether, surfactant, solvent.

## **Introduction**

Professional cleaning is a basic service occupation worldwide, and cleaning products are used daily in different environments, both indoors and outdoors.<sup>1, 2</sup> In recent years, a growing number of scientific studies have strengthened the association of cleaning work with respiratory adverse effects including asthma.<sup>3-5</sup> In addition, skin diseases such as hand dermatitis have also been reported.<sup>6-8</sup> One explanation for the observed respiratory adverse health effects among cleaning workers is chemical exposures deriving from cleaning products.<sup>2, 9-11</sup>

Several studies have investigated the relationship between adverse health effects, cleaning activity, and cleaning products.<sup>12-19</sup> Several risk factors were identified including chemical substances, application of cleaning products and cleaning activities. It was also stated that objective and more accurate estimates of occupational exposure to cleaning products are needed to better understand the adverse effects of cleaning products.<sup>12</sup> One major difficulty in this context is the multitude of cleaning products used, and the large number of chemical substances present in these products. Moreover cleaning products are constantly evolving because of ecological, economic and consumer demands.

Safety Data Sheets (SDSs) for professional cleaning products are made available to provide workers health hazard information regarding their substances or mixtures. The current EU classification system (Directives 1999/45/EC and 67/548/EEC) defines substances and preparations as dangerous if they are: explosive (E), oxidizing (O), extremely or highly flammable (F+, F), very toxic (T+), toxic (T), harmful (Xn), corrosive (C), irritant (Xi), sensitizing (Xn or Xi), carcinogenic (T, Xn), mutagenic (T, Xn), toxic for reproduction (T, Xn) or dangerous for the environment (N). These labels are accompanied by risk phrases (R-phrases), and typical R-phrases used for cleaning products are listed in the methods section.

To better understand the exposures to cleaning products among professional cleaners, we identified frequently used professional cleaning products in Switzerland and through a systematic SDS analysis of these products, hazardous (C, Xn, Xi) substances were identified. We plan to use these results in a future exposure study to better characterize exposures to substances presenting a health hazard among professional cleaning workers.

## **Methods**

### **Selection of cleaning products**

To select a representative group of frequently used cleaning products, we mailed a letter to cleaning companies located in the French and German speaking cantons of Switzerland (n=1476, Figure 1). Cleaning companies

were asked to specify cleaning activity, company size, and cleaning products used. Cleaning companies were identified from the Federal Office of Statistics using the code for cleaning companies (“Nomenclature Générale des Activités économiques” (NOGA code) (2008)). The NOGA data contained estimates about company size by number of employees. Companies were grouped into small (5 to 49 employees), medium (50 to 250 employees) and large ( $\geq 250$  employees). Technical terms (both French and German) used in the cleaning sector were retrieved from the training manual used for professional cleaners in Switzerland.<sup>20</sup> To process the large number of responses we used the TeleForm software (Cardiff TeleForm, Version 10.5.2, San Diego, USA). The letter also included a list of cleaning products (n=488) from four major companies that manufactured, produced and/or supplied products in Switzerland. This list of cleaning products by brand names was finalized after discussions with a professional cleaning association, a medium sized cleaning company, and a training center for professional cleaners. The cleaning companies were asked to mark the cleaning products they used from the provided list, and in the case where the cleaning products they used was not listed, the company was asked to write down these names before mailing the responses back. An Excel spreadsheet was generated from TeleForm, and imported to Stata (Stata 12, Stata Corp Lp, Lakeway Drive, USA). Response rates by company size were calculated. Cleaning products marked by at least 10 cleaning companies were included in the systematic SDS analysis.

### **SDS analysis**

SDSs for cleaning products were obtained from the companies’ web sites. If SDSs were not available, products were excluded from the SDS analysis. Selected products were grouped into ten product categories; floor cleaners (FC), general purpose cleaners (GPC), polishing products (PP), carpet cleaners (CC), scale removing products (SRP), bathroom cleaners (BC), glass cleaners (GC), disinfection products (DP), kitchen cleaners (KC), and other surfaces cleaners (OSC).

A comprehensive table was created listing all substances mentioned in the SDSs under section 3. The frequency of a chemical substance’s occurrence in selected products was recorded. Section 3 of SDSs is titled ‘Composition/information on ingredients’ and provides details about hazardous substances in the mixtures. Names, product identifier (CAS number), concentration or concentration ranges and classifications according to current danger letters and R-phrases (Directives 1999/45/EC and 67/548/EEC) as well as new hazard classes and statements (Regulation (EC) No 1272/2008) are presented in the table. This was possible because Switzerland has from December 1, 2010 to June 1, 2017 to replace the current classification system (Directives 1999/45/EC

and 67/548/EEC) with the new (Regulation (EC) No 1272/2008), meeting the requirements of the Globally Harmonized System of Classification and Labeling of Chemicals (GHS). Therefore, both the current classification and the new GHS labeling were available for this study. Additional regulations (Dir 67/548/EEC, Dir 1999/45/EC, EC 1272/2008) define substance concentration restrictions regarding the listing of substances in this section.

Fragrances often do not meet the criteria to be listed in section 3 'Composition/information on ingredients' of the SDSs. However fragrances, preservatives and others are often mentioned in section 15 'Regulatory Information' as they are subject to other regulations such as substances depleting the ozone layer ((EC) No 2037/2000, persistent organic pollutants (EC) No 850/2004 and export/import of dangerous substances (EC) No 689/2008). Names of fragrances, preservatives and other chemical substances listed under section 15 of SDSs were reported in the result section.

The fractions of cleaning products containing at least one substance listed with these symbols were counted and expressed in percentage for each of the 10 product categories. Similar results were presented for the R-phrases. R-phrases relevant in this study are: harmful by inhalation (R20), harmful in contact with skin (R21), harmful if swallowed (R22), causes burns (R34), causes severe burns (R35), irritating to eyes (R36), irritating to respiratory system (R37), irritating to skin (R38), risk of serious damage to eyes (R41), may cause sensitization by skin contact (R43), danger of serious damage to health by prolonged exposure (R48), possible risk of impaired fertility (R62), possible risk of harm to the unborn child (R63), harmful: may cause lung damage if swallowed (R65), repeated exposure may cause skin dryness or cracking (R66), vapors may cause drowsiness and dizziness (R67). The fractions of cleaning products, with at least one substance listed with the R-phrases R20, R21, R22, R34, R35, R36, R37, R38, R41, R43, R48, R62, R63, R65, R66 and R67, were expressed in percentage.

## **Results**

One hundred and sixteen products were selected for SDS analysis and 11 products were excluded because of missing SDSs. In the 105 remaining selected products, 130 different chemical substances were listed in the SDSs reviewed. In average one cleaning product contained 3.5 ( $\pm 2.8$ ) chemical substances listed in section 3 of the SDSs. The composition of the cleaning products varied depending on their intended use, and substances we identified are listed in Table 1. Although the type of glycol ethers varied greatly across cleaning products, they were often (20% of the products) present in both small and large amounts (0.1-50% in the products). Most glycol ethers were found in polishing products (48%), scale removing products (42%), general purpose cleaners (37%)

and floor cleaners (36%), some (20%) were found in disinfection products and kitchen cleaners, and few (10 to 11%) were found in glass cleaners, bathroom cleaners, and carpet cleaners. The choice of surfactants were diverse but were present in 19 % of the products and their concentration ranges varied greatly (0.1-30% in the products). We particularly focused on ethanolamines, known for their sensitizing properties.<sup>21</sup> Three ethanolamines were identified: monoethanolamine, triethanolamine, and 2-diethylaminoethanol. The most frequently used was monoethanolamine, which was present in 8 products; 5 floor cleaners, (n=8), 2 general purpose cleaners, and 1 kitchen cleaner. 16% of the products contained organic solvents and the concentration ranges varied enormously (0.1-75%) making up 75% of one of the products (polishing product). Other ingredients in lower concentrations made up 18% of our substance list (Table 1): phosphates, salts, detergents, pH-stabilizers, acids, and bases. Quaternary ammonium compounds or “quats”, a substance class known for sensitizing and allergic responses among cleaners, were found in 2 products in 3-10% concentrations.<sup>2, 22</sup>

Fragrances were commonly found in the product (27% of substances found) in low concentrations (0.01-5%), except when it also acted as a solvent (30%). Interestingly up to 91% of selected cleaning products contained at least one substance which was subject to other regulations and listed under section 15 of SDSs. In total 26 substances were found under section 15 of the SDS. Most frequently present fragrances in products were linalool (20%), butylphenyl methylpropional (16%), benzisothiazolione, (16%), hexyl cinnamal (15%), limonene (14%), methylisothiazolione (12%). Aliphatic carbohydrates, amyl cinnamal, benzyl salicylate, citronellol, formaldehyde deposit alpha mixture with 5-Chlor-2-methyl-2H-isothiazol-3-on 2-Methyl-2H-isothiazol-3-on, hydroxycitronellol, hydroxyisohexyl 3-cyclohexene carboxyaldehyde, isoeugenol, sodium hydroxymethylglycinate were present in 9 to 10% of products. Alpha-isomethyl ionone, benzyl alcohol, benzyl benzoate, cinnamal, citral, coumarin, eugenol, geraniol, glutaral, octylisothiazolinone and phenoxyethanol were present in less than 7% of products.

Eleven substances listed in section 3 of SDSs were neither classified with danger, symbol letters and R-phrases nor with hazard classes and categories. The remaining 117 substances were classified with danger, symbol letters and R-phrases as well as with hazard classes and categories. Of these, 82 substances were listed in addition with hazard classifications and statements (GHS). Four substances were found in more than 10 products, 17 substances were found in 5 to 10 products, 38 in 2 to 4 products and 69 in 1 product.

By product categories, usually less than 40% of cleaning products were labeled corrosive (C) in section 3 of SDSs, with exception scale removing products (78%, Figure 2). Usually more than 70% of products were

labeled irritant (Xi), except for polishing products (33%), and more than 50% of products were labeled harmful (Xn), except of product category carpet cleaners (31%).

Fifteen R-phrases regarding human health were identified (Figure 3); corrosive (R34, R35), irritant (R36, R37, R38), harmful (R20, R21, R22), sensitizing (R43) and others (R41, R62, R63, R65, R66, R67). Figure 3 shows the percentages of products (all categories) that have been labeled with these R-phrases in section 3 of SDSs.

The response rate to the letter sent to cleaning companies was highest (50%) for large companies ( $\geq 250$  employees), and to a lesser extent for medium (24%) and small (11%) (Figure 1). These results correspond to 25% (12,500 employees) of the total professional cleaning workforce included in the survey. However, counting the number of employees from the response letters gave  $>40,000$  employees. This could be explained by the NOGA categories, of which the largest only states  $\geq 1000$  employees.

## **Discussion**

Frequently used professional cleaning products contain a multitude of chemical substances with known health effects. Cleaners may therefore be exposed to mixtures of health hazardous substances during their cleaning activity.

It is important to note, that SDSs do not list all chemical substances present in a product, as regulations define substances and concentrations that have to be listed. For example, if only small quantities of common sensitizers such as aldehydes are in the cleaning products, these are not required to be listed in Section 3 in the SDS by current labeling regulations. Only if required by other regulations, these sensitizers were listed as a cleaning product ingredient under section 15 in the SDSs. Interestingly, several substances found under section 15 of SDSs can be involved in sensitizing mechanisms and/or allergic reactions.

In our study, we selected frequently used cleaning products known from cleaning companies with five or more employees. The cleaning products included were the most popular brands (four) and according to a professional association for cleaning companies in Switzerland, estimated to cover  $> 50\%$  of the Swiss professional cleaning products market.

As mentioned above, we estimated that our results concern about 50% of the workforce of companies that were included in this study. This is because the large cleaning companies reported to have high numbers of employees



(above several thousands). As cleaning products of globally active cleaning products suppliers were reviewed, results are probably also relevant for other parts of the world such as the EU and the USA.

Not only is there a great diversity of chemical substances within cleaning products, but numerous companies offer hundreds of different cleaning products, which makes the task of assessing chemical substances used in professional cleaning products complicated. Indeed, responses we received, showed cleaning companies using products from 36 different product companies, and some reported that they produced their own products. Thus when investigating exposures among professional cleaners, a SDS review is a requirement. We believe our results provide important information regarding type of cleaning products used in this industry, common chemical substance classes found in these products and their health hazards. This knowledge should help in monitoring professional cleaners and their exposures to cleaning products and substances with known health effects.

The main challenges in conducting an occupational exposure assessment for professional cleaners are the great number of cleaning products available and the vast number of substances in these products. For further investigations, we recommend to focus on the 21 substances found in  $\geq 5$  products (Table 1). Especially of interest are: the recognized sensitizer monoethanolamine and glycol ethers, frequently found in cleaning products. Substances found in professional cleaning products may likely also be ingredients in cleaning products sold to the general public; however, we did not survey these products.<sup>21</sup> Moreover, the general public does not usually have access to SDSs, only the hazard label on the product packaging is available, thereby limiting their information regarding sensitizing and irritant substances.

Most cleaning products identified in this study were sold by global companies, selling and distributing their products world-wide. We do believe our results hold true for other industrialized countries similar to Switzerland, although the cleaning product might be given a different brand name.

## **Conclusion**

This work contributes to efforts to better understand possible exposures scenarios to pollutants associated with the application of professional cleaning products. We found that hazardous substances in cleaning products are in particular fragrances, glycol ethers, surfactants, solvents and to a lesser extent phosphates, salts, detergents, pH-stabilizers, acids, and bases. Cleaning workers that are handling these products are therefore a group at risk for several occupational exposures. Section 15 in the SDS should be conferred as several substances involved in

sensitizing mechanisms and/or allergic reactions were also listed here. Especially glycol ethers and ethanolamines are frequently used in cleaning products, and could therefore be involved in the development of adverse health effects like irritant or sensitizer induced asthma, which was found to be increased in professional cleaners. Concerning asthma, the presence of different aldehydes as fragrances is also of special interest. The fact that beside some sensitizers like ethanolamines, mainly irritants were found, suggests that pathologies of the skin and the respiratory tract do probably also occur without mechanisms of sensitization. A simultaneous exposure to several hazardous chemical substances might be involved in these pathologies. As professional cleaners represent a large workforce, and cleaning products are ubiquitously used, including private cleaning, it is a major environmental and public health issue to better understand the exposures that may be caused by the use of cleaning products. Our list of substances provides important information about which chemicals and hazards are relevant for further investigations in this field and we plan to use these results for field exposure studies.


### **Conflict of interest**

The authors have declared no conflict of interest.

### **Acknowledgements**

The authors thank the Federal Office for Public Health of Switzerland (Office fédérale de la santé publique [OFSP], Bundesamt für Gesundheit [BAG]) for funding this study.

**Figure 1: Mailing and response rates of the letter to cleaning companies**



**Figure 2: Products that contain at least one substance labeled with a C, Xi or Xn**




Figure 2: Percentages of products by product categories containing at least one substance labeled as corrosive (C), irritant (Xi) and harmful (Xn) in section 3 of SDSs. (FC), general purpose cleaners (GPC), polishing products (PP), carpet cleaners (CC), scale removing products (SRP), bathroom cleaners (BC), glass cleaners (GC), disinfection products (DP), kitchen cleaners (KC), and other surfaces cleaners (OSC).

**Figure 3: Products that contain at least one substance labeled with one of the selected R-phrases**




Figure 3: Percentages of cleaning products that have been labeled with corrosive (R34, R35), irritant (R36, R37, R38), harmful (R20, R21, R22), sensitizing (R43) and other (R41, R62, R63, R65, R66, R67) R-phrases in section 3 of SDSs.

**Table 1: List of substances found in section 3 of SDSs of 105 selected cleaning products**

Substances		EU <sup>1</sup>		GHS <sup>2</sup>		Product <sup>3</sup>	
Name	CAS	L <sup>4</sup>	R <sup>5</sup>	C <sup>6</sup>	S <sup>7</sup>	% <sup>8</sup>	N <sup>9</sup>
(d)-Limonene	5989-27-5	Xi	R10	Flam.Liq3	H226	0.1-1	2
		N	R38	AquaticAcute1	H400		
			R43	AquaticChronic1	H410		
			R50/53	SkinIrrit.2	H315		
				SkinSens.1	H317		
(L)-(-)-Ethyl lactate	687-47-8	Xi	R10	EyeDam.1	H318	3-10	1
			R37	Flam.Liq.3	H226		
			R41	STOTSE3	H335		
1,4-Dioxacycloheptadecane-5,17-dione	105-95-3	N	R10	na <sup>10</sup>	na	<5	1
			R51				
			R53				
1-Methoxy-2-hydroxypropane	107-98-2	na	R10	Flam.Liq.3	H226	0.1- <10	4
1-Penten-3-one, 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-	7779-30-8	N	R51/53	AquaticChronic2	H411	0.1-1	1
1-Propoxy-2-propanol	1569-01-3	na	R10	Flam.Liq3	H226	1-50	2
				EyeIrrit2	H319		
2-cis-3,7-Dimethyl-2,6-octadien-1-ol	106-25-2	F	R12	SkinIrrit2	H315	0.01-10	2
		Xi	R38	Flam.Gas1	H220		
				Press.Gas	H280		
2-Diethylaminoethanol	100-37-8	C	R10	SkinCorr.1B	H314	1-3	1
			R20/21/22	Flam.Liq.3	H226		

			R34	AcuteTox4	[H302 H312 H332]		
2-t-Butylcyclohexyl acetate	88-41-5	N	R51, R53	AquaticChronic2	H411	0.1-1	2
2-trans-3,7-Dimethyl-2,6- octadien-1-ol	106-24-1	na	R38 R41 R43	EyeDam.1 SkinSens.1 SkinIrrit.2	H318 H317 H315	0.01- 0.1	1
3,7-Dimethyl-6-octen-1-ol	106-22-9	Xi N	R38 R43 R51/53	SkinIrrit2 SkinSens1 AquaticChronic2	H315 H317 H411	<0.01	1
6-Octenenitrile, 3,7-dimethyl-	51566-62-2	na	R52/53	AquaticChronic3	H412	0.01- 0.1	1
Acetyl cedrene	32388-55-9	N	R50/53	AquaticAcute1 AquaticChronic1	H400 H410	0.1-1	1
Acid blue 3	3536-49-0	na	na	na	na	<0.01	1
Alanine, N,N- bis(carboxymethyl)-, sodium salt (1:3)	164462-16- 2	na	na	na	na	1-<5	2
Alcohol	64-17-5	F	R11	Flam.Liq.2	H225	1-20	8
Alcohols, C10-12, ethoxylated propoxylated	68154-97-2	N	R51 R53	AquaticCronic2	H411	1-5	3
Alcohols, C10-16, ethoxylated propoxylated	69227-22-1	Xi Xn	R22 R41	na	na	5-15	1
Alcohols, C12-14, ethoxylated	68439-50-9	Xn	R22 R41	EyeDam.1 AcuteTox.4	H318 H302	1-10	5
Alcohols, C13-15-branched and linear, butoxylated ethoxylated	111905-53- 4	Xi	R36/38	SkinIrrit2 EyeIrrit.2	H315 H319	1-30	6

Alcohols, C16-18 and C18-unsatd., ethoxylated	68920-66-1	Xn	R22	EyeDam.1	H318	1-3	1
		N	R38	AquaticAcute1	H400		
			R41	AcuteTox.4	H302		
			R50	SkinIrrit.2	H315		
Alkanes, C9-12-iso-	90622-57-4	Xn	R10	na	na	30-75	2
			R53				
			R65				
			R66				
Alkylalkoholalkoxylat	na	Xi	R36	SkinIrrit2	H315	1-10	4
			R38	Eyelrrit.2	H319		
Alkylalkoholethoxylat	na	Xi	R22	na	na	<5-15	2
		Xn	R41				
Alkyletherphosphatesodiumsalt	na	Xi	R36	na	na	1-5	1
			R38				
alpha-D-Glucopyranoside, 2-ethylhexyl	125590-73-0	Xi	R41	EyeDam.1	H318	3-10	1
Alpha-isomethylionone	127-51-5	Xi	R43	SkinSens1	H317	0.1-1	1
			R52/53	AquaticChronic3	H412		
alpha-Methyl-4-(1-methylethyl)benzenepropanal	103-95-7	Xn	R38	Repr.2	H361	0.1-1	1
		N	R43	SkinIrrit.2	H315		
			R51/53	SkinSens.1	H317		
			R62	AquaticChronic2	H411		
alpha-Terpineol	98-55-5	Xi	R22	SkinIrrit.2	H315	0.01-	3
		Xn	R41			15	
			R38				
Amides, coconut oil, N-(2-((sulfosuccinyl)oxy)ethyl), sodium salts	68784-08-7	Xi	R41	na	na	na	1



Ammonium hydroxide	1336-21-6	C	R34	SkinCorr.1B	H314	0.01-	3
		N	R50	AquaticAcute1	H400	1	
Amyl salicylate	2050-08-0	N	R51/53	na	na	<5	1
Anethole, trans	4180-23-8	N	R51/53	na	na	<5	1
Aromatic naphtha, type I	64742-95-6	Xi	R10	na	na	0.1-1	1
		Xn	R37				
		N	R53				
			R65				
			R66				
			R67				
			R51				
Benzaldehyde	100-52-7	Xn	R22	na	na	na	1
Benzenesulfonic acid, (1-methylethyl)-, sodium salt (1:1)	28348-53-0	Xi	R36	EyeIrrit.2	H319	1-10	6
Benzenesulfonic acid, 4-C10-13-sec-alkyl derivs.	85536-14-7	C	R22	SkinCorr.1C	H314	3-10	1
			R34	AcuteTox4	H302		
Benzenesulfonic acid, mono-C10-13-alkyl derivs., compds. With ethanolamine	85480-55-3	Xn	R22	EyeDam.1	H318	3-10	1
			R38	AcuteTox.4	H302		
			R41	SkinIrrit.2	H315		
Benzenesulfonic acid, mono-C10-13-alkyl derivs., sodium salts	90194-45-9	Xn	R22	EyeDam1	H318	3-10	1
			R38	AcuteTox4	H302		
			R41	SkinIrrit.2	H315		
Benzyl acetate	140-11-4	Xi	R36/37/38	SkinIrrit.2	H315	10-20	1
				EyeIrrit.2	H319		
				STOTSE3	H335		
Benzyl alcohol	100-51-6	Xn	R20/22	EyeIrrit.2	H319	1-20	5
		N	R36	AcuteTox.4	[H302		

H332]

Benzyl benzoate	120-51-4	Xn	R22	AcuteTox4	H302	1-3	1
		N	R51/53	AquaticChronic2	H411		
Benzyl salicylate	118-58-1	Xi	R43	SkinSens1	H317	0.1-1	1
		N	R51/53	AquaticChronic2	H411		
beta-Pinene	127-91-3	Xn	R65	SkinCorr.1B	H314	na	1
		N	R50	EyeDam.1	H318		
			R53	AcuteTox4	H302		
				AquaticAcute1	H400		
Butane	106-97-8	F+	R12	Flam.Gas1	H220	15-75	5
				Press.Gas	H280		
Butanedioic acid, sulfo-, 1-ester with N-(2- hydroxyethyl)dodecanamide , disodium salt	25882-44-4	Xi	R36/38	SkinIrrit2	H315	3-10	1
				EyeIrrit.2	H319		
Butoxypropanol	5131-66-8	Xi	R36/38	SkinIrrit2	H315	1-30	5
				EyeIrrit.2	H319		
C11-15 Pareth-20	68131-40-8	Xi	R22	na	na	1-5	1
		Xn	R41				
C12-15 Pareth-11	68131-39-5	Xi	R22	EyeDam.1	H318	0.1-	5
		Xn	R41	AquaticAcute1	H400	15	
		N	R50	AcuteTox.4	H302		
Camphene	79-92-5	F	R11	SkinCorr.1B	H314	na	1
		Xi	R36	EyeDam.1	H318		
		N	R50	AcuteTox.4	H302		
			R53	AquaticAcute1	H400		
Citral	5392-40-5	Xi	R38	SkinIrrit2	H315	0.01-	1
			R43	SkinSens.1	H317	0.1	

Citric acid	77-92-9	Xi	R36	EyeIrrit.2	H319	1-30	9
Coconut acid	61788-47-4	Xi	R36/38	SkinCorr.1B	H314	na	2
				EyeDam1	H318		
				AcuteTox4	H302		
				AquaticAcute1	H400		
Coumarin	91-64-5	Xn	R22	AcuteTox3	H301	0.1-1	1
			R43	SkinSens.1	H317		
Cyclohexanol, 4-(1,1-dimethylethyl)-, 1-acetate	32210-23-4	N	R51/53	AquaticChronic2	H411	0.1- <5	3
Deceth-4	26183-52-8	Xi	R22	na	na	1-15	9
		Xn	R41				
Decyl D-glucoside	54549-25-6	Xi	R36	na	na	1-5	2
D-Glucopyranose, oligomeric, decyl octyl glycosides	68515-73-1	Xi	R41	na	na	1-5	1
Diethylene Glycol Monoethyl Ether	111-90-0	Xi	R36	SkinCorr.1B	H314	1-10	14
				EyeDam.1	H318		
				AcuteTox4	H302		
				AquaticAcute1	H400		
Diethylene glycol monomethyl ether	111-77-3	na	R63	na	na	na	1
Diethylene glycol mono-n-butyl ether	112-34-5	Xi	R36	EyeIrrit.2	H319	5-30	3
Dimethyl ether	115-10-6	F+	R12	na	na	50-75	1
Diphosphoric acid, tetrapotassium salt	7320-34-5	Xi	R36	EyeIrrit.2	H314	1-3	1
Dipropylene glycol monomethyl ether	34590-94-8	na	na	na	na	1-20	11
Disodium phosphate	7558-79-4	na	na	na	na	0.1-1	1

Ethylene glycol	107-21-1	Xn	R22	AcuteTox.4	H302	1-20	5
Ethylene glycol monomontanate	73138-45-1	na	na	na	na	3-10	1
Ethylene glycol mono-n-butyl ether	111-76-2	Xi	R20	SkinIrrit.2	H315	1-20	4
		Xn	R21	Eyelrrit.2	H319		
			R22	SkinSens.1	[H302		
			R36	AcuteTox.4	H312		
			R38		H332],		
Eugenol	97-53-0	Xi	R36	Eyelrrit.2	H319	0.1-1	1
			R43	SkinSens.1	H317		
Fatty acids, coco, 2-(2-butoxyethoxy)ethyl esters	91031-83-3	Xi	R36	na	na	1-5	1
Fatty acids, coconut oil, potassium salts	61789-30-8	Xi	R36/38	Eyelrrit2	H319	1-5	3
				SkinIrrit.2	H316		
Fatty acid amides	na	Xi	R38	na	na	<5	1
			R41				
Galaxolide	1222-05-5	N	R50/53	AquaticAcute1	H400	0.1-1	1
				AquaticChronic1	H410		
Genapol X 080	9043-30-5	Xi	R22	EyeDam.1	H318	0.1-5	3
		Xn	R41	AcuteTox.4	H302		
			R51				
			R53				
Heptane	142-82-5	F	R11	Flam.Liq.2	H225	5-20	2
		Xn	R38	Asp.Tox.1	H304		
		N	R50/53	AquaticAcute1	H400		
			R65	AquaticChronic1	H410		
			R67	SkinIrrit.2	H315		
				STOTSE*	H336		

Hydrocarbons, terpene	68956-56-9	Xn	R51/53	Asp.Tox.1	H304	0.01-	3
processing by-products		N	R65	AquaticChronic2	H411	1	
Hydroxyacetic acid	79-14-1	C	R34	na	na	1-5	1
Isobutane	75-28-5	F+	R12	Flam.Gas1	H220	3-20	2
				Press.Gas	H281		
Isoeugenol	97-54-1	Xn	R21/22	SkinIrrit.2	H315	0.1-1	1
			R36/38	Eyelrrit.2	H319		
			R43	SkinSens.1	H317		
				AcuteTox.4	[H302		
					H312],		
Isopropyl alcohol	67-63-0	F	R11	Flam.Liq2	H225	1-75	16
		Xi	R20/21/22	Eyelrrit.2	H319		
		Xn	R36	STOTSE	H336		
			R36/38				
			R67				
Laurylamine	2372-82-9	C	R22	AcuteTox.3	H301	0.1-1	1
dipropylenediamine		N	R35	SkinCorr.1A	H314		
			R48/22	STOTRE2	H373		
			R50	AquaticAcute1	H400		
Lilial	80-54-6	Xi	R22	Repr.2	H361	<0.01	1
		Xn	R38	Acute Tox4	H302		
		N	R43	SkinIrrit.2	H315		
			R62	SkinSens.1	H317		
			R51/53	AquaticChronic2	H411		
Linalool	78-70-6	Xi	R38, R43	SkinIrrit2,	H315,	0.01-	2
				SkinSens.1	H317	3	
Lylal	31906-04-4	Xi	R43	SkinSens1	H317	0.1-1	1
			R52/53	AquaticChronic3	H412		

Methanesulfonic acid	75-75-2	C	R34	SkinCorr.1B	H314	3-10	1
Mineral Oil	8012-95-1	Xn	R65			5-15	1
Monoethanolamine	141-43-5	C	R20	SkinCorr.1b	H314	1-15	8
		Xn	R21	STOTSE3	H335		
			R22	EyeDam1	H318		
			R34	AcuteTox4	[H302		
			R37		H312		
					H332]		
Naphtha (petroleum), heavy alkylate	64741-65-7	Xn	R10	AcuteTox.3	H331	>75	1
			R53	Asp.Tox.1	H304		
			R65	Flam.Liq.3	H226		
			R66	AquaticChronic4	[H413		
					EUH006]		
Naphtha (petroleum), hydrotreated heavy	64742-48-9	Xn	R10	na	na	3->30	3
			R65				
			R66				
Natriumlaurylethoxylsulfate	na	Xi	R38	na	na	<5	1
			R41				
Non-ionic tensides	na	Xi	R22	na	na	5-30	2
		Xn	R38				
		N	R50				
n-Octyl-polyoxyethylene	27252-75-1	Xi	R41	na	na	1-5	1
Oxirane,methyl,polymer and oxibane, butyl ether	9038-95-3	Xn	R22	AcuteTox4	H302	3-10	2
PEG-10 Tridecyl ether	24938-91-8	Xi	R41	EyeDam1	H318	1-15	5
		N	R50				
PEG-15 Cocoate	61791-29-5	Xi	R36	na	na	1-5	2
Pentapotassium triphosphate	13845-36-8	Xi	R36/38	na	na	5-15	1

Phenol, 2-methoxy-4-propyl-	2785-87-7	Xi	R36	EyeIrrit.2	H319	0.1-1	1
			R43	SkinSens.1	H317		
Phenoxyethanol	122-99-6	Xi	R22	AcuteTox.4	H302	1-10	5
		Xn	R36	EyeIrrit2	H319		
Phosphoric acid	7664-38-2	C	R34	SkinCorr.1B	H314	5-30	4
				Met.Corr.1	H290		
Poly(oxy-1,2-ethanediyl), alpha-(1-oxo-9-octadecenyl)-omega-hydroxy-,(Z)	na	Xn	R22	AcuteTox.4	H302	1-3%	1
		Xi	R41	EyeDam.1	H318		
Poly(oxy-1,2-ethanediyl), alpha-(2-propylheptyl)-omega-hydroxy-	160875-66-1	Xi	R41	EyeDam1	H318	3-10	1
Poly(oxy-1,2-ethanediyl), alpha-isodecyl-omega-hydroxy-	61827-42-7	Xi	R22	na	na	1-15	3
		Xn	R41				
Poly(oxy-1,2-ethanediyl), alpha-sulfo-omega-hydroxy-, C10-16-alkylethers, sodium salts	68585-34-2	Xi	R38	EyeDam.1	H318	1-15	4
			R41	SkinIrrit.2	H315		
Poly(oxy-1,2-ethanediyl), alpha-tridecyl-omega-hydroxy-, branched	69011-36-5	Xn	R22	EyeDam1	H318	1-20	14
			R41	AcuteTox.4	H302		
Polymerdispersion	na	na	na	na	na	na	2
Polyoxyl 20 cetostearyl ether	68439-49-6	N	R41	na	na	0.1-1	1
			R50				
Potassium hydroxide	1310-58-3	C	R22	na	na	1-5	1
		Xn	R35				
Propane	74-98-6	F+	R12	Flam.Gas1	H220	1-30	6
				Press.Gas	H280		
Quaternary ammonium	68424-85-1	C	R21/22	SkinCorr.1B	H314	3-10	2

compounds, benzyl-C12-16-		N	R34	AquaticAcute1	H400		
alkyldimethyl, chlorides			R50	AcuteTox.4	H302		
Silicic acid, disodium salt,	10213-79-3	C	R34	SkinCorr1B	H314	1-15	3
pentahydrate		Xi	R37	STOTSE3	H335		
Silicon dioxide	7631-86-9	na	na	na	na	0.1-1	1
Sodium 2-butoxyethyl sulphate	67656-24-0	Xi	R36/38	na	na	1-5	1
Sodium benzoate	532-32-1	na	na	na	na	0.1-1	1
Sodium carbonate	497-19-8	Xi	R36	EyeIrrit.2	H319	1-3	1
Sodium chloride	7647-14-5	C	R34	SkinCorr.1B	H314	0.01-	2
			R37	Met.Corr.1	H290	10	
				STOTSE	H335		
Sodium Ethasulfate	126-92-1	Xi	R22	EyeDam.1	H318	1-5	4
		Xn	R38	SkinIrrit.2	H315		
			R41				
Sodium Hydroxide	1310-73-2	C	R35	SkinCorr.1A	H314	0.01-	3
						10	
Sodium sulfate	7757-82-6	na	na	na	na	0.01-	1
						0.1	
Solvent naphtha (petroleum),	64742-94-5	Xn	R51/53	Asp.Tox.1	H304	0.1-1	1
heavy arom.		N	R65	STOTSE3	H336		
			R66	AquaticChronic2	[H411		
			R67		EUH006]		
Solvent naphtha (petroleum),	64742-88-7	Xn	R10	na	na	25-50	1
medium aliph.			R56				
Sulfamic acid	5329-14-6	Xi	R36/38	SkinIrrit2	H315	3-15	5
			R52/53	EyeIrrit.2	H319		
				AquaticChronic3	H412		
Sulfonic acids, C13-17-sec-	85711-69-9	Xi	R38	EyeDam.1	H318	1-15	8



alkane, sodium salts			R41	SkinIrrit.2	H315		
				SkinCorr.1b	H314		
				AcuteTox.4	H302		
				AquaticAcute1	H400		
Sulfuric acid, mono-C10-16-alkyl esters, sodium salts	68585-47-7	Xi	R38	EyeDam.1	H318	3-10	1
			R41	SkinIrrit.2	H315		
Sulfuric acid, mono-C12-14-alkyl esters, sodium salts	85586-07-8	Xi	R38	na	na	na	1
			R41				
Sulfuric acid, mono-C12-16-alkyl esters, sodium salts	73296-89-6	Xi	R38	na	na	5-15	1
			R41				
Terpinolene	586-62-9	Xn	R10	na	na	<5	1
		N	R51/53				
			R65				
Tri(2-butoxyethyl) phosphate	78-51-3	na	na	na	na	1-5	4
Triethanolamine	102-71-6	Xi	R36/38	na	na	1-5	1
Water	7732-18-5	na	na	na	na	50-75	2
Waxmixture	na	na	na	na	na	na	1

Table 1: <sup>1</sup>Directives 1999/45/EC and 67/548/EEC, <sup>2</sup>Regulation (EC) No 1272/2008, <sup>3</sup>Information about amount and frequency in selection of professional cleaning product, <sup>4</sup>Danger letter, <sup>5</sup>Risk-phrase, <sup>6</sup>Hazard class, <sup>7</sup>Hazard statement, <sup>8</sup>Amount of substance in selected professional cleaning products, <sup>9</sup>Number of selected professional cleaning products that contain the listed chemical substance, <sup>10</sup>not available(na).

## 5 References

1. Zock JP. World at work: cleaners. *Occup Environ Med*. 2005; 62(8): 581-4.
2. S Q and P B. Cleaning Agents and Asthma. *Journal of Investigational Allergology and Clinical Immunology*. 2010; 20(7): 542-50.
3. Arif AA, Delclos GL, Whitehead LW, Tortolero SR and Lee ES. Occupational exposures associated with work-related asthma and work-related wheezing among U.S. workers. *American Journal of Industrial Medicine*. 2003; 44(4): 368-76.
4. Zock JP, Vizcaya D and Le Moual N. Update on asthma and cleaners. *Curr Opin Allergy Clin Immunol*. 2010; 10(2): 114-20.
5. Jaakkola JJ and Jaakkola MS. Professional cleaning and asthma. *Curr Opin Allergy Clin Immunol*. 2006; 6(2): 85-90 10.1097/01.all.0000216849.64828.55.
6. Lynde CB, Obadia M, Liss GM, Ribeiro M, Holness DL and Tarlo SM. Cutaneous and respiratory symptoms among professional cleaners. *Occupational Medicine*. 2009; 59(4): 249-54.
7. Diepgen TL and Coenraads PJ. The epidemiology of occupational contact dermatitis. *International Archives of Occupational and Environmental Health*. 1999; 72(8): 496-506.
8. Gawkrödger DJ, Lloyd MH and Hunter JAA. Occupational skin disease in hospital cleaning and kitchen workers. *Contact Dermatitis*. 1986; 15(3): 132-5.
9. Bello A, Quinn M, Perry M and Milton D. Characterization of occupational exposures to cleaning products used for common cleaning tasks-a pilot study of hospital cleaners. *Environ Health*. 2009; 8(1): 11.
10. Medina-Ramón M, Zock JP, Kogevinas M, Sunyer J, Torralba Y, Borrell A, et al. Asthma, chronic bronchitis, and exposure to irritant agents in occupational domestic cleaning: a nested case-control study. *Occup Environ Med*. 2005; 62(9): 598-606.
11. Le Moual N, Kennedy SM and Kauffmann F. Occupational Exposures and Asthma in 14,000 Adults from the General Population. *American Journal of Epidemiology*. 2004; 160(11): 1108-16.
12. Dumas O, Donnay C, Heederik DJJ, Héry M, Choudat D, Kauffmann F, et al. Occupational exposure to cleaning products and asthma in hospital workers. *Occup Environ Med*. 2012; 69(12): 883-9.
13. Zock JP, Plana E, Jarvis D, Anto JM, Kromhout H, Kennedy SM, et al. The use of household cleaning sprays and adult asthma: an international longitudinal study. *American Journal of Respiratory and Critical Care Medicine*. 2007; 176(8): 735-41.
14. Zock JP, Plana E, Anto JM, Benke G, Blanc PD, Carosso A, et al. Domestic use of hypochlorite bleach, atopic sensitization, and respiratory symptoms in adults. *Journal of Allergy and Clinical Immunology*. 2009; 124(4): 731-8 e1.
15. Makela R, Kauppi P, Suuronen K, Tuppurainen M and Hannu T. Occupational asthma in professional cleaning work: a clinical study. *Occup Med (Lond)*. 2011; 61(2): 121-6.
16. Vizcaya D, Mirabelli MC, Antó J-M, Orriols R, Burgos F, Arjona L, et al. A workforce-based study of occupational exposures and asthma symptoms in cleaning workers. *Occup Environ Med*. 2011; 68(12): 914-9.
17. Wieslander G and Norbäck D. A field study on clinical signs and symptoms in cleaners at floor polish removal and application in a Swedish hospital. *International Archives of Occupational and Environmental Health*. 2010; 83(5): 585-91.
18. Arif AA and Delclos GL. Association between cleaning-related chemicals and work-related asthma and asthma symptoms among healthcare professionals. *Occup Environ Med*. 2012; 69(1): 35-40.
19. Zock JP, Kogevinas M, Sunyer J, Almar E, Muniozgueren N, Payo F, et al. Asthma risk, cleaning activities and use of specific cleaning products among Spanish indoor cleaners. *Scand J Work Environ Health*. 2001; 27(1): 76-81.

20. Allpura. Manuel de formation "La technique du nettoyage". edn., Verlag USTER-Info GmbH, 2009.
21. Lessmann H, Uter W, Schnuch A and Geier J. Skin sensitizing properties of the ethanolamines mono-, di-, and triethanolamine. Data analysis of a multicentre surveillance network (IVDK\*) and review of the literature. Contact Dermatitis. 2009; 60(5): 243-55.
22. Purohit A, Kopferschmitt-Kubler MC, Moreau C, Popin E, Blaumeiser M and Pauli G. Quaternary ammonium compounds and occupational asthma. International Archives of Occupational and Environmental Health. 2000; 73(6): 423-7.

## “Hazardous substances in frequently used professional cleaning products”

<sup>1</sup>Fabian Melchior Gerster, <sup>1</sup>David Vernez, <sup>1,2</sup>Pascal Pierre Wild, <sup>1</sup>Nancy Brenna Hopf

<sup>1</sup>IST, Institute for Work and Health, Institut universitaire romand de Santé au Travail, Route de la Corniche 2,  
University of Lausanne and University of Geneva, Switzerland

<sup>2</sup>INRS, National Institute for Research and Security, Institut national de recherche et de sécurité, Department of  
Epidemiology, Vandoeuvre-lès-Nancy, France

Fabian Melchior Gerster:            Fabian.Gerster@chuv.ch

David Vernez:                         David.Vernez@hospvd.ch

Pascal Pierre Wild:                 Pascal.Wild@inrs.fr

Nancy Brenna Hopf:                 Nancy.Hopf@hospvd.ch

Corresponding author:

Fabian M. Gerster

Institute for Work and Health (IST)

Route de la Corniche 2

CH-1066 Epalinges – Lausanne Switzerland

Tel: +41 (0) 21 314 3782

Fax: +41 (0) 21 314 74 30

Email: Fabian.gerster@chuv.ch