



# The year in cardiology 2014: peripheral circulation

**Victor Aboyans<sup>1,2\*</sup>, Marianne Brodmann<sup>3</sup>, Marco De Carlo<sup>4</sup>, Denis Clement<sup>5</sup>,  
Lucia Mazzolai<sup>6</sup>, Lucas van Bortel<sup>7</sup>, and Marc R.H. van Sambeek<sup>8</sup>  
Charalambos Vlachopoulos<sup>9</sup>, On Behalf the ESC Working Group of  
Peripheral Circulation**

<sup>1</sup>Department of Cardiology, Dupuytren University Hospital, 2, Martin Luther King Avenue, Limoges 87042, France; <sup>2</sup>Inserm 1094, Tropical Neuroepidemiology, University of Limoges, Limoges, France; <sup>3</sup>Division of Angiology, Medical University Graz, Graz, Austria; <sup>4</sup>Cardiac Catheterization Laboratory, Cardiothoracic and Vascular Department, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy; <sup>5</sup>Department of the Dean, Gent University Hospital, Gent, Belgium; <sup>6</sup>Division of Angiology, CHUV, Lausanne University Hospital, Lausanne, Switzerland; <sup>7</sup>Unit of Clinical Pharmacology, Heymans Institute of Pharmacology, Gent University, Gent, Belgium; <sup>8</sup>Department of Vascular Surgery, Catharina Hospital, Eindhoven, The Netherlands; and <sup>9</sup>Peripheral Vessels and Hypertension Units, 1st Department Cardiology, Athens Medical School, Athens, Greece

Received 23 November 2014; revised 10 December 2014; accepted 17 December 2014; online publish-ahead-of-print 3 January 2015

## Abstract

In 2014, the debate on the indication of revascularization in case of asymptomatic carotid disease continued, while another one regarding the use of surgery vs. stenting addressed some new issues regarding the long-term cardiac risk of these patients. Renal arteries interventions trials were disappointing, as neither renal denervation nor renal artery stenting was found associated with better blood pressure management or outcome. In contrast, in lower-extremities artery disease, the endovascular techniques represent in 2014 major alternatives to surgery, even in distal arteries, with new insights regarding the interest of drug-eluting balloons. Regarding the aorta, the ESC published its first guidelines document on the entire vessel, emphasizing on the role of every cardiologist for screening abdominal aorta aneurysm during echocardiography. Among vascular wall biomarkers, the aorta stiffness is of increasing interest with new data and meta-analysis confirming its ability to stratify risk, whereas carotid intima-media thickness showed poor performances in terms of reclassifying patients into risk categories beyond risk scores. Regarding the veins, new data suggest the interest of D-dimers and residual venous thrombosis to help the decision of anti-coagulation prolongation or discontinuation after the initial period of treatment for deep vein thrombosis.

## Carotid disease

In a systematic review and meta-analysis (56 studies) for the U.S. Preventive Services Task Force,<sup>1</sup> the authors concluded a lack of sufficient evidence to establish incremental benefit of carotid endarterectomy (CEA), carotid stenting (CAS), or medical therapy intensification beyond current standard medical therapy, leading to a recommendation *against* screening for asymptomatic carotid

artery stenosis in general population.<sup>2</sup> In line with this, a recent analysis of an ongoing trial<sup>3</sup> comparing medical therapy alone vs. CEA vs. CAS in asymptomatic carotid stenosis looked for the clinical significance of progression of asymptomatic carotid stenosis. No statistically significant difference in annual stroke rates can be found between the progression- and non-progression groups, underlining the fact that the stenosis degree is not the sole factor of incident stroke. Further criteria are matter of active research, including ultrasound assessment of microembolic signals or cerebral blood flow reserve, identification of high-risk plaque through ultrasound, MRI or positron emission tomography, and the detection of asymptomatic brain damage.<sup>3</sup> Against the general trend showing safety in case of asymptomatic carotid stenosis treated with current standards of medical therapy, a recent series reported high rates of early ischaemic neurologic events after ultrasound diagnosis, especially in case of >90% stenosis, in favour of intervention.<sup>4</sup> This study suggests that there may be still some place for discussion regarding revascularization of asymptomatic carotid stenosis, mainly because there might be a subgroup of asymptomatic patients with a higher stroke risk, which must be identified by future research.

The preferred strategy to cure carotid stenosis (either asymptomatic or symptomatic) is also still debated. Carotid stenting has become increasingly popular as an alternative to CEA, with less invasiveness, more comfort for the patient and shorter recovery period. Long-term safety and clinical durability of CAS compared with CEA have been established by numerous randomized clinical trials (RCTs), although they differ regarding their reported peri-operative safety. In addition, significant differences in the requested operators' experience were present. A recent publication of the Carotid Stenting Trialists Collaboration highlighted the significant effect of operator's experience on outcome.<sup>5</sup> In 2014, the Cochrane Stroke Group updated their review and concluded that CAS was associated with a

The opinions expressed in this article are not necessarily those of the Editors of the *European Heart Journal* or of the European Society of Cardiology.

\* Corresponding author. Tel: +33 555 056 310, Fax: +33 555 056 384, Email: [vaboyans@live.fr](mailto:vaboyans@live.fr)

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2015. For permissions please email: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

higher risk than CEA for death and any stroke within 30 days of treatment. This statement was supported by the results from a National Hospital Discharge Database.<sup>6</sup> Large registries have obtained data (only from expert centres of excellence) on equivalence, or the non-inferiority, of CAS over CEA.<sup>7</sup> The long-term (10 years) results of a single-centre study comparing CAS with CEA with equivalent risk of death (over 50%) and similar risk of ipsilateral stroke for both procedures has been recently reported.<sup>8</sup>

Based on the most updated literature, the 2014 AHA/ASA Guidelines<sup>9</sup> recommended CAS as an alternative to CEA for symptomatic patients at average or low risk for complications, quite similar to the ESC guidelines in 2011.<sup>10</sup> For patients >70 years, CEA may be associated with improved outcome compared with CAS. Importantly, the trials supporting these guidelines were mainly focused on mortality and stroke. The trial providing 10 years results reported higher long-term risk of myocardial infarction (MI) in case of symptomatic carotid stenosis, especially among patients who underwent CEA, in accordance with several earlier reports showing increased risk of perioperative MI. It is plausible that the coronary risk should also be entered in the equation for selecting appropriate carotid revascularization modality, but this requires further studies.<sup>8</sup>

## Renal arteries

The CORAL trial (*Table 1*), the largest ever performed to assess the benefits of angioplasty in case of renovascular disease put a definite full stop on the extended use of this technique to hypertension and/or limit renal failure.<sup>16</sup> A meta-analysis adding these new data to those of other six trials (a total of 2139 patients with atherosclerotic renal stenosis) confirmed the lack of improvement in any outcome.<sup>18</sup> At this point, even the analysis of subgroups (e.g. renal function class, hypertension severity, stenosis degree, intra-arterial pressure recovery, etc.) have not identified any benefit of renal angioplasty over medical treatment in any setting. Nonetheless, these data are related to atherosclerotic disease, and renal angioplasty should be discussed in case of fibromuscular dysplasia.

The even more disappointing data came from the multicentre randomized single-blind Simplicity-3 trial (*Table 1*), comparing renal denervation with a sham procedure, conducted to validate the prior promising results in favour of major benefits to control resistant hypertension (i.e. uncontrolled hypertension with systolic blood pressure  $\geq 160$  mmHg despite taking  $\geq 3$  anti-hypertensive drugs at maximally tolerated doses).<sup>17</sup> The lack of efficacy, both in office and ambulatory pressure change, halted the big momentum of ongoing (or planned) trials. New technology, pathophysiology insights, and a better selection of patients are potential keys for future success.

## Lower-extremities artery disease

The Global Burden Disease estimated increased LEAD-related death between 1990 and 2010 (*Figure 1*).<sup>19</sup> In the USA, the incidence of CLI was estimated at 0.35% per year but one half of patients declared CLI without previous diagnosis of LEAD.<sup>20</sup>

Two studies identified elevated cardiac biomarkers as a tool for risk stratification in patients with LEAD, as they were tightly associated with worse prognosis. In patients undergoing endovascular

revascularization, pre-procedural cardiac troponin-T (cTnT) levels >0.01 ng/mL were independently associated with 1-year all-cause mortality (HR = 8.1) and amputation (HR = 3.7).<sup>21</sup> In claudicants, high cTnT was associated with increased risk of 7-year all-cause and cardiovascular (CV) death.<sup>22</sup> The clinical relevance of the use of cTnT for risk stratification in LEAD needs further investigation.

The benefits of statins in preventing peripheral outcomes in LEAD patients (–18% at 4 years) were highlighted in the REACH registry.<sup>23</sup>

A recent meta-analysis confirmed the benefits of exercise training on walking distances.<sup>24</sup> Current research is focused on home-based exercise, with favourable effects in terms of walking distance and vascular function.<sup>25,26</sup>

In the IRONIC trial (158 patients), quality of life improved in case of revascularization (vs. medical therapy) despite trivial amelioration of total walking distance, reopening the debate of revascularization in claudicants.<sup>13</sup>

Initial studies comparing drug-eluting balloons (DEBs) with percutaneous transluminal angioplasty (PTA) suggested that DEB may reduce restenosis and reintervention rates and improve wound healing/limb preservation in CLI patients. Two RCTs on DEB have been published in this setting in 2014.<sup>11,12</sup> The LEVANT-I trial randomized 101 Rutherford classes 2–5 femoro-popliteal lesions to Lutonix DEB or PTA, after stratification by intention to stent the lesion.<sup>12</sup> At 6 months, DEB yielded lower late lumen loss (LLL) (*Table 1*). The IN.PACT-DEEP randomized 358 patients with CLI 2 : 1 to DEB or PTA. Despite significantly longer lesions and deeper ulcers in the PTA arm, unexpectedly neither the clinically driven target lesion revascularization (TLR) nor the LLL were significantly better in the DEB arm at 1 year (*Table 1*).<sup>11</sup> While the study met its non-inferiority hypothesis regarding the primary safety endpoint, there was a trend towards increased major amputations (8.8 vs. 3.6%;  $P = 0.080$ ) in the DEB arm. These results should be interpreted as device-specific and should not preclude further research, which should be systematically associated with standardized wound care.

Most recently, the IN.PACT SFA I trial was presented; 331 patients with femoro-popliteal lesions in Rutherford classes 2–4 were randomized 2 : 1 to DEB or PTA.<sup>14</sup> At 12 months, the DEB group showed better primary patency (*Table 1*), clinically driven TLR (2.4 vs. 20.6%), and MACE rate (6.3 vs. 24.3%) ( $P < 0.001$  for all), setting DEB as a primary therapy for atherosclerotic lesions of the superficial femoral artery. This is the first trial showing superiority of DEB over PTA based on clinical endpoints. Balloon materials and coating technologies affect substantially the ability of DEB to deliver therapeutic paclitaxel regimens into the vessel wall. This, along with patients and lesions selection may explain the variability of results obtained, requiring therefore further trials to refine the best techniques and indications.

## Abdominal aorta aneurysm

The ESC guidelines on the management of the whole aorta were published in 2014.<sup>27</sup> They emphasized on the opportunistic screening of abdominal aorta aneurysm (AAA) during echocardiography in men >65 years (Class IIa, Level B), based on a French nationwide study reporting a 5.4% prevalence, at the median cost of 1.7 min to screen AAA.<sup>28</sup> These guidelines emphasized on the high level of CV risk in these patients and the importance of prevention and


**Table 1** Summary of major randomized trials or meta-analyses in peripheral intervention in 2014

Study	Aim	Type	Challenger	Reference	No. of participants: total (challenger/comparator)	Setting (indication)	Primary endpoint	Main objective achieved?
Carotid arteries								
Meta-analysis for the US Preventive Services Task Force <sup>1</sup>	To assess benefits of screening and treating CS (asymptomatic)	Meta-analysis	Screening + revasc. if needed	No screening	78 Reports/56 studies	Population	Stroke or TIA	No sufficient evidence
Lower-extremities arteries								
IN.PACT-DEEP <sup>11</sup>	To compare DEB vs. PTA for BTK lesions	Open	DEB	PTA	358 (239/119)	CLI	Efficacy: clinically driven TLR and LLL at 1 year; safety: composite of death, major amputation, and TLR	Efficacy No: TLR 9.2 vs. 13.1% ( $P = 0.29$ ); LLL $0.61 \pm 0.78$ mm vs. $0.62 \pm 0.78$ mm ( $P = 0.95$ ); safety yes: 17.7 vs. 15.8% ( $P = 0.02^a$ )
LEVANT-I <sup>12</sup>	Compare DEB vs. PTA in fem-pop lesions	Open	DEB ( $\pm$ stenting)	PTA ( $\pm$ stenting)	101 (49/52)	Rutherford classes 2–5 fem-pop lesions	Late lumen loss at 6 months	Yes: 58% lower LLL ( $P = 0.016$ )
IRONIC <sup>13</sup>	Benefits of revascularization on quality of life	Open	Revascularization (surgical or endovascular)	Non-supervised training advice	158 (79/79)	Unselected patients with claudication	Health-related quality of life after 1 year	Yes: improved quality of life ( $P < 0.01$ )
IN.PACT SFA I <sup>14</sup>	DEB vs. PTA for fem-pop lesions	Open	DEB	PTA	331 (220/111)	Rutherford classes 2–4 fem-pop lesions	Efficacy: primary patency at 1 year; safety: freedom MAE <sup>b</sup>	Efficacy yes: primary patency 82.2 vs. 52.4% ( $P < 0.001$ ); safety yes: MAE <sup>b</sup> 6.3% vs. 24.3% ( $P < 0.001$ )
Abdominal aorta								
IMPROVE <sup>15</sup>	EVAR vs. open surgery in ruptured AAA	Open	EVAR	Open surgery	613 (316/297)	Patients referred for ruptured AAA	30-day mortality	No: 30-day death EVAR: 35.4 vs. 37.4% (surgery), $P = 0.62$
Renal arteries								
CORAL <sup>16</sup>	Benefits of renal artery stenting in renovascular HTN	Open	Renal artery stenting	Medical therapy	933 (459/472)	Renal artery stenosis with either HTN or CKD	Death, nonfatal MI, stroke, renal function, and hosp. for CHF.	No: HR = 0.94 ( $P = 0.58$ )
SIMPLICITY-3 <sup>17</sup>	Benefits of renal denervation in resistant HTN	blinded	Denervation (+medical therapy)	Sham (+medical therapy)	535 (364/171)	Resistant HTN ( $\geq 160$ mmHg with $\geq 3$ drugs)	6-months office SBP change	No: $-2.39$ mmHg difference ( $P = 0.25$ )

AAA, abdominal aorta aneurysm; CS, carotid artery stenosis; DEB, drug-eluting balloon; EVAR, endovascular aortic repair; Fem-pop, femoro-popliteal; HTN, hypertension; LLL, late lumen loss; PTA, percutaneous transluminal angioplasty; SBP, systolic blood pressure; TIA, transient ischaemic attack.

<sup>a</sup>For non-inferiority.

<sup>b</sup>Combining death, clinically driven TVR, major amputation, and thrombosis.



**Figure 1** Mean death rates (per 100 000) related to lower-extremities artery disease in the 21 global regions, 1990 and 2010. The Global Burden Disease project. Adapted from Sampson et al.<sup>19</sup>

medical therapy beyond aortic interventions.<sup>27</sup> A *posthoc* analysis of a Dutch randomized trial comparing surgery vs. endovascular aortic repair (EVAR) for AAA showed a significant reduction of long-term total- (by 50%) and CV mortality (by 60%) in case of preoperative statins,<sup>29</sup> in line with data from a large national database (>20 000 interventions), which showed that >50% of these patients were regrettably not under statins.<sup>30</sup> In turn, anti-coagulation, often necessary for cardiac indications in patients undergoing EVAR, could increase the risk of endoleaks<sup>31,32</sup> and reintervention,<sup>31</sup> suggesting closer surveillance.

In most centres, patients with AAAs are usually treated endovascular (EVAR), if the anatomy is suitable. The ESC guidelines put at a same level of performance open surgery and endovascular therapy (EVAR).<sup>27</sup> Clinical evidence based on several RCTs indicates that EVAR is associated with superior peri-operative outcomes and similar long-term outcomes if compared with open surgery, although requiring more often re-intervention. Nevertheless, the mortality rates reported in these trials is lower than those in registries, outside the restrictions of RCTs.<sup>33</sup> Since these RCTs, stent-graft technology has been further developed by profile down-sizing, fixation, and sealing optimization, the use of low porosity fabrics, improved imaging techniques and better intervention planning. For these reasons, the ESC Guidelines recommended that the indication for EVAR be decided on an individual basis, according to anatomy, pathology, comorbidity, and anticipated durability.<sup>27</sup>


In the past few years, fenestrated and branched stent-graft systems have increasingly been used to treat anatomically challenging aneurysms. At the same time, chimney techniques have been used to extend landing zones for EVAR. The results of both techniques demonstrate that the procedure can be implemented with a high degree

of success. On the other hand, there is concern about the major adverse events, including side-branch patency and endoleak.<sup>34,35</sup> Ongoing studies and technological refinement of stent grafts will hopefully continue to broaden the utilization of EVAR.

There is clinical equipoise between open surgical repair and EVAR for patients with a ruptured AAA (rAAA). A systematic review and meta-analysis and an RCT comparing treatment strategy showed that EVAR is not inferior to open repair in patients with an rAAA. This supports the use of EVAR in suitable patients and OR as a reasonable alternative.<sup>15,36</sup>

## Vascular wall biomarkers

Peripheral vascular wall biomarkers show great potential in improving risk prediction beyond scores based on risk factors. An abnormal value of a biomarker would imply closer follow-up and earlier or even intensified preventive therapies. In a recent meta-analysis of 17 635 individual data, a 1 SD increase in carotid-femoral pulse wave velocity (cfPWV) increases CV events, CV mortality and total mortality by 30, 28, and 17%, respectively.<sup>37</sup> Most importantly, this arterial stiffness biomarker fulfils a stringent criterion for qualification as surrogate clinical endpoint: it improves overall 10-year risk classification for intermediate risk subjects by 13%.<sup>37</sup> In patients with erectile dysfunction, a prognostic marker of generalized arterial disease and CV events, cfPWV improves risk prediction and reclassifies 28% of patients to higher or lower risk category.<sup>38</sup> In contrast, cIMT showed a mediocre improvement (5.6%) in net reclassification index in intermediate risk patients with hypertension, currently tempering the expectations for this biomarker.<sup>39</sup> In an attempt to “map” arterial stiffness along the arterial tree, the Hoorn



**Figure 2** Tridimensional bar graphs representing amplification (peripheral—central systolic blood pressures represented by medians) according to sex (females up, males bottom), age, and blood pressure categories. Some categories are not represented because there were <50 observations.<sup>41</sup>

study<sup>40</sup> showed that local carotid and femoral arterial stiffness indices predicted, independently from each other and from cPWV, all-cause mortality and CV events. Central (aortic, carotid) pressures are lower than peripheral (brachial) pressures and they may have a better predictive ability. Difference between central and peripheral pressures

is higher in younger ages, in males, and with increasing levels of blood glucose, while smoking and dyslipidaemia decrease this difference.<sup>41</sup> In an important step towards clinical implementation, reference values for central pressures were determined in a general healthy population and according to CV risk factors (Figure 2).<sup>41</sup>



21. de Lemos JA, Kumbhani DJ. Lessons from the heart: troponin elevations in patients with established peripheral artery disease. *J Am Coll Cardiol* 2014;**63**:1539–1541.
22. Pohlhammer J, Kronenberg F, Rantner B, Stadler M, Peric S, Hammerer-Lercher A, Klein-Weigel P, Fraedrich G, Kollerits B. High-sensitivity cardiac troponin T in patients with intermittent claudication and its relation with cardiovascular events and all-cause mortality – the CAVASIC study. *Atherosclerosis* 2014;**237**:711–717.
23. Kumbhani DJ, Cannon CP, Eagle KA, Smith SC Jr, Goto S, Ohman EM, Elbez Y, Sritara P, Baumgartner I, Banerjee S, Creager MA, Bhatt DL, REACH Registry Investigators. Statin therapy and long-term adverse limb outcomes in patients with peripheral artery disease: insights from the REACH registry. *Eur Heart J* 2014;**35**:2864–2872.
24. Parmenter BJ, Dieberg G, Smart NA. Exercise training for management of peripheral arterial disease: a systematic review and meta-analysis. *Sports Med* 2014 (in press).
25. McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Zhao L, Liu K, Domanchuk K, Spring B, Tian L, Kibbe M, Liao Y, Lloyd Jones D, Rejeski WJ. Home-based walking exercise in peripheral artery disease: 12-month follow-up of the goals randomized trial. *J Am Heart Assoc* 2014;**3**:e000711.
26. Gardner AW, Parker DE, Montgomery PS, Blevins SM. Step-monitored home exercise improves ambulation, vascular function, and inflammation in symptomatic patients with peripheral artery disease: a randomized controlled trial. *J Am Heart Assoc* 2014;**3**:e001107.
27. Erbel R, Aboyans V, Boileau C, Bossone E, Bartolomeo RD, Eggebrecht H, Evangelista A, Falk V, Frank H, Gaemperli O, Grabenwöger M, Haverich A, Lung B, Manolis AJ, Meijboom F, Nienaber CA, Roffi M, Rousseau H, Sechtem U, Sirnes PA, Allmen RS, Vrints CJ; ESC Committee for Practice Guidelines (CPG), Document Reviewers; Authors/Task Force Members. 2014 ESC Guidelines on the diagnosis and treatment of aortic diseases: document covering acute and chronic aortic diseases of the thoracic and abdominal aorta of the adult. The Task Force for the Diagnosis and Treatment of Aortic Diseases of the European Society of Cardiology (ESC). *Eur Heart J* 2014;**35**:2873–2926.
28. Aboyans V, Bataille V, Bliscaux P, Ederhy S, Filliol D, Honton B, Kurtz B, Messas E, Mohty D, Brochet E, Kownator S; investigators of the E2T3A study. Effectiveness of screening for abdominal aortic aneurysm during echocardiography. *Am J Cardiol* 2014;**114**:1100–1104.
29. de Bruin JL, Baas AF, Heymans MW, Buimer MG, Prinssen M, Grobbee DE, Blankensteijn JD; DREAM Study Group. Statin therapy is associated with improved survival after endovascular and open aneurysm repair. *J Vasc Surg* 2014;**59**:39–44.
30. Galiñanes EL, Reynolds S, Dombrovskiy VY, Vogel TR. The impact of preoperative statin therapy on open and endovascular abdominal aortic aneurysm repair outcomes. *Vascular* 2014. (online).
31. De Rango P, Verzini F, Parlani G, Cieri E, Simonte G, Farchioni L, Isernia G, Cao P. Safety of chronic anticoagulation therapy after endovascular abdominal aneurysm repair (EVAR). *Eur J Vasc Endovasc Surg* 2014;**47**:296–303.
32. Lazarides MK, Georgiadis GS, Charalampidis DG, Antoniou GA, Georgarakos EI, Trellopoulos G. Impact of long-term warfarin treatment on EVAR durability: a meta-analysis. *J Endovasc Ther* 2014;**21**:148–153.
33. Malas M, Arhuidese I, Qazi U, Black J, Perler B, Freischlag JA. Perioperative mortality following repair of abdominal aortic aneurysms: application of a randomized clinical trial to real-world practice using a validated nationwide data set. *JAMA Surg* 2014;**149**:1260–1265.
34. Raux M, Patel VI, Cochenne F, Mukhopadhyay S, Desgranges P, Cambria RP, Becquemini JP, LaMuraglia GM. A propensity-matched comparison of outcomes for fenestrated endovascular aneurysm repair and open surgical repair of complex abdominal aortic aneurysms. *J Vasc Surg* 2014;**60**:858–863.
35. Scali ST, Feezor RJ, Chang CK, Waterman AL, Berceli SA, Huber TS, Beck AW. Critical analysis of results after chimney endovascular aortic aneurysm repair raises cause for concern. *J Vasc Surg* 2014;**60**:865–873.
36. van Beek SC, Conijn AP, Koelemay MJ, Balm R. Editor's Choice – endovascular aneurysm repair versus open repair for patients with a ruptured abdominal aortic aneurysm: a systematic review and meta-analysis of short-term survival. *Eur J Vasc Endovasc Surg* 2014;**47**:593–602.
37. Ben-Shlomo Y, Spears M, Boustred C, May M, Anderson SG, Benjamin EJ, Boutouyrie P, Cameron J, Chen CH, Cruickshank JK, Hwang SJ, Lakatta EG, Laurent S, Maldonado J, Mitchell GF, Najjar SS, Newman AB, Ohishi M, Pannier B, Pereira T, Vasan RS, Shokawa T, Sutton-Tyrell K, Verbeke F, Wang KL, Webb DJ, Willum Hansen T, Zoungas S, McEniery CM, Cockcroft JR, Wilkinson IB. Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. *J Am Coll Cardiol* 2014;**63**:636–646.
38. Vlachopoulos C, Ioakeimidis N, Aznaouridis K, Terentes-Printzios D, Rokkas K, Aggelis A, Panagiotakos D, Stefanadis C. Prediction of cardiovascular events with aortic stiffness in patients with erectile dysfunction. *Hypertension* 2014;**64**:672–678.
39. Bots ML, Groenewegen KA, Anderson TJ, Britton AR, Dekker JM, Engström G, Evans GW, de Graaf J, Grobbee DE, Hedblad B, Hofman A, Holewijn S, Ikeda A, Kavousi M, Kitagawa K, Kitamura A, Ikram MA, Lonn EM, Lorenz MW, Mathiesen EB, Nijpels G, Okazaki S, O'Leary DH, Polak JF, Price JF, Robertson C, Rembold CM, Rosvall M, Rundek T, Salonen JT, Sitzer M, Stehouwer CD, Franco OH, Peters SA, den Ruijter HM. Common carotid intima-media thickness measurements do not improve cardiovascular risk prediction in individuals with elevated blood pressure: the USE-IMT collaboration. *Hypertension* 2014;**63**:1173–1181.
40. van Sloten TT, Schram MT, van den Hurk K, Dekker JM, Nijpels G, Henry RM, Stehouwer CD. Local stiffness of the carotid and femoral artery is associated with incident cardiovascular events and all-cause mortality: the Hoorn study. *J Am Coll Cardiol* 2014;**63**:1739–1747.
41. Herbert A, Cruickshank JK, Laurent S, Boutouyrie P; on behalf of The Reference Values for Arterial Measurements Collaboration, on behalf of The Reference Values for Arterial Measurements Collaboration. Establishing reference values for central blood pressure and its amplification in a general healthy population and according to cardiovascular risk factors. *Eur Heart J* 2014;**35**:3122–3133.
42. ISTH Steering Committee for World Thrombosis Day. Thrombosis: a major contributor to the global disease burden. *J Thromb Haemost* 2014;**12**:1580–1590.
43. Simes J, Becattini C, Agnelli G, Eikelboom JW, Kirby AC, Mister R, Prandoni P, Brighton TA; INSPIRE Study Investigators. Aspirin for the prevention of recurrent venous thromboembolism: The INSPIRE Collaboration. *Circulation* 2014;**130**:1062–1071.
44. Baber U, Mastoris I, Mehran R. Balancing ischaemia and bleeding risks with novel oral anticoagulants. *Nat Rev Cardiol* 2014;**11**:693–703.
45. Palareti G, Cosmi B, Legnani C, Antonucci E, De Micheli V, Ghirarduzzi A, Poli D, Testa S, Tassetto A, Pengo V, Prandoni P; DULCIS (D-dimer and Ultrasonography in Combination Italian Study) Investigators. D-dimer to guide the duration of anticoagulation in patients with venous thromboembolism: a management study. *Blood* 2014;**124**:196–203.
46. Napolitano M, Saccullo G, Malato A, Sprini D, Ageno W, Imberti D, Mascheroni D, Bucherini E, Gallucci P, D'Alessio A, Prantera T, Spadaro P, Rotondo S, Di Micco P, Oriana V, Urbano O, Recchia F, Ghirarduzzi A, Lo Coco L, Mancuso S, Casuccio A, Rini GB, Siragusa S. Optimal duration of low molecular weight heparin for the treatment of cancer-related deep vein thrombosis: the Cancer-DACUS Study. *J Clin Oncol* 2014;**32**:3607–3612.
47. Watson L, Broderick C, Armon MP. Thrombolysis for acute deep vein thrombosis. *Cochrane Database Syst Rev* 2014;**1**:CD002783.