

# ESMO Minimum Clinical Recommendations for diagnosis, treatment and follow-up of soft tissue sarcomas

## Incidence

- The crude incidence of soft tissue sarcoma in the European Union is 1.0–3.0/100 000 per year, the mortality 0.6–0.8/100 000 per year. They are diagnosed at any age but are more frequent in older patients with a peak incidence at the age of 50 years.

## Diagnosis

- Histological diagnosis and evaluation of grades are preferentially made on an incisional surgical biopsy or on the completely resected tumor. Tru-cut biopsies are also used but should be restricted to experienced centers.
- Gastrointestinal stromal tumors should be confirmed by staining for CD117.
- Specific types of small round cell tumors (extra-osseous Ewing's sarcoma, embryonal rhabdomyosarcoma) should be identified by immunohistochemistry and cytogenetics and be treated accordingly.

## Staging and risk assessment

- Staging is performed by physical exam and the appropriate radiological techniques. To exclude lung metastases a CT scan is recommended in operable patients.
- For staging according to the UICC/AJCC 2002 system, tumor size is categorized as small ( $\leq 5$  cm, T1) or large (T2) and is complemented by information about location (superficial: Ta or deep: Tb) and histological grade (using either of the grading systems G1-4 or G1-3 or low/high) as further parameters that are required for the stage grouping are shown in Table 1.

## Treatment plan

- Soft tissue sarcoma requires a multidisciplinary approach by an experienced team.

## Surgery

- Surgery is the main treatment for localized disease. The tumor should be removed by wide excision or by compartmental resection including the cutaneous scar and tractus of a previous biopsy. The width of the resection might be decreased in case of resistant anatomic planes such as muscular fasciae, periosteum, perineurium, if not infiltrated.
- After wide excision of high-grade sarcomas, adjuvant radiation therapy is recommended [III, B].

- In case of radical surgery obtained by compartmental excision or amputation at a large distance from the primary tumor, adjuvant radiation therapy is not necessary.
- Re-operation is recommended in case of previous marginal or intralésional resection [III, B].
- In case of completely resectable lung metastases, surgery has to be considered [III, B].

## Radiation therapy

- Radiation therapy should be administered postoperatively at a dose of 60–65 Gy with a shrinking field technique in case of wide resection [II, B].
- In selected patients preoperative radiotherapy may be an option [III, B].

## Chemotherapy

### Localized disease:

- Preoperative chemotherapy is not standard practice for operable patients [III, B]. It can be considered together with radiotherapy in patients with borderline resectable tumors.
- Adjuvant chemotherapy is not standard practice even though it might improve distant and local control. Its impact on overall survival is still debated [II, A]. It may be considered in younger patients with large and high-grade tumors [II, C].
- In non-resectable tumors confined to an extremity, chemotherapy with or without radiotherapy or isolated hyperthermic limb perfusion with chemotherapy and/or cytokines offers an alternative to amputation [III, B].

### Metastatic disease:

- Chemotherapy is the standard treatment for metastatic disease. Doxorubicin with or without ifosfamide is commonly

**Table 1.**

Stage	Primary tumor	Lymph nodes	Distant metastases	Grading G <sub>1-4</sub> (or G <sub>1-3</sub> or low / high)
IA	T <sub>1a</sub> or b	N <sub>0</sub> or x	M <sub>0</sub>	G <sub>1-2</sub> (or G <sub>1</sub> or low)
IB	T <sub>2a</sub> or b	N <sub>0</sub> or x	M <sub>0</sub>	G <sub>1-2</sub> (or G <sub>1</sub> or low)
IIA	T <sub>1a</sub> or b	N <sub>0</sub> or x	M <sub>0</sub>	G <sub>3-4</sub> (or G <sub>2-3</sub> or high)
IIB	T <sub>2a</sub>	N <sub>0</sub> or x	M <sub>0</sub>	G <sub>3-4</sub> (or G <sub>2-3</sub> or high)
III	T <sub>2b</sub>	N <sub>0</sub> or x	M <sub>0</sub>	G <sub>3-4</sub> (or G <sub>2-3</sub> or high)
IV	Any T Any T	N <sub>1</sub> Any N	M <sub>0</sub> M <sub>1</sub>	Any G Any G

used. Doxorubicin alone seems to be equivalent to its combinations with other agents with regard to survival, despite a higher response rate with combination regimens [II, B].

- Imatinib (STI 571, a tyrosine kinase receptor inhibitor) is the treatment of choice for gastro-intestinal stromal tumors (GIST).

## Response evaluation

- Response evaluation of chemotherapy for metastatic disease should be performed after 2 or 3 cycles with the radiological exams that were positive before treatment.

## Follow-up

- Early detection of recurrence might influence the possibility of a curative therapy. The patients should be followed every 3 months with history and physical examination. MRI of the site of resection of the primary tumor is proposed twice a year for the first 2–3 years and then once a year. For patients with high-grade tumors a chest X-ray is recommended every 3 to 4 months in the first 2–3 years, twice a year up to the 5th year, and once a year thereafter [IV, B].

## Note

Levels of Evidence [I–V] and Grades of Recommendation [A–D] as used by the American Society of Clinical Oncology are given in square brackets. Statements without grading were considered justified standard clinical practice by the experts and the ESMO faculty.

## Literature

1. Wylie JP, O'Sullivan B, Catton C, Gutierrez E. Contemporary radiotherapy for soft tissue sarcoma. *Semin Surg Oncol* 1999; 17: 33–46.
2. Sarcoma meta-analysis collaboration. Adjuvant chemotherapy for localized resectable soft-tissue sarcoma: a meta-analysis of individual data. *Lancet* 1997; 350: 1647–1654.
3. Eggermont AMM, Schraffordt Koops H, Liénard D et al. Isolated limb perfusion with high-dose tumor necrosis factor- $\alpha$  in combination with interferon- $\gamma$  and melphalan for non resectable extremity soft tissue sarcomas: a multicentric trial. *J Clin Oncol* 1996; 14: 2653–2665.
4. O'Byrne K, Steward WP. The role of chemotherapy in the treatment of adult soft tissue sarcomas. *Oncology* 1999; 56: 13–23.
5. Frustaci S, Gherlinzoni F, De Paoli A et al. Adjuvant chemotherapy for adult soft tissue sarcomas of the extremities and girdles: results of the Italian Randomized Cooperative Trial. *J Clin Oncol* 2001; 19: 1238–1247.
6. Weiss SW. *Histological typing of soft tissue tumors*. Heidelberg: Springer-Verlag/World Health Organization 1994.
7. Van Unnik JA, Coindre JM, Contesso C et al. Grading of soft tissue sarcomas: experience of the EORTC soft tissue and bone sarcoma group. *Eur J Cancer* 1993; 29 A: 2089–2093.
8. Flugstad DL, Wilke CP, McNutt MA et al. Importance of surgical resection in the successful management of soft tissue sarcoma. *Arch Surg* 1999; 134: 856–861.
9. Lewis JJ, Leung D, Espat J et al. Effect of reresection in extremity soft tissue sarcoma. *Ann Surg* 2000; 231: 655–663.
10. Van Geel AN, Pastorino V, Jauch KW et al. Surgical treatment of lung metastases. The EORTC-STBSG study of 255 patients. *Cancer* 1996; 77: 675–682.

Coordinating authors for the ESMO Guidelines Task Force: S. Leyvraz<sup>1</sup> & S. Jelic<sup>2</sup>

<sup>1</sup>Invited author, Fondation du Centre Pluridisciplinaire d'Oncologie, CHUV, Lausanne, Switzerland; <sup>2</sup>Assigned task force member, Institute of Oncology and Radiology of Serbia, Department of Medical Oncology, Pasterova 14, 11000 Belgrade, Serbia

Approved by the ESMO Guidelines Task Force: August 2003, last update December 2004.

*Correspondence to:*  
ESMO Guidelines Task Force  
ESMO Head Office  
Via La Santa 7  
CH-6962 Lugano  
Switzerland