

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Permissivity of insect cells to *Waddlia chondrophila*, *Estrella lausannensis* and *Parachlamydia acanthamoebae*.

Authors: Kebbi-Beghdadi C, Fatton M, Greub G

Journal: *Microbes and infection*

Year: 2015 Nov-Dec

Volume: 17

Issue: 11-12

Pages: 749-54

DOI: [10.1016/j.micinf.2015.09.014](https://doi.org/10.1016/j.micinf.2015.09.014)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

1 **Permissivity of insect cells to *Waddlia chondrophila*, *Estrella lausannensis* and**
2 ***Parachlamydia acanthamoebae***

3 Carole Kebbi-Beghdadi, Morgane Fatton, Gilbert Greub*

4 Center for Research on Intracellular Bacteria (CRIB), Institute of Microbiology, University
5 Hospital Center and University of Lausanne, Lausanne, Switzerland

6

7 *Corresponding author:

8 Prof. Gilbert Greub

9 Institute of Microbiology

10 Rue du Bugnon 48

11 1011 Lausanne

12 Switzerland

13 Tel: 0041 21 314 4979

14 Fax 0041 21 314 4060

15 e-mail: gilbert.greub@chuv.ch

16

17

18

19 **Abstract**

20 Recent large scale studies questioning the presence of intracellular bacteria of the
21 *Chlamydiales* order in ticks and fleas revealed that arthropods, similarly to mammals,
22 reptiles, birds or fishes, can be colonized by *Chlamydia*-related bacteria with a
23 predominant representation of the *Rhabdochlamydiaceae* and *Parachlamydiaceae*
24 families. We thus investigated the permissivity of two insect cell lines towards *Waddlia*
25 *chondrophila*, *Estrella lausannensis* and *Parachlamydia acanthamoebae*, three bacteria
26 representative of three distinct families within the *Chlamydiales* order, all documented in
27 ticks and/or in other arthropods. We demonstrated that *W. chondrophila* and *E.*
28 *lausannensis* are able to very efficiently multiply in these insect cell lines. *E.*
29 *lausannensis* however induced a rapid cytopathic effect, which somehow restricted its
30 replication. *P. acanthamoebae* was not able to grow in these cell lines even if inclusions
31 containing a few replicating bacteria could occasionally be observed.

32

33 **Keywords:** Intracellular bacteria; *Chlamydia*-related bacteria; Cell permissivity; Host
34 range; Reservoir; Vector

35

36 1. Introduction

37 All bacteria belonging to the *Chlamydiales* order are strict intracellular organisms sharing
38 a biphasic life cycle that involves two distinct bacterial forms, an infectious Elementary
39 Body (EB) capable of attaching to and entering into its host cell, but that cannot replicate
40 before it differentiates into a metabolically active form, called Reticulate Body (RB). At
41 the end of the multiplication phase, RBs differentiate back into infectious particles and
42 lyse their host cells to start a new cycle [32]. *Chlamydiales* bacteria have been isolated
43 from samples of multiple origins such as mammals, birds, fishes, reptiles or protozoa
44 revealing the large and probably still largely underestimated diversity of their ecological
45 niches (reviewed in [22, 36]). Besides the well-studied human and animal pathogens
46 belonging to the *Chlamydiaceae* family (such as *Chlamydia pneumoniae*, *C. trachomatis*
47 or *C. abortus*), this clade also comprises 8 other family-level lineages whose members
48 are very diverse being either emerging pathogens able to grow in cells of various origins
49 (*Waddlia chondrophila*, *Simkania negevensis*) or harmless environmental species only
50 replicating in amoebae (*Protochlamydia amoebophila*, *Neochlamydia hartmanellae*).
51 Recent large scale studies questioning the presence of *Chlamydiales* bacteria in ticks
52 and fleas [13], [Pilloux et al. submitted 2015] revealed that arthropods may be colonized
53 by these bacteria and thus could serve as reservoir and vectors for potential novel
54 pathogens. In this context, we investigated the permissivity of two insect cell lines,
55 *Aedes albopictus* larva cells and *Spodoptera frugiperda* ovary cells (Sf9), towards
56 *Waddlia chondrophila*, *Estrella lausannensis* and *Parachlamydia acanthamoebae*, three
57 *Chlamydia*-related bacteria representative of three different families, all belonging to the

58 *Chlamydiales* order, documented in fleas and/or ticks and whose pathogenic potential
59 for humans and cattle is confirmed or highly suspected [2-6, 9, 15, 16, 23, 34, 37, 14].

60

61 **2. Materials and methods**

62 *2.1. Cell culture and bacterial strains*

63 *Aedes albopictus* clone C6/36 larva cells_(ATCC® CRL-1660™) and *Spodoptera*

64 *frugiperda* ovarian epithelial cells (Sf9) (ATCC® CRL-1711™) were routinely maintained

65 respectively at 28°C and 5%CO₂ in Dulbecco's modified essential medium (DMEM;

66 Gibco Invitrogen, Basel, Switzerland) supplemented with 10% foetal calf serum

67 (Biochrom, Berlin, Germany) or at 27°C in Grace insect medium (GIM; Gibco Invitrogen,

68 Basel, Switzerland) supplemented with 10% foetal calf serum.

69 *W. chondrophila* strain WSU 86-1044 (ATCC VR-1470), *E. lausannensis* strain CRIB 30

70 and *P. acanthamoebae* strain Hall's coccus were grown at 32°C within *Acanthamoeba*

71 *castellanii* strain ATCC 30010, as described in [19].

72

73 2.2. Infection procedure

74 Insect cells were seeded at 1×10^5 cells per well in 24-wells microplates (Corning) the
75 day before infection. Infection was performed at 28°C or 27°C, as described in [24], with
76 a 5 days-old culture of bacteria in *A. castellanii* diluted, if not otherwise described, 1:1000
77 for *W. chondrophila* (MOI: 1-20) and *P. acanthamoebae* (MOI: 0.1-1) and 1:2000 for *E.*
78 *lausannensis* (MOI: 1-10)

79

80 2.3. Immunofluorescence and confocal microscopy

81 At different time points after infection, immunofluorescence was performed on cells
82 cultivated on glass coverslips following precisely the protocol described in [27] except
83 that rabbit anti-*P. acanthamoebae* and mouse anti-*E. lausannensis* were used
84 respectively at a 1:1000 and 1:500 dilution. Secondary antibody was diluted 1:500 and
85 mixed with a 1/50 dilution of Concanavalin A (Molecular Probe) and 150 ng/ml DAPI
86 (Molecular Probe, Eugene, Oregon, USA). Cells were observed under an epifluorescent
87 microscope (Axioplan 2, Zeiss, Feldbach, Switzerland) or a confocal microscope
88 (AxioPlan 2 LSM 710, Zeiss).

89

90 2.4. Electron microscopy

91 One T25 flask of *Aedes albopictus* or Sf9 cells was infected as described above with *W.*
92 *chondrophila* and *E. lausannensis* diluted 1/200. 24 hours post-infection, cells were
93 washed once with PBS and harvested. After one centrifugation step of 10 min at 1'000
94 rpm, cells were resuspended in phosphate buffer (19ml of 0.2 M NaH_2PO_4 + 81ml of
95 0.2M Na_2HPO_4 , pH 7.4) containing 0.2% glutaraldehyde (Fluka Biochemika, Buchs,
96 Switzerland) and 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, USA)

97 and incubated overnight at 4°C. After two more washing steps with phosphate buffer,
98 cells were prepared as described previously [10]. Thin sections on grids were examined
99 with a transmission electron microscope Philips CM 100 (Philips, Eindhoven, The
100 Netherlands).

101

102 *2.5. Quantitative PCR*

103 Cells were harvested at different time points after infection, genomic DNA was extracted
104 following manufacturer's instructions (Wizard SV Genomic DNA purification kit,
105 Promega, Madison, WI) and qPCR was performed as described in [24]. To account for
106 variation in the inocula used for infection, bacterial growth is expressed as a fold
107 increase in the number of bacteria after normalization using time point "0h" as reference.

108

109 *2.6. Cell viability*

110 Cell viability was determined with a propidium iodide assay as described in [14] except
111 that propidium iodide was added immediately after infection. Positive control was
112 obtained by incubating non-infected cells 5 minutes with MetOH at
113 -20°C. Results were normalized considering the positive control as 100% mortality.

114

115 **3. Results**

116 *3.1. Growth kinetic in insect cells*

117 Two insect cell lines, *Aedes albopictus* larva cells and Sf9 (*S. frugiperda* ovarian
118 epithelial cells) were infected with either *W. chondrophila*, *E. lausannensis* or *P.*
119 *acanthamoebae* and bacterial growth was monitored using specific quantitative PCRs
120 (qPCRs) developed in our laboratory [11, 16, 30]. In parallel, bacterial multiplication was

121 also assessed by immunofluorescence and confocal microscopy using in-house
122 polyclonal antibodies and by electron microscopy.

123

124 3.1.1. *W. chondrophila* and *E. lausannensis*

125 Results shown in Fig. 1 indicated that *W. chondrophila* and *E. lausannensis* are both
126 able to efficiently replicate in insect cells, leading to an increase of bacterial genomic
127 DNA copies of about 3 logs in 48 hours for *W. chondrophila* (Fig. 1 panels a and b) and
128 of about 2 logs in 48 hours for *E. lausannensis* (Fig. 1 panels c and d). During the first 8
129 hours following infection, EBs enter their insect cell host, differentiate into RBs and
130 create a replicative niche able to support their exponential multiplication. Inclusions
131 containing dividing bacteria can already be observed by immunofluorescence and
132 confocal microscopy 24 hours post infection (Fig. 2A, panels a,b,c,and d). Similarly to
133 what has been described in other cell lines [24, 25], *W. chondrophila* then exponentially
134 replicate until 48 hours p.i. and ultimately lyse their host to release infectious particles
135 ready to infect new insect cells (data not shown). Replication of *W. chondrophila* and *E.*
136 *lausannensis* was also documented in Sf9 cells by electron microscopy 24 hours post
137 infection (Fig. 2B). The *W. chondrophila*-containing vacuoles were surrounded by tightly
138 associated mitochondria, a feature that was also described in other cell lines [12, 25].

139

140 3.1.2. *P. acanthamoebae*

141 No growth of *P. acanthamoebae* could be detected by qPCR or immunofluorescence in
142 the two insect cell lines tested (Fig.1 and 2A, panels e and f) even after 7 days.

143 However, when cells were infected with a 10 fold higher bacterial load (MOI of 1-10),

144 rare inclusions containing replicating bacteria could be observed by confocal microscopy

145 in *Aedes albopictus* cells (Fig. 2A, panel g). This bacterial replication is very limited and
146 restricted to about 3% of all infected cells.

147

148 3.2. Cytopathic effect

149 Direct examination by confocal microscopy of *E. lausannensis* infection revealed that
150 these bacteria, like *W. chondrophila*, efficiently multiplied in insect cells but these cells
151 were more rapidly lysed than those infected with *W. chondrophila*. This observed
152 cytopathic effect was confirmed in a host cell viability assay based on propidium iodide
153 incorporation that showed, 48 hours post infection, 100% mortality of *Aedes albopictus*
154 cells infected with *E. lausannensis* versus 50% in cells infected with *W. chondrophila*
155 and 23% in non-infected cells (Fig. 3). Results were normalized using methanol-treated
156 cells at time of infection as a positive control of 100% mortality. Mortality rate over 100%
157 are thus explained by an increase, at later time points, of the total number of cells that
158 ultimately died. We could observe a weak cytopathic effect of *P. acanthamoebae*
159 infection that probably resulted more from the lower fitness of infected cells than from
160 the few replicating bacteria.

161 In Sf9 cells, a similar cytopathic effect of *E. lausannensis* was observed by confocal
162 microscopy (data not shown), however, due to interferences of the culture medium with
163 the propidium iodide assay, this increased cell mortality could not be precisely
164 quantified.

165

166 4. Discussion

167 In the present study, we demonstrated that *W. chondrophila* and *E. lausannensis* are
168 able to enter and efficiently multiply in two different insect cell lines. *E. lausannensis*
169 growth is however slightly less efficient than *W. chondrophila* growth, a difference that
170 can be explained by the rapid cytopathic effect of *E. lausannensis*, which restricts the
171 number of replication rounds possible.

172 *W. chondrophila* is an emerging pathogen for humans and animals and it is able to
173 readily grow at temperatures ranging from 25°C to 37°C in mammalian cell lines of
174 diverse origins [12, 21, 25, 28], in fish cells [24] as well as in protozoa [17, 31]. *E.*
175 *lausannensis*, whose pathogenic potential is currently under investigation, was first
176 isolated from an environmental water sample and is also able to grow at different
177 temperatures in protozoa, fishes and mammalian cells [14, 24, 30]. Strikingly, these two
178 organisms that display such a broad host range, both encode an extended family of
179 OmpA proteins with beta-barrel structure that were recently shown to be dominant in the
180 outer membrane of *W. chondrophila* and to play a role in adhesion of this bacteria to its
181 host [1, 7, 8, 26, 29]. In addition, *Simkania negevensis*, another *Chlamydia*-related
182 bacteria that also possess a large family of 37 MOMP-like proteins in its outer
183 membrane similarly displays a wide host range [1, 27]. Based on these observations, we
184 recently hypothesized that the large diversity of the OmpA protein family is linked to the
185 wide host range of these bacteria [26]. In addition, ability to grow in multiple and diverse
186 eukaryotic cells correlates in most cases with virulence towards mammals [27].
187 *P. acanthamoebae* only encodes one homolog of MOMP and its efficient growth seems
188 to be restricted to amoebal hosts [18, 20]. Several studies have reported a limited
189 replication of this bacteria in host cells such as bone-marrow derived macrophages, fish
190 or insect cell lines [24, 33, 35]. This present work also demonstrated that growth of *P.*

191 *acanthamoebae* in insect cells is limited and restricted to a few cells. Sixt *et al.* also
192 reported a limited *P. acanthamoebae* replication in *Aedes albopictus*, Sf9 and
193 *Drosophila* S2 cells and they observed a bacteria-induced programmed cell death in the
194 last two cell lines [35]. We did not observe nuclear fragmentation in *P. acanthamoebae*-
195 infected cells and cell viability was stable during the first 72 hours post infection, which is
196 not in agreement with apoptosis. Furthermore, preliminary experiments performed in
197 *Aedes albopictus* cells in presence of a pan-caspase inhibitor did not demonstrate an
198 enhanced multiplication of *P. acanthamoebae* (A. Croxatto, unpublished).
199 In conclusion, we demonstrated the permissivity of insect cells to *Waddlia chondrophila*
200 and *Estrella lausannensis* and further highlighted the very broad host range of these
201 possible pathogens and the potential role of insects as reservoir or vectors for these
202 strict intracellular bacteria.

203

204 **5. Acknowledgments**

205 We thank S. Aeby, the Electron Microscopy Facility platform of the University of
206 Lausanne and the PFMU at the Medical Faculty of Geneva for assisting with electron
207 microscopy analysis. We also thank the Cellular Imaging Facility platform of the
208 University of Lausanne for confocal microscopy and P. Couleru and L. Pilloux for
209 technical help. This work was partially supported by a grant from the SNSF no:
210 310030_141050

211

212 **6. Conflict of interest**

213 The authors have no conflict of interest.

214

215 **6. References**

216 [1] Aistleitner K, Anrather D, Schott T, Klose J, Bright M, Ammerer G, et al. Conserved
217 features and major differences in the outer membrane protein composition of
218 chlamydiae. *Environ Microbiol* 2015;17:1397-413.

219 [2] Barkallah M, Fendri I, Dhieb A, Gharbi Y, Greub G, Gdoura R. First detection of
220 *Waddlia chondrophila* in Africa using SYBR Green real-time PCR on veterinary samples.
221 *Vet Microbiol* 2013;164:101-7.

222 [3] Barkallah M, Gharbi Y, Hassena AB, Slima AB, Mallek Z, Gautier M, et al. Survey of
223 infectious etiologies of bovine abortion during mid- to late gestation in dairy herds. *PLoS*
224 *One* 2014;9:e91549.

225 [4] Baud D, Goy G, Osterheld MC, Borel N, Vial Y, Pospischil A, et al. *Waddlia*
226 *chondrophila*: From Bovine Abortion to Human Miscarriage. *Clinical infectious diseases*
227 2011;52:1469-71.

228 [5] Baud D, Goy G, Osterheld MC, Croxatto A, Borel N, Vial Y, et al. Role of *Waddlia*
229 *chondrophila* placental infection in miscarriage. *Emerg Infect Dis* 2014;20:460-4.

230 [6] Baud D, Thomas V, Arafa A, Regan L, Greub G. *Waddlia chondrophila*, a potential
231 agent of human fetal death. *Emerg Infect Dis* 2007;13:1239-43.

232 [7] Bertelli C, Aeby S, Chassot B, Clulow J, Hilfiker O, Rappo S, et al. Sequencing and
233 characterizing the genome of *Estrella lausannensis* as an undergraduate project:
234 training students and biological insights. *Front Microbiol* 2015;6:101.

235 [8] Bertelli C, Collyn F, Croxatto A, Ruckert C, Polkinghorne A, Kebbi-Beghdadi C, et al.
236 The *Waddlia* genome: a window into chlamydial biology. *PLoS One* 2010;5:e10890.

- 237 [9] Blumer S, Greub G, Waldvogel A, Hassig M, Thoma R, Tschuor A, et al. *Waddlia*,
238 *Parachlamydia* and *Chlamydiaceae* in bovine abortion. *Vet Microbiol* 2011;152:385-93.
- 239 [10] Casson N, Medico N, Bille J, Greub G. *Parachlamydia acanthamoebae* enters and
240 multiplies within pneumocytes and lung fibroblasts. *Microbes Infect* 2006;8:1294-300.
- 241 [11] Casson N, Posfay-Barbe KM, Gervaix A, Greub G. New diagnostic real-time PCR
242 for specific detection of *Parachlamydia acanthamoebae* DNA in clinical samples. *J Clin*
243 *Microbiol* 2008;46:1491-3.
- 244 [12] Croxatto A, Greub G. Early intracellular trafficking of *Waddlia chondrophila* in
245 human macrophages. *Microbiology* 2010;156:340-55.
- 246 [13] Croxatto A, Rieille N, Kernif T, Bitam I, Aeby S, Peter O, et al. Presence of
247 *Chlamydiales* DNA in ticks and fleas suggests that ticks are carriers of *Chlamydiae*.
248 *Ticks and tick-borne diseases* 2014;5:359-65.
- 249 [14] de Barsy M, Bottinelli L, Greub G. Antibiotic susceptibility of *Estrella lausannensis*, a
250 potential emerging pathogen. *Microbes Infect* 2014.
- 251 [15] Deuchande R, Gidlow J, Caldow G, Baily J, Longbottom D, Wheelhouse N, et al.
252 *Parachlamydia* involvement in bovine abortions in a beef herd in Scotland. *Vet Rec*
253 2010;166:598-9.
- 254 [16] Goy G, Croxatto A, Posfay-Barbe KM, Gervaix A, Greub G. Development of a real-
255 time PCR for the specific detection of *Waddlia chondrophila* in clinical samples. *Eur J*
256 *Clin Microbiol Infect Dis* 2009;28:1483-6.
- 257 [17] Goy G, Greub G. Antibiotic susceptibility of *Waddlia chondrophila* in *Acanthamoeba*
258 *castellanii amoebae*. *Antimicrobial agents and chemotherapy* 2009;53:2663-6.
- 259 [18] Greub G, Raoult D. *Parachlamydiaceae*: potential emerging pathogens. *Emerg*
260 *Infect Dis* 2002;8:625-30.

- 261 [19] Greub G, Raoult D. Crescent bodies of *Parachlamydia acanthamoeba* and its life
262 cycle within *Acanthamoeba polyphaga*: an electron micrograph study. *Appl Environ*
263 *Microbiol* 2002b;68:3076-84.
- 264 [20] Hayashi Y, Nakamura S, Matsuo J, Fukumoto T, Yoshida M, Takahashi K, et al.
265 Host range of obligate intracellular bacterium *Parachlamydia acanthamoebae*. *Microbiol*
266 *Immunol* 2010;54:707-13.
- 267 [21] Henning K, Schares G, Granzow H, Polster U, Hartmann M, Hotzel H, et al.
268 *Neospora caninum* and *Waddlia chondrophila* strain 2032/99 in a septic stillborn calf.
269 *Vet Microbiol* 2002;85:285-92.
- 270 [22] Horn M. *Chlamydiae* as symbionts in eukaryotes. *Annual review of microbiology*
271 2008;62:113-31.
- 272 [23] Hornung S, Thuong BC, Gyger J, Kebbi-Beghdadi C, Vasilevsky S, Greub G, et al.
273 Role of *Chlamydia trachomatis* and emerging *Chlamydia*-related bacteria in ectopic
274 pregnancy in Vietnam. *Epidemiology and infection* 2014;1-4.
- 275 [24] Kebbi-Beghdadi C, Batista C, Greub G. Permissivity of fish cell lines to three
276 *Chlamydia*-related bacteria: *Waddlia chondrophila*, *Estrella lausannensis* and
277 *Parachlamydia acanthamoebae*. *FEMS Immunol Med Microbiol* 2011;63:339-45.
- 278 [25] Kebbi-Beghdadi C, Cisse O, Greub G. Permissivity of Vero cells, human
279 pneumocytes and human endometrial cells to *Waddlia chondrophila*. *Microbes and*
280 *infection* 2011;13:566-74.
- 281 [26] Kebbi-Beghdadi C, Domrose A, Becker E, Cisse OH, Hegemann JH, Greub G.
282 *OmpA* family proteins and *Pmp*-like autotransporter: new adhesins of *Waddlia*
283 *chondrophila*. *Pathogens and disease* 2015;73.

- 284 [27] Kebbi-Beghdadi C, Greub G. Importance of amoebae as a tool to isolate amoeba-
285 resisting microorganisms and for their ecology and evolution: the Chlamydia paradigm.
286 Environmental microbiology reports 2014;6:309-24.
- 287 [28] Kocan KM, Crawford TB, Dilbeck PM, Evermann JF, McGuire TC. Development of a
288 rickettsia isolated from an aborted bovine fetus. J Bacteriol 1990;172:5949-55.
- 289 [29] Lienard J, Croxatto A, Gervais A, Posfay-Barbe K, Baud D, Kebbi Beghdadi C, et al.
290 Undressing of *Waddlia chondrophila* to enrich its outer membrane proteins to develop a
291 new species-specific ELISA. New Microbe New Infect 2014;2:13-24.
- 292 [30] Lienard J, Croxatto A, Prod'hom G, Greub G. *Estrella lausannensis*, a new star in
293 the Chlamydiales order. Microbes Infect 2011;13:1232-41.
- 294 [31] Michel R, Steinert M, Zöller L, Hauröder B, Henning K. Free-living Amoebae may
295 serve as hosts for the *Chlamydia*-like bacterium *Waddlia chondrophila* isolated from an
296 aborted bovine foetus. Acta protozoologica 2004;43:37-42.
- 297 [32] Moulder JW. Interaction of *chlamydiae* and host cells *in vitro*. Microbiological
298 reviews 1991;55:143-90.
- 299 [33] Roger T, Casson N, Croxatto A, Entenza JM, Pusztaszeri M, Akira S, et al. Role of
300 MyD88 and Toll-like receptors 2 and 4 in the sensing of *Parachlamydia acanthamoebae*.
301 Infect Immun 2010;78:5195-201.
- 302 [34] Ruhl S, Casson N, Kaiser C, Thoma R, Pospischil A, Greub G, et al. Evidence for
303 *Parachlamydia* in bovine abortion. Vet Microbiol 2009;135:169-74.
- 304 [35] Sixt BS, Hiess B, König L, Horn M. Lack of effective anti-apoptotic activities restricts
305 growth of *Parachlamydiaceae* in insect cells. PLoS One 2012;7:e29565.
- 306 [36] Taylor-Brown A, Vaughan L, Greub G, Timms P, Polkinghorne A. Twenty years of
307 research into *Chlamydia*-like organisms: a revolution in our understanding of the biology

308 and pathogenicity of members of the phylum *Chlamydiae*. *Pathogens and disease*
309 2015;73:1-15.

310 [37] Wheelhouse N, Howie F, Gidlow J, Greub G, Dagleish M, Longbottom D.
311 Involvement of *Parachlamydia* in bovine abortions in Scotland. *Veterinary journal*
312 2012;193:586-8.

313 [38] Yamane C, Yamazaki T, Nakamura S, Matsuo J, Ishida K, Yamazaki S, et al.
314 Amoebal endosymbiont *Parachlamydia acanthamoebae* Bn9 can grow in immortal
315 human epithelial HEp-2 cells at low temperature; an in vitro model system to study
316 chlamydial evolution. *PLoS One* 2015;10:e0116486.

317

318 **7. Figure legends**

319 **Fig. 1 Bacterial growth within insect cells**

320 *W. chondrophila* (panels a and b), *E. lausannensis* (panels c and d) and *P.*
321 *acanthamoebae* (panels e and f) replication measured by qPCR in *Aedes albopictus*
322 cells (panels a, c and e) and in Sf9 cells (panels b, d and f). Values are normalized to
323 the number of bacteria at 0h post infection (p.i). Results are the mean +/-SD of at least
324 four independent experiments performed in duplicates.

325

326 **Fig.2 Bacterial replication assessed by confocal and electron microscopy**

327 A. Immunofluorescence and confocal microscopy of *W. chondrophila* (panels a and
328 b), *E. lausannensis* (panels c and d) and *P. acanthamoebae* (panels e and f) in
329 *Aedes albopictus* cells (panels a, c and e) and in Sf9 cells (panels b, d and f) 24
330 hours post-infection. Panel g displays one rare inclusion of *P. acanthamoebae* in
331 *Aedes albopictus* cells 4 days post infection. Bacteria (green) are stained with
332 species-specific polyclonal antibodies and insect cells (red) are stained with Texas
333 Red-conjugated Concanavalin A. Scale bar 10 μm .

334 B. Electron micrographs of *W. chondrophila* (panel a) and *E. lausannensis* (panel b)
335 in Sf9 cells 24 hours post infection. Scale bar: 1 μm .

336

337 **Fig. 3 Cell viability**

338 *Aedes albopictus* cells viability was determined with a propidium iodide assay at different
339 time points after infection. Results are the mean +/- SD of 2 independent experiments
340 performed in triplicates.



Figure 1

Figure 2
A



B




Figure 3

