

Elsevier Editorial System(tm) for Journal of  
Structural Biology  
Manuscript Draft

Manuscript Number: JSB-16-119R1

Title: Structural and calorimetric studies demonstrate that the Hepatocyte Nuclear Factor 1 $\beta$  (HNF1 $\beta$ ) transcription factor is imported into the nucleus via a monopartite NLS sequence

Article Type: Short Communication

Keywords: nuclear transport / hepatocyte nuclear factor 1 $\beta$  (HNF1 $\beta$ ) / nuclear localization sequence

Corresponding Author: Dr. Murray Stewart, MA PhD

Corresponding Author's Institution: MRC Laboratory of Molecular Biology

First Author: Mareike M Wiedmann, MChem

Order of Authors: Mareike M Wiedmann, MChem; Shintaro Aibara, PhD; David R Spring, PhD; Murray Stewart, MA PhD; James D Brenton, PhD

Abstract: The transcription factor hepatocyte nuclear factor 1 $\beta$  (HNF1 $\beta$ ) is ubiquitously overexpressed in ovarian clear cell carcinoma (CCC) and is a potential therapeutic target. To explore potential approaches that block HNF1 $\beta$  transcription we have identified and characterised extensively the nuclear localisation signal (NLS) for HNF1 $\beta$  and its interactions with the nuclear protein import receptor, Importin- $\alpha$ . Pulldown assays demonstrated that the DNA binding domain of HNF1 $\beta$  interacted with a spectrum of Importin- $\alpha$  isoforms and deletion constructs tagged with eGFP confirmed that the HNF1 $\beta$  229KKMRRNR235 sequence was essential for nuclear localisation. We further characterised the interaction between the NLS and Importin- $\alpha$  using complementary biophysical techniques and have determined the 2.4 Å resolution crystal structure of the HNF1 $\beta$  NLS peptide bound to Importin- $\alpha$ . The functional, biochemical, and structural characterisation of the nuclear localisation signal present on HNF1 $\beta$  and its interaction with the nuclear import protein Importin- $\alpha$  provides the basis for the development of compounds targeting transcription factor HNF1 $\beta$  via its nuclear import pathway.

*J. Struct. Biol. JSB-16-119 (revised)*

1  
2  
3  
4  
5  
6 **Structural and calorimetric studies demonstrate that the**  
7 **Hepatocyte Nuclear Factor 1 $\beta$  (HNF1 $\beta$ ) transcription factor is**  
8 **imported into the nucleus via a monopartite NLS sequence**  
9  
10  
11  
12

13 Mareike M. Wiedmann<sup>a,b</sup>, Shintaro Aibara<sup>c</sup>, David R. Spring<sup>a</sup>, Murray Stewart<sup>c\*</sup>, James D.  
14 Brenton<sup>b</sup>  
15

16 <sup>a</sup> Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW,  
17 UK  
18

19 <sup>b</sup> Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre,  
20 Robinson Way, Cambridge CB2 0RE, UK  
21

22 <sup>c</sup> MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical  
23 Campus, Cambridge CB2 0QH, UK  
24

25  
26  
27  
28  
29  
30 \* To whom correspondence should be addressed: Dr Murray Stewart, MRC Laboratory of  
31 Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB2  
32 0QH, UK; E-mail: ms@mrc-lmb.cam.ac.uk; Fax: +44 1223 267000; Phone: +44 1223 267074.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Abstract

1 The transcription factor hepatocyte nuclear factor 1 $\beta$  (HNF1 $\beta$ ) is ubiquitously overexpressed  
2 in ovarian clear cell carcinoma (CCC) and is a potential therapeutic target. To explore  
3 potential approaches that block HNF1 $\beta$  transcription we have identified and characterised  
4 extensively the nuclear localisation signal (NLS) for HNF1 $\beta$  and its interactions with the  
5 nuclear protein import receptor, Importin- $\alpha$ . Pull-down assays demonstrated that the DNA  
6 binding domain of HNF1 $\beta$  interacted with a spectrum of Importin- $\alpha$  isoforms and deletion  
7 constructs tagged with eGFP confirmed that the HNF1 $\beta$  <sup>229</sup>KKMRRNR<sup>235</sup> sequence was  
8 essential for nuclear localisation. We further characterised the interaction between the NLS  
9 and Importin- $\alpha$  using complementary biophysical techniques and have determined the 2.4 Å  
10 resolution crystal structure of the HNF1 $\beta$  NLS peptide bound to Importin- $\alpha$ . The functional,  
11 biochemical, and structural characterisation of the nuclear localisation signal present on  
12 HNF1 $\beta$  and its interaction with the nuclear import protein Importin- $\alpha$  provide the basis for the  
13 development of compounds targeting transcription factor HNF1 $\beta$  via its nuclear import  
14 pathway.  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 **Keywords:** Importin- $\alpha$ , nuclear import pathway, nuclear localisation signal sequence (NLS),  
25 Hepatocyte nuclear factor-1 $\beta$  (HNF1 $\beta$ ), site-directed mutagenesis, X-ray crystallography,  
26 isothermal titration calorimetry.  
27  
28  
29  
30

31 **Abbreviations:** CCC - Ovarian clear cell carcinoma; HNF1 $\beta$  - hepatocyte nuclear factor 1 $\beta$ ;  
32 HNF1 $\beta$ <sup>DBD</sup> - HNF1 $\beta$  DNA binding domain; NLS - nuclear localization signal; IBB - Importin- $\beta$ -  
33 binding domain; mImportin- $\alpha$  - mouse Importin- $\alpha$ ; xImportin- $\alpha$  - *Xenopus* Importin- $\alpha$ ; GST -  
34 glutathione S-transferase.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

Ovarian clear cell carcinoma (CCC) accounts for 5-10% of ovarian cancer cases (Anglesio et al., 2011; Kato et al., 2008). Prognosis for patients with advanced stage or for relapsed disease is poor because of intrinsic resistance to platinum based chemotherapy and the lack of targeted therapies available (Tan et al., 2013; Tan and Kaye, 2007), although Bitler *et al.* (2015) have recently discovered a way of targeting cancers with *ARID1A* mutations by targeting EZH2 methyltransferase activity. Common mutations in CCC include loss of function mutations in the chromatin remodeling gene *ARID1A* in 46–57% of cases (Jones et al., 2010; Wiegand et al., 2010), activating mutations in *PIK3CA* (Kuo et al., 2009) in 33–46% of cases, and loss of *PTEN* in 20% of cases (Anglesio et al., 2011; Landen et al., 2008; Tan and Kaye, 2007). Overexpression of the HNF1 $\beta$  transcription factor is the most important clinical immunohistochemical marker for the disease because it is ubiquitously overexpressed in CCC, both at the mRNA and protein level (Hirota Kajihara, 2010; Kato et al., 2006; Tsuchiya et al., 2003; Yamaguchi et al., 2010). In CCC the *HNF1B* gene is upregulated by hypomethylation of its CpG island whereas, in high grade serous ovarian cancer, HNF1 $\beta$  expression is silenced via hypermethylation (Kato et al., 2008; Shen et al., 2013a), suggesting that HNF1 $\beta$  has a loss of function (tumour suppressor) role in high grade serous ovarian cancer but a gain of function (oncogenic) role in CCC (Gounaris et al., 2011; Shen et al., 2013b). This hypothesis is supported by the observation that nearly half of the overexpressed genes identified in CCC are downstream targets of HNF1 $\beta$  (Kobayashi et al., 2009; Yoshida et al., 2009). Evidence that targeting HNF1 $\beta$  might have utility was provided by Liu *et al.* (Liu et al., 2009) who showed that downregulation of HNF1 $\beta$  increased cisplatin- and paclitaxel-mediated cytotoxicity.

Transcription factor HNF1 $\beta$  (also known as vHNF1, vAPF, LF-B3 and Tcf2) is expressed in the liver, digestive tract, pancreas and the kidneys, where it plays a crucial role in early differentiation (Lu et al., 2007). Sequence-specific DNA binding is mediated by a bipartite motif that consists of a POU homeodomain (POU<sub>H</sub>) and a POU specific domain (POU<sub>S</sub>) (Rosenfeld, 1991; Ryan and Rosenfeld, 1997). HNF1 $\beta$  has 70% sequence homology to HNF1 $\alpha$  and both proteins are atypical members of the POU transcription factor family and bind DNA as both homo- and heterodimers (Bach et al., 1991; Rey-Campos et al., 1991). Human HNF1 $\beta$  is constructed from three domains: the dimerization domain, which is further stabilised by the dimerization cofactor of HNF1 (DcoH), the transactivation domain, which is involved in binding transcriptional co-activators [15], and the POU DNA binding domain (HNF1 $\beta$ <sup>DBD</sup>).

Transcription factors together with histones, DNA polymerase, RNA polymerase and many other proteins, have specific amino acid sequences, termed nuclear localisation signals (NLSs), that are recognised by members of the karyopherin family that facilitate their nuclear import (reviewed by Lange et al., 2007). Many NLS sequences are recognised in the

1 cytoplasm by a heterodimeric transport carrier complex composed of Importin- $\beta$  (also known  
2 as Karyopherin- $\beta$ 1) and Importin- $\alpha$  (reviewed by Stewart, 2007). Nuclear pore complexes  
3 (NPCs) are the channels through which macromolecules, such as proteins and RNA, are  
4 transported between the cytoplasm and nucleus (reviewed by Stewart, 2007). Small  
5 molecules and proteins (< 40 kDa) can pass through NPCs by passive diffusion, but larger  
6 proteins require carriers to overcome the NPC physical barrier. The autoinhibitory Importin-  
7  $\beta$  binding (IBB) domain of Importin- $\alpha$  (Kobe, 1999) binds to Importin- $\beta$  in the cytoplasm,  
8 enabling classical NLSs (cNLS) to bind to Importin- $\alpha$  either via a major site, a minor site, or  
9 both (Fontes et al., 2000; Lange et al., 2007). There are two types of cNLS that are  
10 recognised by Importin- $\alpha$  that consist of either a single cluster (monopartite) or two clusters  
11 (bipartite) of positively charged residues, primarily lysines or arginines, that assume an  
12 ordered state once bound by Importin- $\alpha$  (reviewed by Lange et al. 2007; Marfori et al., 2011).  
13 Monopartite cNLSs are exemplified by the simian virus 40 large T-antigen (SV40) NLS  
14  $^{126}\text{PKKKRRV}^{132}$  (Lange et al., 2007). The cargo:carrier heterotrimer is then translocated into  
15 the nucleus in an energy dependent manner powered by RanGTPase (reviewed by Stewart,  
16 2007). In the nucleus, RanGTP then binds Importin- $\beta$  leading ultimately to the release of the  
17 cargo.  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 To function, HNF1 $\beta$  needs to be translocated to the cell nucleus and so we have  
28 investigated and characterized the putative NLS that has been proposed to lie between the  
29 two POU domains of HNF1 $\beta$ . This putative NLS was identified in domain swapping  
30 experiments and studies of nephrogenesis, in which truncated GFP-HNF1 $\beta$  fusion constructs  
31 retaining the POU<sub>H</sub> domain showed exclusive nuclear localisation in transfected HeLa cells  
32 (Bohn et al., 2003; Wu et al., 2004). However, the precise location of the HNF1 $\beta$  NLS has not  
33 been defined. Because of the potential importance of HNF1 $\beta$  as a target in diseases such as  
34 CCC, we have identified and characterized extensively its NLS and its interactions with  
35 Importin- $\alpha$ . We demonstrate that the sequence  $^{229}\text{KKMRRNR}^{235}$  in HNF1 $\beta^{\text{DBD}}$  is responsible  
36 for the nuclear import of the protein. Several eGFP-constructs of HNF1 $\beta$  were generated and  
37 Importin- $\alpha$  binding of the HNF1 $\beta^{\text{DBD}}$  was assessed by both pull-down experiments and ITC.  
38 We also determined the crystal structure of the HNF1 $\beta$  NLS peptide bound to Importin- $\alpha$ . The  
39 identification and structural characterisation of the HNF1 $\beta$  NLS and its interaction with the  
40 nuclear import protein Importin- $\alpha$  provides a basis for the development of inhibitors targeting  
41 the nuclear import of transcription factor HNF1 $\beta$  along the lines suggested by Stelma et al.  
42 (2016).  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 **Materials and Methods**

### 57 ***Mammalian cell culture***

1 HEK293T cells were cultivated in Dulbecco's Modified Eagle medium (DMEM) (1X)  
2 supplemented with 5% foetal bovine serum (FBS) (Invitrogen) and 0.5%  
3 penicillin/streptomycin (P/S). CCC cell lines PEO1, JHOC5, JHOC7, JHOC9, OVISE and  
4 SKOV3 cells were grown in RPMI 1640 medium (1X) supplemented with 10% FBS and  
5 1% P/S. Normal Ovarian Surface Epithelial (IOSE) cells were cultivated in NOSE-CM: MCDB  
6 105/medium 199 (1:1 ratio, Sigma Aldrich), 15% FBS, 10 ng/ml EGF (Invitrogen), 0.5 µg/ml  
7 hydrocortisone (Sigma Aldrich), 5 µg/ml insulin (Sigma Aldrich), 34 µg protein/ml BPE  
8 (Invitrogen). All cell lines were maintained at 37°C in 5% CO<sub>2</sub> and were mycoplasma tested  
9 on a regular basis (Biorepository Core, CRUK CI, Cambridge). Cell counts were conducted  
10 using a Vi-CELL Cell Viability Analyzer.  
11  
12  
13  
14  
15  
16

### 17 ***Protein extraction from mammalian cells***

18 Cell pellets were washed with phosphate buffered saline (PBS) and 200 µl protein lysis buffer  
19 (50 mM Tris pH 8.0, 150 mM NaCl, 5 mM EDTA, 0.5% Igepal, to which 2 tablets/100ml of  
20 protease inhibitor cocktail tablet (Roche)) was added. The mixtures were incubated on ice for  
21 30 min, lysed by syringing four times using a 26 G needle, and centrifuged at 14,800 × g for  
22 10 min at 4°C. Protein concentrations were measured using the DirectDetect IR  
23 spectrometer (Merck Millipore) according to the manufacturer's instructions.  
24  
25  
26  
27  
28

### 29 ***Western blotting***

30 Denatured protein extracts were separated using NuPAGE Novex 4-20% Tris-Glycine gels  
31 and transferred to a Millipore Immobilon FL PVDF membrane (Invitrogen). Primary  
32 antibodies were used as follows: goat anti-HNF1β (sc-7411, polyclonal, Santa Cruz  
33 Biotechnology, 1:1,000) and rabbit anti-GAPDH (G9545, 14C10, 1:5,000, Cell Signalling  
34 Technology). An Odyssey Infrared Imaging System (Li-Cor) and associated secondary  
35 antibodies: donkey anti-goat (800) (1:15,000) and donkey anti-rabbit (800) (1:5,000) were  
36 used to detect material. The expression levels observed with different cells were evaluated  
37 using GraphPad Prism version 6.0 for Mac, GraphPad Software, La Jolla California USA,  
38 [www.graphpad.com](http://www.graphpad.com). Analysis of variance rejected the null hypothesis that the expression  
39 levels were equal (P < 0.002) and Šídák's multiple comparison modification of Student's t-test  
40 was used to evaluate the significance of the levels seen with different cells relative to that  
41 seen with the control HGSOC cell line.  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 ***Confocal microscopy: eGFP imaging and immunofluorescence assay***

52 For immunofluorescence (IF), cells were fixed with 4% paraformaldehyde in PBS for 10 min  
53 at room temperature, rinsed in Tris buffered saline (TBS) for 2 × 5 min and permeabilized in  
54 TBS-0.5% Triton X-100 for 10 min. Fixed cells were then rinsed in TBS-0.1% Triton X-100 for  
55 3 × 3 min and blocked in 10% goat serum in TBS for 30 min. HNF1β was stained with anti-  
56 HNF1β (SAB1406512, mouse polyclonal, 1:300, Sigma Aldrich) overnight at 4°C. Cells were  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 washed with TBS-0.1% Triton X-100 (Fisher Scientific) for 3 × 5 min and the secondary  
2 antibody (AlexaFluor 568, goat anti-mouse IgG (H+L) 1:1,000, Invitrogen) was added and  
3 incubated for 60 min diluted in antibody dilution solution consisting of TBS - 0.1% Triton X-  
4 100, 2% Bovine Serum Albumin (BSA) (Cell Signaling Technology) and 0.1% sodium azide.  
5 All experiments included an unstained control, a “secondary only” control, and a negative  
6 control using the PEO1 cell line. Cells transduced with eGFP-HNF1β, the HNF1β  
7 <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant, or the control eGFP constructs, were fixed with 4%  
8 paraformaldehyde in PBS for 10 min at room temperature, then rinsed in TBS for 2 × 5 min.  
9 Nuclei were stained using DAPI (1 μg/ml in TBS) for 10 min. Cells were washed in TBS-0.1%  
10 Triton X-100, rinsed in TBS and stored in TBS-0.1% Triton X-100. For microscopy imaging,  
11 cover slips were drained, mounted and sealed using Prolong Gold (Invitrogen) and glass  
12 slides (Thermo Specific). Slides were left to dry overnight at room temperature in the  
13 absence of light and were then stored at 4°C. Cells were imaged using a Leica tandem  
14 confocal microscope.  
15  
16  
17  
18  
19  
20  
21  
22

### 23 **Site directed mutagenesis to generate HNF1β NLS deletion construct**

24 Lv103 (EX-F0366-Lv103, Genecopoeia) is a lentiviral transfer vector containing an eGFP-  
25 HNF1β fusion coding sequence. Lv105 (EX-EGFP-Lv105, Genecopoeia) only contains the  
26 eGFP coding sequence. The plasmids were confirmed by sequencing (GATC, Konstanz,  
27 Germany) before and after mutagenesis (Lv103 Fw: 5'-CCGACAACCACTACCTGA-3'; Rv: 5'-  
28 ATTGTGGATGAATACTGCC-3' and Lv105 Fw: 5'-ATCCACGCTGTTTTGACC-3'; Rv: 5'-  
29 AATACTGCCATTTGTCTCG-3').  
30  
31  
32  
33  
34  
35

36 Mutagenesis experiments were conducted using the Q5 site directed mutagenesis kit  
37 (NEB). The <sup>229</sup>KKMRRNR<sup>235</sup> deletion construct was generated using mutagenesis primers  
38 (Fw: 5'-TTCAAATGGGGGCCGC-3'; Rv: 5'-GTTGGTGGGCTCAGAGCAG-3'). Plasmids  
39 were then transformed in *E. coli* (One Shot® Stbl3™, Invitrogen) and streaked on Amp-  
40 containing agar plates. Single colonies were picked and tested for plasmid containing  
41 colonies by PCR. Plasmids were extracted by Plasmid Mini Prep (Qiagen) and quantified  
42 (Qubit).  
43  
44  
45  
46  
47  
48

### 49 **Lentivirus production and transduction**

50 The general protocol devised by Cribbs *et al.* (2013) was used for lentivirus production. The  
51 transfer vectors Lv103, Lv105, the <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant, and plasmids pRVS-Rev,  
52 pVSV-G, and p-MDLg-pRRE were verified by restriction digest. For each transfection  
53 sample, 16 μg transfer vector, 10.4 μg pMDL/pRRE, 4 μg pRSV-Rev and 5.6 μg pVSV-G  
54 were used. After 24 h, expression of GFP protein was observed in the GFP control virus  
55 HEK293T sample. Virus containing supernatant was harvested posttransfection according to  
56 the protocol by Kutner *et al.* (2009). The number of transducing units (TU) was determined by  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 flow cytometry analysis with GFP as the reporter protein. For titration,  $1 \times 10^5$  cells per well  
2 were seeded in a 12-well plate and 0.5 ml DMEM with 5% FBS added. Virus of respective  
3 concentration was pipetted over the HEK293T cells. Duplicate virus dilutions of 1:500,  
4 1:1000, 1:2000, 1:5000 were added. An untreated control was included. The medium was  
5 changed 24 h posttransfection. 72 h posttransfection cells were harvested by trypsinization.  
6 Cells were collected by centrifugation at  $500 \times g$  for 5 min at room temperature. The  
7 supernatant was discarded and cell pellets were resuspended in PBS. A green fluorescent  
8 protein propidium iodide (GFP PI) based assay for flow cytometric measurement of  
9 transfection efficiency and cell viability was performed (LSR-II machine). The titre was  
10 calculated from dilutions that gave 1–40% GFP-positivity and averaged subsequently using  
11 the following formula:  
12  
13  
14  
15

$$16 \text{Titre } \left( \frac{\text{TU}}{\text{ml}} \right) = \frac{17 \text{(frequency of GFP-positive cells)} \times \text{no. cells plated} \times \text{dilution factor}}{18 \text{volume of inoculum}}$$

19 where, frequency of GFP-positive cells is the percentage of cells that are positive for GFP  
20 divided by 100 (acceptable range: 0.01–0.40), dilution factor is the dilution of the virus stock  
21 used and volume of inoculum is the total volume transduced. The titre was calculated to be  
22  $2.26 \times 10^8 \frac{\text{TU}}{\text{ml}}$ .  
23  
24  
25  
26  
27  
28  
29  
30

31 For transduction of the PEO1 cell line,  $1 \times 10^6$  cells per well were seeded in a 12-  
32 well plate to which 0.5 ml of the respective medium was added. Virus was added so that the  
33 desired multiplicity of infection (MOI) was obtained. Polybrene 0.5-10  $\mu\text{g/ml}$  was added to  
34 increase transduction efficiency. Transduction efficiencies were determined by flow  
35 cytometry analysis using an LSR-II machine. For selection of transduced PEO1 lines, a  
36 puromycin kill curve was constructed (data not shown) to determine the optimum  
37 concentration of puromycin required to kill all untransduced cells while minimising toxicity  
38 effects in transduced cells. A puromycin concentration of 0.1  $\mu\text{g/ml}$  was determined which  
39 was used in all experiments.  
40  
41  
42  
43  
44  
45  
46

#### 47 **Quick change mutagenesis to generate GST-tagged HNF1 $\beta$ <sup>DBD</sup> and mImportin $\alpha$ 1** 48 **mutants** 49

50 The cloning of the DNA binding domain of HNF1 $\beta$  into the pGEX-TEV plasmid was conducted  
51 as described by Lu *et al.* (2006). The plasmid was confirmed by sequencing by Source  
52 Bioscience (Cambridge, UK) using PGEX5 and PGEX3 primers (Source Bioscience).  
53  
54  
55

#### 56 **Protein expression in bacterial cells and purification** 57

58 All proteins were expressed in *E. coli* BL21 (DE3) CodonPlus-RIL cells using IPTG induction  
59 over 18 h at 18°C. The cells were harvested by centrifugation and resuspended in 50 mM  
60  
61  
62  
63  
64  
65



1 Tris/HCl pH 8.0, 500 mM NaCl, 5 mM DTT for GST-tagged constructs or 50 mM Tris/HCl pH  
2 8.0, 500 mM NaCl, 20 mM Imidazole pH 8.0 for His<sub>6</sub>-tagged proteins. The *E. coli* were lysed  
3 by two passes through an Emulsiflex C3 system (AVESTIN) running at a pressure of  
4 15,000 psi and the lysates clarified by centrifugation at 48,000 × g. Either Ni-NTA agarose  
5 resin (QIAGEN) or glutathione Sepharose 4B resin (GE Healthcare) was added to the  
6 clarified lysate containing His<sub>6</sub>-tagged proteins or GST-tagged proteins. The resin was  
7 pooled and packed into a gravity filtration column and washed extensively with their  
8 respective lysis buffers. 1 mg of TEV protease (purified in-house) was added to the washed  
9 resin and incubated for 18 h at 4°C. The tag-free protein of interest was then collected from  
10 the flow-through and subjected to the next purification step. HNF1β<sup>DBD</sup> was purified further by  
11 heparin affinity chromatography using a HiTrap Heparin HP column (GE Healthcare Life  
12 Sciences). HNF1β<sup>DBD</sup> was eluted from the column using a 0.1 - 1 M NaCl gradient. All  
13 proteins were purified to homogeneity by size exclusion chromatography using a HiLoad™  
14 26/60 Superdex™ 75 prep grade column (GE Healthcare) pre-equilibrated in 20 mM HEPES  
15 buffer pH 8.0, 200 mM NaCl. Fractions containing pure and homogenous proteins identified  
16 by SDS-PAGE analysis were pooled and concentrated using an Amicon Ultra 15  
17 centrifugation filtration unit. Protein concentrations were measured using a Nanodrop  
18 (Thermo) microspectrophotometer and small aliquots of the protein were flash frozen in liquid  
19 nitrogen and stored at -80°C until required.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

### 30 ***GST pull-downs***

31 After bacterial cell lysis, cell pellets expressing the two proteins of interest were mixed and  
32 pull-down experiments were conducted as described above for glutathione affinity  
33 chromatography and were visualised by SDS-page analysis.  
34  
35  
36  
37  
38

### 39 ***Isothermal calorimetry***

40 The K<sub>d</sub> of mImportin-α1 with the HNF1β NLS peptide (Ac-TNKKMRRNRFK-NH<sub>2</sub>, purchased  
41 from Insight Biotechnology) was determined by isothermal titration calorimetry (ITC) using a  
42 MicroCal™ iTC200 (GE Healthcare Life Sciences). The protein and peptide samples were  
43 made up in the identical buffer (20 mM Hepes pH 8, 200 mM NaCl). 350 μl of protein sample  
44 (0.02 mM) and 200 μl of ligand (0.2 mM) were prepared for each experiment. Sample  
45 concentrations were confirmed by amino acid analysis (PNAC Facility, Biochemistry  
46 Department, Cambridge University). All experiments were conducted at 25°C. The data was  
47 analysed using the Origin™ Software (MicroCal) using a one site binding model.  
48  
49  
50  
51  
52  
53

### 54 ***Protein crystallography***

55 Diffraction quality crystals of the Importin-α1<sup>ΔIBB</sup>:HNF1β<sup>NLS</sup> complex were obtained by sitting  
56 drop vapour diffusion where 200 nl of 10% PEG 8K, 0.09 M NPS, 20% ethylene glycol, 0.1 M  
57 Buffer 1 (Morpheus Screen, Molecular Dimensions) was mixed with 200 nl of Importin-  
58  
59  
60  
61  
62  
63  
64  
65

1  $\alpha 1^{\Delta IBB} : HNF1\beta^{NLS}$ . The Importin- $\alpha 1^{\Delta IBB} : HNF1\beta^{NLS}$  complex was produced by mixing Importin-  
2  $\alpha 1^{\Delta IBB}$  and HNF1 $\beta^{NLS}$  peptide (Ac-TNKKMRRMRFK-NH<sub>2</sub>, Insight Biotechnology) at a 1:1.1  
3 molar ratio and a concentration of 6.7 mg/ml.  
4  
5

6 Crystals were cryoprotected in the mother liquor supplemented with 20% glycerol and  
7 cooled by plunging into liquid nitrogen. X-ray diffraction data were collected on beamline I03  
8 at the Diamond Light Source (Didcot, UK). Reflections were indexed and integrated using  
9 *DIALS* as implemented in *Xia2* (Kabsch, 2010) and then scaled and merged in *AIMLESS*,  
10 ensuring a completeness of > 98% in the outermost shell while maintaining  $CC_{1/2} > 0.3$   
11 (Evans and Murshudov, 2013). The structure was solved by molecular replacement using  
12 *Phaser* with the structure of  $\Delta IBB$ -mImportin- $\alpha 1$  complexed with a minor site small molecule  
13 inhibitor (PDB ID: 4U54 - Holvey et al., 2015) to avoid introducing model bias into the NLS  
14 binding sites. Iterative cycles of rebuilding using *COOT* (Emsley et al., 2010) and refinement  
15 using *PHENIX* (Adams et al., 2010) were used to generate the final model (Table1) that had  
16 an R-factor of 18.4% (R-free = 22.0%) and a MolProbity score of 0.95 (100th percentile).  
17  
18  
19  
20  
21  
22  
23  
24

## 25 **Results and Discussion**

### 26 ***HNF1 $\beta$ is overexpressed in CCC cell lines***

27 HNF1 $\beta$  protein expression in different CCC cell lines was determined by Western blotting  
28 (Figure 1) using the HGSOC cell line PEO1 (which does not express HNF1 $\beta$ ) as a negative  
29 control. The CCC lines OVI5E, JHOC5, JHOC7, JHOC9 and SKOV3 cell lines were used as  
30 models of CCC that overexpress HNF1 $\beta$ . Several splice variants of HNF1 $\beta$  are known to  
31 contain different isoforms of the C-terminal domain that is responsible for activation of  
32 transcription. These isoforms act as transdominant repressors (Bach and Yaniv, 1993). Only  
33 the 61 kDa isoform of HNF1 $\beta$  was detected in our CCC cell lines (Figure 1) (Bach and Yaniv,  
34 1993; Tsuchiya et al., 2003). HNF1 $\beta$  expression levels varied considerably between these  
35 cell lines. Analysis of variance of the data in Figure 1 rejected the null hypothesis that the  
36 expression levels were equal ( $P < 0.002$ ) and the higher level seen in JHOC7 compared with  
37 PEO1 cells was significant at the 0.5% level (using Sidak's modification for multiple  
38 comparisons), consistent with hypomethylation of the HNF1 $\beta$  CpG island (Kato et al., 2008;  
39 Kato et al., 2007).  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 ***Identification of the HNF1 $\beta$ nuclear localisation signal***

52 To confirm the identity of the NLS, the PEO1 cell line was transduced with lentiviruses Lv103  
53 (eGFP-HNF1 $\beta$ ), Lv105 (eGFP) and the <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant. In cells transduced  
54 with eGFP-HNF1 $\beta$ , the fusion protein (eGFP-HNF1 $\beta$  (Lv103) - Figure 2A(Mutant 8)) was  
55 primarily nuclear, whereas cells transduced with Lv105 (eGFP) showed nuclear and  
56 cytoplasmic localisation of the eGFP protein (eGFP (Lv105) - Figure 2B). Deletion of the  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 seven-residue<sup>229</sup>KKMRRNR<sup>235</sup> NLS sequence in HNF1 $\beta$  resulted in its becoming  
2 mislocalized to the cytoplasm (Mutant 8, Figure 2C), confirming the importance of this  
3 sequence for the nuclear import of HNF1 $\beta$ . HNF1 $\alpha$  has an analogous KKGRRN sequence  
4 that differs only at residue 231, where Met is changed to Gly (Wu et al., 2004) and which  
5 probably also functions in a similar manner. Interestingly, all transduced cells (that were  
6 selected using optimised puromycin concentrations) were apoptotic within 24 h. We  
7 speculate that re-expression of a transcription factor that is epigenetically silenced in HGSO  
8 cell line PEO1 (Shen et al., 2013a) may have a negative effect on proliferation, but further  
9 work is necessary to determine the effects of re-expression.  
10  
11  
12  
13

### 14 ***HNF1 $\beta$ <sup>DBD</sup> binds primarily to the major site on Importin- $\alpha$***

15 Because the sequence of the HNF1 $\beta$  NLS is similar to that of classical monopartite NLSs  
16 (Lange et al., 2007), HNF1 $\beta$  is likely to be imported by Importin- $\alpha$  and to interact with its major  
17 NLS-binding site rather than being imported by direct binding to Karyopherin- $\beta$  (Fontes et al.,  
18 2000). To test whether HNF1 $\beta$  and mImportin- $\alpha$  formed a complex, recombinantly expressed  
19 and purified HNF1 $\beta$ <sup>DBD</sup> and mImportin- $\alpha$ 1 in which the IBB domain had been deleted ( $\Delta$ IBB-  
20 mImportin- $\alpha$ 1) were mixed together and analysed by gel filtration chromatography (Figure  
21 3A), in which  $\Delta$ IBB-mImportin- $\alpha$ 1 and HNF1 $\beta$ <sup>DBD</sup> co-eluted as a stoichiometric complex as  
22 observed by Coomassie blue stained SDS-PAGE analysis. To establish whether HNF1 $\beta$   
23 interacts with other Importin- $\alpha$  isoforms, pull-down experiments using different  $\Delta$ IBB-  
24 mImportin- $\alpha$ 1 isoforms were performed (Figure 3B). Crude extracts of GST-tagged HNF1 $\beta$   
25 and  $\Delta$ IBB-mImportin- $\alpha$ 1 isoforms were mixed, then bound to glutathione resin, washed, and  
26 eluted by cleaving between GST and HNF1 $\beta$ <sup>DBD</sup> using TEV protease. Both HNF1 $\beta$ <sup>DBD</sup> and  
27  $\Delta$ IBB-mImportin- $\alpha$ 1 were released, confirming that they were interacting (Figure 3B). HNF1 $\beta$   
28 interacted more strongly with  $\Delta$ IBB-mImportin- $\alpha$ 5 and less strongly with  $\Delta$ IBB-mImportin- $\alpha$ 3, -  
29  $\alpha$ 4 and - $\alpha$ 7 isoforms. The  $K_d$  for the interaction of  $\Delta$ IBB-mImportin- $\alpha$ 1 with the HNF1 $\beta$  NLS  
30 peptide was determined by isothermal calorimetry (ITC) (Figure 4A) to be  $13.6 \pm 1.5$  nM  
31 which was in the range commonly observed (Hodel et al., 2006). In summary, HNF1 $\beta$  binds  
32 at least two  $\Delta$ IBB-mImportin- $\alpha$  isoforms strongly and some other isoforms more weakly.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 Three different *Xenopus* Importin- $\alpha$ 1 (xlImportin- $\alpha$ 1) mutants (Giesecke and Stewart,  
48 2010) were used to define where the HNF1 $\beta$  NLS bound on Importin- $\alpha$  (Figure 4B). These  
49 mutants contained point mutations that prevent binding via the major (D), minor (E), or both  
50 (ED) sites and also have the IBB domain removed to prevent auto-inhibition (Kobe, 1999).  
51 HNF1 $\beta$ <sup>DBD</sup> bound to the E mutant but not the D or ED mutant of  $\Delta$ IBB-xlImportin- $\alpha$ 1 (Figure  
52 4B), indicating that HNF1 $\beta$  binds primarily via the major site, consistent with its similarity to  
53 the SV40 NLS (Marfori et al., 2011).  
54  
55  
56  
57  
58

### 59 ***Structural characterisation of the interaction of mImportin- $\alpha$ 1 with the HNF1 $\beta$ NLS***

1 To complement the functional and biochemical data, we also determined the 2.4 Å resolution  
2 crystal structure of the HNF1β NLS peptide bound to ΔIBB-mImportin-α1 (Figure 5A). After  
3 initial rounds of refinement, strong difference density, corresponding to the peptide was found  
4 in both the major NLS-binding site on Importin-α1 together with weaker density at the minor  
5 site (Figures 5B and C, respectively).  
6  
7

8  
9 At the major site, the HNF1β NLS peptide backbone interacted with Importin-α1 via a  
10 set of conserved Asn residues on the import protein by H-bonding (Figure 6A). The NLS  
11 residues are conventionally assigned to positions P1-P5 (Figure 6). Key residues on  
12 Importin-α involved for the interaction with the NLS were Asn-235, 188 and 146, the latter two  
13 of which are involved in bidentate H-bonding with the NLS peptide backbone at P3 and P5.  
14 The aliphatic portions of the NLS peptide side chains were located in shallow pockets along  
15 the surface of Importin-α, but, as evident from Figure 6A, the P2 residue, Lys230, forms many  
16 critical interactions, mostly by forming salt bridges with negatively charged residues on  
17 Importin-α. There were three interactions of the protonated nitrogen on the P2 lysine side  
18 chain with Asp192, Gly150 and Thr155. Residue P5 interacted with the side-chain of Gln181.  
19 Furthermore, NLS residues P3 and P5 were stacked against the aromatic indole rings of  
20 Trp231, Trp184 and Trp142 on Importin-α, with the latter two being aligned in parallel.  
21 Interestingly, residues <sup>231</sup>MRRNR<sup>235</sup> also form part of the HNF1β DNA binding site identified  
22 by crystallography (Lu et al., 2007; PDB accession code 2H8R), which is analogous to the  
23 overlap between NLS and DNA binding seen in the androgen receptor (Cutress et al., 2008;  
24 PDB accession code 3BTR).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 Weaker difference density was also observed in the minor NLS-binding site on  
37 Importin-α1 and displayed a partly α-helical conformation (Figure 6B). Several atypical, minor  
38 site selective NLSs have previously been characterised structurally to adopt a similar  
39 topology (Chang et al., 2013; Nakada et al., 2015). Our pull-down experiments with  
40 xImportin-α protein mutants have demonstrated that the interaction was primarily via the  
41 major site on Importin-α and so the presence of the peptide density in the minor site was  
42 probably due to the high protein concentration used to produce the crystals together with the  
43 peptide having additional degrees of freedom that may not present in the context of the intact  
44 HNF1β protein. NLS residues Lys229 (P1), Met231 (P3) and Arg233 (P5) bound into apolar  
45 and shallow pockets consisting of aromatic tryptophan residues (Trp231, Trp184 and Trp142)  
46 stacked in parallel.  
47  
48  
49  
50  
51  
52  
53

54 Identifying the HNF1β NLS and its mode of interaction with Importin-α provides a  
55 basis of the design of therapeutic agents along the lines being investigated for other nuclear  
56 factors that aim to impair nuclear import to which cancer cells are often more sensitive  
57 (reviewed by Stelma et al. 2016). For example, Lin *et al.* (1995) developed a 41-residue  
58  
59  
60  
61  
62  
63  
64  
65

1 synthetic peptide (that contained the NLS of transcription factor NF- $\kappa$ B together with a cSN50  
2 cell membrane permeable motif) that inhibited the nuclear translocation of NF- $\kappa$ B and  
3 attenuated gene transcription in intact cells, but was not cytotoxic within the concentration  
4 range of the experiments (Torgerson et al. 1998).  
5  
6

7  
8 In summary, we have confirmed, using eGFP-tagged deletion constructs, that in the  
9 HNF1 $\beta$  transcription factor that is overexpressed in ovarian clear cell carcinoma, the  
10 sequence <sup>229</sup>KKMRRNR<sup>235</sup> is essential for its nuclear localisation in transduced cell lines.  
11 The HNF1 $\beta$ <sup>DBD</sup> was identified to interact with a spectrum of Importin- $\alpha$  isoforms by pull-down  
12 assays and we have further characterised the interaction between the putative NLS peptide  
13 and Importin- $\alpha$  using complementary biophysical techniques. Furthermore, we have  
14 determined the crystal structure of Importin- $\alpha$  in complex with the HNF1 $\beta$  NLS peptide to  
15 2.4 Å resolution. This information should facilitate the development of compounds that target  
16 the nuclear import of the transcription factor. We have shown that HNF1 $\beta$  interacts strongly  
17 with Importin- $\alpha$ 5 and selectively targeting this Importin-HNF1 $\beta$  interaction may open new  
18 avenues for the development of targeted therapeutics for ovarian clear cell carcinoma along  
19 the lines discussed by Stelma et al. (2016).  
20  
21  
22  
23  
24  
25  
26

#### 27 **Data deposition**

28 Co-ordinates and structure factors for the structure of the HNF1 $\beta$  peptides bound to Importin-  
29  $\alpha$  have been deposited at the Protein Data Bank (PDB) with accession code 5K9S.  
30  
31  
32  
33

#### 34 **Acknowledgements**

35 We thank our colleagues in Cambridge for their assistance, comments and criticisms. M.W.  
36 is funded by Cancer Research UK, Department of Chemistry at the University of Cambridge,  
37 School of the Physical Sciences and the Cambridge Cancer Centre. Funding in part was also  
38 provided by Medical Research Council Grant U105178939 to M.S. We would like to thank  
39 the Biorepository, Research Instrumentation, and Microscopy facilities at the Cancer  
40 Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson  
41 Way, Cambridge CB2 0RE, UK for assistance and Matthew Maggiolini for proofreading. We  
42 are grateful for the use of the Diamond Light Source Synchrotron (Harwell Science &  
43 Innovation Campus, Didcot, OX11 0DE, UK) for data collection.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1  
2 Adams, P.D., Afonine, P.V., *et al.* 2010. PHENIX: a comprehensive Python-based system for  
3 macromolecular structure solution. *Acta crystallogr. Section D, Biol. Crystallogr.* 66,  
4 213-221.
- 5 Anglesio, M.S., Carey, M.S., *et al.* 2011. Clear cell carcinoma of the ovary: A report from the  
6 first Ovarian Clear Cell Symposium, June 24th, 2010. *Gynecologic Oncology* 121,  
7 407-415.
- 8 Bach, I. and Yaniv, M. 1993. More potent transcriptional activators or a transdominant  
9 inhibitor of the HNF1 homeoprotein family are generated by alternative RNA  
10 processing. *EMBO J.* 12, 4229-4242.
- 11 Bach, I., Mattei, M.-G., *et al.* 1991. Two members of an HNF1 homeoprotein family are  
12 expressed in human liver. *Nucleic Acids Res.* 19, 3553-3559.
- 13 Bitler, B.G., Aird, K.M., *et al.* 2015. Synthetic lethality by targeting EZH2 methyltransferase  
14 activity in ARID1A-mutated cancers. *Nature Med.* 21, 231-238.
- 15 Bohn, S., Thomas, H., *et al.* 2003. Distinct Molecular and Morphogenetic Properties of  
16 Mutations in the Human HNF1 $\beta$  Gene That Lead to Defective Kidney Development.  
17 *J. American Soc. Nephrology*, 14, 2033-2041.
- 18 Chang, C.W., Counago, R.M., *et al.* 2013. Distinctive conformation of minor site-specific  
19 nuclear localization signals bound to importin-alpha. *Traffic*, 14, 1144-1154.
- 20 Chen, V.B., Arendall, W.B. 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J.,  
21 Murray, L.W., Richardson, J.S. and Richardson D.C.. (2010). MolProbity: all-atom  
22 structure validation for macromolecular crystallography. *Acta Crystallogr. Section D*,  
23 *Biol. Crystallogr.* 66, 12-21.
- 24 Conti, E., Uy, M., *et al.* 1998. Crystallographic Analysis of the Recognition of a Nuclear  
25 Localization Signal by the Nuclear Import Factor Karyopherin- $\alpha$ . *Cell*, 94, 193-204.
- 26 Cribbs, A., Kennedy, A., *et al.* 2013. Simplified production and concentration of lentiviral  
27 vectors to achieve high transduction in primary human T cells. *BMC Biotech.* 13, 98.
- 28 Cutress, M.L., Whitaker, H.C., Mills, I.G., Stewart, M. and Neal, D.E. (2008). Structural basis  
29 for the nuclear import of the human androgen receptor. *J. Cell Sci.* 121, 957-968.
- 30 Emsley, P., Lohkamp, B., *et al.* 2010. Features and development of Coot. *Acta Crystallogr.*  
31 *Section D, Biological Crystallogr.* 66, 486-501.
- 32 Evans, P.R. and Murshudov, G.N. 2013. How good are my data and what is the resolution?  
33 *Acta Crystallogr. Section D, Biological Crystallogr.* 69, 1204-1214.
- 34 Fontes, M.R.M., Teh, T., *et al.* 2000. Structural basis of recognition of monopartite and  
35 bipartite nuclear localization sequences by mammalian importin- $\alpha$ . *J. Molec. Biol.*  
36 297, 1183-1194.
- 37 Geoffrey, M. and Cooper, S.M., 2000. *The Cell, A Molecular Approach*. 2nd ed., Boston  
38 University.
- 39 Giesecke, A. and Stewart, M., 2010. Novel Binding of the Mitotic Regulator TPX2 (Target  
40 Protein for Xenopus Kinesin-like Protein 2) to Importin- $\alpha$ . *J. Biol. Chem.* 285, 17628-  
41 17635.
- 42 Gounaris, I., Charnock-Jones, D.S., *et al.* 2011. Ovarian clear cell carcinoma—bad  
43 endometriosis or bad endometrium? *J. Pathol.* 225, 157-160.
- 44 Hirota, K., Kajihara, Y.Y., Seiji Kanayama, Naoto Furukawa, Taketoshi Noguchi, Shoji Haruta,  
45 Shozo Yoshida, Toshiyuki Sado, Hidekazu Oi, Hiroshi Kobayashi, 2010. Clear cell  
46 carcinoma of the ovary: Potential pathogenic mechanisms (Review). *Oncology Rep.*  
47 23, 1193-1203.
- 48 Hodel, A.E., Harreman, M.T., *et al.* 2006. Nuclear localization signal receptor affinity  
49 correlates with in vivo localization in *Saccharomyces cerevisiae*. *J. Biological Chem.*  
50 281, 23545-23556.
- 51 Holvey, R.S., Valkov, E., *et al.* 2015. Selective Targeting of the TPX2 Site of Importin-alpha  
52 Using Fragment-Based Ligand Design. *Chem. MedChem.* 10, 1232-1239.
- 53 Jones, S., Wang, T.-L., *et al.* 2010. Frequent Mutations of Chromatin Remodeling Gene  
54 ARID1A in Ovarian Clear Cell Carcinoma. *Science*, 330, 228-231.
- 55 Kabsch, W., 2010. XDS. *Acta Crystallogr. Section D, Biol. Crystallogr.* 66, 125-132.
- 56 Kato, N., Sasou, S., *et al.* 2006. Expression of hepatocyte nuclear factor-1beta (HNF-1beta)  
57 in clear cell tumors and endometriosis of the ovary. *Modern Pathol.* 19, 83-89.
- 58 Kato, N., Tamura, G., *et al.* 2008. Hypomethylation of hepatocyte nuclear factor-1beta (HNF-  
59 1beta) CpG island in clear cell carcinoma of the ovary. *Virchows Arch.* 452, 175-180.
- 60  
61  
62  
63  
64  
65

- 1 Kato, N., Toukairin, M., *et al.* 2007. Immunocytochemistry for hepatocyte nuclear factor-1 $\beta$   
2 (HNF-1 $\beta$ ): A marker for ovarian clear cell carcinoma. *Diagnostic Cytopathol.* 35, 193-  
3 197.
- 4 Kobayashi, H., Yamada, Y. *et al.*, 2009. The role of hepatocyte nuclear factor-1beta in the  
5 pathogenesis of clear cell carcinoma of the ovary. *Internat. J. Gynecol. cancer*, 19,  
6 471-479.
- 7 Kobe, B. 1999. Autoinhibition by an internal nuclear localization signal revealed by the crystal  
8 structure of mammalian importin- $\alpha$ . *Nature Struct. Mol. Biol.* 6, 388-397.
- 9 Kuo, K.T., Mao, T.L. *et al.*, 2009. Frequent activating mutations of PIK3CA in ovarian clear  
10 cell carcinoma. *Am. J. Pathol.* 174, 1597-1601.
- 11 Kutner, R.H., Zhang, X.Y., *et al.* 2009. Production, concentration and titration of pseudotyped  
12 HIV-1-based lentiviral vectors. *Nature Protocols*, 4, 495-505.
- 13 Landen, C.N., Birrer, M.J., *et al.*, 2008. Early Events in the Pathogenesis of Epithelial Ovarian  
14 Cancer. *J. Clin. Oncol.* 26, 995-1005.
- 15 Lange, A., Mills, R.E., *et al.* 2007. Classical Nuclear Localization Signals: Definition, Function,  
16 and Interaction with Importin  $\alpha$ . *J. Biol. Chem.* 282, 5101-5105.
- 17 Lanitis, E., Dangaj, D., *et al.* 2012. Primary Human Ovarian Epithelial Cancer Cells Broadly  
18 Express HER2 at Immunologically-Detectable Levels. *PLoS One*, 7, e49829.
- 19 Liu, P., Khurana, A., *et al.* 2009. Regulation of HSulf-1 Expression by Variant Hepatic Nuclear  
20 Factor 1 in Ovarian Cancer. *Cancer Res.* 69, 4843-4850.
- 21 Lu, P., Rha, G.B., *et al.* 2007. Structural Basis of Disease-Causing Mutations in Hepatocyte  
22 Nuclear Factor 1 $\beta$ . *Biochemistry*, 46, 12071-12080.
- 23 Lu, P., Li, Y., *et al.* 2006. Crystallization of hepatocyte nuclear factor 1beta in complex with  
24 DNA. *Acta Crystallogr., Sect. F, Struct. Biol. Cryst. Commun.* 62, 525-529.
- 25 Marfori, M., Lonhienne, T.G., *et al.* 2012. Structural Basis of High-Affinity Nuclear Localization  
26 Signal Interactions with Importin- $\alpha$ . *Traffic*, 13, 532-548.
- 27 Marfori, M., Mynott, A., *et al.* 2011. Molecular basis for specificity of nuclear import and  
28 prediction of nuclear localization. *Biochim. Biophys. Acta*, 1813, 1562-1577.
- 29 Nakada, R., Hirano, H., *et al.* 2015. Structure of importin-alpha bound to a non-classical  
30 nuclear localization signal of the influenza A virus nucleoprotein. *Sci. Rep.* 5, 15055.
- 31 Neduva, V., and Russell, R.B. 2005. Linear motifs: Evolutionary interaction switches. *FEBS*  
32 *Lett.* 579, 3342-3345.
- 33 Rey-Campos, J., Chouard, T., *et al.* 1991. vHNF1 is a homeoprotein that activates  
34 transcription and forms heterodimers with HNF1. *EMBO J.* 10, 1445-1457.
- 35 Rosenfeld, M.G. 1991. POU-domain transcription factors: pou-er-ful developmental  
36 regulators. *Genes Devel.* 5, 897-907.
- 37 Ryan, A.K. and Rosenfeld, M.G. 1997. POU domain family values: flexibility, partnerships,  
38 and developmental codes. *Genes Devel.* 11, 1207-1225.
- 39 Sabherwal, N., Schneider, K.U., *et al.* 2004. Impairment of SHOX nuclear localization as a  
40 cause for Léri-Weill syndrome. *J. Cell Sci.* 117, 3041-3048.
- 41 Shen, H., Fridley, B.L., *et al.* 2013. Epigenetic analysis leads to identification of HNF1B as a  
42 subtype-specific susceptibility gene for ovarian cancer. *Nature Commun.* 4, 1628.
- 43 Stelma, T., Chi, A., van der Watt P.J., Verrico, A., Lavia, P. and Leaner, V.D. (2016).  
44 Targeting nuclear transporters in cancer: Diagnostic, prognostic and therapeutic  
45 potential. *IUBMB Life*, 68, 268-280.
- 46 Stewart, M. (2007). Molecular mechanism of the nuclear protein import cycle. *Nature Rev.*  
47 *Mol. Cell Biol.* 8,195-208.
- 48 Tan, D.S., Miller, R.E., *et al.* 2013. New perspectives on molecular targeted therapy in  
49 ovarian clear cell carcinoma. *British J. Cancer*, 108, 1553-1559.
- 50 Tan, D.S.P. and Kaye, S., 2007. Ovarian clear cell adenocarcinoma: a continuing enigma. *J.*  
51 *Clin. Pathol.* 60, 355-360.
- 52 Tsuchiya, A., Sakamoto, M., *et al.* 2003. Expression Profiling in Ovarian Clear Cell  
53 Carcinoma: Identification of Hepatocyte Nuclear Factor-1 $\beta$  as a Molecular Marker and  
54 a Possible Molecular Target for Therapy of Ovarian Clear Cell Carcinoma. *Am. J.*  
55 *Pathol.* 163, 2503-2512.
- 56 Wiegand, K.C., Shah, S.P., *et al.* 2010. ARID1A mutations in endometriosis-associated  
57 ovarian carcinomas. *New England J. Med.* 363, 1532-1543.
- 58  
59  
60  
61  
62  
63  
64  
65

1 Wu, G., Bohn, S., *et al.* 2004. The HNF1 $\beta$  transcription factor has several domains involved in  
2 nephrogenesis and partially rescues Pax8/lim1-induced kidney malformations.  
3 European J. Biochem. 271, 3715-3728.  
4 Yamaguchi, K., Mandai, M., *et al.* 2010. Identification of an ovarian clear cell carcinoma gene  
5 signature that reflects inherent disease biology and the carcinogenic processes.  
6 Oncogene, 29, 1741-1752.  
7 Yoshida, S., Furukawa, N., *et al.* 2009. Theoretical model of treatment strategies for clear cell  
8 carcinoma of the ovary: focus on perspectives. Cancer Treat. Rev. 35, 608-615.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## Figure Legends

**Figure 1** (A) HNF1 $\beta$  protein expression levels in CCC cell lines OVISE, JHOC5, JHOC7, JHOC9, SKOV3 and a negative control HGSOC cell line, PEO1; (B) Data is the mean (n = 3, three biological replicates) and SEM. Expression levels were normalised to the housekeeping protein GAPDH. The expression levels observed with JHOC5 (P < 0.005, \*\*\*) and OVISE (P < 0.05, \*) cells were significantly higher than PEO1.

**Figure 2** eGFP imaging of PEO1 transduced lines with (a) eGFP-HNF1 $\beta$  (Lv103) showed primarily nuclear localisation of the GFP signal, whereas cells transduced with eGFP (Lv105) alone (b) showed both nuclear and cytoplasmic localisation. In cells transduced with eGFP-HNF1 $\beta$  (Lv103) in which the seven-residue NLS had been deleted (c, Mutant 8) the eGFP signal was mislocalized to the cytoplasm. The scale bar represents 80  $\mu$ m. Images were taken on a Leica tandem confocal microscope using 20 x, 40 x and 60 x objectives.

**Figure 3** (A) Stable complex formation between  $\Delta$ IBB mouse Importin- $\alpha$ 1 (50 kDa) and HNF1 $\beta$  (27 kDa) during size exclusion chromatography. Fractions were analyzed by Coomassie stained SDS-PAGE (\* marks the peak for the complex); (B) GST pull-down with liberated HNF1 $\beta$  showing pull-downs using different  $\Delta$ IBB mouse Importin- $\alpha$  isoforms (red \*).

**Figure 4** (A) ITC titration for the binding of the HNF1 $\beta$  NLS peptide to  $\Delta$ IBB mImportin- $\alpha$ 1. Fitting the data (with  $\chi^2/\text{DoF} = 2.9 \times 10^4$ ) gave a  $K_d$  of  $13.6 \pm 1.5$  nM for  $0.94 \pm 0.13$  sites,  $\Delta H = -7.2 \pm 1.6$  kcal/mole,  $\Delta S = -1.7$  cal/mol/deg. (B) In pull-down assays, GST-HNF1 $\beta$  bound only the E mutant of  $\Delta$ IBB *Xenopus* Importin- $\alpha$ 1 (that impairs NLS binding at the minor site)  $M_r$  50 kD, but not the D mutant (that impairs binding at the major site) or ED double mutant (see Giesecke and Stewart, 2010).

**Figure 5** (A) Overview of the 2.4  $\text{\AA}$  crystal structure of the  $\Delta$ IBB mouse Importin- $\alpha$ 1:HNF1 $\beta^{\text{NLS}}$  complex. Two copies of the NLS peptide were observed where one was bound in the major site (blue) and another in the minor site (yellow). Each ARM repeat of mouse Importin- $\alpha$ 1 has been labelled with alternating colours. (B) Final *2Fo-Fc* electron density map around NLS peptide in major and minor binding sites on mouse Importin- $\alpha$ 1 contoured at  $1\sigma$ .

**Figure 6** Schematic illustration of the interactions of the HNF1 $\beta^{\text{NLS}}$  peptide in the major (A) and minor (B) sites on  $\Delta$ IBB mouse Importin- $\alpha$ 1.




Figure2

[Click here to download high resolution image](#)




Figure3  
[Click here to download high resolution image](#)





Figure 4

[Click here to download high resolution image](#)





Figure5  
[Click here to download high resolution image](#)





Figure6  
[Click here to download high resolution image](#)



*J. Struct. Biol. JSB-16-119 (revised)*

## **Structural and calorimetric studies demonstrate that the Hepatocyte Nuclear Factor 1 $\beta$ (HNF1 $\beta$ ) transcription factor is imported into the nucleus via a monopartite NLS sequence**

Mareike M. Wiedmann<sup>a,b</sup>, Shintaro Aibara<sup>c</sup>, David R. Spring<sup>a</sup>, Murray Stewart<sup>c\*</sup>, James D. Brenton<sup>b</sup>

<sup>a</sup> Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK

<sup>b</sup> Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge CB2 0RE, UK

<sup>c</sup> MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB2 0QH, UK

\* To whom correspondence should be addressed: Dr Murray Stewart, MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB2 0QH, UK; E-mail: [ms@mrc-lmb.cam.ac.uk](mailto:ms@mrc-lmb.cam.ac.uk); Fax: +44 1223 267000; Phone: +44 1223 267074.



## Abstract

The transcription factor hepatocyte nuclear factor 1 $\beta$  (HNF1 $\beta$ ) is ubiquitously overexpressed in ovarian clear cell carcinoma (CCC) and is a potential therapeutic target. To explore potential approaches that block HNF1 $\beta$  transcription we have identified and characterised extensively the nuclear localisation signal (NLS) for HNF1 $\beta$  and its interactions with the nuclear protein import receptor, Importin- $\alpha$ . Pull-down assays demonstrated that the DNA binding domain of HNF1 $\beta$  interacted with a spectrum of Importin- $\alpha$  isoforms and deletion constructs tagged with eGFP confirmed that the HNF1 $\beta$  <sup>229</sup>KKMRRNR<sup>235</sup> sequence was essential for nuclear localisation. We further characterised the interaction between the NLS and Importin- $\alpha$  using complementary biophysical techniques and have determined the 2.4 Å resolution crystal structure of the HNF1 $\beta$  NLS peptide bound to Importin- $\alpha$ . The functional, biochemical, and structural characterisation of the nuclear localisation signal present on HNF1 $\beta$  and its interaction with the nuclear import protein Importin- $\alpha$  provide the basis for the development of compounds targeting transcription factor HNF1 $\beta$  via its nuclear import pathway.

**Keywords:** Importin- $\alpha$ , nuclear import pathway, nuclear localisation signal sequence (NLS), Hepatocyte nuclear factor-1 $\beta$  (HNF1 $\beta$ ), site-directed mutagenesis, X-ray crystallography, isothermal titration calorimetry.

**Abbreviations:** CCC - Ovarian clear cell carcinoma; HNF1 $\beta$  - hepatocyte nuclear factor 1 $\beta$ ; HNF1 $\beta$ <sup>DBD</sup> - HNF1 $\beta$  DNA binding domain; NLS - nuclear localization signal; IBB - Importin- $\beta$ -binding domain; mImportin- $\alpha$  - mouse Importin- $\alpha$ ; xImportin- $\alpha$  - *Xenopus* Importin- $\alpha$ ; GST - glutathione S-transferase.

## Introduction

Ovarian clear cell carcinoma (CCC) accounts for 5-10% of ovarian cancer cases (Anglesio et al., 2011; Kato et al., 2008). Prognosis for patients with advanced stage or for relapsed disease is poor because of intrinsic resistance to platinum based chemotherapy and the lack of targeted therapies available (Tan et al., 2013; Tan and Kaye, 2007), although Bitler *et al.* (2015) have recently discovered a way of targeting cancers with *ARID1A* mutations by targeting EZH2 methyltransferase activity. Common mutations in CCC include loss of function mutations in the chromatin remodeling gene *ARID1A* in 46–57% of cases (Jones et al., 2010; Wiegand et al., 2010), activating mutations in *PIK3CA* (Kuo et al., 2009) in 33–46% of cases, and loss of *PTEN* in 20% of cases (Anglesio et al., 2011; Landen et al., 2008; Tan and Kaye, 2007). Overexpression of the HNF1 $\beta$  transcription factor is the most important clinical immunohistochemical marker for the disease because it is ubiquitously overexpressed in CCC, both at the mRNA and protein level (Hirota Kajihara, 2010; Kato et al., 2006; Tsuchiya et al., 2003; Yamaguchi et al., 2010). In CCC the *HNF1B* gene is upregulated by hypomethylation of its CpG island whereas, in high grade serous ovarian cancer, HNF1 $\beta$  expression is silenced via hypermethylation (Kato et al., 2008; Shen et al., 2013a), suggesting that HNF1 $\beta$  has a loss of function (tumour suppressor) role in high grade serous ovarian cancer but a gain of function (oncogenic) role in CCC (Gounaris et al., 2011; Shen et al., 2013b). This hypothesis is supported by the observation that nearly half of the overexpressed genes identified in CCC are downstream targets of HNF1 $\beta$  (Kobayashi et al., 2009; Yoshida et al., 2009). Evidence that targeting HNF1 $\beta$  might have utility was provided by Liu *et al.* (Liu et al., 2009) who showed that downregulation of HNF1 $\beta$  increased cisplatin- and paclitaxel-mediated cytotoxicity.

Transcription factor HNF1 $\beta$  (also known as vHNF1, vAPF, LF-B3 and Tcf2) is expressed in the liver, digestive tract, pancreas and the kidneys, where it plays a crucial role in early differentiation (Lu et al., 2007). Sequence-specific DNA binding is mediated by a bipartite motif that consists of a POU homeodomain (POU<sub>H</sub>) and a POU specific domain (POU<sub>S</sub>) (Rosenfeld, 1991; Ryan and Rosenfeld, 1997). HNF1 $\beta$  has 70% sequence homology to HNF1 $\alpha$  and both proteins are atypical members of the POU transcription factor family and bind DNA as both homo- and heterodimers (Bach et al., 1991; Rey-Campos et al., 1991). Human HNF1 $\beta$  is constructed from three domains: the dimerization domain, which is further stabilised by the dimerization cofactor of HNF1 (DcoH), the transactivation domain, which is involved in binding transcriptional co-activators [15], and the POU DNA binding domain (HNF1 $\beta$ <sup>DBD</sup>).

Transcription factors together with histones, DNA polymerase, RNA polymerase and many other proteins, have specific amino acid sequences, termed nuclear localisation signals (NLSs), that are recognised by members of the karyopherin family that facilitate their nuclear import (reviewed by Lange et al., 2007). Many NLS sequences are recognised in the

cytoplasm by a heterodimeric transport carrier complex composed of Importin- $\beta$  (also known as Karyopherin- $\beta$ 1) and Importin- $\alpha$  (reviewed by Stewart, 2007). Nuclear pore complexes (NPCs) are the channels through which macromolecules, such as proteins and RNA, are transported between the cytoplasm and nucleus (reviewed by Stewart, 2007). Small molecules and proteins (< 40 kDa) can pass through NPCs by passive diffusion, but larger proteins require carriers to overcome the NPC physical barrier. The autoinhibitory Importin- $\beta$  binding (IBB) domain of Importin- $\alpha$  (Kobe, 1999) binds to Importin- $\beta$  in the cytoplasm, enabling classical NLSs (cNLS) to bind to Importin- $\alpha$  either via a major site, a minor site, or both (Fontes et al., 2000; Lange et al., 2007). There are two types of cNLS that are recognised by Importin- $\alpha$  that consist of either a single cluster (monopartite) or two clusters (bipartite) of positively charged residues, primarily lysines or arginines, that assume an ordered state once bound by Importin- $\alpha$  (reviewed by Lange et al. 2007; Marfori et al., 2011). Monopartite cNLSs are exemplified by the simian virus 40 large T-antigen (SV40) NLS  $^{126}\text{PKKKRRV}^{132}$  (Lange et al., 2007). The cargo:carrier heterotrimer is then translocated into the nucleus in an energy dependent manner powered by RanGTPase (reviewed by Stewart, 2007). In the nucleus, RanGTP then binds Importin- $\beta$  leading ultimately to the release of the cargo.

To function, HNF1 $\beta$  needs to be translocated to the cell nucleus and so we have investigated and characterized the putative NLS that has been proposed to lie between the two POU domains of HNF1 $\beta$ . This putative NLS was identified in domain swapping experiments and studies of nephrogenesis, in which truncated GFP-HNF1 $\beta$  fusion constructs retaining the POU<sub>H</sub> domain showed exclusive nuclear localisation in transfected HeLa cells (Bohn et al., 2003; Wu et al., 2004). However, the precise location of the HNF1 $\beta$  NLS has not been defined. Because of the potential importance of HNF1 $\beta$  as a target in diseases such as CCC, we have identified and characterized extensively its NLS and its interactions with Importin- $\alpha$ . We demonstrate that the sequence  $^{229}\text{KKMRRNR}^{235}$  in HNF1 $\beta^{\text{DBD}}$  is responsible for the nuclear import of the protein. Several eGFP-constructs of HNF1 $\beta$  were generated and Importin- $\alpha$  binding of the HNF1 $\beta^{\text{DBD}}$  was assessed by both pull-down experiments and ITC. We also determined the crystal structure of the HNF1 $\beta$  NLS peptide bound to Importin- $\alpha$ . The **identification and** structural characterisation of the HNF1 $\beta$  NLS and its interaction with the nuclear import protein Importin- $\alpha$  provides a basis for the development of inhibitors targeting the nuclear import of transcription factor HNF1 $\beta$  **along the lines suggested by Stelma et al. (2016).**

## **Materials and Methods**

### ***Mammalian cell culture***

HEK293T cells were cultivated in Dulbecco's Modified Eagle medium (DMEM) (1X) supplemented with 5% foetal bovine serum (FBS) (Invitrogen) and 0.5% penicillin/streptomycin (P/S). CCC cell lines PEO1, JHOC5, JHOC7, JHOC9, OVISE and SKOV3 cells were grown in RPMI 1640 medium (1X) supplemented with 10% FBS and 1% P/S. Normal Ovarian Surface Epithelial (IOSE) cells were cultivated in NOSE-CM: MCDB 105/medium 199 (1:1 ratio, Sigma Aldrich), 15% FBS, 10 ng/ml EGF (Invitrogen), 0.5 µg/ml hydrocortisone (Sigma Aldrich), 5 µg/ml insulin (Sigma Aldrich), 34 µg protein/ml BPE (Invitrogen). All cell lines were maintained at 37°C in 5% CO<sub>2</sub> and were mycoplasma tested on a regular basis (Biorepository Core, CRUK CI, Cambridge). Cell counts were conducted using a Vi-CELL Cell Viability Analyzer.

### ***Protein extraction from mammalian cells***

Cell pellets were washed with phosphate buffered saline (PBS) and 200 µl protein lysis buffer (50 mM Tris pH 8.0, 150 mM NaCl, 5 mM EDTA, 0.5% Igepal, to which 2 tablets/100ml of protease inhibitor cocktail tablet (Roche)) was added. The mixtures were incubated on ice for 30 min, lysed by syringing four times using a 26 G needle, and centrifuged at 14,800 × g for 10 min at 4°C. Protein concentrations were measured using the DirectDetect IR spectrometer (Merck Millipore) according to the manufacturer's instructions.

### ***Western blotting***

Denatured protein extracts were separated using NuPAGE Novex 4-20% Tris-Glycine gels and transferred to a Millipore Immobilon FL PVDF membrane (Invitrogen). Primary antibodies were used as follows: goat anti-HNF1β (sc-7411, polyclonal, Santa Cruz Biotechnology, 1:1,000) and rabbit anti-GAPDH (G9545, 14C10, 1:5,000, Cell Signalling Technology). An Odyssey Infrared Imaging System (Li-Cor) and associated secondary antibodies: donkey anti-goat (800) (1:15,000) and donkey anti-rabbit (800) (1:5,000) were used to detect material. **The expression levels observed with different cells were evaluated using GraphPad Prism version 6.0 for Mac, GraphPad Software, La Jolla California USA, [www.graphpad.com](http://www.graphpad.com). Analysis of variance rejected the null hypothesis that the expression levels were equal ( $P < 0.002$ ) and Šídák's multiple comparison modification of Student's t-test was used to evaluate the significance of the levels seen with different cells relative to that seen with the control HGSOC cell line.**

### ***Confocal microscopy: eGFP imaging and immunofluorescence assay***

For immunofluorescence (IF), cells were fixed with 4% **paraformaldehyde** in PBS for 10 min at room temperature, rinsed in **Tris buffered saline (TBS)** for 2 × 5 min and permeabilized in TBS-0.5% Triton X-100 for 10 min. Fixed cells were then rinsed in TBS-0.1% Triton X-100 for 3 × 3 min and blocked in 10% goat serum in TBS for 30 min. HNF1β was stained with anti-HNF1β (SAB1406512, mouse polyclonal, 1:300, Sigma Aldrich) overnight at 4°C. Cells were

washed with TBS-0.1% Triton X-100 (Fisher Scientific) for 3 × 5 min and the secondary antibody (AlexaFluor 568, goat anti-mouse IgG (H+L) 1:1,000, Invitrogen) was added and incubated for 60 min diluted in antibody dilution solution consisting of TBS - 0.1% Triton X-100, 2% Bovine Serum Albumin (BSA) (Cell Signaling Technology) and 0.1% sodium azide. All experiments included an unstained control, a “secondary only” control, and a negative control using the PEO1 cell line. Cells transduced with eGFP-HNF1 $\beta$ , the HNF1 $\beta$ <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant, or the control eGFP constructs, were fixed with 4% paraformaldehyde in PBS for 10 min at room temperature, then rinsed in TBS for 2 × 5 min. Nuclei were stained using DAPI (1  $\mu$ g/ml in TBS) for 10 min. Cells were washed in TBS-0.1% Triton X-100, rinsed in TBS and stored in TBS-0.1% Triton X-100. For microscopy imaging, cover slips were drained, mounted and sealed using Prolong Gold (Invitrogen) and glass slides (Thermo Specific). Slides were left to dry overnight at room temperature in the absence of light and were then stored at 4°C. Cells were imaged using a Leica tandem confocal microscope.

#### **Site directed mutagenesis to generate HNF1 $\beta$ NLS deletion construct**

Lv103 (EX-F0366-Lv103, Genecopoeia) is a lentiviral transfer vector containing an eGFP-HNF1 $\beta$  fusion coding sequence. Lv105 (EX-EGFP-Lv105, Genecopoeia) only contains the eGFP coding sequence. The plasmids were confirmed by sequencing (GATC, Konstanz, Germany) before and after mutagenesis (Lv103 Fw: 5'-CCGACAACCACTACCTGA-3'; Rv: 5'-ATTGTGGATGAATACTGCC-3' and Lv105 Fw: 5'-ATCCACGCTGTTTTGACC-3'; Rv: 5'-AATACTGCCATTTGTCTCG-3').

Mutagenesis experiments were conducted using the Q5 site directed mutagenesis kit (NEB). The <sup>229</sup>KKMRRNR<sup>235</sup> deletion construct was generated using mutagenesis primers (Fw: 5'-TTCAAATGGGGGCCGC-3'; Rv: 5'-GTTGGTGGGCTCAGAGCAG-3'). Plasmids were then transformed in *E. coli* (One Shot® Stbl3™, Invitrogen) and streaked on Amp-containing agar plates. Single colonies were picked and tested for plasmid containing colonies by PCR. Plasmids were extracted by Plasmid Mini Prep (Qiagen) and quantified (Qubit).

#### **Lentivirus production and transduction**

The general protocol devised by Cribbs *et al.* (2013) was used for lentivirus production. The transfer vectors Lv103, Lv105, the <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant, and plasmids pRVS-Rev, pVSV-G, and p-MDLg-pRRE were verified by restriction digest. For each transfection sample, 16  $\mu$ g transfer vector, 10.4  $\mu$ g pMDL/pRRE, 4  $\mu$ g pRSV-Rev and 5.6  $\mu$ g pVSV-G were used. After 24 h, expression of GFP protein was observed in the GFP control virus HEK293T sample. Virus containing supernatant was harvested posttransfection according to the protocol by Kutner *et al.* (2009). The number of transducing units (TU) was determined by

flow cytometry analysis with GFP as the reporter protein. For titration,  $1 \times 10^5$  cells per well were seeded in a 12-well plate and 0.5 ml DMEM with 5% FBS added. Virus of respective concentration was pipetted over the HEK293T cells. Duplicate virus dilutions of 1:500, 1:1000, 1:2000, 1:5000 were added. An untreated control was included. The medium was changed 24 h posttransfection. 72 h posttransfection cells were harvested by trypsinization. Cells were collected by centrifugation at  $500 \times g$  for 5 min at room temperature. The supernatant was discarded and cell pellets were resuspended in PBS. A green fluorescent protein propidium iodide (GFP PI) based assay for flow cytometric measurement of transfection efficiency and cell viability was performed (LSR-II machine). The titre was calculated from dilutions that gave 1–40% GFP-positivity and averaged subsequently using the following formula:

$$\text{Titre} \left( \frac{\text{TU}}{\text{ml}} \right) = \frac{(\text{frequency of GFP-positive cells}) \times \text{no. cells plated} \times \text{dilution factor}}{\text{volume of inoculum}}$$

where, frequency of GFP-positive cells is the percentage of cells that are positive for GFP divided by 100 (acceptable range: 0.01–0.40), dilution factor is the dilution of the virus stock used and volume of inoculum is the total volume transduced. The titre was calculated to be  $2.26 \times 10^8 \frac{\text{TU}}{\text{ml}}$ .

For transduction of the PEO1 cell line,  $1 \times 10^6$  cells per well were seeded in a 12-well plate to which 0.5 ml of the respective medium was added. Virus was added so that the desired multiplicity of infection (MOI) was obtained. Polybrene 0.5-10  $\mu\text{g/ml}$  was added to increase transduction efficiency. Transduction efficiencies were determined by flow cytometry analysis using an LSR-II machine. For selection of transduced PEO1 lines, a puromycin kill curve was constructed (data not shown) to determine the optimum concentration of puromycin required to kill all untransduced cells while minimising toxicity effects in transduced cells. A puromycin concentration of 0.1  $\mu\text{g/ml}$  was determined which was used in all experiments.

### **Quick change mutagenesis to generate GST-tagged HNF1 $\beta$ <sup>DBD</sup> and mImportin $\alpha$ 1 mutants**

The cloning of the DNA binding domain of HNF1 $\beta$  into the pGEX-TEV plasmid was conducted as described by Lu *et al.* (2006). The plasmid was confirmed by sequencing by Source Bioscience (Cambridge, UK) using PGEX5 and PGEX3 primers (Source Bioscience).

### **Protein expression in bacterial cells and purification**

All proteins were expressed in *E. coli* BL21 (DE3) CodonPlus-RIL cells using IPTG induction over 18 h at 18°C. The cells were harvested by centrifugation and resuspended in 50 mM

Tris/HCl pH 8.0, 500 mM NaCl, 5 mM DTT for GST-tagged constructs or 50 mM Tris/HCl pH 8.0, 500 mM NaCl, 20 mM Imidazole pH 8.0 for His<sub>6</sub>-tagged proteins. The *E. coli* were lysed by two passes through an Emulsiflex C3 system (AVESTIN) running at a pressure of 15,000 psi and the lysates clarified by centrifugation at 48,000 × g. Either Ni-NTA agarose resin (QIAGEN) or glutathione Sepharose 4B resin (GE Healthcare) was added to the clarified lysate containing His<sub>6</sub>-tagged proteins or GST-tagged proteins. The resin was pooled and packed into a gravity filtration column and washed extensively with their respective lysis buffers. 1 mg of TEV protease (purified in-house) was added to the washed resin and incubated for 18 h at 4°C. The tag-free protein of interest was then collected from the flow-through and subjected to the next purification step. HNF1β<sup>DBD</sup> was purified further by heparin affinity chromatography using a HiTrap Heparin HP column (GE Healthcare Life Sciences). HNF1β<sup>DBD</sup> was eluted from the column using a 0.1 - 1 M NaCl gradient. All proteins were purified to homogeneity by size exclusion chromatography using a HiLoad™ 26/60 Superdex™ 75 prep grade column (GE Healthcare) pre-equilibrated in 20 mM HEPES buffer pH 8.0, 200 mM NaCl. Fractions containing pure and homogenous proteins identified by SDS-PAGE analysis were pooled and concentrated using an Amicon Ultra 15 centrifugation filtration unit. Protein concentrations were measured using a Nanodrop (Thermo) microspectrophotometer and small aliquots of the protein were flash frozen in liquid nitrogen and stored at -80°C until required.

### ***GST pull-downs***

After bacterial cell lysis, cell pellets expressing the two proteins of interest were mixed and pull-down experiments were conducted as described above for glutathione affinity chromatography and were visualised by SDS-page analysis.

### ***Isothermal calorimetry***

The K<sub>d</sub> of mImportin-α1 with the HNF1β NLS peptide (Ac-TNKKMRRNRFK-NH<sub>2</sub>, purchased from Insight Biotechnology) was determined by isothermal titration calorimetry (ITC) using a MicroCal™ iTC200 (GE Healthcare Life Sciences). The protein and peptide samples were made up in the identical buffer (20 mM Hepes pH 8, 200 mM NaCl). 350 μl of protein sample (0.02 mM) and 200 μl of ligand (0.2 mM) were prepared for each experiment. Sample concentrations were confirmed by amino acid analysis (PNAC Facility, Biochemistry Department, Cambridge University). All experiments were conducted at 25°C. The data was analysed using the Origin™ Software (MicroCal) using a one site binding model.

### ***Protein crystallography***

Diffraction quality crystals of the Importin-α1<sup>Δ1BB</sup>:HNF1β<sup>NLS</sup> complex were obtained by sitting drop vapour diffusion where 200 nl of 10% PEG 8K, 0.09 M NPS, 20% ethylene glycol, 0.1 M Buffer 1 (Morpheus Screen, Molecular Dimensions) was mixed with 200 nl of Importin-

$\alpha 1^{\Delta IBB} : HNF1\beta^{NLS}$ . The Importin- $\alpha 1^{\Delta IBB} : HNF1\beta^{NLS}$  complex was produced by mixing Importin- $\alpha 1^{\Delta IBB}$  and HNF1 $\beta^{NLS}$  peptide (Ac-TNKKMRRMRFK-NH<sub>2</sub>, Insight Biotechnology) at a 1:1.1 molar ratio and a concentration of 6.7 mg/ml.

Crystals were cryoprotected in the mother liquor supplemented with 20% glycerol and cooled by plunging into liquid nitrogen. X-ray diffraction data were collected on beamline I03 at the Diamond Light Source (Didcot, UK). Reflections were indexed and integrated using *DIALS* as implemented in *Xia2* (Kabsch, 2010) and then scaled and merged in *AIMLESS*, ensuring a completeness of > 98% in the outermost shell while maintaining  $CC_{1/2} > 0.3$  (Evans and Murshudov, 2013). The structure was solved by molecular replacement using *Phaser* with the structure of  $\Delta IBB$ -mImportin- $\alpha 1$  complexed with a minor site small molecule inhibitor (PDB ID: 4U54 - Holvey et al., 2015) to **avoid introducing model bias into the NLS binding sites**. Iterative cycles of rebuilding using *COOT* (Emsley et al., 2010) and refinement using *PHENIX* (Adams et al., 2010) were used to generate the final model (Table1) **that had an R-factor of 18.4% (R-free = 22.0%) and a MolProbity score of 0.95 (100th percentile)**.

## Results and Discussion

### ***HNF1 $\beta$ is overexpressed in CCC cell lines***

HNF1 $\beta$  protein expression in different CCC cell lines was determined by Western blotting (Figure 1) using the HGSOc cell line PEO1 (which does not express HNF1 $\beta$ ) as a negative control. The CCC lines OVISe, JHOC5, JHOC7, JHOC9 and SKOV3 cell lines were used as models of CCC that overexpress HNF1 $\beta$ . Several splice variants of HNF1 $\beta$  are known to contain different isoforms of the C-terminal domain that is responsible for activation of transcription. These isoforms act as transdominant repressors (Bach and Yaniv, 1993). Only the 61 kDa isoform of HNF1 $\beta$  was detected in our CCC cell lines (Figure 1) (Bach and Yaniv, 1993; Tsuchiya et al., 2003). HNF1 $\beta$  expression levels **varied considerably between these cell lines**. Analysis of variance of the data in Figure 1 rejected the null hypothesis that the expression levels were equal ( $P < 0.002$ ) and the higher level seen in JHOC7 compared with PEO1 cells was significant at the 0.5% level (using Sidak's modification for multiple comparisons), consistent with hypomethylation of the HNF1 $\beta$  CpG island (Kato et al., 2008; Kato et al., 2007).

### ***Identification of the HNF1 $\beta$ nuclear localisation signal***

To confirm the identity of the NLS, the PEO1 cell line was transduced with lentiviruses Lv103 (eGFP-HNF1 $\beta$ ), Lv105 (eGFP) and the <sup>229</sup>KKMRRNR<sup>235</sup> deletion mutant. **In cells transduced with eGFP-HNF1 $\beta$ , the fusion protein (eGFP-HNF1 $\beta$  (Lv103) - Figure 2A(Mutant 8)) was primarily nuclear, whereas cells transduced with Lv105 (eGFP) showed nuclear and cytoplasmic localisation of the eGFP protein (eGFP (Lv105) - Figure 2B). Deletion of the**



seven-residue <sup>229</sup>KKMRRNR<sup>235</sup> NLS sequence in HNF1 $\beta$  resulted in its becoming mislocalized to the cytoplasm (Mutant 8, Figure 2C), confirming the importance of this sequence for the nuclear import of HNF1 $\beta$ . HNF1 $\alpha$  has an analogous KKGRRN sequence that differs only at residue 231, where Met is changed to Gly (Wu et al., 2004) and which probably also functions in a similar manner. Interestingly, all transduced cells (that were selected using optimised puromycin concentrations) were apoptotic within 24 h. We speculate that re-expression of a transcription factor that is epigenetically silenced in HGSOc cell line PEO1 (Shen et al., 2013a) may have a negative effect on proliferation, but further work is necessary to determine the effects of re-expression.

### ***HNF1 $\beta$ <sup>DBD</sup> binds primarily to the major site on Importin- $\alpha$***

Because the sequence of the HNF1 $\beta$  NLS is similar to that of classical monopartite NLSs (Lange et al., 2007), HNF1 $\beta$  is likely to be imported by Importin- $\alpha$  and to interact with its major NLS-binding site rather than being imported by direct binding to Karyopherin- $\beta$  (Fontes et al., 2000). To test whether HNF1 $\beta$  and mImportin- $\alpha$  formed a complex, recombinantly expressed and purified HNF1 $\beta$ <sup>DBD</sup> and mImportin- $\alpha$ 1 in which the IBB domain had been deleted ( $\Delta$ IBB-mImportin- $\alpha$ 1) were mixed together and analysed by gel filtration chromatography (Figure 3A), in which  $\Delta$ IBB-mImportin- $\alpha$ 1 and HNF1 $\beta$ <sup>DBD</sup> co-eluted as a stoichiometric complex as observed by Coomassie blue stained SDS-PAGE analysis. To establish whether HNF1 $\beta$  interacts with other Importin- $\alpha$  isoforms, pull-down experiments using different  $\Delta$ IBB-mImportin- $\alpha$ 1 isoforms were performed (Figure 3B). Crude extracts of GST-tagged HNF1 $\beta$  and  $\Delta$ IBB-mImportin- $\alpha$ 1 isoforms were mixed, then bound to glutathione resin, washed, and eluted by cleaving between GST and HNF1 $\beta$ <sup>DBD</sup> using TEV protease. Both HNF1 $\beta$ <sup>DBD</sup> and  $\Delta$ IBB-mImportin- $\alpha$ 1 were released, confirming that they were interacting (Figure 3B). HNF1 $\beta$  interacted more strongly with  $\Delta$ IBB-mImportin- $\alpha$ 5 and less strongly with  $\Delta$ IBB-mImportin- $\alpha$ 3, - $\alpha$ 4 and - $\alpha$ 7 isoforms. The  $K_d$  for the interaction of  $\Delta$ IBB-mImportin- $\alpha$ 1 with the HNF1 $\beta$  NLS peptide was determined by isothermal calorimetry (ITC) (Figure 4A) to be  $13.6 \pm 1.54E-3 \mu\text{M}$  which was in the range commonly observed (Hodel et al., 2006). In summary, HNF1 $\beta$  binds at least two  $\Delta$ IBB-mImportin- $\alpha$  isoforms strongly and some other isoforms more weakly.

Three different *Xenopus* Importin- $\alpha$ 1 (xImportin- $\alpha$ 1) mutants (Giesecke and Stewart, 2010) were used to define where the HNF1 $\beta$  NLS bound on Importin- $\alpha$  (Figure 4B). These mutants contained point mutations that prevent binding via the major (D), minor (E), or both (ED) sites and also have the IBB domain removed to prevent auto-inhibition (Kobe, 1999). HNF1 $\beta$ <sup>DBD</sup> bound to the E mutant but not the D or ED mutant of  $\Delta$ IBB-xImportin- $\alpha$ 1 (Figure 4B), indicating that HNF1 $\beta$  binds primarily via the major site, consistent with its similarity to the SV40 NLS (Marfori et al., 2011).

### ***Structural characterisation of the interaction of mImportin- $\alpha$ 1 with the HNF1 $\beta$ NLS***

To complement the functional and biochemical data, we also determined the 2.4 Å resolution crystal structure of the HNF1β NLS peptide bound to ΔIBB-mImportin-α1 (Figure 5A). After initial rounds of refinement, strong difference density, corresponding to the peptide was found in both the major NLS-binding site on Importin-α1 together with weaker density at the minor site (Figures 5B and C, respectively).

At the major site, the HNF1β NLS peptide backbone interacted with Importin-α1 via a set of conserved Asn residues on the import protein by H-bonding (Figure 6A). The NLS residues are conventionally assigned to positions P1-P5 (Figure 6). Key residues on Importin-α involved for the interaction with the NLS were Asn-235, 188 and 146, the latter two of which are involved in bidentate H-bonding with the NLS peptide backbone at P3 and P5. The aliphatic portions of the NLS peptide side chains were located in shallow pockets along the surface of Importin-α, but, as evident from Figure 6A, the P2 residue, Lys230, forms many critical interactions, mostly by forming salt bridges with negatively charged residues on Importin-α. There were three interactions of the protonated nitrogen on the P2 lysine side chain with Asp192, Gly150 and Thr155. Residue P5 interacted with the side-chain of Gln181. Furthermore, NLS residues P3 and P5 were stacked against the aromatic indole rings of Trp231, Trp184 and Trp142 on Importin-α, with the latter two being aligned in parallel. Interestingly, residues <sup>231</sup>MRRNR<sup>235</sup> also form part of the HNF1β DNA binding site identified by crystallography (Lu et al., 2007; PDB accession code 2H8R), which is analogous to the overlap between NLS and DNA binding seen in the androgen receptor (Cutress et al., 2008; PDB accession code 3BTR).

Weaker difference density was also observed in the minor NLS-binding site on Importin-α1 and displayed a partly α-helical conformation (Figure 6B). Several atypical, minor site selective NLSs have previously been characterised structurally to adopt a similar topology (Chang et al., 2013; Nakada et al., 2015). Our pull-down experiments with xImportin-α protein mutants have demonstrated that the interaction was primarily via the major site on Importin-α and so the presence of the peptide density in the minor site was probably due to the high protein concentration used to produce the crystals together with the peptide having additional degrees of freedom that may not present in the context of the intact HNF1β protein. NLS residues Lys229 (P1), Met231 (P3) and Arg233 (P5) bound into apolar and shallow pockets consisting of aromatic tryptophan residues (Trp231, Trp184 and Trp142) stacked in parallel.

Identifying the HNF1β NLS and its mode of interaction with Importin-α provides a basis of the design of therapeutic agents along the lines being investigated for other nuclear factors that aim to impair nuclear import to which cancer cells are often more sensitive (reviewed by Stelma et al. 2016). For example, Lin *et al.* (1995) developed a 41-residue

synthetic peptide (that contained the NLS of transcription factor NF- $\kappa$ B together with a cSN50 cell membrane permeable motif) that inhibited the nuclear translocation of NF- $\kappa$ B and attenuated gene transcription in intact cells, but was not cytotoxic within the concentration range of the experiments (Torgerson et al. 1998).

In summary, we have confirmed, using eGFP-tagged deletion constructs, that in the HNF1 $\beta$  transcription factor that is overexpressed in ovarian clear cell carcinoma, the sequence <sup>229</sup>KKMRRNR<sup>235</sup> is essential for its nuclear localisation in transduced cell lines. The HNF1 $\beta$ <sup>DBD</sup> was identified to interact with a spectrum of Importin- $\alpha$  isoforms by pull-down assays and we have further characterised the interaction between the putative NLS peptide and Importin- $\alpha$  using complementary biophysical techniques. Furthermore, we have determined the crystal structure of Importin- $\alpha$  in complex with the HNF1 $\beta$  NLS peptide to 2.4 Å resolution. This information should facilitate the development of compounds that target the nuclear import of the transcription factor. We have shown that HNF1 $\beta$  interacts strongly with Importin- $\alpha$ 5 and selectively targeting this Importin-HNF1 $\beta$  interaction may open new avenues for the development of targeted therapeutics for ovarian clear cell carcinoma [along the lines discussed by Stelma et al. \(2016\)](#).

#### **Data deposition**

Co-ordinates and structure factors for the structure of the HNF1 $\beta$  peptides bound to Importin- $\alpha$  have been deposited at the Protein Data Bank (PDB) with accession code 5K9S.

#### **Acknowledgements**

We thank our colleagues in Cambridge for their assistance, comments and criticisms. M.W. is funded by Cancer Research UK, Department of Chemistry at the University of Cambridge, School of the Physical Sciences and the Cambridge Cancer Centre. Funding in part was also provided by Medical Research Council Grant U105178939 to M.S. We would like to thank the Biorepository, Research Instrumentation, and Microscopy facilities at the Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge CB2 0RE, UK for assistance and Matthew Maggolini for proofreading. We are grateful for the use of the Diamond Light Source Synchrotron (Harwell Science & Innovation Campus, Didcot, OX11 0DE, UK) for data collection.

## References

- Adams, P.D., Afonine, P.V., *et al.* 2010. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallogr. Section D, Biol. Crystallogr.* 66, 213-221.
- Anglesio, M.S., Carey, M.S., *et al.* 2011. Clear cell carcinoma of the ovary: A report from the first Ovarian Clear Cell Symposium, June 24th, 2010. *Gynecologic Oncology* 121, 407-415.
- Bach, I. and Yaniv, M. 1993. More potent transcriptional activators or a transdominant inhibitor of the HNF1 homeoprotein family are generated by alternative RNA processing. *EMBO J.* 12, 4229-4242.
- Bach, I., Mattei, M.-G., *et al.* 1991. Two members of an HNF1 homeoprotein family are expressed in human liver. *Nucleic Acids Res.* 19, 3553-3559.
- Bitler, B.G., Aird, K.M., *et al.* 2015. Synthetic lethality by targeting EZH2 methyltransferase activity in ARID1A-mutated cancers. *Nature Med.* 21, 231-238.
- Bohn, S., Thomas, H., *et al.* 2003. Distinct Molecular and Morphogenetic Properties of Mutations in the Human HNF1 $\beta$  Gene That Lead to Defective Kidney Development. *J. American Soc. Nephrology*, 14, 2033-2041.
- Chang, C.W., Counago, R.M., *et al.* 2013. Distinctive conformation of minor site-specific nuclear localization signals bound to importin-alpha. *Traffic*, 14, 1144-1154.
- Chen, V.B., Arendall, W.B. 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S. and Richardson D.C.. (2010). MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr. Section D, Biol. Crystallogr.* 66, 12-21.
- Conti, E., Uy, M., *et al.* 1998. Crystallographic Analysis of the Recognition of a Nuclear Localization Signal by the Nuclear Import Factor Karyopherin- $\alpha$ . *Cell*, 94, 193-204.
- Cribbs, A., Kennedy, A., *et al.* 2013. Simplified production and concentration of lentiviral vectors to achieve high transduction in primary human T cells. *BMC Biotech.* 13, 98.
- Cutress, M.L., Whitaker, H.C., Mills, I.G., Stewart, M. and Neal, D.E. (2008). Structural basis for the nuclear import of the human androgen receptor. *J. Cell Sci.* 121, 957-968.
- Emsley, P., Lohkamp, B., *et al.* 2010. Features and development of Coot. *Acta Crystallogr. Section D, Biological Crystallogr.* 66, 486-501.
- Evans, P.R. and Murshudov, G.N. 2013. How good are my data and what is the resolution? *Acta Crystallogr. Section D, Biological Crystallogr.* 69, 1204-1214.
- Fontes, M.R.M., Teh, T., *et al.* 2000. Structural basis of recognition of monopartite and bipartite nuclear localization sequences by mammalian importin- $\alpha$ . *J. Molec. Biol.* 297, 1183-1194.
- Geoffrey, M. and Cooper, S.M., 2000. *The Cell, A Molecular Approach*. 2nd ed., Boston University.
- Giesecke, A. and Stewart, M., 2010. Novel Binding of the Mitotic Regulator TPX2 (Target Protein for Xenopus Kinesin-like Protein 2) to Importin- $\alpha$ . *J. Biol. Chem.* 285, 17628-17635.
- Gounaris, I., Charnock-Jones, D.S., *et al.* 2011. Ovarian clear cell carcinoma—bad endometriosis or bad endometrium? *J. Pathol.* 225, 157-160.
- Hirota Kajiha, Y.Y., Seiji Kanayama, Naoto Furukawa, Taketoshi Noguchi, Shoji Haruta, Shozo Yoshida, Toshiyuki Sado, Hidekazu Oi, Hiroshi Kobayashi, 2010. Clear cell carcinoma of the ovary: Potential pathogenic mechanisms (Review). *Oncology Rep.* 23, 1193-1203.
- Hodel, A.E., Harreman, M.T., *et al.* 2006. Nuclear localization signal receptor affinity correlates with in vivo localization in *Saccharomyces cerevisiae*. *J. Biological Chem.* 281, 23545-23556.
- Holvey, R.S., Valkov, E., *et al.* 2015. Selective Targeting of the TPX2 Site of Importin-alpha Using Fragment-Based Ligand Design. *Chem. MedChem.* 10, 1232-1239.
- Jones, S., Wang, T.-L., *et al.* 2010. Frequent Mutations of Chromatin Remodeling Gene ARID1A in Ovarian Clear Cell Carcinoma. *Science*, 330, 228-231.
- Kabsch, W., 2010. XDS. *Acta Crystallogr. Section D, Biol. Crystallogr.* 66, 125-132.
- Kato, N., Sasou, S., *et al.* 2006. Expression of hepatocyte nuclear factor-1beta (HNF-1beta) in clear cell tumors and endometriosis of the ovary. *Modern Pathol.* 19, 83-89.
- Kato, N., Tamura, G., *et al.* 2008. Hypomethylation of hepatocyte nuclear factor-1beta (HNF-1beta) CpG island in clear cell carcinoma of the ovary. *Virchows Arch.* 452, 175-180.

- Kato, N., Toukairin, M., *et al.* 2007. Immunocytochemistry for hepatocyte nuclear factor-1 $\beta$  (HNF-1 $\beta$ ): A marker for ovarian clear cell carcinoma. *Diagnostic Cytopathol.* 35, 193-197.
- Kobayashi, H., Yamada, Y. *et al.*, 2009. The role of hepatocyte nuclear factor-1beta in the pathogenesis of clear cell carcinoma of the ovary. *Internat. J. Gynecol. cancer*, 19, 471-479.
- Kobe, B. 1999. Autoinhibition by an internal nuclear localization signal revealed by the crystal structure of mammalian importin- $\alpha$ . *Nature Struct. Mol. Biol.* 6, 388-397.
- Kuo, K.T., Mao, T.L. *et al.*, 2009. Frequent activating mutations of PIK3CA in ovarian clear cell carcinoma. *Am. J. Pathol.* 174, 1597-1601.
- Kutner, R.H., Zhang, X.Y., *et al.* 2009. Production, concentration and titration of pseudotyped HIV-1-based lentiviral vectors. *Nature Protocols*, 4, 495-505.
- Landen, C.N., Birrer, M.J., *et al.*, 2008. Early Events in the Pathogenesis of Epithelial Ovarian Cancer. *J. Clin. Oncol.* 26, 995-1005.
- Lange, A., Mills, R.E., *et al.* 2007. Classical Nuclear Localization Signals: Definition, Function, and Interaction with Importin  $\alpha$ . *J. Biol. Chem.* 282, 5101-5105.
- Lanitis, E., Dangaj, D., *et al.* 2012. Primary Human Ovarian Epithelial Cancer Cells Broadly Express HER2 at Immunologically-Detectable Levels. *PloS One*, 7, e49829.
- Liu, P., Khurana, A., *et al.* 2009. Regulation of HSulf-1 Expression by Variant Hepatic Nuclear Factor 1 in Ovarian Cancer. *Cancer Res.* 69, 4843-4850.
- Lu, P., Rha, G.B., *et al.* 2007. Structural Basis of Disease-Causing Mutations in Hepatocyte Nuclear Factor 1 $\beta$ . *Biochemistry*, 46, 12071-12080.
- Lu, P., Li, Y., *et al.* 2006. Crystallization of hepatocyte nuclear factor 1beta in complex with DNA. *Acta Crystallogr., Sect. F, Struct. Biol. Cryst. Commun.* 62, 525-529.
- Marfori, M., Lonhienne, T.G., *et al.* 2012. Structural Basis of High-Affinity Nuclear Localization Signal Interactions with Importin- $\alpha$ . *Traffic*, 13, 532-548.
- Marfori, M., Mynott, A., *et al.* 2011. Molecular basis for specificity of nuclear import and prediction of nuclear localization. *Biochim. Biophys. Acta*, 1813, 1562-1577.
- Nakada, R., Hirano, H., *et al.* 2015. Structure of importin-alpha bound to a non-classical nuclear localization signal of the influenza A virus nucleoprotein. *Sci. Rep.* 5, 15055.
- Neduva, V., and Russell, R.B. 2005. Linear motifs: Evolutionary interaction switches. *FEBS Lett.* 579, 3342-3345.
- Rey-Campos, J., Chouard, T., *et al.* 1991. vHNF1 is a homeoprotein that activates transcription and forms heterodimers with HNF1. *EMBO J.* 10, 1445-1457.
- Rosenfeld, M.G. 1991. POU-domain transcription factors: pou-er-ful developmental regulators. *Genes Devel.* 5, 897-907.
- Ryan, A.K. and Rosenfeld, M.G. 1997. POU domain family values: flexibility, partnerships, and developmental codes. *Genes Devel.* 11, 1207-1225.
- Sabherwal, N., Schneider, K.U., *et al.* 2004. Impairment of SHOX nuclear localization as a cause for Léry-Weill syndrome. *J. Cell Sci.* 117, 3041-3048.
- Shen, H., Fridley, B.L., *et al.* 2013. Epigenetic analysis leads to identification of HNF1B as a subtype-specific susceptibility gene for ovarian cancer. *Nature Commun.* 4, 1628.
- Stelma, T., Chi, A., van der Watt P.J., Verrico, A., Lavia, P. and Leaner, V.D. (2016). Targeting nuclear transporters in cancer: Diagnostic, prognostic and therapeutic potential. *IUBMB Life*, 68, 268-280.
- Stewart, M. (2007). Molecular mechanism of the nuclear protein import cycle. *Nature Rev. Mol. Cell Biol.* 8,195-208.
- Tan, D.S., Miller, R.E., *et al.* 2013. New perspectives on molecular targeted therapy in ovarian clear cell carcinoma. *British J. Cancer*, 108, 1553-1559.
- Tan, D.S.P. and Kaye, S., 2007. Ovarian clear cell adenocarcinoma: a continuing enigma. *J. Clin. Pathol.* 60, 355-360.
- Tsuchiya, A., Sakamoto, M., *et al.* 2003. Expression Profiling in Ovarian Clear Cell Carcinoma: Identification of Hepatocyte Nuclear Factor-1 $\beta$  as a Molecular Marker and a Possible Molecular Target for Therapy of Ovarian Clear Cell Carcinoma. *Am. J. Pathol.* 163, 2503-2512.
- Wiegand, K.C., Shah, S.P., *et al.* 2010. ARID1A mutations in endometriosis-associated ovarian carcinomas. *New England J. Med.* 363, 1532-1543.

- Wu, G., Bohn, S., *et al.* 2004. The HNF1 $\beta$  transcription factor has several domains involved in nephrogenesis and partially rescues Pax8/lim1-induced kidney malformations. *European J. Biochem.* 271, 3715-3728.
- Yamaguchi, K., Mandai, M., *et al.* 2010. Identification of an ovarian clear cell carcinoma gene signature that reflects inherent disease biology and the carcinogenic processes. *Oncogene*, 29, 1741-1752.
- Yoshida, S., Furukawa, N., *et al.* 2009. Theoretical model of treatment strategies for clear cell carcinoma of the ovary: focus on perspectives. *Cancer Treat. Rev.* 35, 608-615.

## Figure Legends

**Figure 1** (A) HNF1 $\beta$  protein expression levels in CCC cell lines OVISE, JHOC5, JHOC7, JHOC9, SKOV3 and a negative control HGSOC cell line, PEO1; (B) Data is the mean (n = 3, three biological replicates) and SEM. Expression levels were normalised to the housekeeping protein GAPDH. The expression levels observed with JHOC5 (P < 0.005, \*\*\*) and OVISE (P < 0.05, \*) cells were significantly higher than PEO1.

**Figure 2** eGFP imaging of PEO1 transduced lines with (a) eGFP-HNF1 $\beta$  (Lv103) showed primarily nuclear localisation of the GFP signal, whereas cells transduced with eGFP (Lv105) alone (b) showed both nuclear and cytoplasmic localisation. In cells transduced with eGFP-HNF1 $\beta$  (Lv103) in which the seven-residue NLS had been deleted (c, Mutant 8) the eGFP signal was mislocalized to the cytoplasm. The scale bar represents 80  $\mu$ m. Images were taken on a Leica tandem confocal microscope using 20 x, 40 x and 60 x objectives.

**Figure 3** (A) Stable complex formation between  $\Delta$ IBB mouse Importin- $\alpha$ 1 (50 kDa) and HNF1 $\beta$  (27 kDa) during size exclusion chromatography. Fractions were analyzed by Coomassie stained SDS-PAGE (\* marks the peak for the complex); (B) GST pull-down with liberated HNF1 $\beta$  showing pull-downs using different  $\Delta$ IBB mouse Importin- $\alpha$  isoforms (red \*).

**Figure 4** (A) ITC titration for the binding of the HNF1 $\beta$  NLS peptide to  $\Delta$ IBB mImportin- $\alpha$ 1. Fitting the data (with  $\chi^2/\text{DoF} = 2.9 \times 10^4$ ) gave a  $K_d$  of  $13.6 \pm 1.5$  nM for  $0.94 \pm 0.13$  sites,  $\Delta H = -7.2 \pm 1.6$  kcal/mole,  $\Delta S = -1.7$  cal/mol/deg. (B) In pull-down assays, GST-HNF1 $\beta$  bound only the E mutant of  $\Delta$ IBB *Xenopus* Importin- $\alpha$ 1 (that impairs NLS binding at the minor site)  $M_r$  50 kD, but not the D mutant (that impairs binding at the major site) or ED double mutant (see Giesecke and Stewart, 2010).

**Figure 5** (A) Overview of the 2.4  $\text{\AA}$  crystal structure of the  $\Delta$ IBB mouse Importin- $\alpha$ 1:HNF1 $\beta^{\text{NLS}}$  complex. Two copies of the NLS peptide were observed where one was bound in the major site (blue) and another in the minor site (yellow). Each ARM repeat of mouse Importin- $\alpha$ 1 has been labelled with alternating colours. (B) Final *2Fo-Fc* electron density map around NLS peptide in major and minor binding sites on mouse Importin- $\alpha$ 1 contoured at  $1\sigma$ .

**Figure 6** Schematic illustration of the interactions of the HNF1 $\beta^{\text{NLS}}$  peptide in the major (A) and minor (B) sites on  $\Delta$ IBB mouse Importin- $\alpha$ 1.

**Table 1 - Protein Crystallography Parameters for  $\Delta$ IBB-mImportin- $\alpha$ 1 complexed with the HNF1 $\beta$  NLS peptide**

<b>Data collection statistics</b>	
Wavelength (Å)	0.97624
Space group	$C222_1$
Unit cell parameters: a, b, c (Å); $\alpha$ , $\beta$ , $\gamma$ (°)	81.4, 111.4, 129.4; 90.0, 90.0, 90.0
Resolution range (outer shell in brackets; Å)	129.4 - 2.40 (2.49 - 2.40)
Unique reflections	23388 (2414)
Total observations	173271 (18283)
$\langle I / \sigma(I) \rangle$ : all (outer shell)	10.2 (1.8)
Rp.i.m.: all (outer shell)	0.041 (0.60)
Completeness: all (outer shell) (%)	100 (100)
Multiplicity	7.4
Wilson B-factor	53.7
<b>Refinement statistics</b>	
Non-hydrogen atoms	3500
Number of water molecules	53
Bond length deviation from ideal values (Å)	0.008
Bond angle deviation from ideal values (°)	0.78
Ramachandran favoured/outliers (%)	98.9 / 0
All-atom clashscore	1.86
Average protein B factor	67.2
Average water B factor	61.4
Rwork/Rfree (%)	18.4 / 22.0
MolProbity score (percentile)	0.95 (100)



**Supplementary Material (Online Only)**

[Click here to download Supplementary Material \(Online Only\): Supplementary Information.docx](#)