# **Reverse-Engineering the Brain:** The parts are as complex as the whole.

#### Jens Pohl, PhD

Professor of Architecture (Emeritus) California Polytechnic State University (Cal Poly) Vice President, Engineering & Technology Tapestry Solutions (a Boeing Company) San Luis Obispo, California, USA

#### Abstract

The purpose of this paper is to review the current state of neuroscience research with a focus on what has been achieved to date in unraveling the mysteries of brain operations, major research initiatives, fundamental challenges, and potentially realizable objectives. General research approaches aimed at constructing a wiring diagram of the brain (i.e., connectome), determining how the brain encodes and computes information, and whole brain simulation attempts are reviewed in terms of strategies employed and difficulties encountered. While promising advances have been made during the past 50 years due to electron microscopy, the development of new experimental methods, and the availability of computer-enabled high throughput imaging systems, brain research is still greatly encumbered by inadequate monitoring and recording capabilities. Four hypotheses relating to comprehension through the assembly of parts, formation of memories, influence of genes, and synapse formation are described as plausible explanations even though they cannot be validated at this time. By assessing the feasibility of overcoming the principal problems that beleaguer brain research in comparison with the potential benefits that can be derived from even partial achievement of the goals the author concludes that the significant investment of government funding is justified.

## Keywords

chromosomes, comprehension, connectome, digital atlas, DNA, genes, genome, ion channel, measurable brain parameters, memory, microscopy, neuroinformatics, neuron, neuroscience, phrenology, plasticity, protein, RNA, synapse, twins

## Neuroscience Research in Perspective<sup>1</sup>

Although a great deal of progress has been made in the scientific study of the human nervous system over the past 50 years, even today (2016) neuroscience is essentially still a collection of experimental findings and hypothesis rather than an integrating set of theories and validated principles. We know that the brain is a network of networks in which billions of neurons communicate with each other, but we know very little about what they communicate or the role of individual neurons in achieving the end result of the communication. Similarly, although we

<sup>&</sup>lt;sup>1</sup> The author is not a neuroscience researcher but has been involved in the design and development of decisionassistance software with embedded intelligence for the past 40 years. His attraction to neuroscience research stems from his interest in the current level of understanding of how the brain operates and how that knowledge may impact the design of intelligent software. Since this paper is intended for persons with similar background and research interests an appendix describing molecular biology fundamentals has been added.

have some understanding of the molecular elements and their functions within a neuron and its synaptic connections to other neurons, we can only hypothesize how these functions may contribute to our ability to comprehend, memorize and reason (Marcus 2015).

There is general agreement that the brain is an information processor of some kind that takes inputs from the outside, encodes these inputs into some format that allows it to compute a virtual model of the outside, and formulates instructions to the various components that control our physical (e.g., movement) and behavioral (e.g., thinking) functions. However, the fundamental principles of brain computation have yet to be discovered and there is no overarching theory of how networks of neurons contribute to something as complex as human cognition.

The difficulties that are continuing to confront neuroscience research are related to the enormous scope of the undertaking in terms of the number of neurons involved, the complexity of the individual parts and their interactions within the whole, and the relatively limited capabilities of the available research techniques and devices. Although there have been significant advances in microscopy with the development of the electron microscope that can achieve resolutions of objects smaller than 50 picometer<sup>2</sup>, the maximum number of neurons that can be recorded simultaneously is around 100 with a recording speed that lags well behind the neuronal firing rate.

So far most of the recording of neural activity has been confined to the measurement of single neurons. While this provides some insight into the operation of a single node, it provides little reliable information about the behavior of the network of several million of nodes that this neuron is part of. Sporns (2015) draws attention to this problem by pointing to variables such as synaptic weightings and neuron shapes that are likely to have significant influence on the behavior of a network of neurons. It is now generally recognized that the behavior of a network as a whole may differ markedly from the behavior of its nodes under certain conditions. This gives rise to the notion of *emergence* in network theory, where the collective interactions among the nodes of a complex network can lead to new properties that do not exist at the node level. In the case of the human brain such emergent phenomena leading to the coordinated firing of large numbers of neurons is referred to as *neural synchronization*. This is a condition that represents the global outcome of many local events that are orchestrated by the network as a whole rather than any specific chain of causes at the node level (Sporns 2015, 93).

## **Measurable Brain Parameters**

The complexity of the brain is not only due to the enormous number of neuron nodes (around 86 billion in the human brain) and the even much larger number of connections between nodes (around 1000 trillion in the human brain), but also due to the variables that govern the behavior of the nodes. First, there are probably hundreds of thousands of neuron types that appear to play different roles and are therefore likely to function in different ways. The roundworm *C elegans*<sup>3</sup>, whose complete connectome was published in 1986 (White et al. 1986), has only 302 neurons but 100 types of neurons. Therefore, even though researchers have been able to record the behavior of single neurons the role and function of different types of neurons is likely to involve the tracing of synaptic chains (i.e., pathways) involving a large number of neurons. This requires the ability to record thousands if not millions of neurons simultaneously. While increasingly

<sup>&</sup>lt;sup>2</sup> In the metric system of units 1 picometer (pm) is equal to one trillionth of a meter (i.e.,  $10^{-12}$ m)

<sup>&</sup>lt;sup>3</sup> Caenorhabditis elegans (*C elegans*) is a roundworm (nematode) about 1 mm in length that lives in temperate soil environments.

more powerful technologies are becoming available for recording the behavior of groups of neurons, they are limited in both what they can measure and their ability to keep up with the speed at which the state of individual neurons can change.

Second, neurons produce (i.e., express) extremely complex sets of molecules that generate and sense the electrical signals that underlie all neuronal responses and synaptic transmissions. There is evidence that the reweighting of the synapses is influenced by multiple factors that include not only the shape (i.e., morphology) of the neuron but also the rate of excitatory or inhibitory signals and the strengths of those signals. In this way the synapses play a decisive role in how neurons interact to reinforce or diminish, compete or cooperate with each other. Even though the communication at a synapse is a point-to-point connection that is restricted to the two neurons involved, the influence of synapse variables on the complex behavior of large groups of connected neurons will likely continue to be a severe challenge to researchers for the foreseeable future.

Neurons and synapses: Neurons consist of a cell body and branches called neurites. There are two kinds of neurites, namely: a relative long axon along which the neuron passes an electric charge when it *spikes* or *fires*; and, many much shorter and more convoluted *dendrites* (Figure 1). The axon and dendrites of any one neuron connect to many other neurons forming a complex network of equally complex sub-networks. The connection or contact points between neurons are referred to as synapses.



Figure 1: Brain component – the neuron

However, the two sides of a synapse do not actually touch but are separated by a very narrow cleft or *synaptic gap* (Figure 2). The sending side of the cleft stores neurotransmitters in

vesicles, while the receiving side of the cleft contains receptors referred to as postsynaptic density. Triggered by an electrical spike one or more vesicles (the senders) secrete a neurotransmitter into the cleft, which is sensed by the postsynaptic density receiver. While 100 over types there are of neurotransmitters consisting of atoms that are bonded to each other, a neuron typically secretes the same neurotransmitters (often only one) at all of its synapses.



Figure 2: Brain component – the synapse

The receptor senses the neurotransmitter but will accept it only if it is the right kind. The analogy is a lock and key. The receptor is the lock and the neurotransmitter is the key. Each type of neurotransmitter has a distinctive molecular shape that must match the configuration

of the *binding site* (i.e., the receptor lock). Some receptors are a combination of a lock and a door, with a small tunnel threading through the receptor molecule that connects the inside of the neuron with the outside. This tunnel is blocked by a door-like structure most of the time. However, when the neurotransmitter binds to the receptor the door is opened for an instant to allow the electrical current to momentarily flow through the tunnel. This type of protein that contains a tunnel is referred to as an *ion channel*, where *ions* are the electrically charged particles that conduct electricity in aqueous solutions. However, there are many types of ion channels that are not receptors. For example, some of them enable neurons to generate spikes.

How does a synapse deal with the relative slowness of chemical communication, the prevention of potential crosstalk, and timing? Certainly chemical signals are much slower than electrical signals. However, a synaptic gap is very narrow and therefore the neurotransmitter does not have to travel far. Also, since brain tissue is tightly packed with up to a million neurons and a billion synapses per cubic millimeter, the synapse has to ensure that the secretion of a neurotransmitter is not sensed by other nearby connections. To prevent a cascading sequence of nearby neuron activations, the synapse either sucks the neurotransmitter back up for reuse or degrades it into an inert form. Using the same mechanism of recycling and degrading that minimizes crosstalk the synapse is also able to control the timing of the chemical signal by ensuring that the signal does not linger on.

*Synaptic chains:* A synapse is activated when a spike triggers secretion on the sender side. On the receiver side receptors sense the neurotransmitter and then make electrical current flow. In other words, a synapse converts an electrical signal into a chemical signal and then back into an electrical signal. A pathway (i.e., synaptic chain) is created when the electrical signal passes from neuron A to neuron B to neuron C, and so on. These pathways are directional because synapses are one-way communication devices. At any synapse one neuron is always the sender and the other always the receiver. In the case of electrical signals the spike travels from the cell body along the axon away from the neuron and from other neurons through dendrites towards the neuron to the cell body.

In summary, in any neuron electrical signals flow from dendrites to the cell body and once an action potential threshold has been exceeded a new electrical signal flows from the cell body along the axon to the dendrites of connected neurons. Chemical signals jump from the axon of neuron A to the dendrite of neuron B. In neuron B electrical signals again flow from dendrites to the cell body and along the axon if the action potential is exceeded and a spike occurs. These pathways end when axons finally connect to nerves that connect to muscle fibers, which respond by contracting. This causes the muscle to shorten and produce a movement.

The electrical signals (i.e., spikes) that trigger the chemical communication at the synapse can travel long distances by propagating through axons and dendrites much like wires. However, the transmission of electrical signals is isolated to occur only at synapses. In axons the electrical signals are brief pulses (i.e., spikes) lasting about a millisecond and referred to as action potential, much like Morse Code to prevent the signal from becoming corrupted by noise over longer distances. Almost all synapses are weak and far below the level required to produce a spike. However, once a spike occurs and travels along the axon it will be passed onto multiple other neurons through their dendrites. In other words, a single spike becomes many spikes in connected neurons. Accordingly, the stimulation of one sense organ can

cause multiple responses and any one neuron can be activated to generate a spike by receiving weak signals from multiple connected neurons.

*Weighting of synapses:* In most neurons the electrical signals received by the dendrites of that neuron are weighted according to their strength. Strong synapses produce large currents and weak synapses produce small currents. In principle every neuron is different in its behavior due to the unique configuration of its ion channels. Therefore the weighting model, which is common in most neurons, must not imply a common behavior of neurons<sup>4</sup>. However, the classification of neurons into types suggests that neurons of the same type behave in similar ways. It is also possible for a neuron to receive multiple signals from the same external neuron through several synapses. These signals must be received within a brief time span from milliseconds to seconds for them to be combined by the cell body into one signal that is compared with the action potential of that neuron. It follows that there are two mechanisms that prevent neurons from spiking indiscriminately, namely: the threshold of the neuron's action potential; and, the action of inhibitory synapses that will reduce the strength of the combined signal.

A neuron is either excitatory or inhibitory but not both. The existence of inhibitory synapses is crucial to the operation of the nervous system. An excitatory neuron makes only excitatory synapses onto other neurons and an inhibitory neuron makes only inhibitory synapses. However, this is not the case for the synapses received by a neuron from other neurons. Therefore an inhibitory neuron may also receive excitatory signals and vice versa. The analogy is to a conformist stance as opposed to a contrarian stance in society. There is also a type of excitatory synapse that has a direct electrical connection and does not use a neurotransmitter. These synapses communicate more quickly, but are far less in number than the chemical synapses.

The threshold action potential of the neuron spike mechanism has two functions, to either transmit a signal that may precipitate an action or to not send a signal because the combined strength of the signals it has received is insufficient to warrant an action. In this respect the brain is far more sophisticated than a telecommunication network or the wiring of a Cray supercomputer that are designed to essentially support only two states (i.e., *on* or *off*).

*Life cycle of the human brain:* The creation of neurons and synapses proceeds through four stages (Figure 3): neurons are created; neurons migrate to their assigned places in the brain; neurons extend branches (i.e., neurites) throughout the body; and, neurites make connections.

*Neurons are created:* Through the division of progenitor cells neurons are created in the prenatal stage before birth. Similar to stem cells, progenitor cells replicate into a specific type of cell. However unlike stem cells that can replicate indefinitely, progenitor cells can divide only a limited number of times.

*Neurons migrate to the brain:* The migration of neurons to the brain during the prenatal stage can be disrupted with the result of structural abnormalities in the brain, such as the

<sup>&</sup>lt;sup>4</sup> Computer simulations of brain functions such as Henry Markram's *Blue Brain* project (Markram 2006) and Dharmendra Modha's simulation of a cat's brain (Wong et al 2012) rely mostly on the weighted voting model of neurons that assumes that the behavior of all neurons is essentially the same. Also, since there is no known cortical *connectome* these computer models typically assume that connections are formed randomly. However, since *Neural Darwinism* includes the concept of activity-dependent synapse elimination, the surviving connections are not subject to randomness. Finally, computer models do not make allowances for reconnection, rewiring and regeneration.

cortex lacking folds.

Both of these stages occur in the prenatal brain. By the time a baby is born the creation and migration of neurons are virtually complete and there are only a few areas of the brain in which neurons will continue to be created after birth.



Figure 3: Creation of neurons from conception to early childhood

- *Neurons grow neurites:* After birth the neurons continue to grow branches in the form of dendrites and axons. The axons have to grow the most since they are much longer than the dendrites. The growing tip of an axon is called a growth cone, which navigates its way through the body under the guidance of genes. The surfaces of neurons are coated with special guidance molecules and the spaces between neurons contain drifting guidance molecules. The axon's growth cone is equipped with sensor molecules that can detect the guidance molecules much like dogs following a scent. Since both the guidance molecules and the sensor molecules are produced under genetic control we can say that the wiring of the brain is guided by genes. Axons grow fairly straight and do not branch until the growth cones reach their destinations. While the overall shape of an axon appears to be genetically determined, the branching pattern at the tip is likely to be largely random. The analogy is a tree where the need for branches is determined by the genetic plan but the actual location and size of branches is influenced by environmental conditions and a degree of randomness.
- *Neurites make connections:* During the wiring of the brain neurons make connections between neurites by creating synapses. This process is probably driven by a combination of genetic control (i.e., wiring paths and potential affinity between neuron types) and a degree of randomness (i.e., some probability whenever the neurites of different neurons cross each other). Accordingly it appears that the initial connectome at the very early development stage is largely a product of genes and randomness. Genes appear to control the wiring in terms of the shape of each neuron and the region over which it extends branches, while synapses are formed randomly where the neurites of neurons touch. However, as development proceeds sensory stimuli (i.e., experiences) also shape the connectome.

Synapses are created at an astonishing rate in the infant brain. For example, in *Brodmann area 17* alone half a million synapses per second are produced during the third and forth month after birth (Huttenlocher 1990). Another argument in favor of the randomness hypothesis is that while such large numbers of synapses are formed during development many synapses are also eliminated. Pruning of both synapses and dendrites takes place to a significant extent after the age of two years, leaving persons that have just reached adulthood with only about 60% of the synapses that they had when they were two years of age (Huttenlocher and Dabholkar 1997).

Figure 4 (Seung 2013, 108) is an example of the increase in synapses and dendrites from birth to one month, 6 months and two years, followed by the results of pruning at the age of

four years. One could speculate that the early connectome is a rough draft based on gene guidance and randomness, which is subsequently refined based on the survival of the fittest principles of Neural Darwinism<sup>5</sup>. In other words, dendrites and synapses that are not or very infrequently used are weakened (or remain very weak from the time of their creation) and eventually eliminated. However, it appears that the creation and destruction of synapses are not separate processes but occur concurrently.



Figure 4: Pruning of dendrites and synapses

In summary, while the creation of synapses is largely random, the elimination of synapses is based on lack of utilization of synapses. In other words, the dendrites and synapses that are eliminated were not needed in the representation, interpretation and comprehension of the stimuli that were processed by the brain.

# **General Research Approaches**

Recent large-scale research efforts in neuroscience have focused on three related areas, namely: (a) *connectomics*, to determine (map) which neuron is connected to which other neurons (Sporns 2015, Zador 2015, Hawrylycz et al. 2015, Seung 2013); (b) *brain activity mapping*, to observe the electrical discharges along the synaptic pathways (Shenoy 2015, Koch et al. 2015); and, (c) *large-scale brain simulation*, the integration of data of all areas of neuroscience to construct a biophysically realistic model that can be compared with experimental findings (Hill 2015). While each of these large-scale research efforts is certainly valuable none of them on their own is likely to lead to a holistic understanding of how the multi-level brain functions. In addition, it is often difficult to determine how to integrate the findings from each area since each area has missing pieces that the other two areas cannot compensate for. For example, an activity map without a connectome could tell us a great deal about the behavior of larger neural groups

<sup>&</sup>lt;sup>5</sup> The theory of Neural Darwinism first proposed by Edelman in 1978 and extended in 1987 (Edelman 1987) is still considered to be highly speculative (Lichtman and Coleman 2000, Purves and Lichtman 1985). Studies of synapse elimination have been performed on neuron to muscle connections. While early in human development each muscle fiber has synapses with many neurons, over time many of these are eliminated until each fiber receives synapses from just one axon.

(networks) but would not allow us to reconstruct the underlying circuitry. A connectome without an activity map would provide the static circuit architecture but not tell us anything about the dynamic reweighting of excitatory and inhibitory synapses.

Even though the brain of every human is different since both genes and experience shape the human connectome, mapping of experimental data into a common reference framework is a necessary requirement for coordinating research efforts. Not only will it allow the comparison of multiple brains, but it will also facilitate the recognition of unusual features in a particular brain and contribute to an understanding of how these apparent anomalies differ from the norm. Digital atlases can provide an effective framework for building a standard atlas that serves as a single-point-of-access clearing house for accessing multiple databases and related documents (Hawrylycz 2015). However, a standardized brain atlas will require a common three-dimensional coordinate system so that the findings of neuroimaging experiments can be readily analyzed, compared and integrated into the appropriate atlas database. This requires agreement on the specification of an origin (i.e., reference point) and a transformation function that relates each individual three-dimensional brain image from its native coordinates to the coordinates of the atlas<sup>6</sup>. A standard toolset that allows different neuroscience data to be combined is now under development by the International Neuroinformatics Coordinating Facility (INCF) for the mouse, which has remained to this day one of the most important subjects in experimental neuroscience.

*Connected brain:* There is much evidence to suggest that the overall interconnectedness of the brain rather than the simple expansion of specialized regions of the brain is responsible for human cognitive capabilities. For example, it is now recognized that neuropsychiatric disorders such as depression, anxiety, schizophrenia, and obsessive-compulsive behavior, are likely to be the result of complex genetic and environmental factors impacting neural circuitry. The independent proposals in 2005 for a unified approach to defining connectional atlases of the brain by Sporns of Indiana University and Hagmann of Lausanne suggested the term *connectome* to refer to a complete map of the neural connections within the brain (Sporns et al 2005, Hagmann 2005).

Since the study of the first complete neural circuit of any organism in the 1970s by Nobel laureate Sydney Brenner several neuroinformatics databases of connectivity have become available (e.g., CoCoMac <u>www.cocomac.org</u> and Brain Architecture Management System (BAMS) <u>http://brancusi.usc.edu</u>). Several projects to map the connectome of the laboratory mouse have been supported by public and private funding. Of these the Allen Mouse Brain Connectivity Atlas developed by the Allen Institute contains more than 1.5 petabytes of data mapped into a common three-dimensional reference space that allows the identification of neural circuits of the laboratory mouse.

Mapping the connectome of the human brain is one of the great scientific challenges of the 21<sup>st</sup> century. The Human Connectome Project (HCP) (<u>www.humanconnectome.org</u>) is tackling a key aspect by focusing on some of the main neural pathways that underlie brain functions and behavior. Due to the enormous complexity of the human brain HCP is targeting only large-scale circuitry in 1,200 adults using non-invasive neuroimaging techniques (e.g., MRI, fMRI, EEG). However, the current non-invasive imaging techniques cannot capture brain activity at a neural level and mapping the connectome at a cellular level

<sup>&</sup>lt;sup>6</sup> Many of these standardization advantages grew out of the National Institutes of Health's *Decade of the Brain* initiative in the 1990s that established the field of *neuroinformatics* (i.e., the application of computer-based and mathematical technologies for organizing and understanding brain data).

currently requires microscopic postmortem analysis of brain tissues. The required data collection would take years given the current technology when we consider that the human cerebral cortex alone contains  $10^{10}$  neurons and  $10^{14}$  synapses. The inadequacy of existing annotation tools to fully delineate and extract information at a neuronal scale and the incompleteness of the algorithms that are required to map relevant connections and construct connectivity pathways are current obstacles. The open source, crowd-sourcing initiative Open Connectome Project (www.openconnectomeproject.org) was launched in 2011 to meet this challenge.

*Computational brain:* Similar to a computer the brain operates over input and manipulates information systematically. However, both the memory capabilities of the brain and its encoding operations appear to be based on different principles. Marcus (2015) believes that it is very important for the neuroscience community to break away from the notion that the brain is not a computer. He believes that the computational capability of the brain is a fundamental component of the highly plausible and extremely useful notion that perception and comprehension in the brain are constructed progressively and hierarchically from low level feature detectors (e.g., differences in luminance and orientation) into meaningful objects such as cars, animals, persons, and faces. He argues that the unwillingness to consider the brain as a computer stems from the promises of neural networks. The pattern matching capabilities of neural networks (i.e., Parallel Distributed Processing (PDP)) were seen as the first serious alternative to the simplistic understanding of intelligence in terms of stored computer-like programs. However, eventually it became clear that the ability of PDP networks to explain higher level cognition is severely limited. Even though the brain is itself a network of neurons it is far more complex than a PDP network that consists of an array of input nodes that are connected to output nodes through one or more layers of hidden nodes.

The attractiveness of PDP networks was that they did not resemble computer programs. They had no variables or syntax like loops and if-then comparisons. It was assumed at the time that the brain operates in a similar pattern matching manner based on an initial more or less random organization that is adapted and tuned by experience. While the parallel processing capability of PDP networks is mirrored by the brain there is now some evidence that suggests that even at the embryo stage the brain begins to develop intricate (rough drafts) of brain structures in the absence of experience. For example, Nobel laureate Thomas Südhof conducted an experiment with mice whose transsynaptic neurotransmitter secretion was genetically blocked, thereby shutting down most of the brain's internal communication mechanism (Yang et al 2010). If the brain's organization is initially random and then tuned by experience then this laboratory breed of mice would essentially have random brain structures at birth. In fact, these synaptically silent mouse embryos had developed brains that appeared to be normal in respect to folds, different neuron types and the organized structures that one would expect in typical mice. Subsequent studies by others in physiology and behavior further validated that the brain's basic organization is structured in advance of experience.

It can be argued that computers are no longer totally sequential. Since the advent of the personal microcomputer some 30 years ago computers have used input-output controllers, graphic processing units and multi-core co-processors operating in parallel. In other words, computers can no longer be strictly characterized as von Neumann machines that execute a set of stored instructions sequentially. Similarly, Marcus (2015, 209) argues that many pathways in the visual cortex appear to perform transformations of the representation of

visual scenes in parallel, such as extracting edges across a scene. He concludes that at least the brains of vertebrates are computational in the sense that they operate over inputs and manipulate information systematically.

This does not mean that a brain is a digital computer. The memory capabilities of a brain may be based on different principles and a brain may perform different kinds of operations on the information it encodes, but it does encode information. Specifically, brains transduce inputs into chemical and electrical information, operate over that encoded information, and use the resulting output to generate actions and update the internal representation accordingly. Similarly, computers take inputs, encode and manipulate information, and transform their inputs into outputs.

The question then arises how do brains store and encode information and what operations do they perform over the encoded information? If one were to reverse engineer a computer coprocessor one would find that its basic elements are transistors and that these transistors are organized so that they perform a primitive set of instructions. More complex operations would be combinations of these instructions interacting with other components of the environment in which the co-processor functions. Transistors connected in circuits create basic logical operations. In a brain the neuron is the equivalent of the transistor and a circuit-level understanding of the equivalent neurocomputational logic is fundamental to understanding how a brain functions.

*Simulating the brain:* While a computer-based simulation of the brain would be very useful, building the necessary model will require in-depth knowledge of how the brain operates. Should we attempt to build such a model of the brain although we currently do not even have the most superficial understanding of how the brain works? The Human Brain Project (Hill 2015, 113) research group that is endeavoring to create a *Whole Brain Simulation* believes that even identifying the data that are required and systematically categorizing the information that is available is a step in the right direction and a worthwhile contribution to the body of knowledge in the field of neuroscience.

Unfortunately much information is lacking. There are no comprehensive atlases available that identify those parts of the brain that have been mapped and those that have not been mapped. Currently available tools are inadequate for determining whether any particular test data are significant or irrelevant to understanding the functions of the brain. Many fundamental questions are still unanswered. For example, how many classes or types of neurons are there? To date researchers have identified a few hundred but suspect that there are thousands or perhaps even millions. As mentioned previously, the worm *C. elegans* was found to have 100 neuron types out of 302 neurons (White et al 1986). Foremost the Human Brain Project group is relying on advances in information management technology to classify and characterize the enormous volume of data that is being collected in the emerging discipline of neuroinformatics. For example: there are around 100,000 publications per year; a single human brain scan generates 1 terabyte (1000 gigabytes) of data; and, the Allen Institute produced more than 1 petabyte (1,000,000 gigabytes) of data in one year in one study to characterize mouse brain connectivity. At least four formidable challenges confront the Human Brain Project:

*Challenge 1:* Since it is unlikely that the human brain will ever be fully mapped it will be necessary to identify principles to fill in missing data and parameters based on the available data and knowledge. This will require the use of predictive neuroscience

approaches and methodologies.

- *Challenge 2:* For whole brain simulation it will be necessary to build individual models of the components such as proteins, neurons, synapses, circuits, and brain regions. This is more difficult then it seems at face value because the behavior of these individual brain components may vary based on how they are connected. In other words, the behavior of the whole may not be able to be extrapolated from the behavior of the parts.
- *Challenge 3:* During all stages of its construction the whole brain model will need to be validated by comparison with thousands of experimental test results and findings. Following the Open Source approach that has proven itself in computer software, it will be necessary to periodically release new versions of the whole brain model after it has been modified to resolve inconsistencies with recent experimental results and clinical evidence.
- *Challenge 4:* The primary input to the brain is from our senses (i.e., vision, hearing, smell, taste, and touch) and the primary output is the control of muscles and movement. Thus it will be necessary to provide a simulated body to couple with a simulated brain. Neither the European Common Market *Human Brain Project*<sup>7</sup> nor the United States DARPA sponsored *SyNAPSE* project<sup>8</sup> are in their present state able to couple with a human body simulation. Therefore, since these models do not have memory, movement or learning capabilities it is not possible to evaluate them in terms of what is arguably the purpose (i.e., output) of the brain.

# Monitoring and Recording Techniques and Devices

Advances in neuroscience have been largely made possible by developments in experimental laboratory techniques, innovative applications and extensions of existing devices, and the development of entirely new equipment, such as the electron microscope, with orders of magnitude greater capabilities. Existing techniques fall essentially into five categories (Figure 5): staining; imaging; scanning; slicing; and, computing.

Historically, the discovery of the *Golgi Stain* in 1873 by the Italian physician Camillo Golgi made neurons visible for the first time. It allowed the Spanish neuroanatomist Santiago Ramon y Cajal to identify and draw many types of neurons. Both Golgi and Cajal became Nobel laureates in 1906 in recognition of their contributions to neuroscience. The electron microscope invented by the German physicist Ernst Ruska in 1933 used electrons rather than light to produce orders of magnitude greater magnification and sharper images. In the 1950s the first two-dimensional images of synapses under the electron microscope showed that two neurons do not fuse at a synapse. At the same time dense staining methods showed the entangling of the branches of many neurons, confirming that brain tissue is packed full of neurons and their branches. However, these were still two-dimensional images.

Subsequent development of a sequence of innovative techniques, device extensions and the integration of computer processing allowed researchers to transition to three-dimensional imaging. The invention of the ultramicrotome in 1953 by Keith Porter and Joseph Blum (Porter and Blum 1953) allowed brain tissue to be cut into 50 nanometer thick slices that could be imaged and stacked into a three-dimensional model of bundles of neurons. Fifty years later the invention of the SBFSEM (Serial Block Face Scanning Electron Microscope) in 2003 by

<sup>&</sup>lt;sup>7</sup> <u>https://www.humanbrainproject.eu/2016-overview</u>

<sup>&</sup>lt;sup>8</sup> <u>http://www.artificialbrains.com/darpa-synapse-program</u>

German physicist Winfried Denk (Denk and Horstmann 2004) overcame the problem of inevitable distortion during handling of the very thin slices. The SBFSEM consists of an ultramicrotome mounted inside the vacuum chamber of an electron microscope that captures a succession of block face images after each slice is cut.



Figure 5: Experimental recording techniques and devices

In the early 2000s the availability of computers allowed the processing of the large volume of data provided by stacks of two-dimensional brain slices. The invention of the ATUM (Automated Tape-collecting Ultramicrotome) by Ken Hayworth (Hayworth et al 2014) automated the stacking of two-dimensional brain slices on a plastic tape like a film strip. In collaboration with Jeff Lichtman they were able to reduce the thickness of slices from an initial 10 microns to 30 nanometers. While the ATUM is certainly a promising step towards addressing the challenge of processing the enormous volume of data produced by three-dimensional bundles of neurons, it may not be sufficient when we consider that one cubic millimeter of brain tissue can yield a petabyte of image data. For example, the reconstruction of the complete connectome of *C elegans* during the 1970s and 1980s required the manual tracing of each neuron pathway because there was no automated alternative for interpreting the images of the enormous number of 50 nanometer thick slices of brain tissue. The C elegans connectome, which has only 302 neurons, was published in 1986 (White et al. 1986). Based on this experience we can extrapolate that the manual reconstruction of just one cubic millimeter of the cortex of the human brain would take more than a million person-years. Clearly the future of connectomics depends on the ability to automate image analysis. However, this is a difficult task because it essentially relies on the ability of computers to recognize boundaries (i.e., edges) that are only vaguely defined on the two-dimensional image of each slice and even the most advanced computer software has difficulties performing this task (Seung 2015, 160-163).

Devices that are currently available for monitoring the electrical activity of neurons and imaging brain tissue include:

Electroencephalography (EEG) and Magnetoencephalography (MEG): Both EEG and MEG

are used for recording the electrical activity of the brain, mostly non-invasively with electrodes placed along the scalp. They measure voltage fluctuations resulting from the flow of ionic currents in the dendrites of neurons during synaptic transmissions (i.e., neuron spiking). However, since the magnetic fields generated by the brain are very weak approximately 50,000 neurons must be active for a detectable magnetic field. The brain's magnetic field strength of around 10 femtotesla  $(fT)^9$  for cortical activity is far below the ambient magnetic noise of  $10^8$  fT (or  $0.1\mu$ T) in an urban environment.

*Magnetic Resonance Imaging (MRI):* MRI is capable of imaging two-dimensional slices of the brain. A stack of such two-dimensional images provides a means for reconstructing the brain in three dimensions. While it is useful for determining the size of brain regions, it produces poor spatial resolution. Also, MRI's millimeter-scale resolution it is not enough for the discrimination of single neurons and axons (Seung 2013, 209).

*Diffusion Magnetic Resonance Imaging (dMRI):* dMRI maps the diffusion process of molecules (mainly water) in biological tissue both in vivo and non-invasively. The first diffusion MRI images of the normal and diseased brain were made public in 1985. The primary clinical applications are in the study and treatment of neurological disorders such as stroke. Since dMRI allows non-invasive visualization of anatomical connections between different parts of the brain it represents a major breakthrough for human connectome research. While it is useful for tracing neuronal pathways and even computes the direction of the axon, the inherently poor spatial resolution of MRI makes it difficult to follow thin axon tracts if they cross. Nevertheless, dMRI is currently used by the Human Connectome Project to track white-matter pathways (Seung 2013, 209).

While efforts are underway to improve MRI technology, it should be noted that the current resolution of dMRI is 1000 times worse than light microscopy, which is a 100 times worse than electron microscopy. Since microscopy already delivers the resolution required for connectomics research and needs only to be scaled up to handle larger volumes of data and MRI improvements call for a major breakthrough, MRI and microscopy are likely for at least the time being to remain complimentary methods.

*Functional Magnetic Resonance Imaging (fMRI):* Developed in the 1990s, fMRI provides images that indicate brain activity (in color) typically superimposed on a black-and-white MRI image of the brain. While showing which cell networks are active when a person performs a particular physical movement or thought process, it has temporal limitations because brain activation changes from moment to moment. Again, fMRI suffers from the same spatial resolution limitations that govern all current MRI-based imaging techniques.

*Positron Emission Tomography (PET):* The PET scan method is an imaging test that uses a radioactive drug (i.e., tracer) that may be swallowed, injected or inhaled. In clinical applications PET scans with flurodeoxyglucose (FDG) as the tracer are used to indicate tissue metabolic activity in the exploration of cancer metastasis (i.e., spreading of cancer to other sites). In neuroscience research PET neuroimaging is used on the assumption that areas of high radioactivity are associated with brain activity (i.e., actually the flow of blood to different parts of the brain).

Individual-Particle Electron Tomography (IPET): IPET is a technique that allows

<sup>&</sup>lt;sup>9</sup> *Femto* is a prefix in the metric system of units that denotes a factor of  $10^{-15}$ . It is derived from the Danish word *femten* meaning 15.

determination of a high-resolution structure for a single molecule. Each particle from a tilt series of micrographs is tracked and then reconstructed into a three-dimensional density map using an iterative reconstruction algorithm (Zhang and Ren 2012). A University of California (Berkeley) research team led by Gang (Gary) Ren published the first high-resolution three-dimensional image of a DNA segment reconstruction using IPET in April 2016 (Ren et al 2016).

*In-Situ Hybridization (ISH):* ISH is a technique for mapping gene expression that uses probes that bind to RNA within sectioned but intact brain tissue, thereby preserving spatial context. ISH was automated by Gregor Eichele and colleagues at the Max Planck Institute.

*Two-Photon Microscopy:* Two-Photon excitation microscopy is a fluorescence imaging technique that allows imaging of living tissues up to about 1 millimeter in depth. It is a special variant of the multi-photon fluorescence microscope that uses red-shifted excitation light to excite fluorescent dyes. The method was pioneered by Winfried Denk at Cornell University in 1990 by combining the idea of two-photon absorption with the use of a laser scanner.

# **Promising Research Programs**

Neuroscience is no longer a niche field but a major scientific endeavor. The US-based Society for Neuroscience has more than 40,000 members. On-going world-wide research funding is now estimated to be over \$3 billion. Despite the limitations of the available research methods and the daunting scope of the data processing requirements, headway is being made through the application of innovative combinations of MRI imaging, electron microscopy, staining, stacking of nanoscale slices of brain tissue to construct three-dimensional models of neural groupings, and automated (i.e., computer-based) high throughput image interpretation techniques. The major advances are in the molecular area with increased understanding of the structure and function of ionic channels and receptors<sup>10</sup>.

*Digital atlases:* Digital brain atlases are used to represent the spatial organization of neuronal structures, for planning and guidance during neurosurgery, and as a reference for interpreting gene expression (i.e., the structure and functions of proteins). Existing atlases are extensive and include to the extent possible with existing research techniques and devices partial mappings of the mouse, rat, rhesus macaque<sup>11</sup>, and human brain. In addition to atlases based on anatomy, histology, MRI, and PET scans, modern digital atlases use gene expression, connectivity, probabilistic and multi-modal techniques, as well as visualization software (Hawrylycz 2015). The development of large-scale brain atlases is now a major undertaking in neuroscience and there have been recent advances in neuroimaging techniques that are providing maps of remarkable resolution:

*Brainbow:* A combinatorial color labeling technique using fluorescent proteins. Josh Sanes and Jeff Lichtman at Harvard University have been classifying and visualizing microscopic neurons by marking individual neurons with over one hundred distinct colors to trace and reconstruct their cellular structure (Lichtman et al. 2008).

<sup>&</sup>lt;sup>10</sup> The ionic channels and receptors function as switches and modulators in the membrane to control the processing of information by neurons, resulting in spikes along the axon and the release of neurotransmitters in synapses.

<sup>&</sup>lt;sup>11</sup> Rhesus macaque is a monkey species widely used for medical research.

*Array Tomography:* Developed by Stephen Smith at Stanford University to classify diversity in the synaptic code (Micheva and Smith 2007). Can also achieve combinatorial labeling of synaptic connections using an electron microscope.

*High Resolution 3-D Microscopy:* Culmination of the work of Katrin Amunts and the Karl Zilles laboratory at Jülich (Germany) and Evans in Montreal (Canada), *BigBrain* is a dataset 125,000 times larger than a typical MRI. However, still not at the resolution of the finest structures in the brain (Amunts et al. 2013).

*CLARITY:* A method that subjects the brain to a three-dimensional network of hydrophilic polymers and then removes the lipids from the brain by electrophoresis<sup>12</sup>. The brain remains fully intact but optically transparent and macromolecule permeable. CLARITY uses intact tissue in-situ hybridization and immunohistochemistry with multiple rounds of staining and de-staining to visualize gene expression or protein binding.

*EyeWire:* Crowdsourcing project launched by Sebastian Seung of MIT to map the retinal connectome through an interactive game (Johnson 2012).

The transition from printed to digital atlases has been revolutionary, since digital atlases allow navigation, three-dimensional reconstruction and visualization from the smallest nuclei to macro-scale regions of the brain.

*Integrated mapping of the brain:* Ideally projects focused on creating an integrated mapping of the brain should aim to include a sufficiently comprehensive set of data to convey an understanding of the brain's biological structure and functions that interlink to form an integrated system. This would include: cell types (the distinct features of the different neuron types); cell networks (the circuitry architecture within and between neural groups (i.e., networks)); connections (the strengths and types of synaptic excitatory and inhibitory transmissions); pathways (from lowest level feature detectors to highest level neuron functions); histories of electrical activity patterns over time; and, histories of molecular changes over time.

Church et al (2015, 52-56), in drawing an analogy to the Rosetta stone<sup>13</sup>, have proposed in conceptual terms that the construction of a Rosetta Brain could be reduced to the two operations of labeling and counting. Their argument proceeds along logical lines as follows:

- A connectome is at its core a large matrix that indicates whether neuron A is connected to neuron B to neuron n, where n is however a very large number.
- If each neuron could be given a unique barcode so that the children of neuron A have the barcode of their parent (neuron A) included then we could easily trace synaptic chains (i.e., pathways).
- Gene expression (i.e., DNA to messenger RNA (i.e., transcription) to protein (i.e., translation)) could be used to determine cell type by counting the numbers of each

<sup>&</sup>lt;sup>12</sup> In electrophoresis a sample is placed on one end of a gel consisting of a substance with a molecular matrix of evenly spaced opening. When an electric current is passed through the gel it moves the molecules in the sample through the gel. The rate at which the molecules move through the gel depends on both their size and their charge, with small molecules and charged molecules moving more quickly. The gel can be visualized under ultraviolet light to show DNA samples as bands.

<sup>&</sup>lt;sup>13</sup> The Rosetta stone allowed hieroglyphics to be deciphered because it recorded the same statement in three different languages, two of which were known.

messenger RNA in the neuron.

- Gene expression changes due to learning and memory in combination with time stamps, could be used to track the history of molecular changes.
- On the assumption that the number of synapses influences the strength of a connection, synaptic connection strengths and types could be inferred by counting the abundances of different proteins on either side of a synapse in combination with the number of distinct synapses between neuron *A* and neuron *B*.

Recognizing that DNA can be used to store any nitrogen bases string in addition to those used in the biological blueprint genome of an organism, they propose to generate and read barcodes by using the four nitrogen bases of nucleotide in the DNA (i.e., adenine (A), thymine (T), guanine (G), and cytosine (C)) to form barcode sequences (e.g., ATGC). There would be a sufficient number of unique barcodes because a DNA barcode of 25 nitrogen bases has  $4^{25}$  unique combinations and there are only about  $4^{23}$  synapses in the human brain (i.e.,  $10^{14}$  is approximately  $4^{23}$ ). They further argue that to make a test tube with all  $4^{25}$  unique combinations is not difficult and that to randomly insert one sequence into each of the 100 million neurons in the brain of a mouse is potentially feasible. To read these DNA sequences would then require the normal DNA sequencing process to be applied to brain slices using fluorescent dyes to distinguish between A,T,G and C.



Figure 6: Connectome – barcoding solution

The same barcoding concept with a somewhat different implementation approach has been proposed by Zador's team at Cold Spring Harbor Laboratory (2015) for constructing the connectome of sections of the neocortex of laboratory mice. The selection of the neocortex region of the brain is based on its central role in biological intelligence (higher level functions such as sensory perception), the similarity of its basic structure across all mammals (a piece of mouse cortex is not very different from a monkey), its uniformity (portion for processing sound is not very different to the portion processing touch), and its layered

modularity (six layers). The BOINC<sup>14</sup> approach (Figure 6) utilizes the high-throughput DNA sequencing technology originally developed for sequencing the human genome. It proposes to overcome three challenges in very innovative ways:

- Challenge 1: How to express a unique sequence of DNA (i.e., a DNA barcode) in each neuron of the neuronal circuit? Approach: Breeding a laboratory mouse that is engineered with: (a) a specific sequence of DNA inserted into a known location on a chromosome that is scrambled randomly in each neuron; and, (b) a transiently expressed special Recombinase protein.
- Challenge 2: How to induce each neuron to share copies of its barcode with each connected neuron? Approach: Using a virus such as the pseudorabies virus ((PRV) that is essentially a core of DNA wrapped in a protein coat. PRV unlike most viruses propagates from neuron to neuron across the synaptic cleft, thereby carrying the DNA sequence with it. However, the DNA is inverted between any pair of Recombinase sites.
- Challenge 3: How to join pre-synaptic and post-synaptic barcodes into a single molecule for high-throughput DNA sequencing? Approach: Express a specialized protein called an *integrase*, which irreversibly joins the DNA at the sites forming a single piece of DNA out of two. By positioning the barcode sequence near the *integrase* sites it can be ensured that the single piece of DNA contains two barcodes in sequence. The joined barcodes are then sent to a high throughput sequencing machine for processing.

The advantages of the BOINC approach are that it is relatively inexpensive<sup>15</sup> and that its error rate does not increase with the length of the synaptic chain (i.e., pathway) even if there are millions of neurons involved. However on the negative side, it does not include any spatial representation and provides no indication of whether a neuron is excitatory or inhibitory because it cannot differentiate between cell types.

*Whole brain neuroimaging:* A team comprised of Ahrens, Engert, Keller and others at the Howard Hughes Medical Institute (Ashburn, Virginia, USA) has developed an experimental microscopy technique for the holistic study of the sensorimotor system of the zebrafish, with the ability to simultaneously record about 80,000 of the total 100,000 neurons (Ahrens et al. 2012). They partially paralyzed the zebrafish so that only the connections between the neurons and the muscles are inactive. This allows the neural commands to the muscles to be recorded while the brain is stationary and the zebrafish is exposed to a virtual reality environment displayed underneath them using a video projector.

Using the fluorescent protein technique (i.e., staining) the researchers are able to track the activity of some 80,000 neurons at the rate of several times per second. Nevertheless, the question remains how to make sense of the enormous volume of data collected? For example, one neuron may simply reflect the activity level of another. They identified groups of neurons that appeared to be active together at some points in time and inactive at others, and found these groups to have a well defined anatomical structure (e.g., six tightly packed and symmetrically arranged clumps of neurons). However, why these neuron groups appeared to be tightly communicating remains unknown. In a broader sense, how do large groups of neurons coordinate to represent a feature of the visual environment, or how does

<sup>&</sup>lt;sup>14</sup> The BOINC acronym stands for Barcoding Of Individual Neuronal Connections.

<sup>&</sup>lt;sup>15</sup> The cost of sequencing an entire human genome has fallen dramatically in recent years; - from (US)\$1 million in 2007 to around (US)\$1,000 in 2015.

the entire brain respond to a signal that suggests the presence of food and guide the zebrafish to food?

**Recording many neurons simultaneously:** Until now the recording of neuron activity has been restricted to either measuring the activity of one neuron at a time or measuring the aggregate activity of entire networks of thousands to millions of neurons. Recent reports in the literature suggest that researchers may be on the threshold of being able to concurrently measure the individual activities of several hundred neurons and that further significant increases in the number of concurrent recordings may be on the horizon (Shenoy 2015, 78).

While the action potential and chemical activity (e.g., neurotransmitters and ion concentrations) of individual neurons has been measured with intracellular and extracellular electrodes, the aggregate activity of neural networks has been recorded with EEG, MEG and fMRI facilities. Some of these techniques have now scaled up and new techniques are being developed. For example genetically encoded calcium indicators (GCaMP) allow calcium concentration changes due to action potential to be optically imaged in thousands of neurons simultaneously (Ahrens 2015). Traditional electrode approaches for measuring the activity of single neurons have also scaled up to hundreds of neurons with electrode arrays (Shenoy 2015, 79-81). However, there remain at least two serious obstacles that need to be overcome before the simultaneous recording of much larger numbers of neurons can be realized.

First, there is the need to analyze the parallel data from thousands of neurons. As our knowledge of brain operations increases we must expect the discovery of additional variables (e.g., cell type, axon and dendrite projection patterns, synaptic connection strength) that will need to be recorded and a much greater increase in the combinations of these variables due to interdependencies. By comparison, it would be a far easier undertaking to try to gain an understanding of how a computer functions by measuring the electrical activity of its millions of transistors simultaneously. It is doubtful that such an undertaking would lead to more than a superficial understanding of computer operations even though there are no concurrent chemical processes involved in a computer and the number of connections between transistors is far fewer than there are synapses in the brain.

Second, there is the question of degree of importance of the individual variables and their relationships. A further complication could arise if it were to be found that the influence of specific combinations of variables becomes relevant only under certain circumstances that are related to the particular task undertaken by the brain. For example, whether the degree of synaptic strength is an important factor in the movement of a limb may depend on the cause of the movement rather than the accomplishment of the movement task.

# **Hypothetical Explanations**

While the severe limitations of current experimental techniques do not allow what might appear to be plausible hypotheses to be validated into theories, there are four hypotheses that the author would like to briefly discuss; namely, comprehension through the assembly of parts, ability to retain information in memory, influence of genes, and randomness in the formation of synapses.

In general terms the hypotheses are based on the assumption that neurons are assembled into networks and that the brain is in fact a neural network of networks. While any neuron will have a number of connections within its own assembly of cells (i.e., subnetwork) it may also have one or more connections to neurons in other networks. Therefore, a spike in one neuron may activate

several neurons in other networks apart from activating the connected cells in its own assembly of cells.

*Hypothesis 1 - Comprehension through the assembly of parts:* According to this proposition the comprehension of, for example, a visual sensory input such as the scene shown in Figure 7 starts with the activation of neurons that respond to very low level properties of the image such as the luminance of one or more distinct areas or dots or lines or planes in the image.



Figure 7: Comprehension through hierarchical feature detection

The firing of these neurons will activate other connected neurons that respond to other features that may or may not build on the partial interpretations of the previous level neurons, and so on. With each successive level of response the comprehension of the whole image becomes more complete and the contribution of the next level builds more and more on the cumulative results of the previous levels. Throughout this process involving millions of neurons in thousands of subnets there will be not only excitatory but also inhibitory neuron responses. It is in this way that the evolving comprehension of the image is continuously challenged, rejected or validated, and fine tuned. At lower levels these challenges and validations are likely to involve factors that are based more on conditioned (subconsciously acquired) experience, while at higher levels conscious memory might be expected to play an increasing role.

This explanation is based on the proposition that the brain consists of neurons that respond to particular stimuli. At the lowest level(s) these stimuli are very primitive elements like spots of light or sound vibrations that by themselves provide no clues of what the eventual outcome of the perception (i.e., the whole) is going to be, except for the gross distinction between visual, auditory, olfactory, somatic (touch), gustatory (taste), and vestibular (movement) senses. At succeeding levels the stimuli that the neurons respond to become increasingly more indicative in their representation of the whole.

Along the way individual neurons will receive and/or send both excitatory signals and inhibitory signals. While the excitatory signals indicate agreement with the current path to comprehension, the inhibitory signals indicate disagreement. This means that along the entire hierarchy from the lowest to the highest level there are many decision points that will influence the final comprehension of the whole. This process is based on the following hypothetical rules (Figure 8) for which there exists some experimental evidence but not enough for definitive validation (Seung 2013, 62-75):

1. A neuron that detects a 'whole' receives excitatory synapses from neurons that detect its parts.

This rule applies to how higher level neurons in the hierarchy of a network receive signals from lower level neurons. In the opposite direction the following rule would govern how a neuron sends signals to one or more higher level neurons.

2. A neuron that detects a part (i.e., something that it has been conditioned to recognize) sends excitatory synapses to higher level neurons thereby contributing to the comprehension of the 'whole'.

For example, the comprehension of a person in the visual field could start with outlines of one or more limbs such as the head, arms, or legs. At higher levels the limbs are combined into a human body and particular details such as the face, clothing, color, and posture, are added depending on the distance of the person from the beholder and the

nature of the interaction. At the higher levels recognition of the person as a known acquaintance would be triggered by connections to other networks that respond to particular persons or other associated memories. If the person is located close to the beholder then the comprehension sequence start immediately with may primitive features of the face, lips or eyes (e.g., pupil, iris, white, and blue leading to blue eyes).



Figure 8: Hypothesis 1 – Potential Formal Rules

In this case the visual stimuli are likely to occur simultaneously with any auditory stimuli that would be processed concurrently by other networks of neurons from primitive sound elements to speech, to language, to meaning, to relevant knowledge. Seung (2013, 68) suggests that this leads to a third rule:

- 3. The function of a neuron is defined chiefly by its connections to other neurons.
- *Hypothesis* 2 *Ability to retain information in memory:* It has been generally agreed for several decades that long term memory in the human brain is associative. If we have difficulty recalling information about a past event or experience we find it helpful to establish a line of thought by thinking about related information that may eventually lead us to the information that we are seeking. For example, if we are randomly asked to reconstruct where we were and what we were doing on a particular day 20 years ago then we would most likely initially draw a complete blank. However, given enough incentive to try to recollect we would start by broadening the targeted information into where was I

living during that year, what was my principal occupation, who were my closest friends, and so on? One or more of these related information items would likely eventually lead into a line of thought that would allow me to recall some of the particulars about that time period in my life.

It appears that the connectivity of cells in networks of networks is the basis of memory. If

this is true then the focus of our attention in determining how the brain forms memories must be on the plasticity of synapses, in respect to: how are cell assemblies formed; how is the strength of a synapse increased or decreased; how is the action potential of a neuron increased or decreased; what role (if any) does ordering in time play; to what extent is repetition a factor; and, can new connections be formed?



Figure 9: Hypothesis 2 - Memory

In respect to repetition it has been proposed (Figure 9) that we learn to associate ideas when one idea repeatedly accompanies or succeeds another. This has led to the following rule of plasticity:

1. If neuron A and neuron B are repeatedly simultaneously activated, then the connections between them are strengthened in both directions.

A similar rule has been proposed for our ability to learn sequential ideas (Hebb 1949):

2. If neuron A and neuron B are repeatedly activated sequentially, then the connection from neuron A to neuron B is strengthened.

Both of these rules have become known as the Hebbian rules of plasticity and are applicable only to synapses between excitatory neurons. In his 1949 book (The

Organization of Behavior) Hebb also proposed that neurons are connected in cell assemblies. Since the 1950s it has been possible to measure the spiking strength of a neuron by inserting a glass electrode with an extremely sharp tip into a single neuron. In the 1970s it became possible to measure changes in synaptic strength by repeatedly stimulating the spiking of two connected neurons.



Figure 10: Hypothesis 2 – Process of forgetting

While it is difficult to keep two neurons alive after they have been repeatedly penetrated, there is experimental evidence to suggest that the increased synaptic strength can be persistent for time periods of hours to days and weeks (Figure 10).

Seung (2013, 76-95) traces the logic of the corollary as follows. Neurons whose synaptic strengths have increased through some particular kind of sensory stimulation will through their connectivity form an assembly of neurons that have now become sensitive to this type of sensory stimulus. We can call this a synaptic chain based on a pattern of strong connections. Any stimulus that activates the first group of lower level neurons will activate the higher level neurons in this chain (chain A).

It would be reasonable to expect that a stimulation coming from another cell assembly (chain B) through one or more synapses anywhere in chain A could activate the neurons in chain A. If the connections are sufficiently strong then the spiking will propagate through the entire chain without the need for the particular sensory stimulus that was originally responsible for the formation of chain A. This would provide an explanation of how the brain operates in support of associative memory; - i.e., any stimulus that activates chain A will trigger the recollection of the sequence of ideas (memories) that are associated with chain A and every successive recollection of these memories will further strengthen the connections of chain A by Hebbian plasticity.

*Hypothesis* 3 - Influence of genes: The relative influences of genes and life experience on the behavioral characteristics of an individual person is still very much a topic of debate. While there is little doubt about the influence of the genes of the parents on the

physical features of children, it is not clear to what extent the behavioral similarities between parents and child are due to nature (genes) or nurture (upbringing). Studies of monozygotic (MZ) twins with 100% of the same genes and dizygotic (DZ) twins with 50% of the same genes have shown that twins score more similarly on intelligence tests than two randomly selected persons (Figure 11).



Figure 11: Hypothesis 3 – The role of genes

However, twins score less similarly on personality tests than on intelligence tests but still more similarly than two randomly selected persons. Turkheimer (2000) has formulated

three laws of behavior genetics:

- 1. All human behavioral traits are heritable.
- 2. The effect of being raised in the same family is smaller than the effect of genes.
- 3. A substantial portion of the variation in complex human behavioral traits is not accounted for by the effects of genes or families.



Figure 12: Hypothesis 3 – Potential formal rules

Turkheimer's first law holds not only for mental differences but also for mental disorders between persons (Figure 12). In the case of twins it was found that the concordance rate of autism for MZ twins has a 60% to 90% probability, but only 10% to 40% in DZ twins. In the case of schizophrenia it is 40% to 65% for MZ twins and only 0% to 30% for DZ twins. In other words, these studies of twins show that genes matter, but they do not explain why.

*Hypothesis 4 – Randomness in the formation of synapses:* Little is known why certain neurons form connections and others do not. With the high density of neurons hundreds of thousands of dendrites are intertwined with axons in every cubic millimeter of brain tissue. In this thick forest of neurites there are many more opportunities for the formation of synapses than are actually formed. It is interesting to note that although the human brain has an estimated 1000 trillion neurons this very large number is in fact only a small fraction of the number of synapses that would exist if each neuron were to be connected to every other neuron. With 86 billion neurons there would not be nearly enough room in the skull to accommodate the required number of synapses.

It would not seem to be an unreasonable proposition that with so many opportunities for the formation of synapses that there is at least initially a degree of randomness involved. We do know that initially there are more synapses than are required. In comparing the density of neurites in the brain of a two-year old child with a four-year old child we see that noticeable pruning has already taken place (Figure 4). The pruning continues with an approximately 40% reduction in synapses by the time the child reaches adulthood. It could be argued that any new synapse either finds use and becomes part of potentially many synaptic chains or it never gains much strength and is eventually eliminated (Seung 2013, 87-88).

## **Concluding Assessment of Feasibility and Value**

In general terms reverse-engineering means decomposing a system (i.e., a whole) systematically into its component parts without prior knowledge of the model of the whole and the logic of each individual part. The objective of reverse-engineering is to build a system with the same behavior by discovering the model of the whole through discovery of the logic of its component parts. However, with a system as complex as the human brain the behavior of the whole may not be able to be extrapolated from the behavior of the parts. In other words, the interaction of neurons through synapses and possibly other glial connections may not be predictable from an understanding of the fundamental operations of each neuron type.

In this regard, the number and complex nature of the problems confronting neuroscience researchers call into question the feasibility of the undertaking and the advisability of channeling a significant proportion of available government funding into brain research. On the other hand, it can be argued that even if the research findings fall far short of the original objectives they will be of enormous benefit to improving the health of mankind through the treatment of mental and physical disorders. It therefore seems appropriate to conclude this paper with a comparative discussion of the feasibility of the attempt to reverse-engineer the brain and the benefits that may accrue to clinical medicine, psychiatry and psychology.

*Feasibility:* There are many fundamental problems facing neuroscience research today. While continued advances in experimental techniques and the combination of microscopy

with high speed computer analysis are very promising, there is no guarantee that they will be sufficient to gain a full understanding of the operation of the human brain.

Zador (2015, 40-49) suggests that if the functional behavior of the brain is based on a general biological algorithm with many exceptions then the reverse engineering approach may not lead to the kind of overarching theory that neuroscience researchers are seeking. With about 86 billion neurons and 1000 trillion connections there could be a staggering number of exceptions. In fact one of the reasons for the large number of neurons and the complexity of the brain may be that the brain operates according to a relatively small number of principles with a very large number of exceptions. Under these circumstances recording the behavior of single neurons within neural circuits would be an endless task of discovering exceptions and lead to the danger of assuming that some of these exceptions are sufficiently common to be mistaken as an overarching theory.

The challenges confronting neuroscience researchers appear to be insurmountable within the current limitations of laboratory experimentation that cannot take into account the enormous scale of the human brain involving a vast range of stimuli in a dynamically changing environment. As Freemen (2015, 100) and other researchers point out, the experimental context is so limited in scale that it may mislead rather than usefully predict much about the rules that govern cognition, memory, reasoning, planning, and imagination. For example, the calcium fluorescence image from an experiment is often so messy and indirect that the researcher has difficulty inferring what the brain appears to be doing and in particular to separate the expected experimental signals from noise due to other factors.

Even imaging all neurons is not enough to fully understand brain functions, because looking at brain activity in isolation ignores the physical context of the brain. The brain resides inside the human body with its own environment and the body resides within the external natural environment (Chiel and Beer 1997). The brain through its massively connected neurons is part of the sensorimotor loop. Decisions made by the brain lead to actions that produce bodily and environmental changes that feed back to the brain as new sensory input that is again processed by the brain. This loop is important. For example, it appears that aural input from a decision to listen to a speaker is processed differently from the same aural input in the absence of that decision. It follows that understanding the brain requires an understanding of the entire holistic system of brain, body and environment.

Donoghue (2015, 222-3) gives a simple example in which his team recorded the spiking of two neurons in the motor cortex of an animal as it reaches from one point to another. Combined in the motor cortex commands to move are the results of interactions with many other brain structures (i.e., a network embedded in a network). The experimental record shows that one of the two neurons spikes significantly stronger whenever the animal reaches to the left and the other neuron spikes strongly when the animal reaches up and weakly when reaching down. This would suggest that when both neurons spike strongly the animal is reaching upward to the left. However, the experiment recorded only two neurons out of millions and those two neurons were recorded in isolation from other brain structures that they interact with. Therefore, the assumption that there is a direct relationship between the two neurons and the reaching action of the animal may well be erroneous.

In more general terms, identifying the behavior generated by the brain in any particular situation must occur concurrently with how the brain generates the necessary actions. So far it has not been possible to extract principles of brain function from recordings of neuronal

activity. For example, after decades of research the approximately 30-neuron network that controls the stomach system of decapod crustaceans (e.g., crayfish, crabs, lobsters, prawns, etc.) is still not fully understood. This casts considerable doubt on the ability to understand the 100,000 neuron network of the zebrafish, let alone the 86 billion neuron network of the human brain. The situation is exacerbated by the fact that the current methods for recording neuronal activity are relatively slow compared with the millisecond timescale at which neurons communicate. Order of magnitude increases in recording speeds will be necessary to improve upon the current ability to observe each neuron only a few times per second.

There are two additional problems that confront neuroscience researchers. First, the inability to infer higher level human behaviors from animal studies because human behaviors like language and reasoning do not have comparable animal models. Second, the extreme restriction on invasive studies of the human brain, which are essentially limited to the targeted monitoring of specific brain functions during neurosurgery. Therefore, studies of the human connectome are limited to postmortem analysis of human brain images and living animal studies. Non-invasive imaging methods do not allow researchers to discriminate between individual neurons but only brain regions. While phrenologists<sup>16</sup> have attempted to explain mental differences in terms of the functions and sizes of regions of the brain, they are unable to explain how each region performs its function. Seung (2013, xx) points out that the influence of experience on the organization of the connectome is still a hypothesis that has not been validated by conclusive laboratory tests because of the inadequacies of experimental techniques. While the findings of phrenology can be empirically tested, they are too simplistic. At the same time, the much more sophisticated ideas of connectionism cannot be evaluated experimentally.

**Potential benefits:** Several reasons may be offered why research of the brain is worthy of an investment of billions of dollars. First, there is the significant and growing health, social, and economic impact of brain and mental disorders that could be mitigated by the findings of brain research that would help us to compare healthy and unhealthy brains, develop new medical interventions, and provide an understanding of the biological mechanisms that the brain employs and how they fail in diseases, mental disorders, and following injuries. According to Seung (2013, 100-115) an accurate and comprehensive cataloging of the connectome would provide valuable insight into the sources of many mental disorders; - for example, whether autism and schizophrenia are due to macrocephaly during the creation of neurons, or lissencephaly during the migration of neurons to the brain, or connectopathies during the later stages of neurite growth and the creation of synapses.

Second, due to military conflicts and terrorist attacks there are an increasing number of physical injuries leading to amputation of limbs that require prosthetic replacements with the desirability of linkage to the brain's motor control mechanisms and feedback from the prosthesis to the sensory reception mechanisms.

Third, an understanding of how the brain operates is potentially a major aid in the quest for Augmented Intelligence, Artificial General Intelligence and Artificial Superintelligence.

<sup>&</sup>lt;sup>16</sup> Phrenologists have in the past associated the regions of the human brain and their size to the personal traits of an individual. While their belief that character, thoughts and emotions are located in particular regions of the brain was an important historical contribution to neuroscience, further extensions of this notion in respect to the size of any of these regions is now regarded as being obsolete. Their hypothesis that differences in the mind are due to differences in the brain was not unreasonable based on the limited methods available at the time.

There is a justified expectation that a deeper understanding of brain operations will lead to the development of methodologies and algorithms to improve artificial intelligence (i.e., machine learning) and the discovery of new technologies that exploit the physical principles exhibited by neural computation. Specifically in the area of network theory, there is a need for greater insight into the behavior of complex networks, the discovery of new principles underlying network operations, and the development of new theories describing network behavior (Eliasmith 2015).

Finally, there is the innate driving need of human curiosity to gain an understanding of brain functions as a means of unraveling the mystery of one of the most complex problems facing humanity.

It is highly probable that advances in neurotechnology will produce an entirely new set of much more powerful clinical applications. Early examples include: cochlear implants that convert sound into electrical impulses that are delivered to auditory nerves in the ear; deep brain stimulation (i.e., neuromodulation) using electrodes to electrically alter neural circuits to, for example, reduce the tremor in Parkinson's disease; transcranial magnetic stimulation to temporarily relieve symptoms of depression despite the fact that magnetic devices are too coarse to target specific brain circuits; and, brain-computer interfaces to create replacement commands for actions that can be no longer performed by paralyzed persons.

Research aimed at providing an implantable, untethered interface between the brain and a prosthetic device is facing challenges that would be ameliorated by a deeper understanding of brain operations (Maharbiz 2015). The engineering design of such a neural interface that is capable of lasting a lifetime requires knowledge of the nature of the physical substrate, how to avoid the biotic (i.e., organismic) and abiotic (i.e., chemical and physical) factors that lead to performance degradation at the electrode-tissue interface, the spatial coverage and density of the sensing site, the type of signals, and the communication capabilities. Both the neurons that are being recorded and the algorithms that are decoding the neural signals should have the ability to learn to improve the performance of the prosthetic device. The ultimate objective is a degree of control of the prosthetic device that will allow the user to perform tasks in a natural and effortless manner.

The large amount of funding being allocated by governments in support of the substantial research efforts in the field of neuroscience appear to be entirely justified. Even though the ultimate objective of complete knowledge of how the brain works may not be realized for several generations, if ever, the benefits to mankind of even partial understandings are considerable.

## References

Ahrens M. (2015); 'Whole Brain Neuroimaging and Virtual Reality'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 17-24).

Ahrens M., J. Li, M. Orger, D. Robson, A. Schier, F. Engert and R. Portugues (2012); 'Brain-wide neuronal dynamics during motor adaptation in zebrafish'; *Nature*, 485 (7399), (pp. 471-477).

Amunts K., C. Lepage, L. Borgeat, H. Mohlberg, T. Dickscheid, M. Rousseau, S. Bludau, P. Bazin, L. Lewis, A. Oros-Peusquens, N. Shah, T. Lippert, K. Zilles and A. Evans (2013); 'BigBrain: An Ultrahigh-Resolution 3D Human Brain Model'; *Science*, 340(6139), June (pp. 1472-1475).

Brodmann K. (1909) – English translation by Garey L. (2006); 'Brodmann's localization in the cerebral cortex: The principles of comparative localization in the cerebral cortex based on cytoarchitectonics'; Springer, New York, New York.

Chiel H. and R Beer (1997); 'The brain has a body: Adaptive behavior emerges from interactions of nervous system,

body and environment'; Trends in Neurosciences, 20 (12), (pp. 553-557).

Church G., A. Marblestone and R. Kalhor (2015); 'Rosetta Brain: The Multileveled Complexity of the Brain'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 50-64).

Denk W. and H. Horstmann (2004); 'Serial Block-Face Scanning Electron Microscopy to Reconstruct Three-Dimensional Tissue Nanostructure'; *PLoS Biology*, 2(11), October.

Donoghue J. (2015); 'Neurotechnology'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 219-233).

Edelman G. (1987); 'Neural Darwinism: The Theory of Neural Group Selection'; Basic Books, New York, New York.

Eliasmith C. (2015); 'Building a Behaving Brain'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 125-136).

Freeman J. (2015); 'Large-Scale Neuroscience: From Analytics to Insight'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 100-107).

Gosline A. (2005); 'Why your brain has a Jennifer Aniston cell'; New Scientist, Daily News, 22 June.

Hagmann P. (2005); 'From Diffusion MRI to Brain Connectomics'; PhD Thesis, STI School of Engineering, University of Lausanne: EPLF.

Hawrylycz M. (2015); 'Building Atlases of the Brain'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 3-16).

Hayworth K., J. Morgan, R. Schalek, D. Berger, D. Hildebrand and J. Lichtman (2014); 'Imaging ATUM Ultrathin Section Libraries with WaferMapper: A Multi-Scale Approach to EM Reconstruction of Neural Circuits'; *Front Neural Circuits*, 68(8), June.

Hebb D. (1949); 'The organization of behavior: A neuropsychological theory'; Wiley, New York, New York.

Hill S. (2015); 'Whole Brain Simulation'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 111-124).

Huttenlocher P. (1990); 'Morphometric study of human cerebral cortex development'; *Neuropsychologia*, 28 (6) (pp. 517).

Huttenlocher P. and A. Dabholkar (1997); 'Regional differences in synaptogenesis in human cerebral cortex'; *Journal of Comparative Neurology*, 387 (2) (pp. 167-178).

Johnson C. (2012); 'Scientist's Game Helps Map the Brain: MIT researcher creates online game to gather help unlocking brain's mysteries'; *Boston Globe*, 16 July.

Klein M. and E. Kandel (1978); 'Presynaptic modulation of voltage-dependent  $Ca^{2+}$  current: Mechanism for behavioral sensitization in Aplysia califonica'; Proc. National Academy of Sciences of the USA (PNAS), 75 (7) (pp. 3512-7).

Koch C. and G. Marcus (2015); 'Neuroscience in 2064: A look at the Last Century'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 255-269).

Lashley K. (1929); 'Brain mechanisms and intelligence: A quantitative study of injuries to the brain'; University of Chicago Press, Chicago, Illinois.

Lichtman J. and H. Coleman (2000); 'Synapse elimination review and indelible memory'; Neuron, 25 (pp. 269-278).

Lichtman J., J. Livet and J. Sanes (2008); 'A Technicolor Approach to the Connectome'; *Nature Reviews Neuroscience*, 9(6), June (pp. 417-422).

Maharbiz M. (2015); 'Neural Dust: An Untethered Approach to Chronic Brain-Machine Interfaces'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 243-251).

Marcus G. (2015); 'The Computational Brain'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 205-215).

Markram H. (2006); 'The Blue Brain Project'; Nature Reviews Neuroscience, 7, February (pp. 153-160).

Micheva K. and S. Smith (2007); 'Array Tomography'; Neuron, 55(1), July (pp. 25-36).

Moser M. and E. Moser (2015); 'Understanding the Cortex through Grid Cells'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 67-77).

Nguyen T. (2010); 'Total Number of Synapses in the Adult Human Neocortex'; Undergraduate Journal of Mathematical Modeling: One + Two, Vol. 3, Issue 1, Article 26 [http://scholarcommons.usf.edu/ujmm/vol3/iss1/26]

Porter K. and J. Blum (1953); 'A Study in Microtomy for Electron Microscopy'; *The Anatomical Record*, 117(4), December (pp. 685-709).

Purves D. and J. Lichtman (1985); 'Principles of neural development'; Sinauer Associates, Sunderland, Massachusetts.

Ren G. et al. (2016); 'Three-Dimensional Structural Dynamics and Fluctuations of DNA-Nanogold Conjugates by Individual-Particle Electron Tomography'; *Nature Communications*, March.

Seung S. (2013); 'Connectome: How the Brain's Wiring Makes Us What We Are'; Mariner Books, New York, New York.

Shenoy K. (2015); 'Recording from Many Neurons Simultaneously: From Measurement to Meaning'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 78-89).

Smedemark-Marguilies N. and J. Trapani (2013); 'Tools, Methods, and Applications for Optophysiology and Neuroscience'; *Frontiers in Molecular Neuroscience*, 6 (18).

Sporns O. (2015); 'Network Neuroscience'; in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 90-99).

Sporns O., G. Tononi and R. Kötter (2015); 'The Human Connectome: A Structural Description of the Human Brain'; *PLoS Computational Biology*, 1(4): e42.

Turkheimer E. (2000); 'Three laws of behavior genetics and what they mean'; *Current Directions in Psychological Science*'; 9 (5) (pp. 160).

Yang G., F. Pan and W. Gan (2009); 'Stably maintained dendritic spines are associated with lifelong memories'; *Nature*, 462 (7275) (pp. 920-4).

White J., E. Southgate, J. Thomson and S. Brenner (1986); 'The Structure of the Nervous System of the Nematode Caenorhabditis elegans'; *Philosophical Transactions of the Royal Society*, B 314, November (pp. 1-446).

Wong T., R. Preissl, P. Datta, M. Flickner, R. Singh, S. Esser, E. McQuinn, R. Appuswamy, W. Risk, H. Simon and D. Modha (2012); '10<sup>14</sup>'; IBM Research Report, Almaden Research Center, San Jose, California, November.

Yang X., Y. Kaeser-Woo, Z. Pang, W. Xu and T. Südhof (2010); Neuron, 68(5), December (pp. 907-920).

Zador A. (2015); 'The Connectome as a DNA Sequencing Problem' in Marcus and Freeman (eds.) The Future of the Brain, Princeton University Press, Princeton, New Jersey (pp. 40-49).

Zhang L. and G. Ren (2012); 'High-Resolution Single-Molecule Structure Revealed by Electron Microscopy and Individual Particle Electron Tomography'; *Journal of Physical Chemistry & Biophysics*, 2:e103, May.

#### **Appendix: Molecular Biology Fundamentals**

The *genome* of a living organism is made up of *chromosomes* that are made up of *genes*, which are composed of *deoxyribonucleic acid (DNA)*. DNA is made up of atoms that are arranged in a particular order on a double helix structure. Genes are pieces of DNA with particular variations in the order of atoms depending on the type of gene that form into chromosome strands. Every human cell contains 23 pairs of chromosomes in its nucleus (Figure 13). *Ribonucleic acid (RNA)* is the chemical messenger that carries the DNA instructions to the cell cytoplasm for execution.

While a cell such as a *neuron* is built from many types of molecules, one of the main types is a class of molecules known as *proteins*. Some of these proteins have structural functions (i.e., supporting the cell like the frame of a house), others perform functions on other molecules, and many proteins combine both structural and functional roles. Just like DNA is a chain of *nucleotides*, a protein molecule is a chain of smaller molecules called *amino acids*. Since there are 20 types of amino acids each protein is specified by a sequence of 20 letters rather than the four letters used in DNA. This amino acid sequence represents a gene in the genome.

The cell reads the nucleotide sequence of a gene and translates this into an amino acid sequence to synthesize a protein. When a cell reads a gene and constructs a protein, it is said to *express* the

gene. Every cell in a person's body contains the genome. The question then arises: Why do a liver cell and a heart cell look different and perform different functions? The answer is that cells of different type express different genes.



Figure 13: DNA – Chromosomes – Genes

A genome contains tens of thousands of genes, each corresponding to a different kind of protein. Each type of cell expresses some of these genes but not others. Neurons are arguably the most complex type of cell in the body and therefore it is no surprise that many genes encode proteins that are either exclusively or partially devoted to supporting functions in neurons. For this reason alone it would appear that genes are likely to play a role in the operation of the brain.

**DNA:** Deoxyribonucleic acid (DNA) is a molecule that contains the information that governs the way cells function in an organism. The information is encoded in a double helix (ladder like) structure of atoms in the form of a strand. A single human DNA molecule carries a vast amount of information equivalent to 4000 500-page books.

- 1. The double helix structure of DNA is like two ladders twisted around each other.
- 2. DNA is made up of molecules called nucleotides. Each nucleotide contains a phosphate group, a sugar group and a nitrogen base. There are four types of nitrogen bases: adenine (A); thymine (T); guanine (G); and, cytosine (C). Adenine (A) always pairs with thymine (T) and guanine (G) always pairs with cytosine (C). The order or sequence of these nitrogen bases determines genetic code (i.e., instructions).
- 3. Sometimes during cell division the sequence or order of the nitrogen bases is not reproduced exactly resulting in a mutation (i.e., mutant gene).
- 4. In humans half of the DNA comes from the mother and half from the father at conception. Therefore much of the biological blueprint of a person is inherited from the parents.

- 5. Apart from being able to store information DNA can also replicate itself (i.e., exact copy) during cell division.
- 6. DNA instructions determine the production of enzymes, which are proteins. Enzymes ensure that the production of protein in a cell is assembled from the correct sequence of amino acids. Protein is the fundamental building block of cells.

*Chromosomes:* Chromosomes are composed of DNA strands that are made of genes. Thousands of genes make up the thread-like chromosomes that are about six feet long. Every human has 22 pairs of matching chromosomes plus a  $23^{rd}$  pair of sex chromosomes (i.e., XX for female and XY for male). Each cell (except sex cells) contains 23 pairs of chromosomes. Sex cells contain only 23 unpaired chromosomes so that at conception they can create one cell containing a full set of chromosomes. While chromosomes are normally hidden in the opaque nucleus of a cell, they can be chemically stained to become visible during cell division.

*Genes:* Genes are made up of DNA and control heredity. The DNA structures from the two parents can combine in an almost infinite number of ways. Therefore there is almost no probability that any two persons (except identical twins) will have exactly the same genetic profile.

*Genetics:* Genetics is the science of the mechanisms of heredity and the variations that can occur in both animals and plants.

*Genome:* A Genome is all of the genetic material in a living organism. It is the entire set of DNA instructions for building, operating and maintaining an organism. Genome Sequencing is the process of determining the order of DNA nucleotides (i.e., bases) in a genome.

*Cells:* A cell is the basic structural and functional unit of all living organisms. Cells are the smallest unit of life that can replicate independently. Cell cytoplasm is a thick solution that fills each cell and is enclosed by the cell membrane. It includes all of the material inside the cell, but outside of the cell nucleus.

**RNA:** Ribonucleic acid (RNA) acts as a chemical messenger that communicates the DNA's instructions to the cell cytoplasm. There are three types of RNA:

- 1. Ribosomal RNA is made in the cell's nucleus and is a direct copy of the gene that is being exported along messenger RNA (mRNA) pathways through the cytoplasm.
- 2. Transfer RNA (tRNA) acts as an adaptor that allows the information-coded ribosome to attach to successive amino acids and trigger the release of the appropriate enzymes to construct them into a growing protein chain.
- 3. Like DNA, RNA also contains four bases of nucleotides. Since the average protein contains about 400 amino acids and the coded signal is known to be three bases in a row, it follows that each protein message must be about 1,200 nucleotides long.