

This is a repository copy of *Gravitational waves emitted by a particle rotating around a Schwarzschild black hole : A semiclassical approach*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/112435/>

Version: Accepted Version

Article:

Higuchi, Atsushi orcid.org/0000-0002-3703-7021, Bernar, Rafael P and Crispino, Luis C B (2017) Gravitational waves emitted by a particle rotating around a Schwarzschild black hole : A semiclassical approach. *Physical Review D*. 064042. ISSN 1550-2368

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Gravitational waves emitted by a particle rotating around a Schwarzschild black hole: A semiclassical approach

Rafael P. Bernar,^{1,*} Luís C. B. Crispino,^{1,†} and Atsushi Higuchi^{2,‡}

¹*Faculdade de Física, Universidade Federal do Pará, 66075-110, Belém, Pará, Brazil.*

²*Department of Mathematics, University of York, YO10 5DD, Heslington, York, United Kingdom.*

(Dated: February 14, 2017)

We analyze the gravitational radiation emitted from a particle in circular motion around a Schwarzschild black hole using the framework of quantum field theory in curved spacetime at tree level. The gravitational perturbations are written in a gauge-invariant formalism for spherically symmetric spacetimes. We discuss the results, comparing them to the radiation emitted by a particle when it is assumed to be orbiting a massive object due to a Newtonian force in flat spacetime.

PACS numbers: 04.60.-m, 04.62.+v, 04.50.-h, 04.25.Nx, 04.60.Gw, 11.25.Db

I. INTRODUCTION

Black holes are among the most important predictions of General Relativity (GR). Several observations indicate the presence of supermassive black holes in the center of nearly all large galaxies [1, 2]. In addition, there is strong evidence for stellar-mass black holes having an influence on other stars in binary systems [3], emitting X-rays through accretion (see Ref. [4] for a review on observational evidence of stellar-mass and supermassive black holes). Moreover, black holes are believed to play an important part in powerful astrophysical processes, such as gamma-ray bursts [5]. The recent detections of gravitational waves [6, 7] emitted by binary black hole mergers make the study of black holes and radiation-emission scenarios even more appealing, particularly the emission of gravitational waves. Binary black hole systems can provide settings in which the extreme curvature of the black hole generates remarkable signatures which can, in principle, be experimentally detected. It is also interesting to study gravitational radiation emitted by a relatively small object which can be approximated by a point particle in circular orbit around a black hole, in highly relativistic motion, the so-called geodesic synchrotron radiation scenario. The possibility of this mechanism for gravitational synchrotron radiation was raised in Refs. [8, 9]. While studying the scalar radiation emitted by a point source in circular geodesic motion around a Schwarzschild black hole, it was argued that gravitational radiation emitted by the source would be mostly of the synchrotron type, which has frequencies much higher than the angular frequency of the orbit and radiation distributed in narrow angles. This was further investigated in Ref. [10] where the authors computed the high-frequency spectra of electromagnetic and gravitational radiation for a particle orbiting a Schwarzschild black hole. In Refs. [11, 12], using the Regge-Wheeler for-

malism [13–15] and Green’s function techniques [16–19], it was shown that the spectrum of gravitational radiation from a point particle around a Schwarzschild black hole is much broader than the scalar or electromagnetic ones, at least for high- l multipole modes. For stable orbits, a full analysis and numerical computations were done in Refs. [20, 21].

The framework of Quantum Field Theory (QFT) in curved spacetimes [22, 23] has been used at tree level to compute the (massless) scalar radiation of a point source in circular orbit around Schwarzschild [24, 25], Reissner-Nordström [26] and Kerr black holes [27]. The case of massive scalar radiation from a point source orbiting a stellar object or a Schwarzschild black hole was analyzed in Ref. [28]. Electromagnetic radiation from a point charge rotating around an uncharged static black hole was analyzed in Ref. [29], using the same semiclassical approach. Although this approach is found to give the same results as the classical methods (e.g. using Green’s function techniques), it will make the radiative quantum corrections to these results more straightforward. It also allows an alternative interpretation of the radiation processes discussed in this paper. We also find that by regarding the classical fields as quantum particles one can treat several aspects of the radiation phenomena in an unified manner.

In this paper we analyze the gravitational radiation emitted by a point particle in geodesic circular orbit around a Schwarzschild black hole using the framework of QFT in curved spacetimes at tree level. Using numerically obtained solutions, we compute the total emitted power, as well as the power radiated to infinity in both stable and unstable orbits. We also analyze the spectrum of the emitted radiation.

The rest of this paper is organized as follows. In Sec. II we present the formalism developed in Refs. [30, 31] for linear perturbations of the gravitational field around spherically symmetric spacetimes. We give a brief review of the formalism, specializing it to the background of a 4-dimensional Schwarzschild spacetime. In Sec. III we present the framework of QFT for linearized gravity in which we will work, applying this framework to the

* rafael.bernar@icen.ufpa.br

† crispino@ufpa.br

‡ atsushi.higuchi@york.ac.uk

case of a point particle emitting gravitational radiation, in geodesic circular motion around a Schwarzschild black hole. We also obtain numerical results for the emitted power of gravitational radiation. In Sec. IV we compare these results to an analogous case in flat spacetime, namely the radiation emitted by a particle orbiting a Newtonian massive object. We conclude this paper with some remarks in Sec. V. We present the derivation of the normalization factor of one type of the modes, the scalar-type modes, in Appendix A. Throughout this paper we use the metric signature $-+++$ and natural units such that $G = c = \hbar = 1$.

II. GRAVITATIONAL PERTURBATIONS IN SCHWARZSCHILD SPACETIME

In this section we present a brief review of the formalism developed in Refs. [30, 31]. By expanding suitably defined gauge-invariant quantities in terms of harmonic tensors, the perturbed Einstein's equations are reduced to a set of self-adjoint ordinary differential equations, one for each type of perturbation: scalar-, vector- and tensor-type gravitational perturbations. This formalism can be used for background spacetimes in any dimensions with some special isometries. Here we restrict this formalism to Schwarzschild spacetime in 3+1 dimensions.

A. Background Schwarzschild Spacetime

We work in the background spacetime of a chargeless nonrotating black hole of mass M , described by the line element:

$$\begin{aligned} ds^2 &= g_{\mu\nu} dx^\mu dx^\nu \\ &= -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \end{aligned} \quad (1)$$

where

$$f(r) = 1 - \frac{2M}{r}. \quad (2)$$

It is useful for us to define the line element of the orbit spacetime:

$$ds_{orb}^2 = g_{ab} dx^a dx^b = -f(r)dt^2 + \frac{dr^2}{f(r)}. \quad (3)$$

As we will see in Sec. IIB it is basically in the orbit spacetime that the dynamical equations for perturbations have to be solved. It is also useful to define the line element of the two-sphere S^2 :

$$d\sigma^2 = \gamma_{ij} dx^i dx^j = d\theta^2 + \sin^2\theta d\varphi^2. \quad (4)$$

The definitions (1), (3) and (4) above establish part of the notation used. Greek letters are used for spacetime indices running from 0 to 3, the first letters from the Latin

alphabet are used for the t and r components and letters i, j, k, \dots are used for the θ and φ components. Covariant derivatives and Christoffel symbols corresponding to ds^2 , ds_{orb}^2 and $d\sigma^2$ are denoted by ∇_μ , $\Gamma_{\mu\nu}^\alpha$; D_a , Γ_{bc}^a ; and \hat{D}_i , $\hat{\Gamma}_{jk}^i$ respectively.

B. Scalar-type and vector-type perturbations

Gravitational perturbations of the scalar-type are defined as the metric perturbations whose angular dependence is described by the scalar spherical harmonics $Y^{lm}(\theta, \varphi)$, which satisfy

$$(\hat{\Delta}_2 + k_S^2) Y^{lm}(\theta, \varphi) = 0, \quad (5)$$

with eigenvalues

$$k_S^2 = l(l+1), \quad l = 0, 1, 2, \dots, \quad (6)$$

where $\hat{\Delta}_2$ is the Laplace-Beltrami differential operator on S^2 , namely

$$\hat{\Delta}_2 \equiv \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta} \right) + \frac{1}{\sin^2\theta} \frac{\partial^2}{\partial\varphi^2}. \quad (7)$$

The solutions to Eq. (5) are given by

$$Y^{lm}(\theta, \varphi) = C^{lm} P_m^l(\cos\theta) e^{im\varphi}. \quad (8)$$

The normalization constants are [32]:

$$C^{lm} = \sqrt{\frac{(2l+1)(l-m)!}{4\pi(l+m)!}}. \quad (9)$$

The scalar-type metric perturbation modes $h_{\mu\nu}^{(S;lm)}$ can be written as follows [31]

$$h_{ab}^{(S;lm)} = f_{ab}^{(S;l)} Y^{lm}, \quad (10)$$

$$h_{ai}^{(S;lm)} = r f_a^{(S;l)} \mathbb{S}_i^{(lm)}, \quad (11)$$

$$h_{ij}^{(S;lm)} = 2r^2 \left(H_L^{S;l} \gamma_{ij} Y^{lm} + H_T^{S;l} \mathbb{S}_{ij}^{(lm)} \right), \quad (12)$$

where

$$\mathbb{S}_i^{(lm)} = -\frac{1}{k_S} \hat{D}_i Y^{lm}, \quad (13)$$

$$\mathbb{S}_{ij}^{(lm)} = \frac{1}{k_S^2} \hat{D}_i \hat{D}_j Y^{lm} + \frac{1}{2} \gamma_{ij} Y^{lm}. \quad (14)$$

The quantities $f_{ab}^{(S;l)}$, $f_a^{(S;l)}$, $H_L^{S;l}$ and $H_T^{S;l}$ are gauge-dependent quantities and functions of t and r only.

Gauge-invariant quantities can be defined for $l \geq 2$ and written in terms of a master variable $\Phi_l^S(t, r)$, which satisfies the following wave equation, resulting from the perturbed Einstein's equations:

$$\square \Phi_l^S(t, r) - \frac{V_S(r)}{f(r)} \Phi_l^S(t, r) = 0, \quad (15)$$

with the Zerilli effective potential [15]:

$$V_S(r) = f(r) \frac{2\lambda^2(\lambda+1)r^3 + 6\lambda^2Mr^2 + 18\lambda M^2r + 18M^3}{r^3(\lambda r + 3M)^2} \quad (16)$$

where

$$\lambda = \frac{1}{2}(l-1)(l+2). \quad (17)$$

The \square is the d'Alembert operator in the orbit spacetime, namely

$$\square \equiv -f(r)^{-1} \frac{\partial^2}{\partial t^2} + \frac{\partial}{\partial r} \left[f(r) \frac{\partial}{\partial r} \right]. \quad (18)$$

The derivation of Eq. (15) is highly involved and can be found in Refs. [30, 31]. Using the same gauge choice as in Refs. [33, 34], one can write the perturbation modes in terms of the master variable $\Phi_l^S(t, r)$ as

$$h_{ai}^{(S;lm)} = 0, \quad (19)$$

$$h_{ij}^{(S;lm)} = 2r^2 \gamma_{ij} F^l Y^{lm}, \quad (20)$$

$$h_{ab}^{(S;lm)} = F_{ab}^{(l)} Y^{lm}, \quad (21)$$

with

$$F^l = \frac{1}{4r^2 H} [(H - rf')\Omega_l^S + 2rD^a r D_a \Omega_l^S], \quad (22)$$

$$F_{ab}^{(l)} = \frac{1}{H} \left(D_a D_b - \frac{1}{2} g_{ab} \square \right) \Omega_l^S, \quad (23)$$

$$H = 2 \left(\lambda + \frac{3M}{r} \right), \quad (24)$$

and

$$\Omega_l^S = rH\Phi_l^S. \quad (25)$$

The mode with $l=0$ cannot be described in terms of the master variable of Eq. (15). However, it is a spherically symmetric perturbation, which, by Birkhoff's theorem, consists in a shift of the mass parameter of the black hole [16]. Hence, we will not consider this mode, since it is non-radiative. The $l=1$ modes can always be eliminated by a gauge transformation [31].

Gravitational vector-type perturbations are defined as the metric perturbations whose angular dependence is described by vector spherical harmonics satisfying the following equations

$$\left(\hat{\Delta}_2 + k_V^2 \right) Y_i^{(lm)}(\theta, \varphi) = 0, \quad (26)$$

$$\hat{D}^j Y_j^{(lm)}(\theta, \varphi) = 0. \quad (27)$$

The set of eigenvalues takes the form

$$k_V^2 = l(l+1) - 1, \quad l = 1, 2, 3, \dots \quad (28)$$

The solutions to Eqs. (26) and (27) on the unit 2-sphere are [13, 35]

$$Y_i^{(lm)}(\theta, \varphi) = \frac{\epsilon_{ij}}{\sqrt{l(l+1)}} \hat{\Delta}^j Y^{lm}(\theta, \varphi). \quad (29)$$

The Levi-Civita tensor on the S^2 is defined by

$$\epsilon_{\theta\theta} = \epsilon_{\varphi\varphi} = 0, \quad (30)$$

$$\epsilon_{\theta\varphi} = -\epsilon_{\varphi\theta} = \sin\theta. \quad (31)$$

The gravitational perturbation modes of the vector-type $h_{\mu\nu}^{(V;lm)}$ can be written as

$$h_{ab}^{(V;lm)} = 0, \quad (32)$$

$$h_{ai}^{(V;lm)} = r f_a^{(V;l)} Y_i^{(lm)}, \quad (33)$$

$$h_{ij}^{(V;lm)} = 2r^2 H_T^{V;l} \mathbb{V}_{ij}^{(lm)}, \quad (34)$$

with

$$\mathbb{V}_{ij}^{(lm)} = -\frac{1}{2k_V} \left(\hat{D}_i Y_j^{(lm)} + \hat{D}_j Y_i^{(lm)} \right). \quad (35)$$

By defining the gauge-invariant quantity for the modes with $l \geq 2$,

$$F_a^{(V;l)} = f_a^{(V;l)} + \frac{r}{k_V} D_a H_T^{V;l}, \quad (36)$$

we can write it in terms of a master variable $\Phi_l^V(t, r)$ as

$$rF_a^{(V;l)} = \epsilon_{ab} D^b (r\Phi_l^V). \quad (37)$$

The master variable satisfies the following equation:

$$\square \Phi_l^V(t, r) - \frac{V_V(r)}{f(r)} \Phi_l^V(t, r) = 0, \quad (38)$$

with the Regge-Wheeler effective potential [13]:

$$V_V(r) = f(r) \left(\frac{l(l+1)}{r^2} - \frac{6M}{r^3} \right). \quad (39)$$

The $l=1$ vector-type modes correspond to rotational perturbations, i.e. perturbations which give nonzero angular momentum to the background metric [16]. Again, we will not consider these non-radiative modes. In a specific gauge choice [33, 34], one can write the vector-type modes as follows

$$h_{ai}^{(V;lm)} = Y_i^{(lm)} \epsilon_{ab} D^b (r\Phi_l^V), \quad (40)$$

with all other components vanishing, where ϵ_{ab} is the Levi-Civita tensor in the orbit spacetime.

In $n+1$ dimensional spacetime with $n \geq 4$, there are also gravitational tensor-type perturbations, whose angular dependence is given by traceless tensor spherical harmonics. However, they do not exist on S^2 [13, 36]. (A concise proof of this fact can be found in Ref. [35].) Thus, there are no tensor-type modes for gravitational perturbations in $3+1$ dimensions.

III. QUANTIZATION AND GEODESIC SYNCHROTRON RADIATION

We will consider the case of a test¹ point particle in circular orbit emitting gravitational waves as it rotates around the black hole. We compute the emitted power using a semiclassical analysis, i.e. by considering the gravitational perturbations as a quantized field in the background Schwarzschild spacetime.

A. Quantization of gravitational perturbations in Schwarzschild spacetime

We quantize the field $h_{\mu\nu}$ in the same manner as in Refs. [33, 34] (see Ref. [37] for a more complete description.) The Lagrangian density of free linearized gravity in a background spacetime can be written as:

$$\mathcal{L} = \sqrt{-g} \left[\nabla_\mu h^{\mu\lambda} \nabla^\nu h_{\nu\lambda} - \frac{1}{2} \nabla_\lambda h_{\mu\nu} \nabla^\lambda h^{\mu\nu} + \frac{1}{2} (\nabla^\mu h - 2\nabla_\nu h^{\mu\nu}) \nabla_\mu h + R_{\mu\nu\lambda\sigma} h^{\mu\lambda} h^{\nu\sigma} \right], \quad (41)$$

where $h \equiv h^\mu{}_\mu$. The conjugate momentum current is given by

$$p^{\lambda\mu\nu} \equiv \frac{1}{\sqrt{-g}} \frac{\partial \mathcal{L}}{\partial (\nabla_\lambda h_{\mu\nu})}, \quad (42)$$

thus

$$p^{\lambda\mu\nu} = -\nabla^\lambda h^{\mu\nu} + g^{\lambda\mu} \left(\nabla_\kappa h^{\kappa\nu} - \frac{1}{2} \nabla^\nu h \right) + g^{\lambda\nu} \left(\nabla_\kappa h^{\kappa\mu} - \frac{1}{2} \nabla^\mu h \right) + g^{\mu\nu} (\nabla^\lambda h - \nabla_\kappa h^{\lambda\kappa}). \quad (43)$$

Note that we have not yet chosen any gauge condition. For any two solutions to the Euler-Lagrange equations, we define their symplectic product by

$$\Omega(h, h') \equiv - \int_{\Sigma} d\Sigma n_\alpha (h_{\mu\nu} p'^{\alpha\mu\nu} - p^{\alpha\mu\nu} h'_{\mu\nu}), \quad (44)$$

where $p^{\alpha\mu\nu}$ and $p'^{\alpha\mu\nu}$ are the conjugate momentum currents of the two solutions $h_{\mu\nu}$ and $h'_{\mu\nu}$, respectively, and Σ is a Cauchy surface with future-directed unit normal vector n^α . It can be shown that $\Omega(h, h')$ is independent of the choice of Σ [37, 38]. If there were no degeneracy, i.e. if there were no solutions $h_{\mu\nu}^{(null)}$ satisfying $\Omega(h^{(null)}, h) = 0$ for all solutions $h_{\mu\nu}$, one could define an inner product by

$$\langle h, h' \rangle = i\Omega(\bar{h}, h'), \quad (45)$$

where the overbar denotes complex conjugation. Suppose that a complete set of positive-frequency solutions, i.e. solutions whose time dependence is of the form $e^{-i\omega t}$, $\omega > 0$, is given by $\{h_{\mu\nu}^{(n)}\}$, where n represents all (continuous and discrete) labels. Then a positive- and a negative-frequency solution would be orthogonal to one another with respect to the inner product (45), and this inner product would be positive definite on the space of positive-frequency solutions. Then, we could expand the quantum field $\hat{h}_{\mu\nu}(x)$ as

$$\hat{h}_{\mu\nu}(x) = \sum_n \left[\hat{a}_n h_{\mu\nu}^{(n)}(x) + \hat{a}_n^\dagger \overline{h_{\mu\nu}^{(n)}}(x) \right]. \quad (46)$$

The canonical equal-time commutation relations would be equivalent to

$$[\hat{a}_m, \hat{a}_n^\dagger] = (M^{-1})_{mn} \quad (47)$$

and

$$[\hat{a}_m, \hat{a}_n] = [\hat{a}_m^\dagger, \hat{a}_n^\dagger] = 0, \quad (48)$$

where M^{-1} is the inverse of matrix $M^{mn} = \langle h^{(m)}, h^{(n)} \rangle$. However, due to gauge invariance, the symplectic product given by (44) is degenerate: a pure gauge solution $h_{\mu\nu}^{(gauge)} = \nabla_\mu \Lambda_\nu + \nabla_\nu \Lambda_\mu$ has vanishing symplectic product with any other solution, as it is well known. Thus, one needs to modify the quantization procedure described above. One standard way to proceed is to consider only the physical solutions, i.e. solutions satisfying gauge conditions that fix the gauge degrees of freedom completely. When all gauge degrees of freedom are eliminated, the symplectic product is non-degenerate and one quantizes the field by imposing the equal-time commutation relations given by Eqs. (47) and (48). We follow this procedure after fixing the gauge completely as described in Sec. IIB and normalizing the scalar- and vector-type modes, so that $M^{mn} = \delta^{mn}$, which may involve Dirac delta functions. Thus, we expand the quantum gravitational perturbation as in Eq. (46) in terms of positive- and negative-frequency solutions given by Eqs. (19)-(21) and (40), with definite frequencies ω . We require the positive-frequency solutions to be normalized with respect to the inner product (45) as follows:

$$\langle h^{(P;\omega lm)}, h^{(P';\omega' l' m')} \rangle = \delta^{PP'} \delta^{ll'} \delta^{mm'} \delta(\omega - \omega'), \quad (49)$$

where $P = S, V$ labels the type of the perturbations, with S denoting the scalar-type and V denoting the vector-type perturbations.

We write the positive-frequency modes of the master variables as

$$\Phi_{\omega l}^P(t, r) = e^{-i\omega t} u_{\omega l}^P(r), \quad \omega > 0. \quad (50)$$

Then the functions $u_{\omega l}^P(r)$ satisfy the following Schrödinger-like differential equation:

$$-f(r) \frac{d}{dr} \left(f(r) \frac{d}{dr} u_{\omega l}^P(r) \right) + (V_P(r) - \omega^2) u_{\omega l}^P(r) = 0. \quad (51)$$

¹ The word ‘‘test’’ is used here in the sense that the particle does not modify the background metric field.

Close to and far away from the horizon, both effective potentials given by Eqs. (16) and (39) tend to zero. Hence the two independent solutions of Eq. (51) can be written as

$$u_{\omega l}^{P,up} \approx \begin{cases} A_{\omega l}^P (e^{i\omega r^*} + \mathcal{R}_{\omega l}^{P,up} e^{-i\omega r^*}), & r \gtrsim 2M, \\ A_{\omega l}^P \mathcal{T}_{\omega l}^{P,up} e^{i\omega r^*}, & r \gg 2M; \end{cases} \quad (52)$$

$$u_{\omega l}^{P,in} \approx \begin{cases} A_{\omega l}^P \mathcal{T}_{\omega l}^{P,in} e^{-i\omega r^*}, & r \gtrsim 2M, \\ A_{\omega l}^P (e^{-i\omega r^*} + \mathcal{R}_{\omega l}^{P,in} e^{i\omega r^*}), & r \gg 2M, \end{cases} \quad (53)$$

where $r^* \equiv r + 2M \log\left(\frac{r}{2M} - 1\right)$ is the tortoise coordinate.

The modes $u_{\omega l}^{P,up}$ are purely incoming from the past horizon \mathcal{H}^- while the modes $u_{\omega l}^{P,in}$ are purely incoming from the past null infinity \mathcal{I}^- . Using Eq. (49), we determine the asymptotic normalization constants $A_{\omega l}^P$ to be:

$$A_{\omega l}^V = \frac{1}{\sqrt{8\pi\omega(l-1)(l+2)}} \quad (54)$$

and

$$A_{\omega l}^S = \frac{1}{\sqrt{2\pi\omega(l-1)l(l+1)(l+2)}}. \quad (55)$$

We present the calculation of the inner product for the scalar-type modes, which is necessary for finding Eq. (55), in Appendix A.

B. Gravitational radiation emission by a point particle

The point particle will contribute to the action with the interaction term given by

$$\hat{S}_I = \frac{\sqrt{32\pi}}{2} \int d^4x \sqrt{-g} T^{\mu\nu}(x) \hat{h}_{\mu\nu}(x), \quad (56)$$

where $T^{\mu\nu}$ is its energy-momentum tensor. Without loss of generality (due to the spherical symmetry of the problem), we consider the particle orbiting the black hole in the $\theta = \pi/2$ plane, at $r = R$, with angular velocity Ω , as measured by a static asymptotic observer. Its 4-velocity is written as

$$u^\mu = (\gamma, 0, 0, \gamma\Omega), \quad (57)$$

where

$$\gamma = \frac{dt}{d\tau} = \frac{1}{[f(R) - R^2\Omega^2]^{\frac{1}{2}}}. \quad (58)$$

One can write the energy-momentum tensor as

$$T^{\mu\nu} = \mu \frac{u^\mu u^\nu}{\gamma\sqrt{-g}} \delta(r - R) \delta(\theta - \pi/2) \delta(\varphi - \Omega t), \quad (59)$$

where μ is the particle's mass.

We expand the graviton field $\hat{h}_{\mu\nu}(x)$ as:

$$\hat{h}_{\mu\nu}(x) = \sum_{P,\lambda} \sum_{l=2}^{\infty} \sum_{m=-l}^l \int_0^{\infty} d\omega \left[\hat{a}_{lm}^{P,\lambda}(\omega) h_{\mu\nu}^{(P,\lambda;\omega lm)}(x) + \hat{a}_{lm}^{P,\lambda\dagger}(\omega) \overline{h_{\mu\nu}^{(P,\lambda;\omega lm)}}(x) \right]. \quad (60)$$

To first order in perturbation theory, the emission amplitude of a $\lambda = in, up$ graviton of the P -type with quantum numbers l, m and frequency ω is

$$\mathcal{A}_{\text{em}}^{P,\lambda;\omega lm} = \langle P, \lambda; \omega lm | i\hat{S}_I | 0 \rangle, \quad (61)$$

which can be found to be

$$\mathcal{A}_{\text{em}}^{P,\lambda;\omega lm} = \frac{i\sqrt{32\pi}}{2} \int d^4x \sqrt{-g} T^{\mu\nu} \overline{h_{\mu\nu}^{(P,\lambda;\omega lm)}}. \quad (62)$$

Here, the initial state $|0\rangle$ is the one annihilated by all the $\hat{a}_{lm}^{P,\lambda}(\omega)$, i.e. the Boulware vacuum. If we had chosen the Unruh [39] or Hartle-Hawking vacuum states [40] (see also Refs. [41, 42]), then the transition rate calculated from the amplitude given by Eq. (61) would be associated with the *net* radiation emitted by the particle, since the absorption and stimulated emission rates (these two rates are induced by the thermal fluxes) give the same result. The emission amplitude $\mathcal{A}_{\text{em}}^{P,\lambda;\omega lm}$ is proportional to $\delta(\omega - m\Omega)$, and hence the particle will only emit gravitons with the condition $\omega = m\Omega$ satisfied. In particular, since ω and Ω are both positive, only modes with $m \geq 1$ will be emitted.

The emitted power for a graviton with a given type of labels P, λ and quantum numbers l, m reads

$$W_{\text{em}}^{P,\lambda;lm} = \int_0^{\infty} d\omega \omega \frac{|\mathcal{A}_{\text{em}}^{P,\lambda;\omega lm}|^2}{T}, \quad (63)$$

where

$$T = 2\pi\delta(0) = \int_{-\infty}^{\infty} dt \quad (64)$$

is the total time measured by an asymptotic static observer [43]. Thus, the vector-type contributions to the emitted power are given by

$$W_{\text{em}}^{V,\lambda;lm} = 64\pi^2 \mu^2 \gamma^2 f(R)^2 m \Omega^3 \left| Y_{\varphi}^{(lm)}\left(\frac{\pi}{2}, \Omega t\right) \right|^2 \times \left| \frac{d}{dR} \left(R u_{\omega m l}^{V,\lambda}(R) \right) \right|^2, \quad (65)$$

with

$$\omega_m = m\Omega. \quad (66)$$

We note that the vector-type modes only contribute for odd values of $(l + m)$, due to the presence of the

$\left|Y_{\varphi}^{(lm)}\left(\frac{\pi}{2}, \Omega t\right)\right|^2$ factor, which vanishes for even values of $(l+m)$. The scalar-type contributions can be written as

$$W_{\text{em}}^{S,\lambda;lm} = 16\pi^2 \mu^2 \gamma^2 m \Omega \left|Y^{(lm)}\left(\frac{\pi}{2}, \Omega t\right)\right|^2 \times \left|F_{tt}^{(\lambda;\omega_m l)}(R) + 2R^2 \Omega^2 F^{\lambda;\omega_m l}(R)\right|^2, \quad (67)$$

where the functions $F_{tt}^{(\lambda;\omega_m l)}(R)$ and $F^{\lambda;\omega_m l}(R)$ are:

$$F^{\lambda;\omega_m l}(R) = \frac{1}{4HR^2} \left\{ [H - Rf'(R)] [RHu_{\omega_m l}^{S,\lambda}(R)] + 2Rf(R) \frac{d}{dR} [RHu_{\omega_m l}^{S,\lambda}(R)] \right\} \quad (68)$$

and

$$F_{tt}^{(\lambda;\omega_m l)}(R) = \frac{1}{2H} \left\{ f(R)^2 \frac{d^2}{dR^2} [RHu_{\omega_m l}^{S,\lambda}(R)] - \omega_m^2 RHu_{\omega_m l}^{S,\lambda}(R) \right\}. \quad (69)$$

We note that only the scalar-type modes with even values of $(l+m)$ contribute to the emitted power due to the $\left|Y^{(lm)}\left(\frac{\pi}{2}, \Omega t\right)\right|^2$ factor.

Next, we compute numerically the solutions to Eqs. (15) and (38) in the frequency domain. We integrate numerically these differential equations by requiring the boundary conditions given by Eqs. (52) and (53) to be satisfied, choosing suitable values of r . For the value close to the horizon, we have chosen $r/M \geq 2 + \epsilon$, with $\epsilon = 10^{-3}$. As for the ‘‘numerical infinity’’, r_{∞} , we write it as a function of l and ω_m such that the following condition is satisfied

$$\omega_m^2 \gg \frac{l(l+1)}{r_{\infty}^2}. \quad (70)$$

In our computation we have chosen our ‘‘numerical infinity’’ to be:

$$r_{\infty} = \frac{250\sqrt{l(l+1)}}{\omega_m}. \quad (71)$$

With the numerically obtained solutions, we use Eqs. (65) and (67) to compute the total emitted power as

$$W_{\text{em}} = \sum_{\lambda} \sum_{l=2}^{\infty} \sum_{m=1}^l (W_{\text{em}}^{S,\lambda;lm} + W_{\text{em}}^{V,\lambda;lm}). \quad (72)$$

We compute the emitted power as a function of the angular velocity Ω , relating to it the radial coordinate R of the test particle by

$$R = \left(\frac{M}{\Omega^2}\right)^{1/3}, \quad (73)$$

which is required for the particle to be in circular orbit around the black hole, according to GR [44]. We include in our results both stable (up to $\Omega = (6\sqrt{6}M)^{-1} \approx$

$0.068 M^{-1}$) and unstable circular orbits (up to $\Omega = (3\sqrt{3}M)^{-1} \approx 0.192 M^{-1}$). As the circular orbit approaches the orbit of the light ray at $R = 3M$, the emitted power increases rapidly, mainly because its motion becomes ultrarelativistic (the particle’s energy increases with u^t). For this reason we find it more appropriate to plot the emitted power on a logarithmic scale. The particle becomes ultrarelativistic only for unstable orbits and we note that, if backreaction is taken into account, a particle in an unstable orbit starts its plunge into the black hole.




Figure 1. Total power emitted by the test particle rotating around the black hole, given by Eq. (72), plotted as a function of the angular velocity Ω . The summation in l in Eq. (72) is truncated at a certain value of l , which we denote as l_{max} .




Figure 2. Scalar-type power emitted by the test particle, given by Eq. (67), as a function of the angular velocity Ω . We show here the modes with $l = m$, which give the main contributions to the total emitted power.

The results for the emitted power are plotted in Figs. 1, 2 and 3. For stable orbits, the main contribution to the total emitted power are the modes with $l = 2$ ($m = 2$ and $m = 1$ for the scalar- and vector-type radi-




Figure 3. Comparison between the $l = 2$ scalar-type power, given by Eq. (67), and the $l = 2$ vector-type power emitted by the test particle, given by Eq. (65), as a function of the angular velocity Ω .

ation, respectively). We note that the vector-type emitted power, given by Eq. (65), is suppressed by a factor of $R^2\Omega^2 = M/R$, compared to the scalar-type emitted power [20], given by Eq. (67) (see Fig. 3).

We note that the $l = 2$ modes have a dominant contribution to the emitted power for most of the Ω range, as one can see in Fig. 4. The contribution from all the other l modes start to dominate over the $l = 2$ mode contribution at $\Omega \approx 0.164 M^{-1}$. For unstable orbits, the high- l contributions become more important, but the $l = 2$ modes still have a significant contribution (see Fig. 5). For stable orbits, we can see that the contribution of the high multipoles is small. For unstable orbits, high multipoles are enhanced, but the low multipoles (especially the $l = 2$ modes) are still important.




Figure 4. Comparison between the contributions of the $l = 2$ and $l > 2$ modes (we considered contributions up to $l = 20$). For most of the range of Ω (up to $\Omega \approx 0.164 M^{-1}$), the $l = 2$ mode is the dominant contribution.




Figure 5. Ratio $W_{\text{em}}^{l=2}/W_{\text{em}}$ between the emitted power of the $l = 2$ modes and the total emitted power (we have considered contributions to the total emitted power up to $l = 20$). The $l = 2$ modes remain relevant contributions to the emitted power even for unstable orbits.

One can compute the power of emitted gravitons observed at infinity by considering only the modes which are purely outgoing at future null infinity. Since these modes are related to the modes $u_{\omega l}^{P,in}$ by complex conjugation, we can write the power observed at infinity as²

$$W_{\text{em}}^{\text{obs}} = \sum_{P,l,m} W_{\text{em}}^{P,in;lm}. \quad (74)$$

In Figs. 6 and 7, the ratio $W_{\text{em}}^{\text{obs}}/W_{\text{em}}$ is plotted as a function of the angular velocity Ω . For unstable orbits, a considerable amount of emitted power is absorbed by the black hole, as shown in Fig. 7. Approximately 38% of the radiation fails to reach the asymptotically flat region (infinity), for the innermost unstable orbit. In contrast, for stable orbits, more than 99% of the emitted power escapes to infinity, as one can see in Fig. 6. We compared our results for the asymptotic radiation (scalar- and vector-type contributions) in stable orbits with other works [12, 21], resulting in excellent agreement.

Next, we analyze the radiation associated to the modes with large m , and hence with large l . Since only modes with $\omega_m = m\Omega$ are emitted, the total power for a given frequency, $P(\omega_m)$, can be written as a function of m , and hence as a function of ω_m , as [12]

$$P(\omega_m) = \sum_{\lambda,P} \sum_{l \geq |m|}^{\infty} W_{\text{em}}^{P,\lambda;lm}. \quad (75)$$

² The corresponding formulae for the massless scalar radiation in Ref. [24] [Eqs. (35) and (36) of that reference] are incorrect [45]. Similar incorrect formulae were used in Refs. [25–29]. Correction of these formulae does not affect the main conclusions in those references.




Figure 6. Ratio $W_{\text{em}}^{\text{obs}}/W_{\text{em}}$, between the asymptotically observed and the total emitted power, as a function of Ω , plotted for stable orbits. We have considered contributions up to $l = 20$. We see that almost all the energy is radiated away to infinity, in the case of stable orbits.




Figure 7. Ratio $W_{\text{em}}^{\text{obs}}/W_{\text{em}}$, between the asymptotically observed and the total emitted power, as a function of Ω , plotted for stable, as well as for unstable circular orbits. As in Fig. 6, we have considered contributions up to $l = 20$.

The power $P(\omega_m)$ depends on a discrete variable ω_m , but we can regard it as a continuous variable for $m \gg 1$. In this continuum limit, we can write the emitted power W_{em} in terms of the spectral density function, denoted by $P(\omega)/\Omega$, as [46]

$$W_{\text{em}} = \sum_{m=1}^{\infty} \frac{P(\omega_m)}{\Omega} \Delta\omega_m \approx \int_0^{\infty} d\omega \frac{P(\omega)}{\Omega}, \quad (76)$$

with $\Delta\omega_m = \omega_m - \omega_{m-1} = \Omega$. We compute $P(\omega_m)$ and its asymptotically observed counterpart, obtained by summing only $W_{\text{em}}^{P, in; lm}$ in Eq. (75), for a particle orbiting the black hole in a highly relativistic unstable orbit with $R = (3 + \delta)M$, $\delta = 5 \times 10^{-4}$, for frequencies up to $\omega = 2500\Omega$. We neglect the scalar-type modes with $l > m$ and




Figure 8. Total power emitted for a given frequency ω_m . Frequencies up to $m = 30$ are shown in this plot. Since only gravitons with frequencies that are integer multiples of the particle's angular velocity are emitted ($\omega_m = m\Omega$), the spectrum is discrete.




Figure 9. Emitted power for a given frequency ω_m in the high-frequency range ($100 \leq m \leq 2500$).

vector-type modes with $l > m + 1$ since the $l = m$ scalar-type and $l = m + 1$ vector-type modes contribute to more than 99% of the power at a given l . The results are shown in Figs. 8 and 9. These results are in excellent agreement with those for high multipoles in Refs. [10, 12] and show that the contribution of the low frequency modes is still relevant for the total radiation, even for unstable orbits.

IV. COMPARISON WITH FLAT SPACETIME RESULTS

Let us now compare the emitted power W_{em} in Schwarzschild spacetime with its analogues in Minkowski spacetime, W_{em}^M . For the flat spacetime computation, we consider the particle to be in a circular orbit bound to a

stellar object, due to a Newtonian force. When the particle is not very close to the central object, the two cases should give similar results.

In Minkowski spacetime, we use essentially the same procedure as in the Schwarzschild case to obtain the emitted power, with the difference that we set $f(r) = 1$. The perturbed metric $h_{\mu\nu}$ will have the same form as in the Schwarzschild case, but now both master fields in Minkowski spacetime satisfy the same equation, namely

$$\square\Phi_l^{M;P}(t, r) - V_M\Phi_l^{M;P}(t, r) = 0, \quad (77)$$

with $V_M = l(l+1)/r^2$ and $P = S, V$. The d'Alembertian in Eq. (77) is the one compatible with the (flat) orbital spacetime line element $ds_{orb}^2 = -dt^2 + dr^2$. One can write positive-frequency solutions to Eq. (77), which are regular at the origin, as

$$\Phi_{\omega l}^{M;P}(t, r) = C_{\omega l}^P e^{-i\omega t} r j_l(\omega r), \quad (78)$$

where $j_l(\omega r)$ are the spherical Bessel functions of the first kind [47]. Using the inner product given by Eq. (45), we obtain the normalization constants $C_{\omega l}^P$:

$$C_{\omega l}^V = \sqrt{\frac{\omega}{2\pi(l-1)(l+2)}} \quad (79)$$

and

$$C_{\omega l}^S = \sqrt{\frac{2\omega}{\pi(l-1)l(l+1)(l+2)}}. \quad (80)$$

To compute the emitted power in Minkowski spacetime, we simply substitute the master fields of Eq. (78) into Eqs. (65) and (67) to obtain³

$$W_{em}^{M;S;lm} = 16\pi^2 \mu^2 \gamma_M^2 m \Omega \left| Y^{(lm)}\left(\frac{\pi}{2}, \Omega t\right) \right|^2 \times \left| F_{tt}^{(M;\omega_m l)}(R_M) + 2R_M^2 \Omega^2 F^{M;\omega_m l}(R_M) \right|^2 \quad (81)$$

and

$$W_{em}^{M;V;lm} = 64\pi^2 \mu^2 \gamma_M^2 m \Omega^3 \left| Y_{\varphi}^{(lm)}\left(\frac{\pi}{2}, \Omega t\right) \right|^2 \times \left| C_{\omega_m l}^V \frac{d}{dR_M} (R_M^2 j_l(\omega_m R_M)) \right|^2, \quad (82)$$

where $\gamma_M = (1 - R_M^2 \Omega^2)^{-1/2}$ is the Lorentz factor. The quantities $F^{M;\omega_m l}(R_M)$ and $F_{tt}^{(M;\omega_m l)}(R_M)$ are obtained by substituting the master field $\Phi_{\omega_m l}^{M;S}$ into Eqs. (22) and

(23), respectively, with the mass of the black hole set to zero, namely

$$F^{M;\omega_m l}(R_M) = \frac{C_{\omega_m l}^S}{4} \left\{ \omega_m^2 R_M^2 j_l(\omega_m R_M) + \frac{d^2}{dR_M^2} [R_M^2 j_l(\omega_m R_M)] \right\} \quad (83)$$

and

$$F_{tt}^{(M;\omega_m l)}(R_M) = \frac{C_{\omega_m l}^S}{2} \left\{ \frac{d^2}{dR_M^2} [R_M^2 j_l(\omega_m R_M)] - \omega_m^2 R_M^2 j_l(\omega_m R_M) \right\}. \quad (84)$$

We have used the equation of motion given by Eq. (77) to simplify Eq. (83). In Newtonian gravity, for the particle to be in a circular orbit around the stellar object, its radial coordinate is related to its angular velocity by Kepler's third law, namely $R_M(\Omega) = (M\Omega^{-2})^{1/3}$. Since the angular velocity Ω is a quantity measured by an asymptotic static observer and, hence, coordinate independent, it is meaningful to compare the emitted powers from the orbiting particle in Schwarzschild and Minkowski spacetimes with the same value of Ω . This comparison is plotted in Fig. 10.




Figure 10. Ratio W_{em}/W_{em}^M as a function of Ω . We have considered contributions up to $l = 20$. The maximum value considered for ΩM is $(6\sqrt{6})^{-1}$. As Ω increases, the ratio decreases up to approximately 25%, until it starts to increase due to the ultrarelativistic effect.

We note that there is a significant conceptual difference between the radiations in the two cases: in Schwarzschild spacetime, the circular orbit is a geodesics of the spacetime, whereas in Minkowski spacetime, it is a trajectory supported by an external force. In other words, the circular orbit is not a geodesic of Minkowski spacetime. Thus, the total radiation computation in Minkowski spacetime should include the radiation generated by the source of the external force, since the particle's energy-momentum tensor is not conserved by itself. However, it is likely that

³ We adopt the usual Minkowski vacuum, i.e. the vacuum that is annihilated by the annihilation operators corresponding to the positive-frequency mode functions given by Eq. (78).

this additional radiation is negligible. (This is indeed the case if the circular motion is supported by a thin rod connecting the particle and the origin [48].) The comparison in this section has been done primarily to show consistency of our numerical results in Schwarzschild spacetime by comparing them to the solutions obtained analytically in Minkowski spacetime, given by Eq. (78), which should be a good approximation to the master fields far away from the black hole.

V. CONCLUDING REMARKS

We have computed the power of gravitational radiation emitted by a particle in circular orbit around a Schwarzschild black hole. By writing the gravitational perturbations in a gauge-invariant formalism, we used QFT at tree level to obtain numerically the emitted power, for both stable and unstable orbits. The scalar-type gravitational radiation was shown to be dominating over the vector-type gravitational radiation, the latter being suppressed by a factor of $R^2\Omega^2 = M/R$. This result is in agreement with Ref. [20], where the emitted power was computed for stable orbits only. For most of the range of the test particle's angular velocity, we have shown that the main contributions to the emitted power are from the $l = 2$ modes for both scalar- and vector-type gravitational perturbations. In unstable orbits, the contributions from high multipole modes are enhanced. Nevertheless, the $l = 2$ modes still have contributions that are far from negligible.

In stable orbits, almost all of the emitted energy escapes to infinity. For unstable orbits however, a considerable amount of energy is absorbed by the black hole.

Comparing the emitted powers in Schwarzschild and Minkowski spacetimes, we see that the ratio between them (Schwarzschild over Minkowski) approaches unity when the particle is located very far away from the black hole. This ratio decreases as the orbit moves inwards but starts to increase near the innermost stable circular or-

bit ($R = 6M$) because the orbit becomes ultrarelativistic sooner (i.e., for smaller values of Ω) in the Schwarzschild spacetime, than in flat spacetime.

We found that high multipoles of the gravitational radiation are significantly enhanced only for unstable orbits, although the low multipole modes remain as a relevant contribution to the emitted power even in this case. We also have found that no high frequency peaks are present in the power spectrum, unlike the scalar-radiation case [25]. We note that the particle in a stable orbit will emit radiation, gradually losing energy and inspiraling to orbits with smaller radii. When it reaches an unstable orbit, the particle emits a high amount of gravitational radiation and quickly plunges down into the black hole.

ACKNOWLEDGMENTS

We would like to acknowledge Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for partial financial support. A. H. also acknowledges partial support from the Abdus Salam International Centre for Theoretical Physics through the Visiting Scholar/Consultant Programme. A. H. thanks the Universidade Federal do Pará (UFPA) in Belém and R. B. is grateful to the University of York for the kind hospitality while part of the work was carried out.

Appendix A: Computation of the scalar-type inner product

In this Appendix, we compute the inner product for scalar-type perturbations at the future horizon, using Eddington-Finkelstein coordinates. For the scalar-type perturbation, the inner product (45) reads

$$\langle h^S, h'^S \rangle = -i \int_{\Sigma} d\Sigma n_a J^a, \quad (\text{A1})$$

where

$$J^a = \overline{Y^{lm} Y'^{l'm'}} \left[\frac{4}{r} D^d r \left(\overline{F^{(\omega l)ab}} F_{bd}^{(\omega'l')} - F^{(\omega'l')ab} \overline{F_{bd}^{(\omega l)}} \right) - \left(\overline{F^{(\omega l)bc}} D^a F_{bc}^{(\omega'l')} - F^{(\omega'l')bc} D^a \overline{F_{bc}^{(\omega l)}} \right) \right], \quad (\text{A2})$$

and n_a is the (future-pointing) unit vector normal to the Cauchy hypersurface Σ . The inner product given by Eq. (A1) can be rewritten as

$$\langle h^S, h'^S \rangle = \frac{i\delta^{ll'}\delta^{mm'}}{2} \int_{2M}^{\infty} \frac{dr}{f(r)} \times \left(\overline{\Phi_{\omega l}^S} \partial_t \Phi_{\omega'l'}^S - \Phi_{\omega'l'}^S \partial_t \overline{\Phi_{\omega l}^S} \right). \quad (\text{A3})$$

We will derive Eq. (A3) using Eddington-Finkelstein coordinates. To simplify the notation, from now on we omit the labels for the frequency and angular quantum numbers, denoting quantities that depend on ω' , l' and m' with a prime.

Defining a new coordinate by

$$u \equiv t - r^*, \quad (\text{A4})$$

where r^* is the tortoise coordinate, we find

$$du = dt - \frac{dr}{f(r)} \quad (\text{A5})$$

and the orbit spacetime line element (3) becomes

$$ds_{orb}^2 = -f(r)du^2 - 2dudr. \quad (\text{A6})$$

We compute the inner product at the future horizon. The horizon is at $r = 2M$ and $-\infty < u < \infty$. If $r < 2M$, the $r = \text{constant}$ surface is a spacelike surface. A normalized (future-pointing) vector orthogonal to this surface can be written as

$$\begin{aligned} n^a &= -[-f(r)]^{-1/2} D^a r \\ &= [-f(r)]^{-1/2} \left(\frac{\partial}{\partial u} \right)^a + [-f(r)]^{1/2} \left(\frac{\partial}{\partial r} \right)^a. \end{aligned} \quad (\text{A7})$$

For this surface we have

$$d\Sigma = d\Omega_2 du [-f(r)]^{1/2} r^2. \quad (\text{A8})$$

Hence, using Eq. (A7), we obtain

$$d\Sigma n^a = r^2 d\Omega_2 du \left[\left(\frac{\partial}{\partial u} \right)^a - f(r) \left(\frac{\partial}{\partial r} \right)^a \right]. \quad (\text{A9})$$

In the limit $r \rightarrow 2M$, we get

$$\lim_{r \rightarrow 2M} d\Sigma n^a = 4M^2 d\Omega_2 du \left(\frac{\partial}{\partial u} \right)^a. \quad (\text{A10})$$

From one of the equations of motion ($g^{ab}F_{ab} = 0$), we have

$$F_{ur} = \frac{f(r)}{2} F_{rr}, \quad (\text{A11})$$

which means that F_{ur} vanishes at the horizon. Moreover, the first term in Eq. (A2) does not contribute to the inner product because

$$n^a D^d r \left(\overline{F_a^b F'_{bd}} - F_a^b \overline{F_{bd}} \right) = 0. \quad (\text{A12})$$

This follows from the fact that F_{ab} is a symmetric tensor and that $n^a \propto D^a r$ [see Eq. (A7)].

For the second term in Eq. (A2), we have, at the horizon,

$$\begin{aligned} \overline{F^{bc} D_u F'_{bc}} - F'^{bc} D_u \overline{F_{bc}} &= \overline{F_{rr} \partial_u F'_{uu}} + \overline{F_{uu} \partial_u F'_{rr}} \\ &\quad - F'_{rr} \partial_u \overline{F_{uu}} - F'_{uu} \partial_u \overline{F_{rr}} \\ &\quad + \frac{1}{M} \left(\overline{F_{rr} F'_{uu}} - F'_{rr} \overline{F_{uu}} \right). \end{aligned} \quad (\text{A13})$$

We integrate Eq. (A13) by parts with respect to u , indicating with the symbol “ \approx ” the equivalence under integration by parts. We find

$$\begin{aligned} \overline{F^{bc} D_u F'_{bc}} - F'^{bc} D_u \overline{F_{bc}} &\approx 2 \left(\overline{F_{rr} \partial_u F'_{uu}} + \overline{F_{uu} \partial_u F'_{rr}} \right) \\ &\quad + \frac{1}{M} \left(\overline{F_{rr} F'_{uu}} - F'_{rr} \overline{F_{uu}} \right). \end{aligned} \quad (\text{A14})$$

At the horizon, one can write the components of the gauge-invariant quantity $F_{ab}^{(\omega l)}$ as

$$F_{uu} = \left(2M \partial_u^2 + \frac{1}{2} \partial_u \right) \Phi^S \quad (\text{A15})$$

and

$$F_{rr} = \left[2M \partial_r^2 + \left(2 - \frac{6}{H} \right) \partial_r \right] \Phi^S. \quad (\text{A16})$$

Here, $H = H(r = 2M)$ [see Eq. (24)]. Thus,

$$H = l^2 + l + 1. \quad (\text{A17})$$

We note that we set $r = 2M$ after all differentiation is done. Now, we substitute Eqs. (A15) and (A16) into Eq. (A14), and integrate by parts with respect to u to obtain

$$\overline{F^{bc} D_u F'_{bc}} - F'^{bc} D_u \overline{F_{bc}} \approx \partial_u \Phi'^S \hat{O} \overline{\Phi^S} - \partial_u \overline{\Phi^S} \hat{O} \Phi'^S, \quad (\text{A18})$$

where the fourth order differential operator \hat{O} reads

$$\begin{aligned} \hat{O} &= 8M^2 \partial_u^2 \partial_r^2 - 6M \partial_u \partial_r^2 + 8M \left(1 - \frac{3}{H} \right) \partial_u^2 \partial_r \\ &\quad - 6 \left(1 - \frac{3}{H} \right) \partial_u \partial_r + \partial_r^2 + \frac{1}{M} \left(1 - \frac{3}{H} \right) \partial_r. \end{aligned} \quad (\text{A19})$$

We write \hat{O} as the following linear combination

$$\hat{O} = A \square \left[\frac{f(r)}{V_S(r)} \square \right] + (B + C \partial_u) \square, \quad (\text{A20})$$

with A , B and C being suitably chosen constants. We can use the equations of motion to write

$$\hat{O} \Phi_{\omega l}^S = \underbrace{\frac{V_S(r)}{f(r)}}_{\text{constant at } r=2M} [(A + B) \Phi_{\omega l}^S + C \partial_u \Phi_{\omega l}^S]. \quad (\text{A21})$$

We note that the term containing C does not contribute to the inner product. However, its presence is needed to write the fourth order operator (A19) in the form (A20). We may use a symbolic computation software to obtain

$$A = \frac{[3 + (l-1)l(l+1)(l+2)]^2}{2[3 + l(l+1)(l^4 + 2l^3 - l + 1)]}, \quad (\text{A22})$$

$$B = 2 - \frac{3}{l^2 + l + 1} - \frac{3}{2} \frac{l^2 + l + 1}{(l-1)l(l+1)(l+2) + 3}, \quad (\text{A23})$$

and

$$C = 6M \frac{l^2 + l + 1}{(l-1)l(l+1)(l+2) + 3}. \quad (\text{A24})$$

Then, at the horizon, the inner product (A1) can be written as

$$\begin{aligned}
\langle h^S, h'^S \rangle &= i \frac{(l-1)l(l+1)(l+2)}{2} \int d\Omega_2 du \overline{Y^{lm}} Y^{l'm'} \left(\overline{\Phi_{\omega l}^S} \partial_u \Phi_{\omega' l'}^S - \Phi_{\omega' l'}^S \partial_u \overline{\Phi_{\omega l}^S} \right) \\
&= \lim_{r \rightarrow 2M} i \frac{(l-1)l(l+1)(l+2)}{2} \int_{\Sigma} d\Sigma n^a \overline{Y^{lm}} Y^{l'm'} \left(\overline{\Phi_{\omega l}^S} \partial_a \Phi_{\omega' l'}^S - \Phi_{\omega' l'}^S \partial_a \overline{\Phi_{\omega l}^S} \right). \tag{A25}
\end{aligned}$$

This can be evaluated in a $t = \text{constant}$ Cauchy surface in tr coordinates as

$$\langle h^{(S;\omega lm)}, h^{(S;\omega' l' m')} \rangle = i \frac{(l-1)l(l+1)(l+2)}{2} \int_{S^2} d\Omega_2 \overline{Y^{lm}} Y^{l'm'} \int_{2M}^{\infty} \frac{dr}{f(r)} \left(\overline{\Phi_{\omega l}^S} \partial_t \Phi_{\omega' l'}^S - \Phi_{\omega' l'}^S \partial_t \overline{\Phi_{\omega l}^S} \right), \tag{A26}$$

which leads to Eq. (A3).

Another way to obtain Eq. (A3) is by computing the inner product (A1) directly using tr coordinates. However, this method is much more involved, requiring several cumbersome integration by parts, although we can still use a computational software to do all the algebraic

computations. We have done so and the same result as the Eddington-Finkelstein one has been obtained, as expected. In addition, since computing the inner product in these coordinates does not require the presence of a horizon, we can also use this method in flat spacetime, in spherical coordinates.

-
- [1] D. Richstone, E. A. Ajhar, R. Bender, G. Bower, A. Dressler, S. M. Faber, A. V. Filippenko, K. Gebhardt, R. Green, L. C. Ho, J. Kormendy, T. R. Lauer, J. Magorrian, and S. Tremaine, Supermassive black holes and the evolution of galaxies. *Nature (London)* **395**, A14 (1998), astro-ph/9810378.
 - [2] J. Kormendy and L. C. Ho, Coevolution (or not) of supermassive black holes and host galaxies, *Annual Review of Astronomy and Astrophysics* **51**, 511 (2013), <http://dx.doi.org/10.1146/annurev-astro-082708-101811>.
 - [3] W. Lewin, J. van Paradijs, and E. van den Heuvel, *X-ray Binaries*, Cambridge Astrophysics (Cambridge University Press, Cambridge, 1997).
 - [4] A. Celotti, J. C. Miller, and D. W. Sciama, Astrophysical evidence for the existence of black holes, *Classical and Quantum Gravity* **16**, A3 (1999).
 - [5] A. I. MacFadyen and S. E. Woosley, Collapsars: Gamma-ray bursts and explosions in “failed supernovae”, *The Astrophysical Journal* **524**, 262 (1999).
 - [6] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
 - [7] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), GW151226: Observation of gravitational waves from a 22-solar-mass binary black hole coalescence, *Phys. Rev. Lett.* **116**, 241103 (2016).
 - [8] C. W. Misner, Interpretation of gravitational-wave observations, *Phys. Rev. Lett.* **28**, 994 (1972).
 - [9] C. W. Misner, R. A. Breuer, D. R. Brill, P. L. Chrzanowski, H. G. Hughes, and C. M. Pereira, Gravitational synchrotron radiation in the Schwarzschild geometry, *Phys. Rev. Lett.* **28**, 998 (1972).
 - [10] D. M. Chitre and R. H. Price, Nature of gravitational synchrotron radiation, *Phys. Rev. Lett.* **29**, 185 (1972).
 - [11] M. Davis, R. Ruffini, J. Tiomno, and F. Zerilli, Can synchrotron gravitational radiation exist? *Phys. Rev. Lett.* **28**, 1352 (1972).
 - [12] R. A. Breuer, R. Ruffini, J. Tiomno, and C. V. Vishveshwara, Vector and tensor radiation from Schwarzschild relativistic circular geodesics, *Phys. Rev. D* **7**, 1002 (1973).
 - [13] T. Regge and J. A. Wheeler, Stability of a Schwarzschild singularity, *Phys. Rev.* **108**, 1063 (1957).
 - [14] L. A. Edelman and C. V. Vishveshwara, Differential equations for perturbations on the Schwarzschild metric, *Phys. Rev. D* **1**, 3514 (1970).
 - [15] F. J. Zerilli, Effective potential for even-parity Regge-Wheeler gravitational perturbation equations, *Phys. Rev. Lett.* **24**, 737 (1970).
 - [16] F. J. Zerilli, Gravitational field of a particle falling in a Schwarzschild geometry analyzed in tensor harmonics, *Phys. Rev. D* **2**, 2141 (1970).
 - [17] M. Davis and R. Ruffini, Gravitational radiation in the presence of a Schwarzschild black hole. A boundary value search, *Lett. Nuovo Cim.* **2**, 1165 (1971).
 - [18] M. Davis, R. Ruffini, W. H. Press, and R. H. Price, Gravitational radiation from a particle falling radially into a Schwarzschild black hole, *Phys. Rev. Lett.* **27**, 1466 (1971).
 - [19] M. Davis, R. Ruffini, and J. Tiomno, Pulses of gravitational radiation of a particle falling radially into a Schwarzschild black hole, *Phys. Rev.* **D5**, 2932 (1972).
 - [20] E. Poisson, Gravitational radiation from a particle in circular orbit around a black hole. I. Analytical results for the nonrotating case, *Phys. Rev. D* **47**, 1497 (1993).
 - [21] C. Cutler, L. S. Finn, E. Poisson, and G. J. Sussman, Gravitational radiation from a particle in circular orbit around a black hole. II. Numerical results for the nonrotating case, *Phys. Rev. D* **47**, 1511 (1993).
 - [22] N. D. Birrell and P. C. W. Davies, *Quantum fields in Curved Space* (Cambridge University Press, Cambridge, 2013).

- [23] L. Parker and D. Toms, *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity* (Cambridge University Press, Cambridge, 2009).
- [24] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, Scalar radiation emitted from a source rotating around a black hole, *Classical and Quantum Gravity* **17**, 19 (2000).
- [25] L. C. B. Crispino, Synchrotron scalar radiation from a source in ultrarelativistic circular orbits around a Schwarzschild black hole, *Phys. Rev. D* **77**, 047503 (2008).
- [26] L. C. B. Crispino, A. R. R. da Silva, and G. E. A. Matsas, Scalar radiation emitted from a rotating source around a Reissner-Nordström black hole, *Phys. Rev. D* **79**, 024004 (2009).
- [27] C. F. B. Macedo, L. C. B. Crispino, and V. Cardoso, Semiclassical analysis of the scalar geodesic synchrotron radiation in Kerr spacetime, *Phys. Rev. D* **86**, 024002 (2012).
- [28] J. Castiñeiras, L. C. B. Crispino, and D. P. M. Filho, Source coupled to the massive scalar field orbiting a stellar object, *Phys. Rev. D* **75**, 024012 (2007).
- [29] J. Castiñeiras, L. C. B. Crispino, R. Murta, and G. E. A. Matsas, Semiclassical approach to black hole absorption of electromagnetic radiation emitted by a rotating charge, *Phys. Rev. D* **71**, 104013 (2005).
- [30] H. Kodama, A. Ishibashi, and O. Seto, Brane world cosmology: Gauge-invariant formalism for perturbation, *Phys. Rev. D* **62**, 064022 (2000).
- [31] H. Kodama and A. Ishibashi, A master equation for gravitational perturbations of maximally symmetric black holes in higher dimensions, *Prog. Theor. Phys.* **110**, 701 (2003), arXiv:hep-th/0305147 [hep-th].
- [32] C. Cohen-Tannoudji, B. Diu, and F. Laloe, *Quantum Mechanics (2 vol. set)* (Wiley-Interscience, New York, 2006).
- [33] R. P. Bernar, L. C. B. Crispino, and A. Higuchi, Infrared-finite graviton two-point function in static de Sitter space, *Phys. Rev. D* **90**, 024045 (2014).
- [34] R. P. Bernar, L. C. B. Crispino, and A. Higuchi, Graviton two-point function in $3 + 1$ static de Sitter spacetime, *Int. J. Mod. Phys. D* **25**, 1641016 (2016).
- [35] A. Higuchi, Symmetric tensor spherical harmonics on the N -sphere and their application to the de Sitter group $SO(N,1)$, *J.Math.Phys.* **28**, 1553 (1987), **43**, 6385(E) (2002).
- [36] M. A. Rubin and C. R. Ordóñez, Eigenvalues and degeneracies for n -dimensional tensor spherical harmonics, *Journal of Mathematical Physics* **25**, 2888 (1984).
- [37] C. J. Fewster and D. S. Hunt, Quantization of linearized gravity in cosmological vacuum spacetimes, *Rev. Math. Phys.* **25**, 1330003 (2013).
- [38] J. L. Friedman, Generic instability of rotating relativistic stars, *Commun. Math. Phys.* **62**, 247 (1978).
- [39] W. G. Unruh, Notes on black-hole evaporation, *Phys. Rev. D* **14**, 870 (1976).
- [40] J. Hartle and S. Hawking, Path integral derivation of black hole radiance, *Phys. Rev. D* **13**, 2188 (1976).
- [41] W. Israel, Thermo-field dynamics of black holes, *Physics Letters A* **57**, 107 (1976).
- [42] B. S. Kay and R. M. Wald, Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on spacetimes with a bifurcate Killing horizon, *Physics Reports* **207**, 49 (1991).
- [43] R. A. Breuer, *Gravitational perturbation theory and synchrotron radiation* (Springer-Verlag, Berlin, 1975).
- [44] R. Wald, *General Relativity* (The University of Chicago Press, Chicago, 1984).
- [45] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, Corrigendum: Scalar radiation emitted from a source rotating around a black hole (2000 *Class. Quantum Grav.* **17** [<http://dx.doi.org/10.1088/0264-9381/17/1/303>] 19), *Classical and Quantum Gravity* **33**, 209502 (2016).
- [46] R. A. Breuer, P. L. Chrzanowski, H. G. Hughes, and C. W. Misner, Geodesic synchrotron radiation, *Phys. Rev. D* **8**, 4309 (1973).
- [47] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables* (Dover, New York, 1964).
- [48] R. H. Price and V. D. Sandberg, Role of constraining forces for ultrarelativistic particle motion as a source of gravitational radiation, *Phys. Rev. D* **8**, 1640 (1973).