

## Using remote sensing and Detection of Early Season Invasives (DESI) to analyze the temporal dynamics of invasive cheatgrass (*Bromus tectorum*).

Bishop, Tara B.B.<sup>1</sup>, St. Clair, S.B.<sup>1</sup>, Belnap, J.<sup>2</sup>, Munson, S.<sup>3</sup>, Gill, R.A.<sup>1</sup>

<sup>1</sup>Brigham Young University, <sup>2</sup> Southwest Biological Science Center, USGS, <sup>3</sup> Colorado Plateau Research Station, USGS

### Introduction

The invasion of exotic annual grasses during the last century has transformed plant habitats and communities worldwide.

Cheatgrass (*Bromus tectorum*) is a winter annual grass that has invaded over 100 million acres of the western United States (Pellant and Hall, 1994. Pellant, 1996). Cheatgrass quickly utilizes available resources especially after a disturbance to the landscape. A major impact of invasion is the increased frequency in fires (D'Antonio and Vitousek, 1992). As cheatgrass is highly successful at invading open and disturbed landscapes at a rapid pace it increases the frequency and severity of fires in arid landscapes (Brooks, 2005). Cheatgrass' prolific seed production and flammability allows it to competitively exclude native plant species (Seabloom et al., 2003). The successful life strategy of cheatgrass gives a unique spectral image reflectance that can allow the use of remote sensing platforms to track and locate invasions.

Cheatgrass invasion is particularly worrisome in eastern and southern Utah as it spreads and degrades much of Utah's wildlands. Utah has 13 national parks and monuments with over 10 million visitors annually. Within those parks there are over 18 threatened and endangered species and pristine habitat for over 200 endemic plant species. With an economic benefit of over \$725,00,000 annually (National Park Service, 2014) the increasing invasion of

cheatgrass puts all national parks at risk of altering valuable visitor experiences and economic benefit in the future.

Increasing invasion, and thus potential and actual fire frequency, also has serious ecological impacts as the native plants have a decreased ability to re-establish after a fire. This leads to the degradation of the native plant community as the cheatgrass continues to replace the native perennials and/or shrubs (Zouhar, 2003). This change in the native plant community can lead to negative impacts on the surrounding wildlife habitat and changes in the surrounding physical environment.

Restoration and rehabilitation of areas that have been invaded are a top priority of land managers. But large scale surveying of the land is timely and can have high cost association. Using a geographic information system (GIS, ESRI) with Detection of Early Season Invasive (DESI) software (Kokaly, 2011) landscape level analysis can be done of invasive annual grasses. Understanding landscape controls and the temporal dynamics of large, full scale invasions may be critical to controlling, managing and even preventing loss of natural habitat to the conversion of invasive grasslands. Our primary objectives to achieve this understanding are to (1) Measure areas of permanent cheatgrass invasion and areas that are temporally dynamic; (2) Develop a statistical model that explains the controls over the spatial and temporal distribution of cheatgrass and (3) Identify areas that are

sensitive to and conditions that will promote cheatgrass expansion in the future.

**METHODS**

*Detection of Early Season Invasives*

The study sites, in collaboration with USGS, will be conducted at seven national parks and monuments: Arches National Park, Bryce Canyon National Park, Canyonlands National Park, Capitol Reef National Park, Dinosaur National Monument, Glen Canyon National Recreation Area, and Natural Bridges National Monument all located in the state of Utah.

The United States Geological Survey (USGS) developed a software algorithm that uses remote sensing data from Landsat TM/ETM imagery to detect locations and populations of cheatgrass called Detection of Early Season Invasives (DESI) (Kokaly, 2011).

Using ENVI (Exelis Visual Information Solutions) software the Normalized Difference Vegetation Index (NDVI) values for reflectance of red and near-infrared radiation by plants are extracted from the Landsat TM/ETM images. NDVI is taken for early spring and summer to capture the

values in early spring and summer (Figure 1), and including masks for cloud cover and other climatic conditions, the software can detect locations for early seasons invasives, specifically cheatgrass. The image produced is a map with 30m x 30m pixels designating locations at which cheatgrass meets high and low thresholds. The thresholds are determined by examining the value (minimum dNDVI values) of a specific pixel and then the surrounding pixels and their corresponding value.

DESI images were produced for each study site for years 1999-2009 (Figure 2). Not all parks had a complete 10-year data set as some images were not acceptable for proper analysis due to cloud cover and other environmental factors.




Figure 2: Left is a DESI output image for Landsat imagery encompassing central and southeast Utah. Right is Arches National Park DESI output (clipped from larger image). Red indicates the high threshold for cheatgrass growth and yellow indicates the low threshold for cheatgrass growth.




Figure 1: Seasonal trends of dNDVI for plots in Canyonlands National Park in 2001.

senescence of early season invasives. By taking the difference of NDVI (dNDVI)

*Extracting CORE and VARIABLE populations*

Analysis of the DESI images to identify areas of permanent core populations versus the transient temporally dynamic populations required building models in GIS software, ArcMap 10x (ESRI, 2011). To determine whether or not a core population existed in a park all of the DESI images were overlaid each other. Then using the Raster Calculator tool syntax was derived to pull out any pixel that was positive for

cheatgrass presence every year at the same location (Figure 3). The end result was a raster layer that only had pixels where cheatgrass was deemed present every year, which became the core population. All thresholds were included (low and high).




Figure 3: Graphical concept for adding all DESI years for Canyonlands NP together to extract a CORE and VARIABLE population.

Every other pixel from the individual DESI images that was present for cheatgrass became part of the temporal population or the “variable” population. Meaning, a pixel was present for cheatgrass for at least one but not all of the years of the individual park’s DESI images.

### *Landscape and Climate Models*

We initially began our work by focusing on Arches National Park, evaluating lags between precipitation in preceding seasons and DESI estimates of annual grass cover. We found inconclusive results, leading us to believe that there are other heavily weighing factors that will determine the locations and predict growth or decline in certain areas of the park.

Factors that are currently being considered are climate, topographic, and cultural in nature (Figure 4).




Figure 4: Data layers used for DESI output analysis acquired by remote sensing and satellite imagery

Topographic data include: DEM (digital elevation model), slope, and soil texture and percent clay.

Digital Elevation Model (DEM) for elevation (USGS) has been collected from Utah Automated Geographic Reference Center (UT AGRC). Tiles were mosaicked using ArcMap 10x to encompass all areas of each park. Slope was calculated using the ArcMap 10x Slope Tool with the DEM layer.

Soil texture and percent clay data was downloaded from the NRCS Web Soil Survey (Soil Survey Staff, 2015).

Climate data include precipitation for the preceding fall DESI year. If the DESI output image was for 2003 then fall 2002 precipitation was used. Maximum and minimum temperatures for the preceding fall year were also included with the climate data. Climate data has been collected from PRISM climate datasets (PRISM Climate Group, 2004). PRISM data was resampled from a 90m x 90m pixel using a cubic convolution to match the 30m x 30m pixel size of the DESI output image.

The cultural data was gathered from existing GIS databases as well as digitizing trail maps and other sources gathered from the National Park Service. Trails, visitor centers, campgrounds and any other high traffic use areas were located and combined into one layer. A 100m buffer zone was created (ESRI) around all locations. This buffer zone is used as an error buffer as well as to account for growth that may occur near but not on these specific locations. For instance, cheatgrass would not grow on a road but on the shoulder or adjacent land to the road.

All data preparation was done using ArcMap 10x to ensure quality and compatibility of the multiple data layers. An example model that was used for these adjustments can be seen in Figure 5.




Figure 5: One of the models used to adjust and transform the various data layers to all align with the DESI output images. Input raster (2) will be target data layer for manipulation. Output raster will be the new data transformed and clipped to

Transformations were required to ensure accuracy of the data for spatial and statistical analysis. Not all databases were found to be in the same datum or geographic coordinate system. Once the data was aligned, spatial autocorrelation and extrapolation was done using Multi-Value to Point Tool to build a statistical model that explains the control over the spatial and temporal distribution of cheatgrass. This statistical modeling and analysis will be done using Program R (R Core Team, 2012).

## RESULTS

### *Preliminary*

Arches NP, Bryce Canyon NP, Capitol Reef NP, and Glen Canyon NRA were the only parks found to have a core population. Canyonlands NP, Dinosaur NM, and Natural

Bridges NM did not have any of the same location of pixels present for cheatgrass that were present every year within the years of dataset available.

Initial descriptive statistics were performed to see general trends between a CORE and VARIABLE population. Analysis was done only on those parks containing a core and variable population.

When all parks and monuments were combined for soil texture there is a spike in the CORE population for higher clay content whereas the VARIABLE population had a higher sand content (Figure 6).




Figure 6: All parks that had a core and variable population. When a circle is above the bar, it indicates the probability of cheatgrass presence is higher than would be expected randomly for that bin.

When all parks and monuments were combined for preceding fall precipitation there is a clustering of moderate precipitation ranges in the variable population but higher than expected in the extreme ranges for core population (Figure 7).




Figure 7: All parks that had a core and variable population. When a circle is above the bar, it indicates the probability of cheatgrass presence is higher than would be expected randomly for that bin.

No analyses have been done using any other covariate at this point in the project.

	Soil Texture	Precipitation
Core Population	Constrained by soil texture; 15-30% clay (loamy)	Broad range
Variable Population	Wide range and abundant in 0-15% clay (rock outcrops and sandy)	Mostly found in wet range. Arches dry range anomaly is exception

Table 1: A summary of the soil texture and precipitation findings for core vs variable populations

## DISCUSSION

Cheatgrass is highly dynamic and temporally variable from year to year. Based on preliminary results we know that there are other factors other than preceding fall precipitation and clay content in soils that will determine whether cheatgrass will become established or not.

When analyzing the datasets of core populations compared to variable populations a map was created to visually validate the core versus variable theory. Based on this visual validation it was determined that the core populations were in fact, ecologically irrelevant. There was no

spatial clustering of the core population pixels to indicate that there was a large scale invasion that is well established enough to be present every year within the datasets. Because of this, the initial descriptive results of soil texture and precipitation trends may be invalid.

It is important that this large landscape level work be ecologically relevant as the primary goal of this work is to be useful in land management and conservation goals.

Other findings of the visual validation show that there is spatial clustering of cheatgrass present pixels for most years, which indicate there are landscape controls and characteristics that would determine cheatgrass growth. But in certain years there are incredibly large diebacks of cheatgrass populations. These large-scale diebacks highly reduce the probability of pixels being present in certain locations every year consistently. Another visual inspection of clustering indicates that the distance to the park boundary may be an important factor. This would make sense logically as the surrounding land is not under control of National Park Service and will have highly variable levels of conservation and rehabilitation for and of invasion of annual grasses.

Further research will need to be conducted in order to identify areas that are sensitive to and conditions that will promote cheatgrass expansion. Once the research has been completed, a set of criteria should be possible to model as control factors that indicate locales that either are sensitive to or promote the invasion of cheatgrass. As cheatgrass is an aggressive invader, the models are most likely going to be much more intricate than original sought after at the initial stages of this project.

Characterizing conditions and locations of cheatgrass populations will give land



managers insight into areas that should be of high priority for conservation. It will also give NPS an understanding if the park has been fully invaded or if there are areas that meet the criteria for invasion but have not yet experience large-scale cheatgrass growth. If the factors that control cheatgrass invasion are controllable then these areas would mostly likely set apart has high priority for conservation. Understanding this system will also make it possible to focus the restorative efforts on areas that have an increased likelihood of success in those endeavors.

The novelty of this work is it will give the scientific community, including land managers, the ability to monitor invasions at an unprecedented landscape scale.

Since it is known that annual alien grass species contribute to increasing fire cycles and is an aggressive invader it will be crucial to maintain the integrity of the wildlands in Utah on a large scale. This technology of using free open source Landsat imagery will allow for this continued large scale monitoring to occur.

All data collected for this project utilizes existing data and remote sensing platforms and is available in free, open-source databases, reducing the costs directly to land managers. It reduces the need for large field crews to be extensively sampling remote areas and reduces human bias in the collection process based on conditions of the landscape (Peterson, 2008).

## References

1. Brooks, M. L., D'Antonio, C. M., Richardson, D. M., Grace, J. B., Keeley, J. E., DiTomaso, J. M., ... & Pyke, D. (2004). Effects of invasive alien plants on fire regimes. *BioScience*, 54(7), 677-688.
2. D'Antonio, C.M., Vitousek, P. (1992). Biological invasions by exotic grasses, the grass/firecycle, and global change. *Annual Review of Ecological Systems*. 25, 63-87
3. ESRI 2011. ArcGIS Desktop: Release 10.3. Redlands, CA: Environmental Systems Research Institute.
4. Exelis Visual Information Solutions, Boulder, Colorado
5. Kokaly, R.F..(2011). Detecting cheatgrass on the Colorado Plateau using Landsat data: A tutorial for the DESI software: *U.S. Geological Survey Open-File Report 2010–1327*, 88
6. NASA (2014) Strategic Plan. [http://www.nasa.gov/sites/default/files/files/FY2014\\_NASA\\_SP\\_508c.pdf](http://www.nasa.gov/sites/default/files/files/FY2014_NASA_SP_508c.pdf)
7. National Park Service (2014) Working with Utah: By the numbers.
8. Pellant, M.. (1996) Cheatgrass: The invader that won the west. *Bureau of Land Management Interior Columbia Basin Ecosystem Management Project*
9. Peterson, E. B. (2008). Invasion of the exotic grasses: mapping their progression via satellite. *Shrublands under fire: disturbance and recovery in a changing world. Usda Forest Service Rocky Mountain research station proceedings*, 52, 33-39.
10. PRISM Climate Group, Oregon State University, <http://prism.oregonstate.edu>, created 4 Feb 2004.
11. R Core Team (2012). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>
12. Seabloom, E.W., Harpole, W.S., Reichman, O.J., Tilman, D.. (2003). Invasion, competitive dominance and resources use and native California grassland species. *Proceedings of the National Academy of Sciences of the United States of America*. 100(23), 13384-13389
13. Sieberg, M., Howery, L., Thwaites, A.. (2007) Western rangeland weeds. *Learning Modules; University of Arizona College of Agriculture and Life Sciences*.
14. Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov/>. Accessed [07/30/2015].
15. Zouhar, K. 2003. Bromus tectorum. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2015, April 20].