



Prionus Root Borer (*Prionus californicus*)

Diane Alston, Entomologist • Shawn Steffan, Entomologist¹ • Michael Pace, Box Elder County Extension Agent

Do You Know?

- This long-horned beetle is native to western North America and lives for 3 years or more underground, feeding on tree roots.
- Severe infestations can cause the death of stone fruit trees.
- In northern Utah, the rootborer is found most commonly in sweet cherry and peach orchards growing in sandy soils.
- Insecticide applications have done little to suppress prionus beetle populations.
- An attractive pheromone lure is under evaluation for monitoring, and may also suppress populations in an orchard.

Prionus root borers belong to a family of beetles (Order Coleoptera) commonly known as long-horned beetles (Cerambycidae). The larvae are often referred to as round-headed borers because their body shape is cylindrical.

Several species in the *Prionus* genus are large, root-boring beetles, and they are widely distributed across the U.S. One species, the California prionus (*Prionus californicus*), appears to have a particular affinity for stone fruit trees growing in sandy soils of Utah.

HOSTS

sweet cherry, peach, apricot, other stone fruits, cottonwood and other native/ornamental trees and shrubs

LIFE HISTORY

The adult California prionus is a large beetle, ranging in size from 1¾ - 2¼ inches long (45-60 mm). Adults are reddish-brown in color, relatively smooth and shiny with long, deeply notched antennae (Fig. 1). The larvae can be as long as 4¼ inches (108 mm) with a diameter of approximately ¾ inch (18 mm) at the widest point of its body (Fig. 2).



Fig. 1. Adult prionus root borer, female (left) with ovipositor extended, and male (right) with larger antennae for detecting female sex pheromone.

Adults emerge from pupae in the soil in July in northern Utah. The beetles fly at night in search of mates, and eggs are laid in the soil soon thereafter. Young larvae tunnel into the soil to seek out tree roots. Research in a Utah sweet cherry orchard suggests that the younger larvae begin feeding on smaller diameter roots and ultimately reach the tree crown as mature larvae as they move inward and upward along larger roots. The "crown" refers to the region of the trunk (usually at or near the soil level) that represents the transition between below-ground (roots) and above-ground (trunk) growth. In cherries, a greater proportion of the larvae found at the crown were large, mature larvae, while most of the



Fig. 2. Prionus larvae of various ages, ¾ - 4¼ inches long.

¹Washington State University

smaller, younger larvae were found in roots. Following 3 or more years of root and crown feeding and upward movement, the insect pupates close to the soil surface. The adults then emerge and seek out mates.

HOST INJURY



Fig. 3. Furrows in sweet cherry roots made by prionus larvae.

Larvae will chew deep furrows in the roots (Fig. 3), often causing severe reduction in a tree's ability to take up water and nutrients. The feeding injury is often associated with rot and decay, particularly in wet soils. The colonization of feeding wounds by soil microbes will compound the damage caused by larval tunneling. Larvae feeding in the crown form spiraling furrows which girdle the crown and upper roots (Fig. 4).

Sandy soils appear to favor prionus infestations. Not only is sand likely to increase the ease with which larvae can move about, but it allows rapid percolation of irrigation water. The loss of water from the soil profile combined with the tree's reduced ability to uptake water (due to prionus feeding) can become major stressors, even for well-established trees.



Fig. 4. Cross-section of crown of a sweet cherry tree showing circular pattern of prionus tunnels.

MONITORING

Adult prionus can be monitored using pheromone and light traps. The *Prionus californicus* sex pheromone has been identified and found to be highly attractive to male beetles. Studies in northern Utah have found that hanging the pheromone lure over a funnel placed in a buried bucket can be an attractive trap to the male beetles (Fig. 5 A and B). In a sweet cherry orchard in 2009, males were captured in traps during July and August; more males were caught in bucket than panel traps, and more were caught in pheromone-bated (Phero) than non-baited (Untrt) traps (Fig. 6.) In 2010, males were caught in pheromone-bated traps from July through September. A commercial pheromone lure is expected to be available soon. If pheromone traps are attractive enough, they may be successful in reducing the local *Prionus californicus* population (i.e., mass trapping). The beetles typically fly soon after sunset, and they are attracted to lights. Trap-catch may decline after midnight, presumably due to colder temperatures. Adults begin to emerge in early July in northern Utah, and probably 2-4 weeks earlier in southern Utah.



Fig. 5A. Bucket trap with experimental prionus pheromone lure.



Fig. 5B. Prionus males caught in bucket trap.




Fig. 6. Weekly capture of prionus males in pheromone- and non-baited bucket and panel traps.

Larvae can be monitored by digging into the soil (6-10 inches deep) around a trunk and looking for darkened, soft areas around the tree crown. Probing into the dark, soft material within the crown or lower trunk may expose

larvae. Alternatively, more soil can be unearthed to search for infestations on roots. Usually, infested trees will show signs of limb dieback or a marked decline in overall vigor during hot spells (Fig. 7).



Fig. 7. Sweet cherry tree decline due to prionus infestation.

To prevent prionus infestations, maintain tree health and prevent stress. Watch for canopy dieback or sudden loss of tree vigor. Following an infested field for two or more years, planting annual crops that will be tilled under each year, and avoiding planting stone fruit trees into infested sites are the best options for managing prionus root borer in infested soil. When the pheromone becomes commercially available, mass-trapping males in bucket traps may be an additional control option.

MANAGEMENT

Currently, there are no registered insecticides for prionus root borers. Several insecticides registered for stone fruit may provide incidental suppression of the adult. Soil fumigation is a pre-planting option where infestations are known to exist, but its cost-effectiveness is questionable. Systemic materials, such as imidacloprid (Admire, Merit, and Bayer Advanced Garden Tree and Shrub Insect Control) may be options, but recent research indicated that this material may be effective only on the younger larvae on roots, and not against the older larvae in the crown or lower trunk. Imidacloprid may suppress the local population if used annually over several years. Some imidacloprid formulations are not registered on bearing fruit trees, so always read the label carefully. Other neurotoxic insecticides, such as chlorpyrifos, permethrin, or carbaryl will kill adults and may inhibit egg-laying when fresh residue is present on the lower trunk and soil, but will not suppress larval populations on the roots.

Precautionary Statement: Utah State University Extension and its employees are not responsible for the use, misuse, or damage caused by application or misapplication of products or information mentioned in this document. All pesticides are labeled with ingredients, instructions, and risks, and not all are registered for edible crops. "Registered use" pesticides may only be applied by a licensed applicator. The pesticide applicator is legally responsible for proper use. USU makes no endorsement of the products listed herein.

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions. Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities. This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Dept. of Ag., Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.