

Human–Wildlife Interactions 8(1):78–87, Spring 2014

Morbidity and mortality in reptiles presented to a wildlife care facility in Central Illinois

ANNE E. RIVAS, Department of Veterinary Clinical Medicine, University of Illinois College of Veterinary Medicine, 1008 West Hazelwood Drive, Urbana, IL 61802, USA arivas2@illinois.edu

MATTHEW C. ALLENDER, Department of Comparative Biosciences, University of Illinois College of Veterinary Medicine, 2001 South Lincoln Avenue, Urbana, IL 61802, USA

MARK MITCHELL, Department of Veterinary Clinical Medicine, University of Illinois College of Veterinary Medicine, University of Illinois College of Veterinary Medicine, 1008 West Hazelwood Drive, Urbana, IL 61802, USA

JULIA K. WHITTINGTON, Department of Veterinary Clinical Medicine, University of Illinois College of Veterinary Medicine, 1008 West Hazelwood Drive, Urbana, IL 61802, USA

Abstract: We examined morbidity and mortality of 200 reptiles, representing 13 different species that were presented to the University of Illinois Wildlife Medical Clinic (WMC) from 2003 to 2010. Snapping turtles (*Chelydra serpentina*; $n = 46$), box turtles (*Terrapene* sp.; $n = 43$), painted turtles (*Chrysemys picta*; $n = 37$), and red-eared slider turtles (*Trachemys scripta elegans*; $n = 33$) were the most frequently seen species. Turtles were significantly more likely to be presented to the WMC following collision with a motor vehicle ($n = 73$) than any other reason, including idiopathic trauma (i.e., trauma of unknown origin; $n = 25$) or infectious disease ($n = 18$). The findings from this cross-sectional study suggest a potential for community education in limiting reptile traumas resulting in presentations to a wildlife hospital.

Key words: anthropogenic, human–wildlife conflicts, morbidity, mortality, reptile, trauma

REPTILE POPULATIONS are in decline worldwide (Gibbons et al. 2000), and many of the declines are attributed to human encroachment on natural habitat along with other anthropogenic activities (Gardner et al. 2007). However, there are likely regional differences for reptile population declines, too, and studies are required to characterize morbidity and mortality of local reptile populations and their effect on biodiversity to effectively develop management plans to protect these animals. One means of monitoring local causes of morbidity and mortality of native wildlife, including reptiles, is using case information obtained from wildlife rehabilitation facilities (Brown et al. 2003, Joyner et al. 2006).

The Wildlife Medical Clinic (WMC; Urbana, Il.) at the University of Illinois College of Veterinary Medicine is a wildlife hospital that provides medical care to free-living wildlife species in east central Illinois. Although WMC will accept native wildlife from anywhere within the state of Illinois, most wildlife patients originated in Champaign County, with the surrounding counties serving as the second most common source of animals (Figure 1). This region encompasses 1,307,892 ha, divided into cropland (1,048,316 ha), grassland (157,974

ha), forest (40,154 ha), urban and built-up areas (34,055 ha), open water (15,338 ha), wetland (10,790 ha), and barren land (454 ha; Illinois Department of Natural Resources 2004). The most recent census reported the population of these counties as 521,644 (U.S. Census Bureau 2010).

As a point-of-care facility for sick or injured wildlife, the WMC serves as an investigative source for wildlife pathologies in this region. A few previous investigations have evaluated trends in morbidity and mortality among reptiles presented to wildlife hospitals (Hartup 1996, Brown and Sleeman 2002, Schrader et al. 2010). However, these studies have been in regions with different habitats, species distributions, and land-use breakdowns. To our knowledge, morbidity and mortality trends in reptiles from central Illinois have not been evaluated.

This study reviews medical records of reptiles presented to the WMC. Our goal was to determine causes of morbidity and mortality to assess risk factors that can be used to guide future case management and preventive measures against anthropogenic injury of reptiles in Illinois.

Methods

This cross-sectional study was conducted using data from reptile cases presented to the WMC from 2003 to 2010. All cases were presented by good Samaritans who presumed that the animals were in need of assistance. We examined original medical records from 249 reptiles admitted to the WMC within this time period. Exclusion criteria consisted of non-native species, previously owned animals, animals that were presented for research transmitter implantation, and those with incomplete medical records. Based on these criteria, 200 of these cases were included for analysis.

Diagnosis was assigned with the following criteria: hit by car = trauma due to known automobile collision or after being found traumatized in the vicinity of a roadway ($n = 73$); lawn equipment = trauma due to lawn mower or plastic fencing ($n = 9$); animal attack = trauma inflicted by another animal as indicated by bite wounds ($n = 7$); fish hook = imbedded fish hook ($n = 16$); idiopathic trauma = trauma of unknown origin ($n = 25$); infectious = pneumonia and aural abscesses ($n = 18$); disturbed hibernation = reptiles collected by finders during hibernation ($n = 14$); healthy = no discernible medical problem ($n = 30$); idiopathic = conditions of unknown etiology ($n = 5$); and miscellaneous trauma = trauma from being closed in a door and entrapped in a glue trap ($n = 3$). Diagnoses of hit by car, lawn equipment, fish hook, disturbed hibernation, and miscellaneous trauma were collectively considered as human-related problems ($n = 115$). Only the primary problems were interpreted for diagnosis. The medical diagnoses were determined by the original attending clinician and were based on case history, physical examination findings, and any ancillary diagnostic tests.

Age classes were assigned to each animal based on subjective criteria that included mass and length. Sex was determined based on plastron morphology (chelonians), cloaca position (chelonians), radiographic evidence of eggs (chelonians), or cloacal probing (serpentes). Final dispositions were classified as released, died, or euthanized.

Frequencies were determined for species, age, sex, diagnosis, and final disposition. Associations between each outcome variable




Figure 1. Map of Illinois with the study area consisting of Champaign and its surrounding counties shaded gray. The approximate location of the Wildlife Medical Clinic is indicated with an asterisk.

(diagnosis and final disposition) and each independent variable (species, age, and sex) were initially evaluated using Fisher's exact test and Pearson's Chi-square. Categorical variables were assessed for equal probabilities using the 1 sample Chi-square test. We used a Chi-square test to determine associations between month animals were presented to the WMC and the 4 most commonly presented species: box turtles (*Terrapene* sp.), painted turtles (*Chrysemys picta*), snapping turtles (*Chelydra serpentina*), and red-eared slider turtles (*Trachemys scripta elegans*). Ordinal regression modeling was then done to evaluate these independent variables (species, age, sex) simultaneously by each dependent variable (final disposition and diagnosis). A $P < 0.05$ was used to determine statistical significance. We used SPSS 19.0 (SPSS Inc., Chicago, Ill.) to analyze the data.

Results

Seventeen reptile species were presented




Figure 2. Number of individuals presented by month of the 4 most common reptile species (eastern box turtle, painted turtle, snapping turtle, slider turtle) seen at the Wildlife Medical Clinic, 2003 to 2010.

to the WMC for care. However, due to inconsistent classification in the records, box turtles (*Terrapene carolina* and *T. ornata*), garter snakes (*Thamnophis radix* and *T. sirtalis*), king snakes (*Lampropeltis calligaster* and *L. getula*), and softshell turtles (*Apolone mutica* and *A. spinifera*) were each reported as a single species for the purposes of this study (Table 1).

All animals that were not classified as healthy received appropriate medical treatment (fracture repair, wound care, antimicrobial therapy, analgesia, fluid therapy, supplemental feedings, or euthanasia). The majority of patients were released or transferred to a licensed wildlife rehabilitator for ongoing care prior to release ($n = 114$). The others were euthanized ($n = 58$) or died while in captivity ($n = 27$). Final disposition varied significantly ($P < 0.001$) based on diagnosis and is presented in Table 2.

Table 1: Reptiles presented to the Wildlife Medical Clinic at the University of Illinois College of Veterinary Medicine, 2003 to 2010.

| Species | Number of animals |
|---|-------------------|
| Snapping turtle (<i>Chelydra serpentina</i>) | 46 |
| Box turtle (<i>Terrapene carolina</i> , <i>T. ornata</i>) | 43 |
| Painted turtle (<i>Chrysemys picta</i>) | 37 |
| Slider turtle (<i>Trachemys scripta</i>) | 33 |
| Garter snake (<i>Thamnophis radix</i> , <i>T. sirtalis</i>) | 17 |
| Fox snake (<i>Elaphe vulpina</i>) | 6 |
| Softshell turtle (<i>Apolone mutica</i> , <i>A. spinifera</i>) | 6 |
| King snake (<i>Lampropeltis calligaster</i> , <i>L. getula</i>) | 4 |
| Eastern massasauga rattlesnake (<i>Sistrurus catenatus</i>) | 2 |
| Black rat snake (<i>Elaphe obsoleta</i>) | 2 |
| Bullsnake (<i>Pituophis melanoleucus</i>) | 2 |
| Brown snake (<i>Storeria dekayi</i>) | 1 |
| Eastern hognose snake (<i>Heterodon platirhinos</i>) | 1 |
| Total | 200 |




Figure 3. Morbidity and mortality category by month of the 4 most common reptile species (eastern box turtle, painted turtle, snapping turtle, slider) presented to the Wildlife Medical Clinic, 2003 to 2010.

Patients were significantly more likely to be adult animals ($n = 149$) than juveniles ($n = 35$) or unknown age ($n = 16$) animals. Eighteen animals were female, 17 animals were male, and 165 animals had sex listed as unknown.

Human-related problems predominated among causes of morbidity and mortality ($n = 115$; Table 3). Hit-by-car was both the most frequently encountered human-related problem and most frequently encountered problem overall ($n = 73$). The next most common diagnosis was healthy ($n = 30$).

Causes of morbidity and mortality varied significantly ($P < 0.001$) among species (Table 4). Hit-by-car was the most frequent diagnosis among the 4 most commonly encountered species: common snapping turtles, box turtles, painted turtles, and red-eared slider turtles. Box turtles constituted nine of the 18 infectious disease cases. The most commonly recorded species was the snapping turtle ($n = 46$), of

which animals were healthy with no medical problems.

The 4 most common species were not presented to the WMC with equal probabilities throughout the year ($P < 0.001$), but animals

Table 2: Final disposition of reptiles by diagnosis presented to the Wildlife Medical Clinic at the University of Illinois College of Veterinary Medicine, 2003 to 2010.

| Diagnosis | Released | Died | Euthanized |
|-----------------------|----------|------|------------|
| Hit by car | 25 | 12 | 35 |
| Lawn equipment | 4 | 0 | 5 |
| Animal attack | 4 | 0 | 3 |
| Fish hook | 14 | 1 | 1 |
| Idiopathic trauma | 11 | 6 | 8 |
| Infectious | 11 | 5 | 2 |
| Disturbed hibernation | 13 | 0 | 1 |
| Healthy | 30 | 0 | 0 |
| Idiopathic | 1 | 1 | 3 |
| Miscellaneous trauma | 1 | 2 | 0 |
| Total | 114 | 27 | 58 |

were seen each month (Figure 2). Hit-by-car was the most common ($P < 0.001$) presented in every month except from November through February (Figure 3). There were significant differences in the diagnosis of infectious disease by month ($P = 0.003$). There was no significant association in month presented and disposition ($P = 0.399$).

Discussion

Compared to previous studies conducted at other wildlife health centers, the results of this cross-sectional study were unique in identifying snapping turtles as the most frequently encountered species (Hartup 1996, Brown and Sleeman 2002). There are a few reasons that could account for this. The WMC is located in the Wabash River Watershed, where snapping turtles are the most abundant aquatic turtle species (Rizkalla and Swihart 2006). Snapping turtles also are known to be more vagrant than other species (Christens and Bider 1987) when searching for suitable hibernation sites and bodies of water (Brown and Brooks 1994, Burke et al. 1995). Galbraith (2008) speculated that snapping turtles’ limited carapace and plastron allow for greater mobility, enabling them to migrate greater distances in comparison to other aquatic turtles. In our 1,307,892-ha study area, 1,082,372 ha are dedicated to agriculture

or urban and built-up use (Illinois Department of Natural Resources 2004). The limited amount of natural habitat that remains, combined with snapping turtles’ tendency to travel greater distances, could result in an increased likelihood that these turtles would encounter humans, leading to both the high number of

Table 3: Morbidity and mortality of reptiles presented to the Wildlife Medical Clinic at the University of Illinois College of Veterinary Medicine, 2003 to 2010.

| Morbidity and mortality category | Number |
|----------------------------------|--------|
| Human-related problem | 115 |
| Hit by car | 73 |
| Fish hook | 16 |
| Disturbed hibernation | 14 |
| Lawn equipment | 9 |
| Miscellaneous trauma | 3 |
| Healthy | 30 |
| Idiopathic trauma | 25 |
| Infectious | 18 |
| Animal attack | 7 |
| Idiopathic | 5 |
| Total | 200 |

Table 4: Morbidity and mortality in most frequently presented reptile species at the Wildlife Medical Clinic at the University of Illinois College of Veterinary Medicine, 2003 to 2010.

| | Box turtle (<i>Terrapene carolina</i> , <i>T. ornata</i>) | Painted turtle (<i>Chrysemys picta</i>) | Slider turtle (<i>Trachemys scripta</i>) | Snapping turtle (<i>Chelydra serpentina</i>) |
|-----------------------|---|--|---|---|
| Human related problem | 23 | 29 | 18 | 24 |
| Hit by car | 18 | 19 | 12 | 18 |
| Fish hook | 0 | 3 | 5 | 5 |
| Disturbed hibernation | 1 | 6 | 1 | 1 |
| Lawn equipment | 4 | 1 | 0 | 0 |
| Miscellaneous trauma | 0 | 0 | 0 | 0 |
| Healthy | 4 | 1 | 3 | 16 |
| Idiopathic trauma | 4 | 4 | 5 | 5 |
| Infectious | 9 | 1 | 5 | 1 |
| Animal attack | 2 | 2 | 1 | 0 |
| Idiopathic | 1 | 0 | 1 | 0 |
| Total | 43 | 37 | 33 | 46 |

vehicular traumas and collection by people who found healthy animals that were assumed to be in need of care.

Overall, most of the reptiles seen were adults, although there was a potential for misclassification bias in age characterization because morphometrics on these patients was not recorded. Adult wild turtles tend to be more vagrant than juveniles (Sexton 1959, Browne et al. 2006), possibly due to the greater physiologic costs and predation risks faced by juvenile turtles during migration (Lefevre and Brooks 1995, Finkler 2001). As with the snapping turtles, increased terrestrial exposure both makes adult wild reptiles more susceptible to vehicular trauma and collection of healthy animals by humans, the 2 most frequently recorded diagnoses of animals in this study.

Trauma after being hit by a car was the most common reason reptiles were presented to the WMC. Subjectively, these findings follow trends seen in similar studies, but, due to differences in data reporting, there are limitations to comparing findings (Hartup 1996, Brown and Sleeman 2002, Schrader et al. 2010). This high prevalence of vehicular trauma is not surprising. Regions with increased road density are known to have higher mortality and subsequent alterations in population structure for wild reptiles (Steen and Gibbs 2004, Patrick and Gibbs 2010). In addition to a higher likelihood of encountering roads during terrestrial migrations, increased vehicular trauma in such habitats may be due to reptiles spending more time basking on paved roads. Herpetofauna in distressed habitats are subject to sublethal environmental stressors that may suppress their immune systems (Carey 1993, Amo et al. 2007). Once ill, some reptiles combat some pathogens by behaviorally selecting and maintaining higher than normal body temperatures (Mongas and Gatten 1983, Burns et al. 1996).

Infectious etiologies were the second leading diagnosis for box turtles, which comprised the majority of the infectious disease cases. This was also the case among box turtles in another study (Schrader et al. 2010). There were seasonal trends in the number of infectious disease cases of animals presented to the WMC (Figure 3). All cases in December and half of the cases in November and April were infectious,

while there were no infectious disease cases in October and few in all other months. The high number of infectious disease cases seen in these months may be related to decreased immune function during hibernation periods. Immune data from reptiles suggests that immune function is generally depressed during hibernation, though heightened in autumn just prior to hibernation (Muñoz and de la Fuente 2004, Nelson and Demas 1996, and Schwanz et al. 2011).

Despite the frequent occurrence of infectious etiologies in box turtles, the true prevalence of infectious disease among reptiles presented to WMC is difficult to ascertain given that many trauma cases are allotted only a brief physical exam prior to euthanasia. Ancillary testing may have shown that reptiles suffering trauma were impacted by underlying infectious disease etiologies. Limited funding prohibits advanced testing in many instances, but the importance of testing for infectious disease pathogens should not be overlooked. Infections with herpes virus, *Mycoplasma* or ranavirus, are without cure in reptiles, and animals harboring these pathogens could spread them upon release back to wild populations (Schumacher 2006, Johnson et al. 2008). To prevent introduction and propagation of pathogens in wild populations, it is recommended to release reptiles only that have been shown to be disease-free through appropriate screening protocols (Jacobson et al. 1999).

There were temporal trends in the presentation of the 4 most common species (Figure 2), which correlated with the reproductive behavior of these species. May and June were the most common months for snapping turtles and box turtles to be brought to the WMC. Illinois populations of snapping turtles are known to nest mid-May to mid-June, and box turtles nest in June (Phillips et al. 1999). Slider turtles and painted turtles, which nest in Illinois during May to July (Phillips et al. 1999), were most commonly seen at the WMC during these same months. It is likely that the increase in number of animals seen during these months is due to increased migration by female turtles in search of nesting sites. Previous studies suggest that there is disproportionate road mortality of aquatic turtle females on nesting migrations in areas with increased road density (Patrick

and Gibbs 2010; Steen and Gibbs 2004). This coincides with our findings, which show a peak number of animal presentation during nesting months, as well as the highest number of hit-by-car diagnoses in May and June (Figure 3).

Overall, most (57%) animals were released or transferred to a licensed wildlife rehabilitator in preparation for release. Of the healthy animals, 49% of the patients were released or transferred to a wildlife rehabilitator. In the same time period at the WMC, the overall release rate for all species including reptiles was 41.2%. The higher release rate of reptile species may be due to reptiles being better able to withstand and recover from severe traumatic events than free-living avian and mammalian species.

The effects of human activities on the reptile patients in this study were not limited to vehicular trauma. Overall, human-related problems constituted >50% of all cases seen. Admittedly, the disease frequencies seen in the patient population at this clinic may not be representative of the entire disease scope of the natural population, as patient presentation is dependent on good Samaritans in the community. There is a suspected bias toward human-related problems in the animals presented to the clinic, and an underrepresentation of infectious disease compared with the natural population. Similar results have been found in other retrospectives assessing morbidity and mortality of reptiles presented to wildlife centers (Spalding and Forrester 1993, Hartup 1996, Brown and Sleeman 2002).

The reliance on good Samaritans may also contribute to a bias in the types of species that present. Chelonians appear over-represented, whereas snakes appear under-represented. This may be due to differences in natural history, with chelonians being less secretive or more likely to travel greater distances. Due to the protective nature of their shells, chelonians are also more likely to survive a traumatic impact when compared with snakes, resulting in a greater chance of being recovered by good Samaritans. This is supported by reptile road mortality surveys from another region of central Illinois that found snakes greatly outnumbered chelonians among vehicular fatalities (Shepard et al. 2008a). Additionally, greater species

charisma might make chelonians more likely to encounter people willing to transport these animals for care. How the patient demographics compare to actual species abundance in the area is difficult to ascertain due to a lack of recent fieldwork on wild reptile populations in the region (Phillips et al. 1999).

In summary, most of known causes of morbidity and mortality among reptiles during the study period resulted from human activities. Many more healthy animals were relocated to new populations when insufficient information on their origin was available. This indicates that human activities are having an adverse effect on wild reptiles, though the true ramifications for the wild populations cannot be determined without prospective study. However, such studies for reptiles tend to be rare given the time and cost constraints in following these long-lived animals. As a result, retrospective evaluations of medical records provide valuable insight to possible trends in population health.

Although more study is required to comprehend the scope of human impacts on wild reptile populations, what is known is that reptiles are one of the most threatened taxonomic groups worldwide (International Union for the Conservation of Nature [IUCN] Red List 2011). In Illinois alone, 18 reptile species are listed as threatened or endangered (Illinois Endangered Species Protection Board 2011). Community education offers a potential to mitigate the impact of human activities that contribute to reptile population decline. Education may also limit the number of healthy animals that are unnecessarily relocated. Information from this study has been incorporated in the WMC community education programs. Future analysis will hopefully indicate that these education efforts are having an effect to reduce the number of reptiles presented to the WMC for human-related morbidity and mortality.

Acknowledgments

We thank the many students from the University of Illinois Wildlife Medical Clinic and the University of Illinois College of Veterinary Medicine for their support of the Wildlife Medical Clinic and the care of these patients.

Literature cited

- Amo, L., P. Lopez, J. Martin. 2007. Habitat deterioration affects body condition of lizards: a behavioral approach with *Iberolacerta cyreni* lizards inhabiting ski resorts. *Biological Conservation* 135:77–85.
- Brown, G. P., and J. R. Brooks. 1994. Characteristics of and fidelity to hibernacula in a northern population of snapping turtles, *Chelydra serpentina*. *Copeia* 1994:222–226.
- Brown, J. D., and J. M. Sleeman. 2002. Morbidity and mortality of reptiles admitted to the Wildlife Center of Virginia, 1991 to 2000. *Journal of Wildlife Disease* 38:699–705.
- Brown, J. D., J. M. Sleeman, and F. Elvinger. 2003. Epidemiologic determinants of aural abscessation in free-living eastern box turtles (*Terrapene Carolina*) in Virginia. *Journal of Wildlife Disease* 39:918–921.
- Browne, D. R., M. A. Bowers, and J. E. Hines. 2006. Connectivity in an agricultural landscape as reflected by interpond movements of a freshwater turtle. *Conservation Biology* 20:780–791.
- Burke, V. J., J. L. Greene, and J. W. Gibbons. 1995. The effect of sample size and study duration on metapopulation estimates for slider turtles (*Trachemys scripta*). *Herpetologica* 51:451–456.
- Burns, G., A. Ramos, A. Muchlinski. 1996. Fever response in North American snakes. *Journal of Herpetology* 30:133–139.
- Carey C. 1993. Hypothesis concerning the causes of the disappearance of boreal toads from the mountains of Colorado. *Conservation Biology* 7:355–362.
- Christens, E., and J. R. Bider. 1987. Nesting activity and hatching success of the painted turtle (*Chrysemys picta marginata*) in southwestern Quebec. *Herpetologica* 43:55–56.
- Finkler, M. S. 2001. Rates of water loss and estimates of survival time under varying humidity in juvenile snapping turtles (*Chelydra serpentina*). *Copeia* 2001:521–525.
- Galbraith, D. A. 2008. Population biology and population genetics. Pages 168–180 in A. C. Steyermark, M. S. Finkler, and R. J. Brooks, editors. *Biology of the snapping turtle (Chelydra serpentina)*. John Hopkins University Press, Baltimore, Maryland, USA.
- Gardner, T. A., J. Barlow, and C. A. Peres. 2007. Paradox, presumption and pitfalls in conservation biology: the importance of habitat change for amphibians and reptiles. *Biological Conservation* 138:166–179.
- Gibbons, J. W., S. E. Scott, T. J. Ryan, K. A. Buhlmann, T. D. Tuberville, B. S. Metts, J. L. Greene, T. Mills, Y. Leiden, S. Poppy, and C. T. Winne. 2000. The global decline of reptiles, *deja-vu* amphibians. *Bioscience* 50:653–667.
- Hartup, B. K. 1996. Rehabilitation of native reptiles and amphibians in DuPage County, Illinois. *Journal of Wildlife Diseases*. 32:109–112.
- Illinois Department of Natural Resources. 2004. Illinois land cover statistical summary: 1999–2000. Springfield, Illinois, USA.
- Illinois Endangered Species Protection Board. 2011. Checklist of endangered and threatened animals and plants of Illinois. Illinois Department of Natural Resources, Springfield, Illinois, USA.
- International Union for the Conservation of Nature Red List. 2011. IUCN Red List summary statistics: number of species in each Red List category in each major animal taxonomic group, <http://www.iucnredlist.org/documents/summarystatistics/2011_2_RL_Stats_Table3a.pdf>. Accessed November 12, 2013.
- Johnson, A. J., A. P. Pessier, J. F. X. Wellehan, A. Childress, T. M. Norton, N. L. Stedman, D. C. Bloom, W. Belzer, V. R. Titus, R. Wagner, J. W. Brooks, J. Spratt, and E. R. Jacobson. 2008. Ranavirus infection of free-ranging and captive box turtles and tortoises in the United States. *Journal of Wildlife Diseases* 44:851–863.
- Jacobson, E. R., J. L. Behler, and J. L. Jarchow. 1999. Health assessment of chelonians and release into the wild. Pages 232–242 in M. E. Fowler and R. E. Millers, editors. *Zoo and wild animal medicine and current therapy*. Saunders, Philadelphia, Pennsylvania, USA.
- Joyner, P. H., S. Kelly, A. A. Shreve, S. E. Snead, J. M. Sleeman, and D. A. Pettit. 2006. West Nile virus in raptors from Virginia during 2003: clinical, diagnostic and epidemiologic findings. *Journal of Wildlife Disease* 42:335–344.
- Lefevre, K., and R. J. Brooks. 1995. Effects of sex and body size on basking behavior in a northern population of the painted turtle, *Chrysemys picta*. *Herpetologica* 51:217–224.
- Mongas, W. R., and R. E. Gatten. 1983. Behavioural fever in the turtles *Terrapene carolina* and *Chrysemys picta*. *Journal of Thermal Biology* 8:285–288.
- Muñoz, F. J., and M. de la Fuente. 2004. Seasonal


- changes in lymphoid distribution of the turtle *Mauremys caspica*. *Copeia* 2004:178–183.
- Nelson, R. J., and G. E. Demas. 1996. Seasonal changes in immune function. *Quarterly Review of Biology* 71:511–548.
- Patrick, D. A., and J. P. Gibbs. 2010. Population structure and movements of freshwater turtles across a road-density gradient. *Landscape Ecology* 25:791–801.
- Phillips, C. A., R. A. Brandon, and E. O. Moll. 1999. Field guide to amphibians and reptiles of Illinois. Manual 8. Illinois Natural History Survey, Urbana, Illinois, USA.
- Rizkalla, C. E., and R. K. Swihart. 2006. Community structure and differential responses of aquatic turtles to agriculturally induced habitat fragmentation. *Landscape Ecology* 21:1361–1375.
- Schrader, G. M., M. C. Allender, and A. Odoi. 2010. Diagnosis, treatment, and outcome of eastern box turtles (*Terrapene carolina carolina*) presented to a wildlife clinic in Tennessee, USA, 1995–2007. *Journal of Wildlife Disease* 45:1079–1055.
- Schumacher, J. 2006. Selected infectious diseases of wild reptiles and amphibians. *Journal of Exotic Pet Medicine* 15:18–24.
- Schwanz, L., D. A. Warner, S. McGaugh, R. Di Terlizzi, and A. Bronikowski. 2011. State-dependent physiological maintenance in long-lived ectotherm, the painted turtle (*Chrysemys picta*). *Journal of Experimental Biology* 214:88–97.
- Sexton, O. J. 1959. Spatial and temporal movements of a population of the painted turtle, *Chrysemys picta marginata*. *Ecological Monographs* 29:113–140.
- Shepard, D. B., M. J. Dreslik, B. C. Jellen, and C. A. Phillips. 2008a. Reptile road mortality around an oasis in the Illinois corn desert with emphasis on the endangered eastern massasauga. *Copeia* 2:350–359.
- Shepard, D. B., A. R. Kuhns, M. J. Dreslik, and C. A. Phillips. 2008b. Roads as barriers to animal movement in fragmented landscapes. *Animal Conservation* 11:288–296.
- Spalding, M. G., and D. J. Forrester. 1993. Disease monitoring of free-ranging and released wildlife. *Journal of Zoo and Wildlife Medicine* 24:271–280.
- Steen, D. A., and J. P. Gibbs. 2004. Effects of roads on the structure of freshwater turtle populations. *Conservation Biology* 18:1143–1148.
- U.S. Census Bureau. 2010. 2010 census population compared to 2000: Illinois counties. U.S. Census Bureau, Spauldings, Maryland, USA.
-



ANNE E. RIVAS received dual bachelor degrees in integrative biology and animal sciences from the University of Illinois in 2009. While attending veterinary school, she served as manager of the Wildlife Medical Clinic. Upon graduating from the University of Illinois College of Veterinary Medicine in 2013, she began a 1-year clinical internship in small animal medicine and surgery at VCA Emergency Animal Hospital and Referral Center in San Diego, California. She is currently working toward the goal of being boarded in zoological medicine. Her interests include free-ranging, as well as captive, wildlife health and medicine.



MATTHEW C. ALLENDER received his B.S. degree in ecology, ethology, and evolution in 2000 and D.V.M. degree in 2004 from the University of Illinois. He received his M.S. degree for his thesis, "Health and disease assessments in two reptiles: an ophidian and a chelonian" in 2006 from the University of Illinois. He completed a clinical residency in zoological medicine at the University of Tennessee–Knoxville Zoological Gardens in 2009 and became a Diplomate of the American College of Zoological Medicine in 2011. He received his Ph.D. degree for his dissertation, "Characterizing the epidemiology of ranaviruses in North American chelonians: diagnosis, surveillance, pathogenesis, and treatment," from the University of Illinois in 2012. He is currently a clinical assistant professor in the Department of Comparative Biosciences at the University of Illinois, where he directs the wildlife epidemiology laboratory. He is a clinical veterinarian for the Miller Park Zoo and an assistant editor for the *Journal of Wildlife Diseases*. His research interests focus on the health epidemiology of diseases of free-ranging wildlife, specifically reptiles.



MARK A. MITCHELL received his B.S. degree in veterinary science in 1990 and D.V.M. degree in 1992 from the University of Illinois. He received his M.S. degree for his thesis, "Assessment of hematologic parameters and infectious and parasitic agents in western Illinois raccoons" in 1997 from the University of Illinois. He received his Ph.D. degree for his dissertation, "Epidemiology of *Salmonella* spp. in the green iguana (*Iguana iguana*)" from Louisiana State University in 2001. He is currently a professor in the Department of Veterinary Clinical Medicine at the University of Illinois. He is the editor of the *Journal of Herpetological Medicine and Surgery* and co-editor of the *Journal of Exotic Pet Medicine*. His research interests focus on the epidemiology of diseases in both captive and free-ranging wildlife.



JULIA K. WHITTINGTON received her bachelor's degree from the University of Illinois with a background in ecology, ethology, and evolution. In 1997, she received her D.V.M. degree from the University of Illinois College of Veterinary Medicine and later worked in small animal and exotic animal practice. In 2001, she returned to the College of Veterinary Medicine and currently serves as a clinical associate professor in zoological medicine and surgery, overseeing the Wildlife Medical Clinic. She also serves as head of the zoological medicine service at the University of Illinois Veterinary Teaching Hospital. Her interests include avian, small mammal, and wildlife medicine.