

Utilising daily diaries to examine oral health experiences associated with dentine hypersensitivity

PORRITT, Jenny, SUFI, Farzana and BAKER, Sarah

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/13306/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

PORRITT, Jenny, SUFI, Farzana and BAKER, Sarah (2016). Utilising daily diaries to examine oral health experiences associated with dentine hypersensitivity. *BMC Oral Health*, 16 (1).

Repository use policy

Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in SHURA to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

1 **Utilising daily diaries to examine oral**
2 **health experiences associated with dentine**
3 **hypersensitivity**

4 Jenny M Porritt^{1*}, Farzana Sufi², Sarah R Baker³

5
6 ¹Senior Lecturer in Psychology

7 Department of Psychology, Sociology & Politics

8 Sheffield Hallam University

9 Collegiate Crescent

10 Sheffield

11 S10 2BQ

12 Corresponding author. Email: j.porritt@shu.ac.uk, (+44 (0)114 225 6535)

13
14 ² Director of Clinical Research (Oral Care)

15 Consumer Healthcare GlaxoSmithKline

16 St Georges Avenue

17 Weybridge, Surrey

18 KT13 0DE

19 Email: farzana.x.sufi@gsk.com

20
21 ³ Professor in Psychology

22 School of Clinical Dentistry

23 19 Claremont Crescent

24 University of Sheffield

25 Sheffield, UK

26 S10 2TA

27 Email: s.r.baker@sheffield.ac.uk

Abstract

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

Background: The current investigation examined the determinants of oral health experiences associated with dentine hypersensitivity using prospective diary methodology.

Methods: Staff and students from a large UK university who had self-diagnosed dentine hypersensitivity completed an online daily diary and text survey for two weeks recording their mood, oral health-related coping behaviours, coping and pain appraisals, pain experiences and functional limitations. Cross sectional and lagged path analyses were employed to examine relationships.

Results: 101 participants took part in the diary study. Participants had a mean age of 26.3 years (range=18-63) and most were female (N=69). Individuals who used more oral health-related coping behaviours predicted and experienced greater levels of pain on subsequent days. Negative mood also predicted worse pain outcomes. The daily diary method provided a useful avenue for investigating variations in oral health experiences and relationships between variables that can fluctuate daily.

Conclusions: Psychological variables such as coping and mood play an important role in the pain experiences of people with dentine hypersensitivity. The study highlights the benefits of using prospective methods to elucidate the experiences of people with oral conditions.

Keywords: pain, dentine hypersensitivity, coping, diary, oral impacts

Background

Dentine hypersensitivity affects approximately half of the population and the pain associated with this condition can limit oral functions such as eating and tooth brushing, which negatively affect individual oral health related quality of life [1-5]. Previous research has revealed that key psychological factors can impact on the quality of life of people with dentine hypersensitivity [6], however, the symptoms of this condition are transient and fluctuate frequently. Moreover, even the self-management of dentine hypersensitivity can vary frequently as people may only use products occasionally. For this reason, study designs that enquire about dentine hypersensitivity need to allow adequate investigation of the everyday experience of the condition and factors that may influence this experience.

Coping may play an important role in an individual's dynamic response to health conditions [7, 8]. People may not have consistent styles of coping (e.g. traits) but appear to respond differently in response to specific situations, such as pain [9, 10]. People with dentine hypersensitivity use a number of dentine hypersensitivity-related coping behaviours, which are problem focused (to prevent pain sensations from occurring) such as warming cold food/drinks up before consuming them and the use of toothpaste specially formulated for sensitive teeth [4, 11]. However, there is little research examining how these adaptations are associated with health outcomes in this group of the population. Understanding how well people feel these behaviours help them cope with this oral health condition on a daily basis (e.g. perceived coping efficacy) is also extremely important because positive appraisals of coping are associated with less frequent subsequent pain [12, 13]. Such research would require frequent assessments given the specificity of stimuli and responses.

Another possible predictor of daily pain experience is mood [14, 15]. There is convincing evidence that individual mood and pain experience are intrinsically linked. Low mood both predicts future pain experience [16-18] and may be a consequence of prior pain experience [19]. Clearly this is relevant to a condition with recurring symptoms but to date no research has investigated the relationship between mood, oral health-related coping behaviours and the oral health experiences of dentine hypersensitivity, or indeed any other acutely-experienced oral health condition. Research of this nature, which can expose both within and between-person variation in the oral health experiences related to dentine hypersensitivity, could help clinicians and patients develop a better understanding of the impact and effective management of the condition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

Therefore, to deepen our understanding of how pain experiences are associated with a particular health condition it has been proposed that studies should be prospective and provide more reliable data on how psychological variables and health outcomes may be inter-related and the day-to-day impacts of health conditions [14, 20-22]. Clearly such research requires frequent data collection. Daily diaries offer a unique opportunity to capture day-to-day experiences as they occur thus overcoming the recall bias arising when data are collected retrospectively [14]. Therefore, the current investigation had two aims; firstly, to identify the determinants of the daily experiences of oral health in people with dentine hypersensitivity and secondly, to explore how daily diaries can be used to examine the daily impacts caused by this oral condition. The specific research questions were as follows:

- Do oral health-related coping, perceived coping efficacy and pain predictions successfully predict following day health outcomes in individuals with dentine hypersensitivity?
- Are daily mood and oral health behaviours associated with dentine hypersensitivity-related pain experience?
- What are participants' experiences of using daily diary methodology to examine dentine hypersensitivity?

Method

This prospective daily diary study formed part of a larger research project and longitudinal data obtained from a retrospective questionnaire-based study, which examined how key clinical and psychological factors (e.g. illness beliefs) influence the oral health-related and health-related quality of life outcomes in individuals with dentine hypersensitivity, have been published elsewhere [6].

An advertisement for the study was placed on the intranet page of a large UK University available for staff and students to view. Individuals interested in participating in the study completed a pre-study screening questionnaire and a strict exclusion criteria based on Holland et al.'s guidelines [23] were employed. Only those individuals who experienced dentine hypersensitivity on a frequent basis (a minimum of 'several times a week') were invited to participate in the study. Individuals were excluded from participating if they were pregnant/breastfeeding, taking pain medication, experiencing serious and/or painful health conditions, had experience of a dental professional hygiene visit within the past 14 day (or periodontal surgery within the past six months), were suffering from xerostomia or if they

1 had a recent history of substance misuse. Individuals who did not regularly access routine dental
2 appointments were also excluded. The screening questionnaire therefore aimed to identify and
3 exclude individuals whose sensitivity could have been caused by alternative factors/clinical pathology.

4
5 Participants who met the inclusion criteria were provided with additional study information. Upon
6 obtaining informed consent, participants were asked to complete a text survey at 2pm every day for
7 two weeks. Participants also completed an online daily diary every evening.

8
9 The online diary was used to assess dentine hypersensitivity-related coping behaviours, oral health
10 behaviours, perceived coping efficacy, pain predictions and mood on a daily basis.

11
12 Dentine hypersensitivity-related coping behaviours were assessed using the mean score from the 12
13 item 'adaptation' subscale from the revised Dentine Hypersensitivity Experience Questionnaire [11].
14 An example item is 'Today, when I ate some foods I have made sure they didn't touch certain teeth'
15 (1=strongly disagree to 7=strongly agree). The participant's use of sensitivity toothpaste was assessed
16 using the item: 'Have you used sensitivity toothpaste designed for sensitive teeth today (or do you
17 expect to have by the end of the day?)' (No=1, Yes=2).

18
19 Perceived efficacy of pain coping was assessed on a 5-point scale (1=not at all, 2=slightly,
20 3=somewhat, 4=moderately, 5=very much) in response to the item 'Considering all the things you did
21 or thought today to contend with your pain, how much were you able to alleviate your pain by doing or
22 thinking these things?' [14]. Higher scores reflected greater perceived coping efficacy. Pain frequency
23 prediction was assessed via the online diary by asking respondents 'How often do you expect you will
24 experience these sensations tomorrow?'

25
26 The Short Positive and Negative Affect Scale (PANAS) assessed daily mood [24] on two primary
27 dimensions: positive (PANAS-P, 5 items) and negative affect (PANAS-N, 5 items). Items are rated on
28 a 5-point scale (1=very slightly or not at all to 5=extremely). Total scores for each mood dimension
29 range from 5 to 25. In the current study state mood was assessed by asking participants to rate how
30 they had felt over the course of the day.

31

1 Pain frequency was recorded at two points throughout the day by enquiring 'How many times have
2 you experienced sensations up until 2pm today?' (text survey) and 'How many times have you
3 experienced sensations since 2pm today?' (online diary). At the end of each day the online survey
4 assessed overall pain intensity, bothersomeness and tolerability for that day using three visual
5 analogue scales (scored 1-10) (11). Functional limitations were assessed using mean scores from the
6 4 item 'restrictions' subscale of the DHEQ [11].

7

8 On completion of the study participants were asked to provide feedback on their experiences (see
9 Table 5). Items included 'Has taking part in the study changed your experience of sensitivity?' and
10 'How easy or difficult was it to complete the online daily diaries?'. Participants were paid a small
11 financial incentive for each diary entry submitted.

12

13 Models and analysis

14 Model 1 aimed to examine the relationship between coping, appraisal, pain experience and functional
15 impacts associated with dentine hypersensitivity using lagged analysis (e.g. examining how coping
16 and appraisal from day 1 influenced outcomes on day 2). Model 2 aimed to examine the relationship
17 between mood, coping and pain experience using longitudinal data using a cross sectional method of
18 analysis (mean scores across the 14 days were calculated for each variable). Based on sample size
19 concerns a pre-selection criteria was employed for Model 2 and only significant baseline predictors of
20 pain experience ($p < 0.20$) were entered into the model (pre-selection based on Spearman & Pearson
21 correlations).

22

23 The statistical modelling procedure of Path Analysis was used (AMOS 18.0) to test the extent to which
24 the models fitted the dataset. Bootstrapping was conducted and bias corrected 95% confidence
25 interval (CI) bootstrap percentiles were used to interpret the results. This approach is recommended
26 for sample sizes of less than 20 [25, 26]. Due to the inclusion of a dichotomous variables (use of
27 sensitivity toothpaste: 1=no; 2=yes) in both models, the ADF estimation method was used.

28

29

Results

Two-hundred and eighty individuals expressed an interest in the study, of whom 101 respondents fulfilled the inclusion criteria. Most were female (N=69) and ranged between 18 years and 63 years (mean age 26.3 years, SD=8.6).

Scores for the daily variables are summarised in Table 1. Dentine hypersensitivity-related coping behaviours reported most frequently on day one included 'today, when I ate some foods I have made sure they don't touch certain teeth' and 'today, I had to change the way I drank or ate certain things' (54% and 51%, respectively). Just over a third of participants used sensitivity toothpaste but daily use varied over the two-week daily diary study (range 34% to 43%, average across 14 days=37%). When asked 'considering all the things you did or thought today to contend with your pain how much were you able to alleviate pain by doing or thinking these things?' 18% reported 'not at all', 26% reported 'slightly', 20% reported 'somewhat', 16% reported 'moderately' and 19% reported 'very much' (on day one) revealing a wide variety of responses related to coping efficacy.

The functional limitations reported most frequently on day one of the diary study was 'today, having sensations in my teeth took a lot of pleasure out of eating and drinking' (33%) and the least frequent limitation reported was 'today, there have been times when I couldn't finish my meal because of the sensations' (7%). Mean pain frequency over the two-week diary study was 2.8 (SD=2.3) and the median pain frequency value ranged between 2 and 3 pain sensations per day. However, the number of pain sensations individuals reported ranged from 0-11 (day 1) to 0-43 (day 8) highlighting the between-person variability in dentine hypersensitivity pain over the two-week period. Examination of individual trajectories also highlighted considerable within-person variation in pain frequency over the same period (Figure 1).

1 *Do oral health-related coping, perceived coping efficacy and pain predictions successfully*
2 *predict following day health outcomes in dentine hypersensitivity?*

3

4 *Model 1: Coping strategies, appraisal and following day impacts*

5 The model hypothesised 10 direct pathways based on previous findings from the stress and coping
6 literature. To examine the relationship between oral health coping and outcomes for dentine
7 hypersensitivity, pathways between dentine hypersensitivity-related coping behaviours and the
8 following factors were examined: use of sensitivity toothpaste (path 1); perceived coping efficacy (path
9 2); pain predictions (path 3) and following day functional limitations and pain frequency (paths 4 & 5).
10 Relationships between use of sensitivity toothpaste and pain predictions (path 6) and the primary
11 outcome of pain were also examined (paths 6 & 7). To examine the role cognitive appraisals play in
12 dentine hypersensitivity relationships between perceived coping efficacy and pain predictions (path 8)
13 and between pain predictions and following day pain frequency (path 9) were examined. Finally, the
14 relationship between pain frequency and functional limitations was also investigated to examine
15 whether these health outcomes are inter-related in dentine hypersensitivity sufferers (path 10).

16

17 In order to examine whether pathways identified in Model 1 were 'stable' over the 2-week period,
18 seven identical validation models were examined using 'pairings' of discrete daily diary data (e.g. day
19 1 to 2, days 3 to 4 etc.). Model 1 did not differ significantly from the data across the seven daily
20 pairings ($p > 0.05$), suggesting that the relationships between the key variables remained stable over
21 the study period. Goodness of fit indices were all acceptable (Table 2) and significant pathways for the
22 different days of interest are presented in Table 3. Variables included within the models accounted for
23 between 23% & 66% of the variance in pain frequency and 30% & 54% of the variance for functional
24 limitations.

25

26 Three significant direct pathways and one indirect pathway remained significant across all daily
27 pairings (Table 3). Individuals using more dentine sensitivity-related coping behaviours were more
28 likely to predict that they would experience more painful sensations the following day and were more
29 likely to experience more functional limitations the following day. The prediction of more
30 sensations was also associated with more pain the following day. Significant total effects, which
31 existed when combining both direct and indirect pathways within the model can be seen in Figure 2.

1 *Are daily mood and oral health behaviours associated with dentine hypersensitivity-related*
2 *pain experience?*

3

4 *Model 2: Mood, coping and pain experience*

5 Positive mood was unrelated to the other variables so was not entered into the Path Analysis. The
6 nine direct pathways hypothesised within the model were based on previous findings from the stress
7 and coping literature. To examine the relationship between mood, oral health-related coping and oral
8 health experiences five direct pathways between negative mood and use of sensitive toothpaste (path
9 1), pain frequency (path 2), pain intensity (path 3), pain tolerability (path 4) and pain bothersomeness
10 (path 5) were examined. The relationship between the use of sensitivity toothpaste and pain was also
11 examined (path 6). Finally, to examine how pain experiences of dentine hypersensitivity are inter-
12 related, relationships between pain frequency and pain intensity (pain 7), bothersomeness (path 8)
13 and tolerability (path 9) were all examined.

14

15 Model 2 did not differ significantly from the observed data ($\chi^2=1.95$, $df=3$, $p= 0.58$) and the goodness
16 of fit indices were all excellent (Table 2). The variables included within the model accounted for 12% of
17 variance in pain frequency, 21% of the variance for pain intensity, 13% of the variance for pain
18 tolerability and 18% of the variance in pain bothersomeness.

19

20 The model contained five significant direct pathways and six indirect pathways (Table 4). More
21 negative mood throughout the two-week period was associated with more frequent painful sensations.
22 Frequent sensations were associated with more intense pain, bothersomeness and less tolerability.
23 Individuals who typically used sensitivity toothpaste reported more frequent pain. Significant total
24 effects, which existed when combining both direct and indirect pathways can be seen in Figure 3.

25

26 *Participants' experiences of using daily diary methodology to examine dentine*
27 *hypersensitivity*

28

29 The number of participants who completed daily diaries on each of the study days ranged from 98 to

1 101. Almost half of participants felt that taking part in diary study had changed their experience of
2 sensitivity, largely because it changed their awareness of how frequently they experienced their
3 sensations:

4 :

5 *“It has made me realise that I do not feel sensations as often as I thought”*

6

7 *“I think I notice any sensations in my teeth more, and am probably generally more aware of*
8 *my oral health as a result”*

9

10 Some perceptions and understanding of the oral condition had changed since participating in
11 the study (e.g. understanding the triggers of dentine sensitivity):

12

13 *“I’ve begun to notice that it varies with my mood, I’m not sure how much one influences the*
14 *other!”*

15

16 For some individuals, participating had helped them develop ways of coping with the pain or accepting
17 their condition:

18

19 *“Made me think about the frequency and methods of dealing with it”*

20

21 Most participants felt that receiving the text survey mid-afternoon helped them recall
22 information on their pain experiences and found the daily diary study easy to complete. However, they
23 found it more difficult to complete the following day.

24

25 **Discussion**

26

27 This research revealed how psychological factors may play an important role in the day-to-day oral
28 experiences of people with dentine hypersensitivity. These factors, coping strategies and experiences,
29 fluctuated within individuals throughout the study period and thus the daily diary approach allowed
30 sufficient precision to detect associations between these fluctuations.

31

1 Previous diary research has found mixed results in relation to the use of coping and subsequent pain
2 levels. For example, the use of cognitive reframing strategies in the morning predicted less evening
3 pain experience, whereas morning use of active problem solving predicted more evening pain [27].
4 Within the current study, the use of dentine hypersensitivity-related coping behaviours predicted worse
5 pain and functional limitations the following day. However, there was some evidence that greater use
6 of specific coping adaptations were positively associated with higher levels of perceived coping
7 efficacy. Therefore, an explanation for the relationship between coping and following day pain
8 experience could be that those individuals who felt the need to use dentine sensitivity related-coping
9 behaviours were already experiencing more pain and found the coping adaptations did help them
10 control this pain. This highlights the importance of examining individuals' coping appraisals as a
11 mechanism for aiding our understanding into the complex, and probable bi-directional, nature of the
12 relationship between coping and pain.

13

14 Pain predictions were strongly associated with subsequent pain experience, indicating that despite its
15 fluctuating nature, pain is predictable in dentine hypersensitivity. The mechanisms through which pain
16 predictions influence subsequent pain experience warrants further examination [28]. Individuals with
17 worse negative daily mood were also more likely to report more pain, of a higher intensity, less
18 tolerability and greater bothersomeness. This is the first research to highlight the role of mood in
19 individuals' experiences of this specific type of dental pain. These findings are compatible with
20 previous research which has found similar effects in relation to chronic pain [17, 18]. Whilst the
21 causality of the relationship between mood and pain is subject to debate, state negative affect can
22 increase symptom perception and reporting [29, 30].

23

24 The explained variance in pain frequency and functional limitations differed considerably across the
25 two-week period (23%-66% and 30%-54%, respectively). This variation in the explanatory power of
26 coping and appraisal factors on dentine hypersensitivity experiences provides evidence that the
27 factors and processes that influence these daily oral health experiences may also fluctuate on a daily
28 basis. It is also important to recognise that there were only small to moderate associations between
29 pain frequency and the other subjective elements of pain experience reported (intensity,
30 bothersomeness and intolerability). This suggests that only measuring the frequency with which

1 someone experiences pain may fail to provide a reliable and holistic account of the individual's pain
2 experience.

3

4 A major strength of the current study was its prospective design of the research with frequent data
5 collection, which is less likely to be subject to recall error or bias. This success was reflected in the
6 high response rate throughout the daily data collection and the positive feedback from participants
7 regarding different aspects of the study's design which is consistent with findings from studies using
8 similar methods [31, 32]. However, participants' response times became increasingly delayed during
9 the study, supporting previous research describing the cumulative burden of completing daily diaries
10 and decreased reliability of responses over time [31]. The data revealed that most individuals felt the
11 text surveys had acted as a useful aid to their pain recall throughout the day, highlighting the
12 usefulness of signal-contingent methods in daily diaries [33]. Incentives were used in the current study
13 and this could have also contributed to the high response rate. The use of incentives are associated
14 with higher levels of engagement levels in research studies which employ diary methods [34].

15

16 The potential value of utilising e-diaries (e.g. via a phone app) as a clinical assessment tool could also
17 be examined in future research. Prospective diary methods could be used to collect accurate data on
18 patient's dentine hypersensitivity-related pain and coping, this information could aid the patient's and
19 clinician's understanding of the patient's own unique experience of the condition and the possible
20 factors that may help them effectively manage their symptoms. However, further evaluation of the
21 feasibility and utility of using this type of assessment tool in a clinical setting is required.

22

23 Whilst these data start to explain the experience of dentine hypersensitivity, the research was not
24 without limitations. Individuals were university students and staff and may not represent the general
25 population and this could affect the generalisability of the findings. For example, the high response
26 rate and level of engagement with the daily diaries may not be replicated in studies that use different
27 groups of the population. The methods employed (e.g. internet diaries and text surveys) also excluded
28 individuals who did not have daily access to the internet and a mobile phone, which could have further
29 biased the sample.

30

1 It is important to recognise that participants self-diagnosed their dentine hypersensitivity and no
2 clinical examinations were undertaken to identify alternative clinical pathologies which could have
3 been responsible for pain experiences. Oral conditions, which have symptoms similar to dental
4 hypersensitivity, should be excluded to ensure an accurate diagnosis of dentine hypersensitivity can
5 be made [23]. Within the current study it is possible that some individuals could have been
6 experiencing oral pain caused by a different pathology. However, a pre-study screening questionnaire
7 and strict exclusion criteria based on Holland et al.'s guidelines [23] were employed. Irregular dental
8 attendance was one of the exclusion criteria employed to minimise the likelihood that individuals with
9 untreated dental disease/conditions (e.g. cavities or trauma) could participate in the study. It is also
10 possible that participants self-monitoring effects could have influenced the data obtained. Self-
11 monitoring is an intervention in its own right and several psychological models highlight the impact
12 selective attention to bodily symptoms can have on the experiencing of symptoms [35, 36]. Therefore,
13 the issues associated with self-monitoring should be recognised by researchers and clinicians who
14 employ daily diary methods.

15

Conclusions

1
2
3
4
5
6
7
8
9
10

These data offer unique insights into the experiences of people with dentine hypersensitivity. The study revealed how dentine hypersensitivity experiences fluctuate on a daily basis, which has significant implications for the how these oral health experiences can be measured in a meaningful and reliable way. The use of prospective methods of data collection, such as daily diaries, can develop understanding of the day-to-day impacts caused by oral conditions and the complex relationships that exist between transient states and experiences such as psychological factors and dental pain. This information can place clinicians in a stronger position to manage their patients' oral health conditions.

List of abbreviations

- ADF: Asymptotically Distribution-Free
- DHEQ: Dentine Hypersensitivity Experience Questionnaire
- PANAS: Positive and Negative Affect Scale

Declarations

Ethics approval and consent to participate: Informed written consent was obtained from participants prior to their involvement in the research. Ethical approval for the study was obtained from the University of Sheffield Research Ethics Committee.

Availability of data and materials: The dataset analysed during the current study is available from the corresponding author on reasonable request.

Funding: This research was funded by GlaxoSmithKline Consumer Healthcare (grant number: R/125710-1).

Competing interests: FS is employed by GlaxoSmithKline Consumer Healthcare. JP and SB declare they have no competing interests.

Authors' contributions: SB, JP and FS contributed to the conception and design of the study. JP was responsible for the collection, analysis and interpretation of data. JP and SB were involved in drafting and revising the manuscript. All authors have given approval for the manuscript to be published.

Consent for publication: Not applicable.

Acknowledgements: The authors would like to thank the individuals who shared their time to participate in this study and Professor Peter Robinson for his guidance and feedback that greatly improved the manuscript.

References

1. Cummins D. Dentin Hypersensitivity: From Diagnosis to a Breakthrough Therapy for Everyday Sensitivity Relief. *Journal of Clinical Dentistry*. 2009;1-9.
2. Docimo R, Montesani L, Maturo P, DeVizio W, Zhang YP, Cummins D, Dibart S, Mateo LR. Comparing the Efficacy in Reducing Dentin Hypersensitivity of a New Toothpaste Containing 8/0% Arginine, Calcium Carbonate, and 1450ppm Fluoride to a Commercial Sensitive Toothpaste Containing 2% Potassium Ion: An Eight-Week Clinical Study in Rome, Italy. *Journal of Clinical Dentistry*. 2009;17-22.
3. Gibson B, Boiko OV, Baker SR, Robinson PG, Barlow APS, Player T, Locker D. The everyday impact of dentine sensitivity: personal and functional aspects. *Social Science and Dentistry*. 2010; 1:11-21.
4. Gillam DG, Seo HS, Bulman JS, Newman HN. Perceptions of dentine hypersensitivity in a general practice population. *Journal of Oral Rehabilitation*. 1999; 26:710-714.
5. Irwin CR, McCusker P. Prevalence of dentine hypersensitivity in a general dental population. *Journal of Irish Dental Association*. 1997; 43:7-9.
6. Porritt JM, Sufi F, Barlow A, Baker SR. The role of illness beliefs and coping in the adjustment to dentine hypersensitivity. *Journal of Clinical Periodontology*. 2014; 41:60-69.
7. Leventhal H, Diefenbah MA, Leventhal EA. Illness Cognition: Using common sense to understand treatment adherence and affect cognition interactions. *Cognitive Therapy and Research*. 1992; 16:143-163.
8. Leventhal H. Findings and theory in the study of fear communications. *Advances in Experimental Social Psychology*. 1970; 5:119-186.
9. Brown GK, Nicassio PM. Development of a questionnaire for the assessment of active and passive coping strategies in chronic pain patients. *Pain*. 1987; 31:53-64.
10. Norman SA, Thompson R, Vedhara K. A prospective examination of illness beliefs and coping in patients with type 2 diabetes. *British Journal of Health Psychology*. 2007; 12:621-638.
11. Boiko OV, Baker SR, Gibson BJ, Locker D, Sufi F, Barlow APS, Robinson PG. Construction and validation of the quality of life measure for dentine hypersensitivity (DHEQ). *Journal of Clinical Periodontology*. 2010; 37:973-980.
12. Hagger MS, Orbell S. A Meta-analytic review of the common-sense model of illness representations. *Psychology and Health*. 2003; 18:141-184.
13. Evans DD, Carter M, Panico R, Kimble L, Morlock JT, Spears MJ. Characteristics and predictors of short-term outcomes in individuals self-selecting yoga or physical therapy for treatment of chronic low back pain. *American Academy of Physical Medicine and Rehabilitation*. 2010; 2:1006-1015.
14. Tennen H, Affleck G, Zautra A. Depression History and Coping With Chronic Pain: A Daily Process Analysis. *Health Psychology*. 2006; 25:370-379.
15. Williamson GM, Schultz R. Physical illness and symptoms of depression among elderly patients. *Psychology and Aging*. 1992; 7:343-351.
16. Zautra AJ, Marbach JJ, Raphael KG, Dohenwend BP, Lennon MC, Kenny DA. The examination of myofascial face pain and its relationship to psychological distress. *Health Psychology*. 1995; 14:223-231.
17. Feldman SI, Downey G, Schaffer-Neitz R. Pain, Negative Mood, and Perceived Support in Chronic Pain Patients: A Daily Diary Study of People With Reflex Sympathetic Dystrophy Syndrome. *Journal of Consulting and Clinical Psychology*. 1999; 67:776-785.

- 1 18. Tang NK, Salkovski PM, Hodges A, Wright KJ, Hanna M, Hester J. Effects of
2 mood on pain responses and pain tolerance: an experimental study in chronic
3 back pain patients. *Pain*. 2008; 138:392-401.
- 4 19. Glover J, Dibble SL, Dodd MJ, Miaskowski C. Mood states of oncology
5 outpatients: Does pain make a difference? *Journal of Pain and Symptom*
6 *Management*. 1995; 10:120-128.
- 7 20. Galli U, Ettlin DA, Palla S, Ehlert U, Gaab J. Do illness perceptions predict
8 pain-related disability and mood in chronic orofacial patients? A 6-month
9 follow-up study. *European Journal of Pain*. 2010; 14:550-558.
- 10 21. Hobro N, Weinman J, Hankins M. Using the self-regulatory model to cluster
11 chronic pain patients: the first step towards identifying relevant treatments?
12 *Pain*. 2004; 108:276-283.
- 13 22. Tennen H, Affleck G. Daily processes in coping with chronic pain: methods
14 and analytic strategies. In: *Handbook of coping*. edn. Edited by Zeidner M,
15 Endler N. New York: Wiley; 1996: 151-180.
- 16 23. Holland GR, Narhi MN, Addy M, Gangarosa L, Orchardson R. Guidelines for
17 the design and conduct of clinical trials on dentine hypersensitivity. *Journal of*
18 *clinical periodontology*. 1997; 24(11):808-813.
- 19 24. Mackinnon A, Jorm AF, Christensen H, Korten AE, Jacomb PA, Rodgers B. A
20 short form of the Positive and Negative Affect Schedule: evaluation of factorial
21 validity and invariance across demographic variables in a community sample.
22 *Personality and Individual Differences*. 1999; 27:405-416.
- 23 25. Efron B, Tibshirani R. An introduction to the bootstrap. New York: Chapman &
24 Hall; 1993.
- 25 26. Hoyle RH, Panter AT. Writing about structural equation models. In: *Structural*
26 *equation modelling*. edn. Edited by RH R. Thousand Oaks, CA: Sage; 1995:
27 158-176.
- 28 27. Holtman S, Newth S, DeLongis A. The Role of Social Support in Coping with
29 Daily Pain among Patients with Rheumatoid Arthritis. *Journal of Health*
30 *Psychology*. 2004; 9:677-695.
- 31 28. Finan PH, Zautra AJ, Tennen H. Daily diaries reveal influence of pessimism
32 and anxiety on pain prediction patterns. *Psychology and Health*. 2008; 23:551-
33 568.
- 34 29. Charles ST, Almeida DM. Daily reports of symptoms and negative affect: Not
35 all symptoms are the same. *Psychology and Health*. 2006; 21:1-17.
- 36 30. Watson D, Pennebaker JW. Health complaints, stress, and distress: Exploring
37 the central role of negative affectivity. *Psychological Review*. 1989; 96:234-
38 254.
- 39 31. Lyons KE, Pahwa R. Electronic motor function diary for patients with
40 Parkinson's disease: A feasibility study. *Parkinsonism and Related Disorders*.
41 2007; 13:304-307.
- 42 32. Quinn P, Goka J, Richardson H. Assessment of an electronic daily diary in
43 patients with overactive bladder. *BJU Int*. 2003; 91:647-652.
- 44 33. Affleck G, Tennen H, Armeli S, Zautra A. Multilevel Daily Process Designs for
45 Consulting and Clinical Psychology: A Preface for the Perplexed. *Journal of*
46 *Consulting and Clinical Psychology*. 1999; 67:746-754.
- 47 34. Lynn P. The Impact of Incentives on Response Rates to Personal Interview
48 Surveys: Role and Perceptions of Interviewers. *International Journal of Public*
49 *Opinion Research*. 2001; 13(3):326-336.
- 50 35. Goodwin L, Fairclough SH, Poole HM. A cognitive-perceptual model of
51 symptom perception in males and females: the roles of negative affect,
52 selective attention, health anxiety and psychological job demands. *J Health*
53 *Psychol*. 2013; 18(6):848-857.
- 54 36. Kolk AM, Hanewald GJFP, Schagen S, Gijsbers van Wijk CMT. A symptom
55 perception approach to common physical symptoms. *Social Science &*
56 *Medicine*. 2003; 57(12):2343-2354.
- 57

Figure Legends

1

2

3 **Figure 1.** Random sample (10%) of individuals' pain frequency trajectories

4

5 **Figure 2.** Significant total pathways between oral health coping behaviours, appraisal and following
6 day impacts proposed within Model 1

7

8 **Figure 3.** Significant total pathways between mood, oral health coping behaviours and pain
9 experience in individuals with dentine hypersensitivity (Model 2)

Table 1. Daily mean scores

Day	Positive mood mean (SD)	Negative mood mean (SD)	Coping mean (SD)	Efficacy mean (SD)	Pain frequency prediction mean (SD)	Pain frequency mean (SD)	Pain intensity mean (SD)	Pain bothersomeness mean (SD)	Pain (in)tolerability mean (SD)	Functional limitations mean (SD)
1	15.1 (4.3)	8.4 (4.0)	2.8 (1.0)	2.9 (1.4)	2.8 (2.9)	3.0 (3.0)	3.9 (1.7)	3.4 (1.8)	2.7 (1.7)	2.7 (1.3)
2	14.6 (4.9)	8.3 (4.1)	2.8 (1.3)	2.9 (1.4)	2.5 (1.9)	2.9 (3.2)	3.5 (2.0)	3.3 (2.2)	3.0 (2.1)	2.8 (1.6)
3	14.4 (5.1)	8.3 (4.1)	2.8 (1.2)	2.9 (1.4)	2.3 (1.9)	2.7 (3.5)	3.4 (2.0)	3.1 (2.1)	2.9 (2.1)	2.8 (1.6)
4	14.0 (5.0)	8.8 (4.5)	2.8 (1.2)	3.0 (1.4)	2.3 (1.8)	2.6 (2.6)	3.3 (1.7)	3.0 (1.7)	2.7 (1.7)	2.8 (1.2)
5	13.8 (5.0)	7.4 (3.4)	2.9 (1.5)	3.0 (1.5)	2.3 (1.8)	2.8 (2.6)	3.6 (2.0)	3.3 (2.1)	3.1 (2.0)	3.0 (1.6)
6	14.0 (5.3)	7.1 (3.0)	2.9 (1.2)	3.0 (1.3)	2.3 (1.8)	2.9 (3.7)	3.3 (1.8)	3.1 (1.8)	2.9 (1.7)	3.0 (1.5)
7	14.0 (5.3)	7.8 (3.5)	2.9 (1.2)	3.0 (1.4)	2.3 (1.9)	2.7 (2.7)	3.3 (1.8)	3.0 (1.9)	2.7 (1.7)	3.1 (1.6)
8	14.3 (4.8)	8.0 (4.1)	3.0 (1.4)	3.0 (1.4)	2.5 (2.4)	3.2 (5.0)	3.5 (2.0)	3.1 (2.0)	2.9 (1.9)	3.1 (1.7)
9	14.1 (5.0)	8.0 (4.3)	3.0 (1.4)	2.8 (1.3)	2.4 (1.9)	2.6 (2.3)	3.3 (2.0)	3.3 (2.1)	3.0 (2.0)	3.1 (1.6)
10	14.2 (5.1)	7.7 (4.0)	3.0 (1.4)	2.8 (1.4)	2.3 (1.8)	2.4 (2.4)	3.3 (1.8)	3.0 (1.8)	2.9 (1.8)	2.9 (1.6)
11	15.0 (5.1)	7.4 (3.0)	3.0 (1.4)	3.1 (1.3)	2.4 (2.0)	3.0 (2.6)	3.2 (1.7)	3.0 (1.8)	3.0 (1.9)	2.9 (1.6)
12	15.0 (5.1)	6.9 (3.3)	2.9 (1.4)	2.9 (1.3)	2.2 (1.8)	2.5 (2.7)	3.1 (1.9)	2.9 (1.9)	2.7 (1.4)	2.7 (1.5)
13	13.1 (4.8)	7.3 (3.5)	3.1 (1.3)	2.9 (1.3)	2.4 (1.9)	2.7 (2.9)	3.1 (1.7)	2.8 (1.6)	2.7 (1.6)	2.9 (1.5)
14	13.4 (5.1)	7.7 (4.0)	3.1 (1.3)	2.8 (1.3)	2.4 (1.9)	3.1 (3.2)	3.3 (1.9)	2.9 (1.8)	2.7 (1.9)	3.0 (1.6)
14 day Mean	14.2 (3.3)	7.8 (2.5)	2.9 (1.1)	2.9 (1.1)	2.4 (1.7)	2.8 (2.3)	3.3 (1.3)	3.1 (1.4)	2.8 (1.4)	2.9 (1.1)

Table 2. Goodness of fit indices for Model 1 and Model 2

Model 1 (N=101)	Goodness of fit indices					
	χ^2 / df ($p > .05$)	CMIN/df (< 2.0)	IFI (> 0.95)	RMSEA (< 0.08)	SRMR (< 0.08)	Criteria fitted
Day 1-2	3.93/5 (p=0.56)	0.79	1.02	0.00	0.07	5
Day 3-4	2.11/5 (p=0.83)	0.42	1.09	0.00	0.03	5
Day5-6	3.16/5 (p=0.68)	0.63	1.04	0.00	0.03	5
Day7-8	4.57/5 (p=0.47)	0.91	1.01	0.00	0.04	5
Day9-10	2.99/5 (p=0.70)	0.60	1.02	0.00	0.03	5
Day 11-12	6.33/5 (p=0.28)	1.26	0.99	0.05	0.06	5
Day13-14	4.09/5 (p=0.54)	0.82	1.02	0.00	0.05	5
Model 2 (N=101)	1.95/3 (p=0.58)	0.65	1.00	0.00	0.03	5

Note: Figures in bold are those which meet the model-fitting criteria.

Table 3. Significant pathways between oral health coping, appraisal and following day impacts (Model 1)

Daily Models (N=101)	Significant pathways	Total β value (bootstrap bias corrected 95% CI)	Indirect β value (bootstrap bias corrected 95% CI)	Direct β value (bootstrap bias corrected 95% CI)
Days 1-2	D1 coping [^] → D1 perceived efficacy	.27** (.06 to .45)	--	.27** (.06 to .45)
	D1 coping → D1 pain prediction	.27** (.07 to .70)	n/s	.30* (.06 to .72)
	D1 coping → D2 pain frequency	.45** (.24 to .65)	.18* (.05 to .84)	.27* (.03 to .41)
	D1 coping → D2 functional	.53** (.32 to .67)	.18* (.05 to .35)	.35* (.16 to .54)
	D1 pain prediction → D2 pain frequency	.68** (.46 to .83)	--	.68** (.46 to .83)
Days 3-4	D3 coping → D3 pain prediction	.31** (.10 to .50)	--	.31** (.10 to .50)
	D3 coping → D4 pain frequency	.23* (.03 to .41)	.20** (.07 to .34)	n/s
	D3 coping → D4 functional	.32** (.08 to .50)	.10* (.02 to .20)	.22* (.02 to .38)
	D3 pain prediction → D4 pain frequency	.61** (.35 to .76)	--	.61** (.35 to .76)
	D3 pain prediction → D4 functional	.28** (.11 to .44)	.28** (.11 to .44)	-
Days 5-6	D4 pain frequency → D4 functional	.46** (.24 to .62)	--	.46** (.24 to .62)
	D5 coping → D5 pain prediction	.34** (.15 to .50)	n/s	.33** (.13 to .50)
	D5 coping → D6 pain frequency	--	.22** (.06 to .35)	n/s
	D5 coping → D6 functional	.38** (.17 to .54)	n/s	.31** (.14 to .46)
	D5 pain prediction → D6 pain frequency	.63* (.21 to .74)	--	.63** (.21 to .74)
Days 7-8	D5 pain predictions → D6 functional	.35* (.23 to .43)	.35* (.23 to .43*)	--
	D6 pain frequency → D6 functional	.56** (.41 to .65)	--	.56** (.41 to .65)
	D7 coping → D7 pain prediction	.30** (.13 to .46)	--	.29** (.11 to .44)
	D7 coping → D8 functional	.45** (.24 to .60)	--	.42** (.25 to .56)
	D7 coping → pain frequency	--	.15* (.03 to .26)	n/s
Days 9-10	D7 pain prediction → D8 pain frequency	.46* (.33 to .67)	--	.46* (.33 to .67)
	D7 pain predictions → D8 functional	.16* (.06 to .23)	.16* (.06 to .23)	--
	D8 pain frequency → D8 functional	.34** (.15 to .46)	--	.34** (.15 to .46)
	D9 coping → D9 pain prediction	.37** (.18 to .53)	n/s	.34** (.14 to .51)
	D9 coping → D10 pain frequency	.26** (.07 to .42)	.24** (.09 to .38)	n/s
Days 11-12	D9 coping → D10 functional	.40** (.40 to .71)	.06** (.02 to .15)	.52** (.33 to .66)
	D9 pain prediction → D10 pain frequency	.69* (.40 to .80)	--	.69* (.40 to .80)
	D9 pain predictions → D10 functional	.17** (.05 to .28)	.17** (.05 to .28)	--
	D10 pain frequency → D10 functional	.25** (.09 to .41)	--	.25** (.09 to .41)
	D11 coping → D11 pain prediction	.41** (.22 to .56)	n/s	.35** (.18 to .51)
Days 13-14	D11 coping → D12 pain frequency	.36** (.18 to .53)	.31** (.18 to .44)	n/s
	D11 coping → D12 functional	.68** (.42 to .82)	.11* (.02 to .19)	.57** (.31 to .77)
	D11 use of toothpaste → D12 pain frequency	.29** (.09 to .47)	n/s	.19* (.02 to .37)
	D11 use of toothpaste → D12 functional	.09* (.02 to .20)	.09* (.02 to .20)	--
	D11 pain prediction → D12 pain frequency	.62* (.41 to .80)	n/s	.64* (.41 to .80)
	D12 pain frequency → D12 functional	.31* (.01 to .50)	n/s	.31* (.01 to .50)
	D13 coping → D13 pain prediction	.41** (.24 to .57)	n/s	.37** (.18 to .55)
Days 13-14	D13 coping → D14 pain frequency	.40** (.22 to .61)	.28** (.17 to .44)	n/s
	D13 coping → D14 functional	.61** (.40 to .76)	.14* (.00 to .27)	.47** (.24 to .72)
	D13 pain prediction → D14 pain frequency	.69** (.56 to .83)	--	.69** (.56 to .83)

[^]dentine hypersensitivity-related coping behaviours * $p < 0.05$, ** $p < 0.01$

Table 4. Significant direct, indirect and total pathways between mood, toothpaste use and pain experience proposed within Model 2

Significant pathways	Direct pathways		Indirect pathways		Total pathways	
	β value	Bootstrap bias corrected 95% CI	β value	Bootstrap bias corrected 95% CI	β value	Bootstrap bias corrected 95% CI
Negative mood → Use of sensitivity toothpaste	-.08	-.29 to .18	--	--	-.08	-.29 to .18
Negative mood → Pain frequency	.24*	.05 to .42	-.02	-.09 to .05	.23*	.02 to .43
Negative mood → Pain intensity	.12	-.06 to .37	.09*	.02 to .20	.21*	.02 to .54
Negative mood → Pain intolerability	.17	-.07 to .45	.06*	.01 to .16	.23	-.02 to .51
Negative mood → Pain bothersomeness	.18	-.00 to .42	.08*	.01 to .17	.26*	.06 to .50
Use of sensitivity toothpaste → Pain frequency	.26*	.04 to .47	--	--	.26*	.04 to .47
Use of sensitivity toothpaste → Pain intensity	--	--	.11**	.02 to .25	.11**	.02 to .25
Use of sensitivity toothpaste → Pain intolerability	--	--	.07**	.01 to .18	.07**	.01 to .18
Use of sensitivity toothpaste → Pain bothersomeness	--	--	.09**	.02 to .20	.09**	.02 to .20
Pain frequency → Pain intensity	.41**	.22 to .58	--	--	.41**	.22 to .58
Pain frequency → Pain intolerability	.28*	.07 to .45	--	--	.28*	.07 to .45
Pain frequency → Pain bothersomeness	.35**	.18 to .51	--	--	.35**	.18 to .51

* $p < 0.05$, ** $p < 0.01$

Table 5. Participant methodology feedback

Feedback item within follow-up questionnaire	Number of people (N=99)
Has taking part in the study changed your experience of sensitivity? Yes Don't know No	43 20 36
How easy or difficult was it to complete the daily text survey? Very difficult Difficult Neither easy not difficult Easy Very easy	0 1 1 11 86
Did completing the text survey mid afternoon help you remember how many times you had experienced sensitive teeth that day? Yes Don't know No	83 7 9
How easy or difficult was it to complete the onl Very difficult Difficult Neither easy not difficult Easy Very easy	0 5 8 34 52
If, on occasions, you completed the daily diary in the evening how easy or difficult was it to remember the experiences related to your sensitive teeth that day? Very difficult Difficult Neither easy not difficult Easy Very Easy	0 4 13 44 38
If, on occasions, you completed the daily diary following day but before 10am how easy or difficult was it to remember the experiences related to your sensitive teeth the previous day? Very difficult Difficult Neither easy not difficult Easy Very Easy Not applicable	2 10 15 37 15 20
If, on occasions, you completed the daily diary after 10am the following day how easy or difficult was it to remember the experiences related to your sensitive teeth the previous day? Very difficult Difficult Neither easy not difficult Easy Very Easy Not applicable	5 7 17 18 9 43
How did you feel about the duration of the study? I felt the duration of the study was about right I felt two-weeks was too long a period of time to be completing diary entries each day I would have been willing to complete more daily diaries over a longer period of time	46 3 50