

Measuring wearing time of knee-ankle-foot orthoses in children with cerebral palsy: comparison of parent-report and objective measurement

MAAS, Jose, DALLMEIJER, Annet J., OUDSHOORN, Bodil, BOLSTER, Eline A. M., HUIJING, Peter A., JASPERS, Richard T. and BECHER, Jules G.

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/14122/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

MAAS, Jose, DALLMEIJER, Annet J., OUDSHOORN, Bodil, BOLSTER, Eline A. M., HUIJING, Peter A., JASPERS, Richard T. and BECHER, Jules G. (2016). Measuring wearing time of knee-ankle-foot orthoses in children with cerebral palsy: comparison of parent-report and objective measurement. *Disability and Rehabilitation*. (In Press)

Repository use policy

Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in SHURA to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

1 **Observational study**

2

3 **Measuring wearing time of knee-ankle-foot orthoses in children with**
4 **cerebral palsy: Comparison of parent-report and objective measurement.**

5

6 **Josina C Maas^{1*}, Annet J Dallmeijer¹, Bodil Y Oudshoorn¹, Eline AM Bolster¹, Peter A Huijting²,**
7 **Richard T Jaspers² and Jules G Becher¹**

¹Department of Rehabilitation Medicine, MOVE Research Institute Amsterdam and the EMGO+ Institute for Health and Care Research, VU University Medical Center, Amsterdam, the Netherlands

²Laboratory for Myology, MOVE Research Institute Amsterdam, Faculty of Human Movement Sciences, VU University, Amsterdam, the Netherlands

8 *Corresponding author: De Boelelaan 1117,1081 HV Amsterdam, the Netherlands;

9 Email : jc.maas@vumc.nl; Phone: +31 204443063; Fax: +31 20 4440787

10

11

12

13

14

15

16

17

18

19

20

21 **Keywords**

22 Orthotic devices, Self-report, Parent-report, Questionnaires, Wearing time, Cerebral palsy,

1 **Abstract**

2 Background Orthotic wearing time may be an important confounder in efficacy studies of treatment in
3 children with spastic cerebral palsy (SCP). Most studies measure parent-reported wearing time with
4 questionnaires, but it is questionable whether this yields valid results. This study aims to compare
5 parent-reported wearing time (WT_{parent}) with objectively measured wearing time (WT_{obj}) in children
6 with SCP receiving orthotic treatment.

7 Methods Eight children with SCP participated in this observational study. For one year, they received
8 knee-ankle-foot orthosis (KAFO) treatment. WT_{parent} was measured using questionnaires. WT_{obj} was
9 measured using temperature sensor-data-loggers that were attached to the KAFOs. The 2.5th and 97.5th
10 percentiles and median of differences between methods (per participant) were used to calculate limits
11 of agreement and systematic differences.

12 Results There was no systematic difference between WT_{parent} and WT_{obj} (0.1 hours per week), but high
13 inter-individual variation of the difference was found, as reflected by large limits of agreement (lower
14 limit/2.5th percentile: -1.7. hours/week; upper limit/97.5th percentile: 11.1 hours/week).

15 Conclusions Parent-reported wearing time of a knee-ankle-foot orthosis differs largely from
16 objectively measured wearing time using temperature sensors. Therefore, parent-reported wearing
17 time of KAFOs should be interpreted with utmost care.

18 Clinical relevance Low wearing time of orthoses may be a cause of inefficacy of orthotic treatment
19 and incorrect reported wearing time may bias results of efficacy studies. Results of this study show
20 that parent-reported wearing time is not in agreement with objectively measured wearing time,
21 Therefore, objective methods are recommended for measuring orthotic wearing time.

22

1 **Introduction**

2 According to the International Society for Prosthetics and Orthotics consensus document^{1;2}, orthotic
3 treatment is applied in efforts to prevent deformities and muscle shortening in children with spastic
4 cerebral palsy (SCP). Applying a high strain onto muscles by stretching the muscle tendon complex
5 using orthoses is presumed to increase joint range of motion or prevent its loss². Wearing time is an
6 important factor in determining effectiveness, as efficacy of treatment with orthoses is suggested to be
7 dependent on duration that muscles are strained³. Two recent studies⁴⁻⁶, investigating the efficacy of
8 ankle-foot orthoses (AFO) and knee-ankle-foot orthoses (KAFO) straining the calf muscles, used
9 questionnaires filled in by the participant's parents to assess wearing time. However, it is questionable
10 whether such parent-reported wearing time is a valid measure.

11 Research focusing on objective methods to measure wearing time has been reported for
12 patients with scoliosis, wearing spine orthoses. Several objective methods are available, such as data
13 acquisition using temperature sensors⁷⁻¹⁰ or using force transducers^{9;11} attached to the orthoses. These
14 methods yielded reliable and valid wearing time estimates⁷⁻¹¹. Temperature sensors have been
15 mentioned as most feasible with good agreement with actual wearing time, with no reported
16 mechanical problems and the sensors being easy to use⁹.

17 Objective measures have been used for comparison of self-reported wearing time and actual
18 wearing time (further referred to as objectively measured wearing time). Previous studies assessing the
19 reliability and validity of self-reported wearing time measurements in patient with spine orthoses
20 showed that, in general, self-reports by patients or parents led to overestimation^{8;12-16}. In particular
21 cases, patient-reported wearing time was twice the objectively measured wearing time^{14;16}. Also
22 children with clubfoot deformities, who were wearing a foot abduction orthosis, patient-reported
23 wearing time (by parents) was overestimated¹⁷. In efficacy studies, such overestimation may lead to
24 erroneous conclusions if orthoses were not worn long enough to be effective.

25 To the best of our knowledge, in children with SCP who wear KAFOs or other orthoses to
26 prevent loss of ankle-foot range of motion, parent-reported wearing time has not been compared to
27 objectively measured wearing time. Because wearing instructions, aim of treatment and design of
28 orthoses are different from patients wearing spine orthoses, validity of parent-reported orthotic
29 wearing time may vary from the above described studies. Therefore, we aim to compare parent-
30 reported wearing time in children with SCP, collected using questionnaires, and objective wearing
31 time, measured using temperature sensor-data-loggers attached to the KAFO. It is hypothesised that
32 parent-reported wearing time differs from objective measurements.

1 **Methods**

2 This observational study is part of a randomized controlled trial that tested the efficacy of KAFO
3 treatment. Approval for this study was obtained from the Medical Ethics Committee of VU University
4 Medical Center^{4,5}. All participants (if older than 12 years) and their parents signed an informed
5 consent form.

6 **Participants and study design**

7 All children participating in this study were also participants in the experimental group of a multi-
8 center study investigating the efficacy of KAFOs to prevent equinus in children with SCP (Splint
9 study^{4,5}). In the Splint study, 30 children were recruited and 17 of them were randomized to the KAFO
10 group. They were recruited between February 2010 to April 2012. The KAFOs of children who were
11 recruited in Amsterdam or Venlo (8 children, allocation was determined by the child's residence) were
12 equipped with a temperature sensor-data-logger to measure wearing time (Figure 1). These 8 children
13 were included in the present study. Although the sensor-data-logger was visible for the participants,
14 they were not informed about the actual aim of the device. Besides that wearing time was measured
15 objectively, parents of participants were asked to report KAFO wearing time for their child. The
16 participants were aged between 4 and 16 years and were able to walk independently (gross motor
17 function classification system (GMFCS) level of I-II) or with walking aids (GMFCS level III)¹⁸. All
18 participants were instructed to wear KAFOs for at least 6 hours every other night.

19

20

<<Figure 1>>

21

22 **Measurements**

23 Parent-reported wearing time (WT_{parent}) in hours/week, was collected using questionnaires. For all
24 days of the 4th week of each month during the treatment period, parents of participants were asked to
25 complete online questionnaires. These questionnaires were sent to the parents by email
26 (Netquestionnaire Nederland B.V., Utrecht, The Netherlands). A reminder was sent after two weeks if
27 parents did not respond. The following multiple choice question was asked: 'How many hours did
28 your child wear his/her KAFO?'. Answers were given as seven categories, ranging from 0-1 hour to 6
29 hours or more.

30 Objectively measured wearing time (WT_{obj}) in hours/week, was measured with a temperature sensor-
31 data-logger (UTBI-001, Onset Computer Corporation, Bourne, MA, USA) during the treatment
32 period. The sensor-data-logger was embedded in the shelf of the KAFO, touching upper-leg skin of
33 the participant. Temperature was recorded every 15 minutes. KAFO wearing time was derived from
34 time-temperature data. An increase of at least 3°C in temperature within 45 minutes, indicated that the

1 KAFO was put on. As KAFO temperature is not expected to increase over normal body temperature
2 (37°), periods with higher temperatures were ignored, because it was assumed that these temperature
3 increases were artefacts caused by external heating (e.g. KAFO exposed to direct sunlight or heater
4 radiation). The KAFO was assumed to be removed if temperature decreased to values similar to
5 previous periods that KAFO was not worn.

6 Analyses

7 For each participant, WT_{parent} and WT_{obj} per week were calculated for all weeks that data of
8 both methods were available. A week was only analysed if WT_{parent} and WT_{obj} measurements were
9 available for at least 4 days. If only 4-6 days of a week were available, wearing time of those days was
10 used to estimate total wearing time of 7 days. For WT_{parent} , the centre value of each time category (i.e.
11 0.5 for 0-1 hours, 1.5 for 1-2 hours up to 6.5 for 6 hours or more) was used to calculate total wearing
12 time of the week.

13 Agreement between the two methods was assessed according to the non-parametric Bland and
14 Altman method¹⁹. The non-parametric method was used because of violation of normal distribution of
15 data due to some extreme values for differences in wearing time between measurement methods. For
16 each participant, median of the differences between both measurement methods per week was
17 determined. In addition, the group median of differences and 2.5th and 97.5th percentile score were
18 determined, reflecting the systematic difference between methods and limits of agreement
19 respectively.

20 Results

21 Temperature sensor data were available for 8 participants of the Splint study (mean age: 8.7 ± 1.9 years;
22 gender: 6 males, 2 females; GMFCS: level I (n=2), level II (n=4), III (n=2)). The period of KAFO
23 treatment of 4 participants was shorter than one year because other treatment (e.g. botulinum toxin
24 treatment) was indicated clinically (n=3) or the KAFO caused too much discomfort (n=1) and
25 therefore the protocol was not finished. As a consequence, KAFO treatment varied from 2 months to 1
26 year. Individual response to questionnaires ranged from 22 to 100% with a median of 87%. For about
27 25% of the sent questionnaires reminders had to be sent. Acceptable time-temperature data that was
28 available per participant ranged from 10 to 100% (median: 66%). Missing time-temperature data was
29 caused by 1) too high air temperature because of weather conditions, leading to a too small increase in
30 temperature if the KAFO was worn (3% of total missing time-temperature data), or 2) technical
31 problems with sensor-data-loggers (due to sensor failure that was not related to KAFO wearing),
32 leading to empty data-files (97% of total missing time-temperature data). The period that data of both
33 WT_{parent} and WT_{obj} were available ranged from 1 to 11 weeks per participant, with a median of 3
34 weeks.

1 << Figure 2 >>

2 Individual median, interquartile range, minimum and maximum values of WT_{parent} and WT_{obj}
3 are presented in Fig. 2. Temperature data show that the KAFO wearing time was lower than prescribed
4 and in one case, KAFO wearing time was 0 hours. In addition, data show that, in some participants,
5 there were large differences between WT_{parent} and WT_{obj} . The median of these differences per week are
6 shown per participant in Fig. 3. This figure shows that there was no systematic difference between
7 both methods (median: 0.13 hours per week), but that differences between methods show large
8 variability between participants (ranging from -2.0 to 12.9 hours per week), resulting in large limits of
9 agreement (lower limit/2.5th percentile: -1.7 hours per week; upper limit/97.5th percentile: 11.1 hours
10 per week), This indicates that agreement between methods is low.

11 << Figure 3 >>

12 Comparing measurements within participants, a high variability in differences between the
13 two methods was shown as well. Individual data of weekly measurements of two participants is shown
14 in Fig. 4. For participant 102, WT_{parent} ranged from 0 to 12 hours per week, while WT_{obj} ranged from 0
15 to 8.5 hours per week (Fig. 3A). Differences between WT_{parent} and WT_{obj} ranged from -5.5 to 5.5 hours
16 per week with a median of 0 hours per week (Fig 3B). For participant 207, WT_{parent} ranged from 0 to
17 16.5 hours per week, while WT_{obj} range from 0 to 8.5 hours per week). Differences between WT_{parent}
18 and WT_{obj} ranged from -1 to 10 hours per week with a median of 2.5 hours per week. The wide range
19 of within participant difference between WT_{parent} and WT_{obj} was also expressed in high individual
20 limits of agreement (see figure 3B and 3D, participant 102: lower limit/2.5th percentile: -5.4 hours per
21 week, upper limit/97.5th percentile: 5.2 hours per week; participant 207: lower limit/2.5th percentile: -
22 -0.9 hours per week, upper limit/2.5th percentile: 9.4 hours per week).

23 << Figure 4 >>

24

25 Discussion

26 The present study shows that parent-reported wearing time of KAFOs differs from objective
27 measurements. The differences in reported wearing time between methods vary considerably per
28 participant: some parents reported higher wearing times than what was measured by temperature
29 sensor-data-loggers while others reported lower wearing times. In literature regarding spine orthoses,
30 mostly overestimation of self/parent-reported wearing time is reported (differences range from 0-
31 150%)^{8;12-16}. It should be noted that these conclusions are often based on means of all participants. In
32 accordance with the reported range of differences between wearing time measurement methods, the
33 present study found no systematic difference or at most a small overestimation of parent-reported

1 wearing time. The large range of overestimation reported in literature may indicate a similar trend as
2 found in this study that self/parent-reported wearing time and objectively measured wearing time
3 varies considerably between participants. Therefore it may be concluded that parent-reported wearing
4 time and objectively measured wearing time are not comparable.

5 In the present study, high values of limits of agreement between parent-reported and
6 objectively measured wearing times were found. If the objectively measured wearing time method is
7 considered as a golden standard, the limits of agreement reflect the error of parent-reported wearing
8 time. In the present study, the limits of agreement indicate that only if parents reported that the KAFO
9 was worn for more than 11 hours per week, it can be concluded with certainty that the KAFO was
10 worn for at least a short period. However, participants often did not wear their KAFO for such long
11 period in general (i.e. Fig. 2 shows that participant 102 and 207 did wear their KAFO not more than
12 8.5 hours per week). This indicates that parent reported wearing time cannot distinguish whether
13 participants wore their KAFO or not. It should be noted that one outlying value (participant 109, Fig.
14 1) affected error of parent-reported outcome considerably. However, sub-analyses excluding this
15 outlier still showed an upper limit of agreement of 2.4 hours per week. Taking into account that the
16 median of objectively measured wearing time of the remaining 7 participants was about 5 hours per
17 week, agreement between methods is still very low.

18 The response rate of questionnaires was relatively high (approximately 85%). Additional
19 analysis showed that there was no correlation between response rate and agreement between parent-
20 reported and objectively measured wearing times (Spearman's $\rho=0.252$, $p=0.5$). Another factor
21 possibly affecting agreement was delayed response to questionnaires: it was often necessary to send
22 reminders to parents after 2 weeks. This may indicate that parents did not complete questionnaires on
23 the day of KAFO wearing and may explain the variability in response. However, uncertainty about
24 delayed responses and therefore variable recall time will always be part of any parent report method.
25 Overestimation was larger compared to under estimation: some parents reported KAFO wearing for
26 periods that the temperature-data-logger did not register any higher temperatures. This may be the
27 result of a more general phenomenon that people adjust their answers according to socially desired
28 answers. It is unknown whether other methods of self-reporting may yield more accurate results.

29 Because of validity problems of parent-reported wearing time, we recommend the use of
30 objective techniques to measure KAFO wearing time. We agree with Hunter⁹ et al, that it is feasible to
31 use temperature sensor-data-loggers to measure KAFO wearing time. However, the high percentage of
32 missing data with the current device indicates that technical failures can occur and need to be solved.
33 In this study, we were able to collect data for each participant after replacing the non-functioning
34 sensors. In most datasets, it was straightforward to distinguish KAFO wearing from non-KAFO
35 wearing using criteria described above. However, at high ambient temperatures (about 25°C), it is

1 more difficult to determine KAFO wearing time because differences in temperature with or without a
2 leg in the KAFO was smaller. Comparison of data from temperature sensors inside the KAFO and
3 additional outside temperature sensors²⁰ may solve this problem.

4 Limitations of the study

5 A limitation of this study is the low number of participants. Non-parametric Bland Altman
6 plots are less reliable in small sample sizes¹⁹. However, because of the large intra-individual variation,
7 we believe that research using larger samples will yield similar conclusions.

8 A second limitation is that parent-reported wearing time was assessed by only one method and
9 cannot be generalized to other self/parent-reporting wearing time methods. We chose to send online
10 questionnaires during each day. This was only done for every 4th week of the month to decrease
11 burden on participants.

12 Applications to clinical practice

13 Measuring parent-reported orthotic wearing time with online daily questionnaires for one
14 week per month does not show to agree with objective measurements using a temperature sensor-data-
15 logger. Outcome of parent-reported questionnaires has to be interpreted with extreme care.

16 **Conclusion**

17 Wearing time of a knee-ankle-foot orthosis assessed by questionnaires is not in agreement with of
18 wearing time assessed by temperature sensor-data-loggers and should therefore be interpreted with
19 utmost care. Using objective measurement methods are preferred. Regarding temperature sensors, we
20 recommend to use a more sophisticated method then used in this study to measure orthotic wearing
21 time with temperature sensor-data-loggers, like the method described by Bus e.a.²⁰, in order to
22 overcome difficulties in determining orthotic wearing time at high ambient temperatures.

23

24 **Acknowledgement**

25 We thank all participants and their parents for their participation in this study. We also thank
26 OIM/Noppe orthopedie, Noordwijkerhout, the Netherlands; Kamer orthopedie, Amsterdam, the
27 Netherlands and Orthopedie Heckert & van Lierop, Venlo, the Netherlands for their coöperation.

28

29 **Conflicts of Interest and Source of Funding**

30 The Splint study is financially supported by a grant from the Phelps Stichting voor Spastici, Bussum,
31 The Netherlands [grant number: 2008032]. In addition to funding, the Splint study is financially
32 supported by a donation of Ultraflex Europe, Ternat, Belgium on behalf of Ultraflex Systems, Inc,

1 Pottstown, Pennsylvania, USA. The “temperature sensor-data-loggers” were partly sponsored by
2 PEDAK Meettechniek BV Heythuysen, The Netherlands. The authors declare that there is no conflict
3 of interest. Although we received financial support from Ultraflex Europe, Ultraflex Systems Inc. and
4 PEDAK Meettechniek, they did not have authority over this study.

5

6

1 **References**

2

3 (1) Condie DN, Meadows CB. Report of a Consensus Conference on the Lower Limb Orthotic
4 Management of Cerebral Palsy. 1995. Copenhagen, International society of Prosthetics &
5 Orthotics.

6

7 (2) Morris C, Condy D. Recent Developments in Healthcare for Cerebral Palsy: Implications and
8 Opportunities for Orthotics. Copenhagen, Denmark: International Society for Prosthetics and
9 Orthotics, 2009.

10 (3) Tardieu C, Lespargot A, Tabary C, Bret M. For how long must the soleus muscle be stretched
11 each day to prevent contractures. *Dev Med and Child Neurol* 1988;30:3-10.

12 (4) Maas J, Dallmeijer A, Huijing P et al. A randomized controlled trial studying efficacy and
13 tolerance of a knee-ankle-foot orthosis used to prevent equinus in children with spastic
14 cerebral palsy. *Clin Rehabil* 2014;28:1025-1038.

15 (5) Maas JC, Dallmeijer AJ, Huijing PA et al. Splint: the efficacy of orthotic management in rest
16 to prevent equinus in children with cerebral palsy, a randomised controlled trial. *BMC Pediatr*
17 2012;12:38.

18 (6) Zhao X, Xiao N, Li H, Du S. Day vs. Day-Night Use of Ankle-Foot Orthoses in Young
19 Children with Spastic Diplegia: A Randomized Controlled Study. *Am J Phys Med Rehabil*
20 2013;92:905-911.

21 (7) Benish BM, Smith KJ, Schwartz MH. Validation of a miniature thermochron for monitoring
22 thoracolumbosacral orthosis wear time. *Spine (Phila Pa 1976)* 2012;37:309-315.

23 (8) Helfenstein A, Lankes M, Ohlert K et al. The objective determination of compliance in
24 treatment of adolescent idiopathic scoliosis with spinal orthoses. *Spine (Phila Pa 1976)*
25 2006;31:339-344.

26 (9) Hunter LN, Sison-Williamson M, Mendoza MM et al. The validity of compliance monitors to
27 assess wearing time of thoracic-lumbar-sacral orthoses in children with spinal cord injury.
28 *Spine (Phila Pa 1976)* 2008;33:1554-1561.

29 (10) Rahman T, Borkhuu B, Littleton AG et al. Electronic monitoring of scoliosis brace wear
30 compliance. *J Child Orthop* 2010;4:343-347.

31 (11) Havey R, Gavin T, Patwardhan A et al. A reliable and accurate method for measuring orthosis
32 wearing time. *Spine (Phila Pa 1976)* 2002;27:211-214.

33 (12) Donzelli S, Zaina F, Negrini S. In defense of adolescents: They really do use braces for the
34 hours prescribed, if good help is provided. Results from a prospective everyday clinic cohort
35 using thermobrace. *Scoliosis* 2012;7.

36 (13) Morton A, Riddle R, Buchanan R, Katz D, Birch J. Accuracy in the prediction and estimation
37 of adherence to bracewear before and during treatment of adolescent idiopathic scoliosis. *J*
38 *Pediatr Orthop* 2008;28:336-341.

39 (14) Nicholson GP, Ferguson-Pell MW, Smith K, Edgar M, Morley T. The objective measurement
40 of spinal orthosis use for the treatment of adolescent idiopathic scoliosis. *Spine (Phila Pa*
41 *1976)* 2003;28:2243-2250.

- 1 (15) Takemitsu M, Bowen JR, Rahman T, Glutting JJ, Scott CB. Compliance monitoring of brace
2 treatment for patients with idiopathic scoliosis. *Spine (Phila Pa 1976)* 2004;29:2070-2074.
- 3 (16) Vandal S, Rivard CH, Bradet R. Measuring the compliance behavior of adolescents wearing
4 orthopedic braces. *Issues Compr Pediatr Nurs* 1999;22:59-73.
- 5 (17) Morgenstein A, Davis R, Talwalkar V, Iwinski H, Jr., Walker J, Milbrandt TA. A randomized
6 clinical trial comparing reported and measured wear rates in clubfoot bracing using a novel
7 pressure sensor. *J Pediatr Orthop* 2015;35:185-191.
- 8 (18) Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and
9 reliability of a system to classify gross motor function in children with cerebral palsy. *Dev*
10 *Med Child Neurol* 1997;39:214-223.
- 11 (19) Bland JM, Altman DG. Measuring agreement in method comparison studies. *Stat Methods*
12 *Med Res* 1999;8:135-160.
- 13 (20) Bus SA, Waaijman R, Nollet F. New monitoring technology to objectively assess adherence to
14 prescribed footwear and assistive devices during ambulatory activity. *Arch Phys Med Rehabil*
15 2012;93:2075-2079.
16
17

1 **Tables**

2 **Figures**

3

4 Figure 1: the temperature sensor-data-logger attached to the KAFO

5 The KAFO as shown in the figure is configured with a fixed knee angle and a variable ankle angle
6 using a spring that provides variable ankle dorsiflexion. The temperature sensor-data-logger was
7 attached to the KAFO with a nut (upper inset) and bolt (lower inset). The flat side (lower inset) of the
8 sensor-data-logger touched the participants skin during wearing of the KAFO.

9

10 Figure 2: Box-and-whisker plots of wearing time per week per participant

11 Comparison of parent-reported wearing time per week (WT_{parent}) and objectively measured wearing
12 time per week (WT_{obj}) per participant. Numbers of weeks included in analyses are for each participant
13 noted on the x-axis. For participant 107, WT_{parent} and WT_{obj} was reported and measured as 0 hours per
14 week. The middle line of each box shows the median wearing time. The outer lines of each box show
15 the 25th and 75th percentile and the end of the whiskers show the maximum and minimum values.

16

17 Figure 3: Bland-Altman plot: median differences per participant

18 Bland-Altman plot of parent-reported wearing time per week (WT_{parent}) and objectively measured
19 wearing time per week (WT_{obj}). Upper and lower dotted lines indicate limits of agreement (2.5th and
20 97.5th percentile). Dotted line in middle indicates median of the difference between WT_{parent} and WT_{obj}
21 for all participants.

22

23 Figure 4: Individual data: examples

24 A+C: Comparison of parent-reported wearing time per week (WT_{parent}) and objectively measured
25 wearing time per week (WT_{obj}) for two participants. If no column is visible, wearing time is 0 hours.

26 B+D: Bland-Altman plot of parent-reported wearing time per week (WT_{parent}) and objectively
27 measured wearing time per week (WT_{obj}) for the same participants as in A and C. Upper and lower
28 dotted lines indicate limits of agreement (2.5th and 97.5th percentile). Dotted line in middle indicates
29 median of the difference between WT_{parent} and WT_{obj} for all measurements.