

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Liangcheng Du Publications

Published Research - Department of Chemistry

---

5-31-2016

## A TonB-dependent receptor regulates antifungal HSAF biosynthesis in *Lysobacter*

Ruping Wang

*Nanjing Agricultural University*

Huiyong Xu

*Nanjing Agricultural University*

Liangcheng Du

*University of Nebraska-Lincoln, ldu3@unl.edu*

Shan-Ho Chou

*National Chung Hsing University, Taipei*

Hongxia Liu

*Nanjing Agricultural University*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/chemistrydu>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#), and the [Chemistry Commons](#)

---

Wang, Ruping; Xu, Huiyong; Du, Liangcheng; Chou, Shan-Ho; Liu, Hongxia; Liu, Youzhou; Liu, Fengquan; and Qian, Guoliang, "A TonB-dependent receptor regulates antifungal HSAF biosynthesis in *Lysobacter*" (2016). *Liangcheng Du Publications*. 12.

<https://digitalcommons.unl.edu/chemistrydu/12>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Liangcheng Du Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Ruping Wang, Huiyong Xu, Liangcheng Du, Shan-Ho Chou, Hongxia Liu, Youzhou Liu, Fengquan Liu, and Guoliang Qian

# SCIENTIFIC REPORTS



OPEN

## A TonB-dependent receptor regulates antifungal HSAF biosynthesis in *Lyso bacter*

Ruping Wang<sup>1</sup>, Huiyong Xu<sup>1</sup>, Liangcheng Du<sup>2</sup>, Shan-Ho Chou<sup>3</sup>, Hongxia Liu<sup>1</sup>, Youzhou Liu<sup>4</sup>, Fengquan Liu<sup>4</sup> & Guoliang Qian<sup>1</sup>

Received: 04 March 2016

Accepted: 25 April 2016


Published: 31 May 2016

*Lyso bacter* species are Gram-negative bacteria that are emerging as new sources of antibiotics, including HSAF (Heat Stable Antifungal Factor), which was identified from *L. enzymogenes* with a new mode of action. LesR, a LuxR solo, was recently shown to regulate the HSAF biosynthesis via an unidentified mechanism in *L. enzymogenes* OH11. Here, we used a comparative proteomic approach to identify the LesR targets and found that LesR influenced the expression of 33 proteins belonging to 10 functional groups, with 9 proteins belonging to the TBDR (TonB-Dependent Receptor) family. The fundamental role of bacterial TBDR in nutrient uptake motivates us to explore their potential regulation on HSAF biosynthesis which is also modulated by nutrient condition. Six out of 9 TBDR coding genes were individually in-frame deleted. Phenotypic and gene-expression assays showed that TBDR7, whose level was lower in a strain overexpressing *lesR*, was involved in regulating HSAF yield. TBDR7 was not involved in the growth, but played a vital role in transcribing the key HSAF biosynthetic gene. Taken together, the current *lesR*-based proteomic study provides the first report that TBDR7 plays a key role in regulating antibiotic (HSAF) biosynthesis, a function which has never been found for TBDRs in bacteria.

TonB-dependent receptors (TBDRs) are a family of proteins that are located in the outer membrane of Gram-negative bacteria<sup>1</sup>. These receptors share a common structural feature of two domains; a C-terminal membrane-embedded  $\beta$ -barrel domain that is sealed by a conserved N-terminal globular domain (plug domain)<sup>2,3</sup>. TBDRs typically act as channels that open in response to outside ligands to allow import of extracellular nutrients, such as iron-siderophore complexes or non-Fe compounds (e.g. vitamin B12), into the periplasmic space<sup>4,5</sup>. The best-characterized examples include FecA, FhuA, FepA and BtuB, which are necessary for the active transport of the corresponding iron siderophores of ferric citrate, ferrichrome, or enterobactin, as well as vitamin B12, respectively<sup>1–3,6</sup>. The TBDR-dependent substrate transport is an active process that requires energy input from the proton motive force across the cytoplasmic membrane. Such a process requires that the ligand-loaded TBDRs interact with the TonB protein complex consisting of three inner membrane proteins (TonB/ExbB/ExbD)<sup>5,7</sup>. Although the basic role of TBDRs is believed mainly in nutrient transport, some TBDRs are also shown to trigger pathogenesis in several animal and plant bacterial pathogens<sup>8,9</sup>. Nevertheless, TBDR is never reported to play a role in regulating bacterial antibiotic biosynthesis to our knowledge.

LuxR solo is defined as a group of LuxR-family proteins possessing a classical AHL (*N*-acyl-homoserine lactones)-binding domain at the N terminus and a HTH DNA-binding domain at the C terminus, as other LuxR proteins in the canonical LuxI/R system<sup>10</sup>. However, LuxR solo lacks any cognate LuxI protein synthesizing the QS (Quorum sensing) signal AHL<sup>10,11</sup>. Bacterial LuxR solo thus potentially responds to signals produced by the bacteria itself, the neighboring bacteria, or the eukaryotes (e.g. plants) to exert the corresponding regulations, such as biofilm formation, virulence and biocontrol activity in several bacteria<sup>10,12–18</sup>. The genus *Lyso bacter*, belonging to the Xanthomonadaceae family, is a group of Gram-negative bacteria with several conserved features, such as high genomic G+C content (approximately 70%), flagella-independent twitching motility, production of abundant lytic enzymes, as well as generation of bioactive natural products<sup>19–21</sup>. These distinct characteristics differentiate

<sup>1</sup>College of Plant Protection (Key Laboratory of Integrated Management of Crop Diseases and Pests), Nanjing Agricultural University, Nanjing 210095, P.R. China. <sup>2</sup>Department of Chemistry, University of Nebraska-Lincoln, Lincoln, Nebraska 68588, United States. <sup>3</sup>Institute of Biochemistry, and NCHU Agricultural Biotechnology Center, National Chung Hsing University, Taichung, Taiwan, ROC. <sup>4</sup>Institute of Plant Protection, Jiangsu Academy of Agricultural Science, Nanjing 210014, P.R. China. Correspondence and requests for materials should be addressed to F.L. (email: fqliu20011@sina.com) or G.Q. (email: glqian@njau.edu.cn)



**Figure 1. Identification of proteins affected by the *lesR* overexpression through 2-D gel proteome analysis of *Lysobacter enzymogenes* OH11.** (A) Monitoring and comparison of the growth curve between the *lesR* overexpression and control strains in 1/10 TSB broth. The time point used for cell collection was set at OD<sub>600</sub> 1.0 (indicated by the dotted line). Data are from three independent biological experiments. Each experiment involved three replicates for each strain. (B) Overexpression of *lesR* almost impaired the HSAF production in strain OH11. The HSAF production (indicated by the red arrow) from the collected cells, as noted in part (A), was extracted and determined by HPLC. (C) Functional classification of the identified 33 proteins affected by the *lesR* overexpression. The detailed information of each gene described in this figure was provided in the Table 1. OH11(*lesR*), the *lesR* overexpression strain; OH11(pBBR), the wild-type strain containing an empty expressing vector.

*Lysobacter* from its ecological/taxonomic related species, such as *Xanthomonas*<sup>19,22</sup>. A well-characterized species of this genus is *L. enzymogenes*, which is emerging as a biological control agent against fungal pathogens of crop plants, such as *Bipolaris sorokiniana* and *Rhizoctonia solani*<sup>23,24</sup>. *L. enzymogenes* is also currently recognized as a new source of antibiotics<sup>20</sup>, including HSAF (Heat Stable Antifungal Factor) that belongs to the distinct PTM (polycyclic tetramate macrolactam) antifungal antibiotic with a new mode of action<sup>20,25,26</sup>. Furthermore, our previous reports show that the *hsaf* *pks/nrps* gene, encoding a hybrid polyketide synthase-nonribosomal peptide synthetase, is responsible for the HSAF biosynthesis in *L. enzymogenes*<sup>27,28</sup>. However, the yield of HSAF in *L. enzymogenes* is relatively low even in the HSAF inducing medium<sup>26,27</sup>. Therefore, elucidation of the regulatory mechanism(s) of HSAF biosynthesis in *L. enzymogenes* is necessary for improving the HSAF yield by genetic engineering or molecular biotechnology. Recent advancements have started to shed light into this issue<sup>21,22,29</sup>. Intriguingly, we previously found that overexpression, but not deletion of *lesR*, the only LuxR solo coding gene in *L. enzymogenes*, almost entirely impaired the HSAF production<sup>29</sup>. However, how the overexpressed *lesR* performed this critical control on the HSAF biosynthesis remains to be investigated.

To further understand the mechanism used by *LesR* in regulating HSAF biosynthesis, we have endeavored to identify the *LesR* targets that contribute to HSAF biosynthesis. By a combination of proteomics, bioinformatics and genetic approaches, we discovered that a certain TBDR protein, whose level was affected by the *lesR* overexpression, played an important role in regulating the antibiotic HSAF biosynthesis in *L. enzymogenes*. Our findings represent the first report about a novel functionality of TBDR proteins in bacteria.

## Results

**The levels of 9 TBDR proteins were affected by the *lesR* overexpression.** Given that overexpression, but not deletion, of *lesR* in the wild-type OH11 was found to almost shut down the HSAF biosynthesis<sup>29</sup>, we therefore selected the *lesR* overexpression strain, as well as the wild-type OH11 possessing an empty vector as a control for 2-D gel proteome analysis to identify potential *LesR* targets in *L. enzymogenes*. To achieve this point, the growth ability of the *lesR* overexpression strain and control strain was first determined and compared in the HSAF-inducing medium (1/10 TSB broth). As shown in Fig. 1A, overexpression of *lesR* in the wild-type OH11 did not alter its growth ability in comparison to that of the control strain. On the basis of this result, a good time point at 12.5 h after initial inoculation that corresponds to the logarithmic phase of both strains was chosen for

cell collection (Fig. 1A). We showed that regulation of *lesR* in the HSAF biosynthesis was functional at this designated point, because the control strain produced HSAF, whereas no HSAF was detected in the *lesR* overexpression strain at this time (Fig. 1B).

Next, total proteins were extracted from the collected cells of the *lesR* overexpression strain and control strain. After purification and quantification, these proteins were separated by 2-D gel electrophoresis. In this way, a total of 98 differentially expressed protein spots (with a threshold of larger than 1.5-fold change) were excised from silver-stained gels and subject to MALDI-TOF-TOF analysis; 33 of them were confidentially identified in the genome of strain OH11 (Table 1). *In silico* analysis further divided these 33 proteins into 10 groups (Fig. 1C and Table 1). The largest percentage of annotated proteins (27%) affected by *lesR* overexpression corresponds to the group of ‘inorganic ion transport and metabolism’ (Fig. 1C), which comprised 9 TBDR proteins (Table 1). These TBDR proteins were then further investigated for their potential roles in regulating HSAF biosynthesis, because the basic role of TBDR in nutrient uptake in *L. enzymogenes* is speculated to be correlated with the nutrient-dependent property of HSAF biosynthesis<sup>25,26</sup>. After a detailed sequence analysis of these 9 TBDR proteins as shown in Fig. 2A, we found that each TBDR possessed two conserved domains, a C-terminal membrane-embedded  $\beta$ -barrel domain (ligand\_gated\_channel) and an N-terminal plug domain (Plug domain; ~150–200 residues) that is similar to the well-characterized TBDR protein BtuB<sup>30,31</sup>. Furthermore, TBDR2, TBDR4 and TBDR9 also contained an additional domain, the TonB\_dep\_Rec domain (TonB dependent Receptor). Notably, all 9 TBDR proteins had a TonB-box region at their corresponding N terminus as BtuB (Fig. 2B).

The analysis also showed that all detected TBDR proteins did not contain a long N-terminal signaling domain, a distinct structural feature of the TBDT (TonB-dependent transducers) family that differentiates them from the conventional TBDRs<sup>32</sup>. Furthermore, none of all detected TBDR proteins was adjacent to a ECF (extracytoplasmic function) sigma factor and anti-sigma factor in their respective genetic organization (Supplementary Fig. S2), which is another typical characteristic of the TBDT-based CSS (cell-surface signaling) system in bacteria<sup>33</sup>. All these results strongly suggest that the 9 detected TBDR proteins belong to the conventional TBDR family, but not TBDT. Collectively, this 2-D proteomic study indicates that the levels of all 9 TBDR proteins were influenced by the *lesR* overexpression in *L. enzymogenes*.

### Systematic mutation revealed that only TBDR7 played a key positive role in controlling HSAF biosynthesis.

To test whether all identified TBDR proteins contribute to HSAF biosynthesis, each *TBDR* gene in *L. enzymogenes* was mutated by an in-frame deletion. In this way, 6 gene-deletion mutants, including the  $\Delta TBDR1$ , 2, 4, 7, 8 and 9, were generated and further validated (Supplementary Table S2). The TBDR3, 5 and 6 coding genes appear to be essential for bacterial survival under the test condition, because these gene knockout bacteria failed to grow under a similar condition. Next, the HSAF yield was quantified in each *TBDR* mutant. As shown in Fig. 3, only 2 out of the 6 *TBDR* mutants were found to change the HSAF level, and knockout of the gene *TBDR2*, 4, 8 or 9 had no effect on the HSAF yield (Fig. 3). In particular, inactivation of *TBDR7* almost abolished the HSAF production, whereas missing of *TBDR1* significantly enhanced the HSAF level (Fig. 3). We also generated a  $\Delta TBDR1$ &7 double mutant (Table 2 and Supplementary Table S2), and found that it produced approximately 55% HSAF to that of the wild type (Fig. 3), suggesting that *TBDR1* and *TBDR7* played opposing roles in regulating HSAF biosynthesis in *L. enzymogenes*. Collectively, the above results suggest that both *TBDR1* and *TBDR7* potentially controlled the HSAF biosynthesis in *L. enzymogenes*.

In the following study, we focused our efforts on *TBDR7*, because it seems to play a greater role in HSAF production than *TBDR1*. To verify the role of *TBDR7*, its complemented strain of  $\Delta TBDR7$  was constructed and verified by RT-PCR (Fig. 4A). As shown in Fig. 4B, the *in trans* *TBDR7* complementation restored the HSAF production of the  $\Delta TBDR7$  mutant to the wild-type level, whereas the  $\Delta TBDR7$  mutant was deficient when complemented with an empty vector (Fig. 4B). In addition, we also created a point mutation (V74A) at the predicted TonB-box region of *TBDR7* (Fig. 2B), because this amino acid (Val74) was previously shown to be important in transporting vitamin B12 in BtuB<sup>34</sup>. As shown in Fig. 4B, the *TBDR7* containing the V74A indeed failed to complement the HSAF deficiency of the  $\Delta TBDR7$  mutant, revealing the importance of the *TBDR7* TonB-box region in controlling HSAF biosynthesis. Collectively, these results suggest that *TBDR7* played a vital role in regulating the HSAF production in *L. enzymogenes*.

### TBDR7 positively regulated the *hsaf pks/nprs* transcription.

To investigate whether the deficiency of HSAF production of the  $\Delta TBDR7$  mutant in 1/10 TSB is due to the different growth rates, the growth capacity of the  $\Delta TBDR7$  mutant and the wild-type OH11 in this medium was examined. As shown in Fig. 4C, deletion of *TBDR7* did not seem to alter the bacterial growth rate under the similar test condition, suggesting that *TBDR7* was not involved in the growth, but controlled the HSAF production in *L. enzymogenes*. To further address this point, we determined the transcriptional level of *hsaf pks/nprs*, the key gene for HSAF biosynthesis in *L. enzymogenes*<sup>27</sup>. As shown in Fig. 4D, transcription of *hsaf pks/nprs* was shut down almost entirely in the  $\Delta TBDR7$  mutant compared to that of the wild-type OH11. This finding was consistent with the decreasing HSAF level in the  $\Delta TBDR7$  mutant (Fig. 3), and further suggests that the contribution of *TBDR7* on the HSAF biosynthesis was, at least partially, due to decreasing transcription of the key HSAF biosynthetic gene in *L. enzymogenes*.

## Discussion

In the present study, an omics-based strategy was utilized to investigate how *LesR*, the LuxR solo from a biological control agent *L. enzymogenes*, is able to regulate the antibiotic HSAF biosynthesis. A comparative proteomic analysis led to the finding that *LesR* affects the expression of 98 protein spots when the threshold was set at 1.5-fold change. For these proteins, we have paid attention most to a series of TBDR proteins, because they are closely associated with the nutrient-dependent trait of HSAF biosynthesis. By using a combination of systematic mutation, phenotypic analysis and quantitative gene expression methods, we have further found that *TBDR7* was not

Cat- alog no.	Spot no.	Fold change OH11(PBBR)/ OH11( <i>lesR</i> )	Sequence coverage (%)	Gene acces- sion no.	Gene name <sup>a</sup> /abbrev- iation	Functional catalog <sup>b</sup>	Function/Similarity	pI (cal)	M <sub>r</sub> (cal) KD
1	7-03	+1000000	22	KP293905	TBDR9	Inorganic ion trans- port and metabolis	TonB-dependent sidero- phore receptor	5.26	83.5
2	7-05	+3.71567	10	KP293900	TBDR4		TonB-dependent receptor	4.95	94.9
3	7-12	-2.42089	23	KP293903	TBDR7		TonB-dependent outer membrane receptor	4.86	103.9
	7-13	-1000000	15						
	7-15	-1000000	12						
	7-16	-2.95065	9						
	7-18	-1000000	19						
	7-30	-1000000	13						
4	7-19	+2.36877	18	KP293903	TBDR8		TonB-dependent receptor	4.89	102.7
	7-28	-6.99916	21						
5	7-20	+6.49466	15	KP293902	TBDR6		TonB-dependent receptor	5.06	105.5
	7-21	-6.21763	10						
6	7-38	+3.28287	9	KP293901	TBDR5		TonB-dependent receptor	5.24	102.3
7	8-03	-1000000	24	KP293901	TBDR3		TonB-dependent receptor	5.39	99.0
8	8-06	-2.06331	27	KP293898	TBDR2		TonB-dependent receptor domain protein	5.60	86.2
9	7-49	+1000000	6	KP293897	TBDR1	putative tonb-dependent outer membrane receptor	5.64	118.6	
10	7-07	+1000000	28	KP293926	OH11GL004315/ le4315	Energy production and conversion	Dihydropolyl dehydro- genase	6.03	50.4
11	8-22	-2.28895	30	KP293925	OH11GL000895/ le0895		catalytic domain of components of various dehydrogenase complexes	6.32	46.5
12	7-14	+3.75737	18	KP293921	OH11GL002176/ le2176	Lipid transport and metabolism	FadL family outer mem- brane protein	5.27	48.4
	7-42	-1000000	26						
13	7-25	+1000000	22	KP293928	OmpA2	Cell wall/mem- brane/envelope biogenesis	OmpA family outer membrane protein	4.8	39.1
14	7-34	+1000000	28	KP293927	OmpA1		OmpA family outer membrane protein	4.8	39.1
15	7-26	+2.82758	19	KP293917	OH11GL000050/ le0050	Posttranslational modification, protein turnover, chaperones	trigger factor	4.96	48.8
	7-45	-1000000	16						
	7-51	-8.98225	8						
16	7-33	+1000000	38	KP293918	OH11GL001285/ le1285		glutaredoxin-like protein	4.98	32.7
	7-46	-3.51678	44						
17	8-44	-2.28179	13	KP293918	OH11GL002659/ le2659		chaperonin GroEL	5.2	57.3
18	7-22	-1000000	22	KP293907	OH11GL001810/ le1810	Function unknown	DNA binding domain-containing protein	4.85	38.1
19	7-23	+1000000	53	KP293910	OH11GL002922/ le2922		hypothetical protein	5.73	35.1
20	7-28	+6.99916	21	KP293912	OH11GL004158/ le4158		No hit	5.22	86.7
21	7-39	-2.05056	58	KP293908	OH11GL002552/ le2552		hypothetical protein	5.34	29.8
22	7-43	+1000000	17	KP293913	OH11GL004940/ le4940		No hit	8.35	26.6
23	7-44	+1000000	24	KP293906	OH11GL002473/ le2473		No hit	5.15	40.8
24	8-30	-2.4328	34	gi 189474077	<i>hmgA</i>		homogentisate 1, 2-dioxygenase	5.93	50.1
25	8-38	+2.00608	40	KP293909	OH11GL002024/ le2024		putative secreted protein	6.77	31.8
26	8-41	+2.54476	39	KP293911	OH11GL004856/ le4856		No hit	5.2	26.6

Continued

Cat-alog no.	Spot no.	Fold change OH11(PBBR)/OH11( <i>lesR</i> )	Sequence coverage (%)	Gene accession no.	Gene name <sup>a</sup> /abbreviation	Functional catalog <sup>b</sup>	Function/Similarity	pI (cal)	M <sub>r</sub> (cal) KD
27	7-32	+3.73628	17	KP293915	OH11GL003474/le3474	General function prediction only	hypothetical protein	4.69	24.4
28	7-35	+1000000	29	KP293916	OH11GL004311/le4311		hypothetical protein	5.78	34.2
29	8-35	-2.79579	47	KP293914	OH11GL002539/le2539		NADP-dependent alcohol dehydrogenase	5.43	38.06
30	7-36	+2.63927	38	KP293922	OH11GL002141/le2141	Amino acid transport and metabolism	spermidine synthase	5.05	31.8
31	8-02	+3.82444	18	KP293920	OH11GL000430/le0430	Transcription	DNA-directed RNA polymerase subunit beta	5.73	155.2
32	8-34	-2.02788	33	KP293920	OH11GL003264/le3264	Translation, ribosomal structure and biogenesis	glutamyl-tRNA synthetase	5.56	51.6
33	8-47	+1000000	18	KP293924	OH11GL005061/le5061		No hit	5.4	43.2

**Table 1. Identification of 33 proteins affected by the *lesR* overexpression in *Lysobacter enzymogenes*.** <sup>a</sup>Gene name was based on the genome sequence of *L. enzymogenes* strain OH11<sup>27</sup>, which could be found with the accession number 1784099 in NCBI database. <sup>b</sup>Functional catalog was performed by using protein blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

involved in the growth, and acted as a key protein in controlling the HSAF production. Although bacterial TBDRs have been reported to play a key fundamental role in various nutrient transport, and some of them also have an important role in pathogen-host interaction in several pathogenic bacteria<sup>8,9</sup>, to our knowledge, no TBDR has been reported to control antibiotic biosynthesis in bacteria<sup>35</sup>. In the present study, we have used *L. enzymogenes* as a model bacterium and provide the first result that TBDR7 was involved in generating HSAF, a unique antifungal antibiotic. The present results therefore reveal a novel function of TBDR in bacteria.


Although the contribution of TBDR7 to the HSAF biosynthesis is well revealed in the present manuscript, the mechanism(s) is still unclear at this moment. It is well accepted that the fundamental function of TBDR is to uptake nutrient in nearly all bacterial species in an energy-dependent way, which requires direct interaction between the periplasmic domain of TonB and the TonB-box region of the ligand-loaded TBDRs<sup>33</sup>. In the present study, we did not yet know the nature of the TBDR7-loading ligand, and also lack the data for the direct interaction between TBDR7 and TonB. However, we did also find that TonB can regulate the HSAF biosynthesis, as mutation of *tonB* (Supplementary Table S2) almost completely impaired the HSAF yield (Supplementary Fig. S3). This result was correlated with the phenotype change of the  $\Delta$ TBDR7 mutant on HSAF production. It will be intriguing to identify the potential TBDR7-loading ligand(s) and to explore the possible TBDR7-TonB interaction for a better understanding on how TBDR7 regulates the HSAF biosynthesis.

TBDR7 is one of the 9 TBDR proteins that were identified from a *lesR*-based proteomic study presented in this work. These 9 TBDR proteins account for approximately 16% of all TBDR proteins (55) distributed in the genome of strain OH11 which belong to the group of 'inorganic ion transport and metabolism'. The functionality of TBDR has been shown to be directly related to different nutrient uptake, such as iron<sup>34</sup>. Competition for iron has long been known to be an important trait for beneficial rhizosphere colonization and for antagonism of plant deleterious microorganisms<sup>36</sup>. This finding has also been reported by PsoR, a LuxR solo of *Pseudomonas fluorescens* that responds to plant compounds<sup>17</sup>. As reported previously, LesR is a LuxR solo, but does not belong to the novel subgroup of plant-responding LuxR solo regulators (e.g. PsoR, OryR and XccR)<sup>29</sup>. However, we found that overexpression of either LesR or PsoR in the background of the relevant wild-type strain affected the gene/proteins involved in iron acquisition<sup>17</sup> (Fig. 1C). This suggests that different types of LuxR solos from different bacterial biological control agents might share a similar role in controlling expression of the certain genes/proteins, such as those relating to 'inorganic ion transport and metabolism'.


Since LesR is a transcription factor, one possible mechanism of LesR, therefore, is to regulate transcription of *TBDR7* via a direct or indirect manner. In partial support of this hypothesis, we found that overexpression of *lesR* significantly increased the transcript of *TBDR7* compared to that of the control strain (Supplementary Fig. S4), although the protein level of TBDR7 was decreased in the *lesR* overexpression strain. These results suggest that a post-transcriptional modification may occur in influencing the protein level of TBDR7 in the overexpressed *lesR* strain. We have also attempted to use gel shifting assay to test the direct interaction between LesR and the promoter of *TBDR7* *in vitro*, but failed, due to the great difficulty in getting purified recombinant LesR protein. This situation is consistent with the previous report on the preparation of representative LuxR solo OryR from *X. oryzae* pv. *Oryzae*<sup>14,37</sup>. In a future study, we will try to fuse the LesR with the MBP tag to obtain soluble fused LesR to check its binding with the *TBDR7* promoter.

## Materials and Methods

**Strains, plasmids and culture conditions.** The bacterial strains and plasmids used in this study are listed in Table 2. *Escherichia coli* strain DH5 $\alpha$  was used for plasmid constructions, and was grown in LB medium at 37 °C. *Lysobacter enzymogenes* OH11 (CGMCC No. 1978) and its derivative strains were grown in LB or 1/10 TSB (Tryptic Soy Broth, Sigma) at 28 °C, shaking at 200 rpm. When required, antibiotics were added into the medium



**Figure 2. Sequence analysis of nine TBDR proteins identified from the *lesR*-based proteomics in *L. enzymogenes*.** (A) The domain analyses of nine TBDR (TonB-Dependent Receptor) proteins. The well-characterized vitamin B12 receptor, BtuB from *E. coli* (gi|948468), served as a reference TBDR. (B) Multiple alignment of the TonB-box region of TBDR1 to TBDR9 with that of BtuB. The predicted TonB-box region was highlighted with a red box, similar to that of BtuB<sup>30</sup>. The conserved amino acid (Val74) that was marked with a red asterisk was selected for point mutation in further study.



**Figure 3. Quantification of HSAF yield from the wild-type OH11 of *L. enzymogenes* and its mutants.** Peak area indicated the area of HSAF determined by HPLC method, while the OD<sub>600</sub> represents the growth status of tested strains at the time points used for the extraction of HSAF. ΔTBDR (number) indicated the deletion mutant of target TBDR gene; ΔTBDR1&7, the double mutant of TBDR1 and TBDR7. Three replicates for each treatment were used, and the experiment was repeated three times. Vertical bars represent standard errors. The asterisk above the bars indicate a significant difference between the wild-type strain OH11 and the tested strains (\**p* < 0.05).

at final concentrations: kanamycin (Km) 100 μg/ml and gentamicin (Gm) 150 μg/ml for *L. enzymogenes* strains; Gm 25 μg/mL for *E. coli* strains.

**Protein extraction and purification, and 2-D gel analyses.** Total protein was extracted from bacterial cells according to the previous description<sup>38</sup>. In brief, cells were harvested from the culture (10% TSB broth) with an OD<sub>600</sub> (Optical Density at 600nm) of 1.0 by centrifugation (10,000 × *g* at 4 °C for 10 min). The cell pellets were re-suspended in 20-mL washing buffer (50 mM Tris-HCl, pH 7.2), and this step (centrifugation and washing) was




Strains and plasmids	Properties or characteristics <sup>a</sup>	Source
<i>Lysobacter enzymogenes</i>		
OH11	The wild-type strain, Km <sup>R</sup>	CGMCC No. 1978
OH11(pBBR)	OH11 harboring plasmid pBBR1-MCS5, Gm <sup>R</sup> , Km <sup>R</sup>	29
OH11( <i>lesR</i> )	OH11 harboring plasmid pBBR- <i>lesR</i> , Gm <sup>R</sup> , Km <sup>R</sup>	29
Δ <i>TBDR1</i>	<i>TBDR1</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR2</i>	<i>TBDR2</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR4</i>	<i>TBDR4</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR7</i>	<i>TBDR7</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR8</i>	<i>TBDR8</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR9</i>	<i>TBDR9</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR1</i> ϕ7	A double in-frame mutation in <i>TBDR1</i> and <i>TBDR</i> , Km <sup>R</sup>	This study
Δ <i>tonB</i>	<i>tonB</i> in-frame deletion mutant of strain OH11, Km <sup>R</sup>	This study
Δ <i>TBDR7</i> ( <i>TBDR7</i> )	Δ <i>TBDR7</i> harboring plasmid pBBR- <i>TBDR7</i> , Gm <sup>R</sup> , Km <sup>R</sup>	This study
Δ <i>TBDR7</i> (pBBR)	Δ <i>TBDR7</i> harboring plasmid pBBR-MCS5, Gm <sup>R</sup> , Km <sup>R</sup>	This study
Δ <i>TBDR7</i> ( <i>TBDR7</i> -V74A)	Δ <i>TBDR7</i> harboring plasmid pBBR- <i>TBDR7</i> , Gm <sup>R</sup> , Km <sup>R</sup>	This study
<i>Escherichia coli</i>		
DH5a	F <sup>-</sup> , ϕ80 <i>dlacZ</i> Δ <i>M15</i> , Δ( <i>lacZYA-argF</i> )U169, <i>deoR</i> , <i>recA1</i> , <i>endA1</i> , <i>hsdR17</i> (r <sub>k</sub> <sup>-</sup> , m <sub>k</sub> <sup>+</sup> ), <i>phoA</i> , <i>supE44</i> , λ <sup>-</sup> , <i>thi-1</i> , <i>gyrA96</i>	Lab collection
Plasmids		
pEX18GM	Suicide vector with a <i>sacB</i> gene, Gm <sup>R</sup>	42
pBBR1-MCS5	Broad-host-range vector with a P <sub><i>lac</i></sub> promoter, Gm <sup>R</sup>	43
pBBR- <i>lesR</i>	pBBR1-MCS5 cloned with a 1525-bp fragment containing the coding region of <i>lesR</i> and its predicted promoter region	29
pEX18- <i>TBDR1</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR1</i> flanking fragments	This study
pEX18- <i>TBDR2</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR2</i> flanking fragments	This study
pEX18- <i>TBDR4</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR4</i> flanking fragments	This study
pEX18- <i>TBDR7</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR7</i> flanking fragments	This study
pEX18- <i>TBDR8</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR8</i> flanking fragments	This study
pEX18- <i>TBDR9</i>	Km <sup>R</sup> , pEX18GM with two <i>TBDR9</i> flanking fragments	This study
pEX18- <i>tonB</i>	Km <sup>R</sup> , pEX18GM with two <i>TonB</i> flanking fragments	This study
pBBR- <i>TBDR7</i>	pBBR1-MCS5 cloned with a 3266-bp fragment containing the coding region of <i>TBDR7</i> and its predicted promoter	This study
pBBR- <i>TBDR7</i> (V74A)	pBBR- <i>TBDR7</i> , where the amino acid V74 within the TonB-box region was substituted by A74	This study

**Table 2. Bacterial strains and plasmids used in this study.** <sup>a</sup>Km<sup>R</sup> and Gm<sup>R</sup> = Kanamycin-, Gentamicin-, respectively.

repeated twice. The final washed cells were resuspended in 400-μl alkyls buffer (addition of 0.1 mM PMSF), and fragmented by an ultrasonic machine. Then the lysate was discarded by centrifugation at 10,000 × g at 4 °C for 20 min. The supernatant was transferred into a fresh 2-mL reaction tube. Total protein of each strain was purified by CKEAN UP kit (GE, USA). Protein samples were subsequently used for 2-D gel analyses, which was described in details in a previous study from our laboratory<sup>39</sup>.

**Trypsin digestion, MALDI-TOF MS analysis and sequence analysis.** Protein spots were excised from the 2-D gel. Removal of silver ions in gels, trypsin digestion and peptide extraction were performed as described previously<sup>39,40</sup>. In brief, after the peptides were cocrystallized with CHCA (alpha cyano-4-hydroxy cinnamic acid) by evaporating organic solvents, tryptic-digested peptide masses were measured using a MALDI-TOF-TOF mass spectrometer (ABI4700 System, USA). All mass spectra were recorded in positive reflector mode and generated by accumulating data from 1000 laser shots. The following threshold criteria and settings were used: detected mass range of 700–3200 Da (optimal resolution for the quality of 1500 Da), using a standard peptide mixture [des-Arg1-Bradykinin Mr904.468, Angiotensin I Mr1296.685, Glu-1-Fibrinopeptide B Mr1570.677, ACTH (1–17) Mr2093.087, ACTH(18–39) Mr2465.199; ACTH (7–38) Mr3657.929] as an external standard calibration, with laser frequency of 50 Hz, repetition rate of 200 Hz, UV wavelength of 355 nm, and accelerated voltage of 20,000 V. Peptide mass fingerprint data were matched to the comprehensive non-redundant sequence database NCBIInr using Profound program under 50 ppm mass tolerance.

Data was processed via the Data Explorer software and proteins were unambiguously identified by searching against the database NCBIInr using the MASCOT software search engine (<http://www.matrixscience.com/cgi/searchform.pl?FORMVER=2&SEARCH=MIS>). The search parameters were as follows: (1) peptide quality of 800–4000 Da, mass tolerance for the fragment ion of 0.25 Da; (2) a minimum of seven matching peptides; (3) one missed cleavage; (4) taxonomy: *L. enzymogenes* (bacterium); and (5) allowed modifications, alkylation of cysteine by carbamidomethylation of Cys (complete) acetylation of the N-terminus and oxidation of methionine (partial).



**Figure 4. TBDR7 modulated the HSAF production in *L. enzymogenes*.** (A) RT-PCR assay validated the transcription of *TBDR7* in the corresponding transformed strain. The size of expected DNA fragment was 388 bp. The gene, *16S rRNA* (abbreviation for *16s*) was used as an internal control, as described previously<sup>22</sup>. (B) Quantification of the HSAF level from the complemented strain of the *TBDR7* deletion mutant. (C) Growth curves of the wild type and *TBDR7* mutant in 1/10 TSB medium. (D) Mutation of *TBDR7* almost entirely shut down the transcription of the key HSAF biosynthetic gene, *hsaf pks/nrps*. OH11, the wild-type strain of *L. enzymogenes*;  $\Delta$ *TBDR7*, the *TBDR7* mutant;  $\Delta$ *TBDR7* (pBRR),  $\Delta$ *TBDR7* containing an empty vector;  $\Delta$ *TBDR7* (*TBDR7*), the complemented strain of  $\Delta$ *TBDR7*;  $\Delta$ *TBDR7* (*TBDR7*-V74A),  $\Delta$ *TBDR7* containing the plasmid-born *TBDR7*, where the amino acid Val74 within the TonB-box region was substituted by Ala74. Three replicates for each treatment were used, and the experiment was repeated three times. Vertical bars represent standard errors. The asterisk above the bars indicate a significant difference between the wild-type strain OH11 and the tested strains (\* $p < 0.05$ ).

Moreover, in order to evaluate protein identification, we considered the percentage of sequence coverage, the observation of distribution of matching peptides (authentic hit is often characterized by peptides that are adjacent to one another in the sequence and that overlap), the distribution of error (distributed around zero), the gap in probability and score distribution from the first to other candidate; only matches with over 90% sequence identity and a maximum  $e$ -value of  $10^{-10}$  were considered. Domain and functional analyzing of proteins identified was carried out using NCBI (<http://www.ncbi.nlm.nih.gov/>) and Pfam (<http://pfam.sanger.ac.uk/>).

**In-frame deletion, complementation and site-directed mutagenesis.** The protocol of gene in-frame deletion in *L. enzymogenes* was utilized as described previously<sup>41</sup>. In brief, two flanking regions of target gene were generated by PCR using a set of primer pairs (Supplementary Table S1), and cloned into the corresponding sites of the suicide vector pEX18Gm (Table 2). The final constructs were transformed into the wild-type OH11 or its derivatives by electroporation. Then *Lysobacter* transformants were selected on LB plates without sucrose and with Km (100  $\mu$ g/ml) and Gm (150  $\mu$ g/ml). The positive colonies were plated on LB plates containing 1/10 (w/v) sucrose and Km (100  $\mu$ g/ml) to select for resolution of the construct by a second cross-over event. The resultant mutants were confirmed by a PCR method (Supplementary Table S2). For generation of complemented strains, the target gene, *TBDR7* together with its native promoter region (Supplementary Table S1) was amplified from strain OH11 and cloned into the broad-host vector pBRR1-MCS5 (Table 2). The final construct was transformed into the *TBDR7*-deletion mutant for complementation. The expression of *TBDR7* in the transformed strains was determined by reverse transcription PCR (RT-PCR), which was described as below. Site-directed mutagenesis of the TonB-box region of *TBDR7* was carried out by using the Fast Mutagenesis System kit (FM111-01, Transgen Biotech, China) according to the product manufacturer. The point-mutated DNA was validated by sequencing.

**HSAF extraction and quantification.** Extraction and HPLC (High-Performance Liquid Chromatography) based quantification of HSAF from various *Lysobacter* strains was performed as described previously<sup>22,25,27</sup>. Three replicates for each treatment were used, and the experiment was performed three times.

**Growth determination.** Various *Lysobacter* strains were cultured in LB liquid at 28 °C overnight. Then 500 µl of the overnight culture for each strain was transferred into the fresh 1/10 TSB broth (50 ml) to grow until the OD<sub>600</sub> reached to 1.0 (the logarithmic phase of growth). Next, 1 ml of each culture was transferred again into the fresh 1/10 TSB broth (50 ml) to start the detection of growth curve. All inoculation broths were grown at 28 °C with shaking at 200 rpm, and the OD<sub>600</sub> value was determined every 2 h until bacterial growth reached the stationary phase. Each sample involves three technical replicates and the experiment was performed three times.

**RNA extraction, qRT-PCR and RT-PCR.** The wild-type OH11 and its derivatives, including the *TBDR7* deletion mutant, OH11 containing an empty vector and the *lesR* overexpression strain, were grown on 1/10 TSB. Cells of the wild-type OH11 and the *TBDR7* mutant were collected at the time point 10 h and 11 h, respectively, corresponding to the same cell density (OD<sub>600</sub> = 1.0). Similarly, OH11 containing the empty vector and the *lesR* overexpression strain were collected at the time point 12.5 h (OD<sub>600</sub> = 1.0). Then the total RNA was extracted from the collected cells of each strain using a kit from OMIGA Company (China). Next, qRT-PCR (quantitative RT-PCR) and RT-PCR assays, including cDNA synthesis and PCR amplification were performed as described previously<sup>22,29</sup>. Primer sequences used in this assay were listed in Supplementary Table S1.

**Data submission.** The sequence data of the present study has been submitted to the NCBI Genbank. The details were provided in Table 1.

**Data analysis.** All analyses were conducted using SPSS 14.0 (SPSS Inc., Chicago, IL, USA). The hypothesis test of percentages (*t*-test, \**p* < 0.05) was used to determine significant differences in production of antibiotic metabolite and gene expression between the control and treatment of the present study.

## References

1. Ferguson, A. D. *et al.* Siderophore-mediated iron transport: crystal structure of FhuA with bound lipopolysaccharide. *Science* **282**, 2215–2220 (1998).
2. Buchanan, S. K. *et al.* Crystal structure of the outer membrane active transporter FepA from *Escherichia coli*. *Nat. Struct. Biol.* **6**, 56–63 (1999).
3. Ferguson, A. D. *et al.* Structural basis of gating by the outer membrane transporter FecA. *Science* **295**, 1715–1719 (2002).
4. Moeck, G. S. & Coulton, J. W. TonB-dependent iron acquisition: mechanisms of siderophore-mediated active transport. *Mol. Microbiol.* **28**, 675–681 (1998).
5. Postle, K. & Kadner, R. J. Touch and go: tying TonB to transport. *Mol. Microbiol.* **49**, 869–882 (2003).
6. Cadieux, N. & Kadner, R. J. Site-directed disulfide bonding reveals an interaction site between energy-coupling protein TonB and BtuB, the outer membrane cobalamin transporter. *Proc. Nat. Acad. Sci.* **96**, 10673–10678 (1999).
7. Ferguson, A. D. & Deisenhofer, J. Metal import through microbial membranes. *Cell* **116**, 15–24. (2004).
8. Tauseef, I. *et al.* Influence of the combination and phase variation status of the haemoglobin receptors HmbR and HpuAB on meningococcal virulence. *Microbiology* **157**, 1446–1456 (2011).
9. Blavillain, S. *et al.* Plant carbohydrate scavenging through TonB-Dependent receptors: A feature shared by phytopathogenic and aAquatic bacteria. *PLoS ONE* **2**, e224 (2007).
10. Gonzalez, J. F., Myers, M. P. & Venturi, V. The inter-kingdom solo OryR regulator of *Xanthomonas oryzae* is important for motility. *Mol. Plant. Pathol.* **14**, 211–221 (2013).
11. Fuqua, C., Parsek, M. R. & Greenberg, E. P. Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu. Rev. Genet.* **35**, 439–468 (2001).
12. Volf, J. *et al.* Role of SdiA in *Salmonella enterica* serovar *Typhimurium* physiology and virulence. *Arch. Microbiol.* **178**, 94–101 (2002).
13. Ahmer, B. M. Cell-to-cell signalling in *Escherichia coli* and *Salmonella enterica*. *Mol. Microbiol.* **52**, 933–945 (2004).
14. Ferluga, S. *et al.* A LuxR homologue of *Xanthomonas oryzae* pv. *oryzae* is required for optimal rice virulence. *Mol. Plant. Pathol.* **8**, 529–538 (2007).
15. Zhang, L. *et al.* A proline iminopeptidase gene upregulated in planta by a LuxR homologue is essential for pathogenicity of *Xanthomonas campestris* pv. *campestris*. *Mol. Microbiol.* **65**, 121–136 (2007).
16. Lee, J. *et al.* Reconfiguring the quorum-sensing regulator SdiA of *Escherichia coli* to control biofilm formation via indole and N-acyl homoserine lactones. *Appl. Environ. Microbiol.* **75**, 1703–1716 (2009).
17. Subramoni, S. *et al.* Bacterial subfamily of LuxR regulators that respond to plant compounds. *Appl. Environ. Microbiol.* **77**, 4579–4588 (2011).
18. Chatnaparat, T. *et al.* XagR, a LuxR homolog, contributes to the virulence of *Xanthomonas axonopodis* pv. *glycines* to soybean. *Mol. Plant. Microbe. Interact.* **25**, 1104–1117 (2012).
19. Christensen, P. & Cook, F. D. *Lysobacter*, a new genus of nonfruiting, gliding bacteria with high base ratio. *Int. J. Syst. Bacteriol.* **28**, 367–393 (1978).
20. Xie, Y. X. *et al.* Bioactive natural products from *Lysobacter*. *Nat. Prod. Rep.* **29**, 1277–1287 (2012).
21. Zhou, X. *et al.* PilG is involved in the regulation of twitching motility and antifungal antibiotic biosynthesis in the biological control agent *Lysobacter enzymogenes*. *Phytopathology* **105**, 1318–1324 (2015).
22. Qian, G. L. *et al.* *Lysobacter enzymogenes* uses two distinct cell-cell signaling systems for differential regulation of secondary-metabolite biosynthesis and colony morphology. *Appl. Environ. Microbiol.* **79**, 6604–6616 (2013a).
23. Zhang, Z. & Yuen, G. Y. Biological control of *Bipolaris sorokiniana* on tall fescue by *Stenotrophomonas maltophilia* strain C3. *Phytopathology* **89**, 817–822 (1999).
24. Yuen, G. Y. & Zhang, Z. Control of brown patch using the bacterium *Stenotrophomonas maltophilia* C3 and culture fluid. *Int. Turfgrass. Soc. Res.* **9**, 742–747 (2001).
25. Yu, F. G. *et al.* Structure and biosynthesis of heat-stable antifungal factor (HSAF), a broad-spectrum antimycotic with a novel mode of action. *Antimicrob. Agents. Ch.* **51**, 64–72 (2007).
26. Li, S. J. *et al.* Distinct ceramide synthases regulate polarized growth in the filamentous fungus *Aspergillus nidulans*. *Mol. Biol. Cell.* **17**, 1218–1227 (2006).
27. Lou, L. L. *et al.* Biosynthesis of HSAF, a tetramic acid-containing macrolactam from *Lysobacter enzymogenes*. *J. Am. Chem. Soc.* **133**, 643–645 (2011).

28. Li, Y. Y. *et al.* Iterative assembly of two separate polyketide chains by the same single-module bacterial polyketide synthase in the biosynthesis of HSAF. *Angew. Chem. Int. Ed. Engl.* **53**, 7524–7530 (2014).
29. Qian, G. L. *et al.* Roles of a solo LuxR in the biological control agent *Lysobacter enzymogenes* strain OH11. *Phytopathology* **104**, 224–231 (2014).
30. Schöffler, H. & Braun, V. Transport across the outer membrane of *Escherichia coli* K12 via the FhuA receptor is regulated by the TonB protein of the cytoplasmic membrane. *Mol. Gen. Genet.* **217**, 378–383 (1989).
31. Peacock, R. S. *et al.* The solution structure of the C-terminal domain of TonB and interaction studies with TonB Box peptides. *J. Mol. Biol.* **345**, 1185–1197 (2005).
32. Braun, V. & Endriss, F. Energy-coupled outer membrane transport proteins and regulatory proteins. *Biometals* **20**, 219–231 (2007).
33. Ferguson, A. D. *et al.* Signal transduction pathway of TonB-dependent transporters. *Proc. Nat. Acad. Sci.* **104**, 513–518 (2007).
34. Gudmundsdottir, A. *et al.* Point mutations in a conserved region (TonB box) of *Escherichia coli* outer membrane protein BtuB affect vitamin B12 transport. *J. Bacteriol.* **171**, 6526–6533 (1989).
35. Hu, Y. H. *et al.* A TonB-dependent outer membrane receptor of *Pseudomonas fluorescens*: virulence and vaccine potential. *Arch. Microbiol.* **194**, 795–802 (2012).
36. O'Sullivan, D. J. & O'Gara, F. Traits of fluorescent *Pseudomonas* spp. involved in suppression of plant root pathogens. *Microbiol. Rev.* **56**, 662–676 (1992).
37. Ferluga, S. & Venturi, V. OryR is a LuxR-family protein involved in interkingdom signaling between pathogenic *Xanthomonas oryzae* pv. *oryzae* and rice. *J. Bacteriol.* **191**, 890–897 (2009).
38. Watt, S. A. *et al.* Comprehensive analysis of the extracellular proteins from *Xanthomonas campestris* pv. *campestris* B100. *Proteomics* **5**, 153–167 (2005).
39. Qian, G. L. *et al.* Proteomic analysis reveals novel extracellular virulence-associated proteins and functions regulated by the diffusible signal factor (DSF) in *Xanthomonas oryzae* pv. *oryzicola*. *J. Proteome. Res.* **12**, 3327–3341 (2013b).
40. Joo, W. A. *et al.* Monitoring protein expression by proteomics: Human plasma exposed to benzene. *Proteomics* **3**, 2402–2411 (2003).
41. Qian, G. L. *et al.* Selection of available suicide vectors for gene mutagenesis using *chiA* (a chitinase encoding gene) as a new reporter and primary functional analysis of *chiA* in *Lysobacter enzymogenes* strain OH11. *World. J. Microbiol. Biotechnol.* **28**, 549–557 (2012).
42. Hoang, T. T. *et al.* A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**, 77–78 (1998).
43. Kovach, M. E. *et al.* Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **166**, 175–176 (1995).

## Acknowledgements

Dr. Guoliang Qian is a visiting scholar in Dr. Mark Gomelsky's laboratory of University of Wyoming, USA, supported by China Scholarship Council. This study was supported by Jiangsu Agriculture Science and Technology Innovation Fund [CX(15)1023], National Basic Research (973) program of China (2015CB150600), National Natural Science Foundation of China (31371981; 31572046; 31329005), the grants from Jiangsu Academy of Agricultural Sciences and Jiangsu Province (ZX(15)1006; BE2014386; BE2015354), Program for New Century Excellent Talents in University of Ministry of Education of China (NCET-13-0863), Fundamental Research Funds for the Central Universities (KYTZ201403) and Special Fund for Agro-Scientific Research in the Public Interest (No. 201303015). Research in the Du lab is supported in part by NIH R01AI097260 and a UNL Redox Biology Center pilot grant. Dr. Chou is supported by the Ministry of Education, Taiwan, under the ATU plan, and by the National Science Council, Taiwan (grants 102-2113-M-005-006-MY3).

## Author Contributions

R.W., H.X. and G.Q. conceived and designed experiments. R.W., H.X. and Y.L. carried out experiments. R.W., H.X., G.Q., F.L. and S.C. analyzed data and prepared figures. R.W., H.X., Y.S., F.L. and G.Q. wrote the manuscript. L.D., Y.S. and S.C. reviewed and revised the manuscript. All authors reviewed the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Wang, R. *et al.* A TonB-dependent receptor regulates antifungal HSAF biosynthesis in *Lysobacter*. *Sci. Rep.* **6**, 26881; doi: 10.1038/srep26881 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

1 **Supplementary Materials**

2 **A TonB-dependent receptor regulates antifungal HSAF biosynthesis in *Lysobacter***

3 Ruping Wang<sup>a</sup>, Huiyong Xu<sup>a</sup>, Liangcheng Du<sup>c</sup>, Shan-Ho Chou<sup>d</sup>, Hongxia Liu<sup>a</sup>, Youzhou Liu<sup>b</sup>,  
4 Fengquan Liu<sup>\*b</sup> and Guoliang Qian<sup>\*a</sup>

5 <sup>a</sup> College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China/Key Laboratory of Integrated  
6 Management of Crop Diseases and Pests (Nanjing Agricultural University), Ministry of Education

7 <sup>b</sup> Institute of Plant Protection, Jiangsu Academy of Agricultural Science, Nanjing 210014, P.R. China

8 <sup>c</sup> Department of Chemistry, University of Nebraska-Lincoln, Lincoln, Nebraska 68588, United States

9 <sup>d</sup> Institute of Biochemistry, and NCHU Agricultural Biotechnology Center, National Chung Hsing University,  
10 Taichung, Taiwan, ROC

11 <sup>\*</sup>To whom correspondence should be addressed. Tel: +86-25-84396109. Fax: +86-25-84396109. E-mail:

12 [glqian@njau.edu.cn](mailto:glqian@njau.edu.cn) or [fqliu20011@sina.com](mailto:fqliu20011@sina.com)

13

14

## Outlines

15 **Figure S1.** 2-D gel proteome analysis of the *lesR* overexpression strain.

16 **Figure S2.** The genomic organization of each *lesR*-controlled TBDR coding genes in *L.*  
17 *enzymogenes*.

18 **Figure S3.** Quantification of the HSAF yield from the *tonB* mutant of *L. enzymogenes*.


19 **Figure S4.** Overexpression of *lesR* caused an approximately twice increase on the transcription of  
20 *TBDR7* in comparison to the control strain.

21 **Table S1.** Primers used in this study

22 **Table S2.** Mutant confirmation by PCR in this study

23

24



25

26

27

28 **Figure S1. 2-D gel proteome analysis of the *lesR* overexpression strain.** The up-expressed


29 protein spots with the threshold ratio of 1.5 were numbered in green either in OH11 (pBBR) or

30 OH11 (*lesR*) compared to each other. Detailed information of the identified *lesR*-controlled31 proteins was provided **Table 2**. OH11(pBBR), the control strain, representing the wild-type OH1132 of *L. enzymogenes* containing an empty expressing vector; OH11(*lesR*), the *lesR* overexpression

33 strain.

34

35



36

37 **Figure S2. The genomic organization of each *lesR*-controlled TBDR coding genes in *L.***

38 ***enzymogenes*.** Each TBDR was colored, whereas other genes were shown in white arrows. The


39 predicted gene products were provided below each arrow. The scale indicates the amino-acid size

40 was identical to 100 aa.

41

42





43

44

45 **Figure S3. Quantification of the HSAF yield from the *tonB* mutant of *L. enzymogenes*.**

46 Mutation of *tonB* (GU121672) almost completely abolished the HSAF production. Three

47 replicates for each treatment were used, and the experiment was repeated three times. Vertical bars

48 represent standard errors. The asterisk above the bars indicate a significant difference between the

49 wild-type strain OH11 and the tested strain (\* $p < 0.05$ ).

50

51

52

53

54

55

56

57

58

59

60


61

62

63

64

65



66

67 **Figure S4. Overexpression of *lesR* caused an approximately twice increase on the**  
68 **transcription of *TBDR7* in comparison to the control strain.** The control strain is OH11  
69 (pBBR), representing the wide-type OH11 containing an empty vector; OH11 (*lesR*), the *lesR*  
70 overexpressing strain. Each column indicates the mean of three biologically independent  
71 quantitative RT-PCR experiments. Vertical bars represent standard errors. The asterisk above the  
72 bars indicate a significant difference between the wild-type strain OH11 and the tested strains (\**p*  
73 < 0.05).

#### 74 **Supplementary reference**

- 75 1. Qian, G. L. *et al.* *Lysobacter enzymogenes* uses two distinct cell-cell signaling systems for  
76 differential regulation of secondary-metabolite biosynthesis and colony morphology. *Appl.*  
77 *Environ. Microbiol.* **79**, 6604-6616 (2013a).
- 78 2. Schöffler, H. & Braun, V. Transport across the outer membrane of *Escherichia coli* K12 via the  
79 FhuA receptor is regulated by the TonB protein of the cytoplasmic membrane. *Mol. Gen.*  
80 *Genet.* **217**, 378-383 (1989).

81

82

83

84

85  
86  
87  
88  
89

**Table S1 Primers used in this study**

Primer	Sequence <sup>a</sup>	Purpose	Source
<i>TBDR1-1F</i>	5'-GGGGTACCCGACTCCGCCACGCATTCCA-3'	To amplify a 314-bp upstream homologue arm of <i>TBDR1</i>	This study
<i>TBDR1-1R</i>	5'-CCCAAGCTTTCGCCTTCCTTGGTGATGTTGA-3'		
<i>TBDR1-2F</i>	5'-CCCAAGCTTCCGGCTGGAACACTACAACACTACCT-3'	To amplify a 1054-bp downstream homologue arm of <i>TBDR1</i>	This study
<i>TBDR1-2R</i>	5'-GCTCTAGA CGGCGGAAATCACGAAAA-3'		
<i>TBDR2-1F</i>	5'-CCC <del>CGT</del> ACCGAGGGCGGTGCGGAGTTC-3'	To amplify a 788-bp upstream homologue arm of <i>TBDR2</i>	This study
<i>TBDR2-1R</i>	5'-CCCTCTAGACAGCAGCAGGCTGGCGAC-3'		
<i>TBDR2-2F</i>	5'-CCCTCTAGACGCAACATCACCAACGAGAA GA-3'	To amplify a 999-bp downstream homologue arm of <i>TBDR2</i>	This study
<i>TBDR2-2R</i>	5'-CGGAATTCAGCGAGTACAGCAGCCACAGC-3'		
<i>TBDR4-1F</i>	5'-CGGAATTCGGTGCCAGTGGAAGGTGTTGA G-3'	To amplify a 1062-bp upstream homologue arm of <i>TBDR4</i>	This study
<i>TBDR4-1R</i>	5'-CCCAAGCTTAGCGTGGCCTGGAACATCG-3'		
<i>TBDR4-2F</i>	5'-CCCAAGCTTCGTTGGTGTGGCGAACTTGC-3'	To amplify a 985-bp downstream homologue arm of <i>TBDR4</i>	This study
<i>TBDR4-2R</i>	5'-GCTCTAGAGCGGACATGCTGCGTTTGGT-3'		
<i>TBDR7-1F</i>	5'-GGGGTACCCGGCTACTCGCACATCCACG-3'	To amplify a 503-bp upstream homologue arm of <i>TBDR7</i>	This study
<i>TBDR7-1R</i>	5'-GCTCTAGACGCCAACACCTTCCACGACG-3'		
<i>TBDR7-2F</i>	5'-GCTCTAGACGGTGGCGGTGATGTTCTGG-3'	To amplify a 554-bp downstream homologue arm of <i>TBDR7</i>	This study
<i>TBDR7-2R</i>	5'-CGGGATCCAACGCACGAACCCGCATCCG-3'		
<i>TBDR8-1F</i>	5'-GGGGTACCAACGGCGACAAGGACGAAGG-3'	To amplify a 806-bp upstream homologue arm of <i>TBDR8</i>	This study
<i>TBDR8-1R</i>	5'-CCCAAGCTTCGTTGCCGCCGTTGTTGTAG-3'		
<i>TBDR8-2F</i>	5'-CCCAAGCTTGGCAACTGGTCGGAGGGCTT-3'	To amplify a 681-bp downstream homologue arm of <i>TBDR8</i>	This study
<i>TBDR8-2R</i>	5'-GCTCTAGAGACCGAGCCCAGTTCACAGT-3'		
<i>TBDR9-1F</i>	5'-GGGGTACCAGTTCGTCGTGTCGCCGCTC-3'	To amplify a 739-bp upstream homologue arm of <i>TBDR9</i>	This study
<i>TBDR9-1R</i>	5'CCCAAGCTTGGGGTGTGTCGTCGCTCTG-3'		
<i>TBDR9-2F</i>	5'-CCCAAGCTTGCAAGAACTACAAGGTCGCC-3'	To amplify a 1052-bp downstream homologue arm of <i>TBDR9</i>	This study
<i>TBDR9-2R</i>	5'-GCTCTAGATTGCCACCGTCGTATTCTGT-3'		
<i>tonB-1F</i>	5'-ACGAATTC AACACCAGCGAGCAGTTGTT-3'	To amplify a 706-bp downstream homologue arm of <i>tonB</i>	This study
<i>tonB-1R</i>	5'-CCAAGCTTGTGCTGTTCGGCTTCAATTC-3'		
<i>tonB-2F</i>	5'-AAAAGCTTGTGCGCACAGCGTGTCGA-3'	To amplify a 738-bp downstream homologue arm of	This study

<i>tonB-2R</i>	5'-AA <u>TCTAGA</u> ATGCCGCCCTGGCGATCCG-3'	<i>tonB</i>	
<i>tonB-F</i>	5'-GATTTCGTCCCGCCGAAGCTG-3'	Mutant confirmation: to amplify a 991- and 250-bp DNA band from the wild-type OH11 and <i>tonB</i> -deletion mutant, respectively	This study
<i>tonB-R</i>	5'-TGATCAGCAGCACCAGCATC-3'		
<i>TBDR7-F</i>	5'-GCT <u>CTAGAG</u> GGTGTCTGGCACTGGTCCCGGC GAT-3'	To amplify 3266-bp fragment of <i>TBDR7</i> with its predicted promoter	This study
<i>TBDR7-R</i>	5'-GGGGTACCTCAGAAGCGCTGGGTGTACTT C-3'		
<i>TBDR7-Fm</i>	5'-TCTCGATCGCATCGAG <sup>a</sup> CGACCGGTTTCGC G-3'	To amplify pBBR- <i>TBDR7</i> (site directed mutagenesis), the box indicated the target point mutation (V74 was changed into A74)	This study
<i>TBDR7-Rm</i>	5'-GC <sup>a</sup> TCGATGCGATCGAGAGTGGTGGGGGT C-3'		
16S rDNA-1	5'-ACGGTCGCAAGACTGAAACT-3'	The internal control for RT-PCR and quantitative RT-PCR	1
16S rDNA-2	5'-AAGGCACCAATCCATCTCTG-3'		
q <i>pks/nrps</i> -F	5'-CATCACATCATCTCCGATGC-3'	To determine the transcriptional level of the gene <i>pks/nrps</i>	1
q <i>pks/nrps</i> -R	5'-CAGTTCCACCTTCTCCTTGC-3'		
q <i>TBDR7</i> -F	5'-ACAACCAGAACAAGGGCAAC-3'	To determine the transcriptional level of the gene <i>TBDR7</i>	This study
q <i>TBDR7</i> -R	5'-GCTGAGGTTGAGGCTGTAGA-3'		

<sup>a</sup> Restricted digestion enzyme site was underlined.

90  
91

Table S2 Mutant confirmation by PCR in this study

Mutant	Selected primers <sup>a</sup>	Expected size from wild type	Expected size from mutant	PCR validation <sup>b</sup>
$\Delta TBDR1$	<i>TBDR1</i> -1F/2R	3629 bp	1368 bp	
$\Delta TBDR2$	<i>TBDR2</i> -1F/2R	3944 bp	1787 bp	
$\Delta TBDR4$	<i>TBDR4</i> -1F/2R	4267 bp	2047 bp	
$\Delta TBDR7$	<i>TBDR7</i> -1F/2R	2604 bp	1057 bp	
$\Delta TBDR8$	<i>TBDR8</i> -1F/2R	3016 bp	1487bp	
$\Delta TBDR9$	<i>TBDR9</i> -1F/2R	2789 bp	1791 bp	
$\Delta TBDR1\&7$	<i>TBDR1</i> -1F/2R	3629 bp	1368 bp	
<i>tonB</i>	<i>tonB</i> -F/R	991bp	250bp	

93 <sup>a</sup> Primers sequence used here were provided in Supplementary Table S1.

94 <sup>b</sup> Red arrow and green box indicated the expected size from the wild-type OH11 or deletion mutant amplified by  
 95 selected primers, respectively; + and – means the positive control (the genomic DNA of strain OH11) and negative  
 96 control (ddH<sub>2</sub>O), respectively.