

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Center for Plant
Science Innovation

Plant Science Innovation, Center for

2015

Modifications of membrane lipids in response to wounding of *Arabidopsis thaliana* leaves

Hieu Sy Vu

University of Nebraska-Lincoln

Rebecca Roston

University of Nebraska- Lincoln, rroston@unl.edu

Sunitha Shiva

University of Nebraska- Lincoln

Manhoi Hur

Iowa State University

Eve Syrkin Wurtele

Iowa State University

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/plantscifacpub>

 Part of the [Plant Biology Commons](#), [Plant Breeding and Genetics Commons](#), and the [Plant Pathology Commons](#)

Vu, Hieu Sy; Roston, Rebecca; Shiva, Sunitha; Hur, Manhoi; Wurtele, Eve Syrkin; Wang, Xuemin; Shah, Jyoti; and Welti, Ruth, "Modifications of membrane lipids in response to wounding of *Arabidopsis thaliana* leaves" (2015). *Faculty Publications from the Center for Plant Science Innovation*. 126.

<http://digitalcommons.unl.edu/plantscifacpub/126>

This Article is brought to you for free and open access by the Plant Science Innovation, Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Center for Plant Science Innovation by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Hieu Sy Vu, Rebecca Roston, Sunitha Shiva, Manhoi Hur, Eve Syrkin Wurtele, Xuemin Wang, Jyoti Shah, and Ruth Welti

Plant Signaling & Behavior

Plant Signal Behav. 10(9): e1056422

Modifications of membrane lipids in response to wounding of *Arabidopsis thaliana* leaves

Hieu Sy Vu^{1,2}, Rebecca Roston², Sunitha Shiva¹, Manhoi Hur³, Eve Syrkin Wurtele³, Xuemin Wang⁴, Jyoti Shah⁵, Ruth Welti^{1*}

1. Kansas Lipidomics Research Center; Division of Biology; Kansas State University; Manhattan, KS USA

2. Department of Biochemistry and Center for Plant Science Innovation; University of Nebraska-Lincoln; Lincoln, NE USA

3. Department of Genetics, Development, and Cell Biology; Iowa State University; Ames, IA USA

4. Department of Biology; University of Missouri; Donald Danforth Plant Science Center; St. Louis, MO USA

5. Department of Biological Sciences; University of North Texas; Denton, TX USA

*. Correspondence to: Ruth Welti; Email: welti@ksu.edu

Addendum to: Vu HS, Shiva S, Roth MR, Tamura P, Zheng L, Li M, Sarowar S, Honey S, McElhiney D, Hinkes P, Seib L, Williams TD, Gadbury G, Wang X, Shah J, Welti R. Lipid changes after leaf wounding in *Arabidopsis thaliana*: Expanded lipidomic data form the basis for lipid co-occurrence analysis. *Plant J.* 2014; 80, 728–43.

© 2015 Taylor & Francis Group, LLC

DOI: 10.1080/15592324.2015.1056422

Published online: 7 August 2015

See the addendum "[Lipid changes after leaf wounding in *Arabidopsis thaliana*: expanded lipidomic data form the basis for lipid co-occurrence analysis.](#)" in *Plant J*, volume 80 on page 728.

Abstract

Mechanical wounding of *Arabidopsis thaliana* leaves results in modifications of most membrane lipids within 6 hours. Here, we discuss the lipid changes, their underlying biochemistry, and possible relationships among activated pathways. New evidence is presented supporting the role of the processive galactosylating enzyme SENSITIVE TO FREEZING2 in the wounding response.

Abbreviations

ASG Acyl sterol glucoside
acMGDG Acylated monogalactosyldiacylglycerol
acPG Acylated phosphatidylglycerol
DAD1 Defective in anther dehiscence 1
DAG Diacylglycerol
DGDG Digalactosyldiacylglycerol
DGMG Digalactosylmonoacylglycerol
dnOPDA Dinor-oxophytodienoyl
GL Galactolipid
Glc Glucoside
GlcCer Glycosylceramide
GIPC Glycosylinositolphosphoceramide
LPC Lysophosphatidylcholine
LPE Lysophosphatidyl-ethanolamine
MGDG Monogalatosyldiacylglycerol
oxDGDG Oxidized digalactosyl- diacylglycerol
oxMGDG Oxidized monogalactosyldiacylglycerol
OPDA Oxophytodienoyl
PA Phosphatidic acid

PC	Phosphatidylcholine
PE	Phosphatidylethanolamine
PG	Phosphatidylglycerol
PG	Phosphatidylglycerol
PI	Phosphatidylinositol
PS	Phosphatidylserine
PLA2	Phospholipase A2
PLD	Phospholipase D
PL	Phospholipid
PDAT1	Phospholipid:diacylglycerol acyltransferase 1
SFR2	SENSITIVE TO FREEZING 2
SG	Sterol glucoside
SQDG	Sulfoquinovosyldiacylglycerol
TeGDG	Tetra-galactosyldiacylglycerol
TAG	Triacylglycerol
TrGDG	Tri-galactosyldiacylglycerol.


Wounding (i.e., mechanical damage) of plant tissues can be caused by wind, rain, or hail or can occur during insect attack, grazing, or pathogen infection. In the laboratory, mechanical wounding of plants provides a controlled experimental treatment to elicit chemical responses, including changes in gene expression, protein expression, and metabolite composition. Changes in lipids occur rapidly, with many examples of changes observed in 5 min or less.¹⁻⁴

In a recent study,⁵ complex lipids were analyzed in leaves of unwounded plants and mechanically wounded plants, sampled 45 min and 6 h after wounding, with each treatment replicated 31 times. The levels of 254 out of

the total of 264 complex lipids detected changed significantly in response to mechanical wounding. Here, the observed changes in complex lipids in wound response are examined further and considered in relation to known and proposed lipid metabolic pathways.

Overview

Vu et al.⁵ found that, after wounding, the total levels of membrane and membrane-derived lipids dropped. The mass spectral intensity of all measured leaf lipids fell 6% from the level in unwounded plants at 45 min after wounding and a further 22% at 6 h after wounding ([Fig. 1A](#)). The drop in total lipid level was mainly due to decreases in the levels of normal-chain, diacyl polar membrane lipids including phospholipids (PLs), sulfolipid (sulfoquinovosyldiacylglycerol, SQDG), and galactolipids (GLs) ([Fig. 1A and B](#)). On the other hand, many less abundant lipids increased ([Fig. 1A, C and D](#)). Lipids with increased levels include molecular species with oxidized fatty acyl chains, head-group acylated membrane lipids, monoacyl polar lipids, phosphatidic acids, oligogalactolipids, sterol glucosides, acyl sterol glucosides, and triacylglycerols. Additionally there were subtle increases in lipid unsaturation in PC and PA. In the following sections, we detail and discuss these changes and discuss possibilities for the underlying biochemistry ([Fig. 2](#)).



[View larger version](#)

Figure 1. Changes in leaf lipid classes following pressure wounding on leaves of wild-type *Arabidopsis thaliana* across the mid-vein with a hemostat. Leaves were harvested from unwounded or wounded plants 45 min and 6 h after wounding.⁵ This figure is based on data in ref. [5](#). **(A)** Normalized mass spectral intensity for lipids in the panels **(B, C, and D)**. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. Values over the bars indicate the percent of the total lipids represented by the bar. **(B)** Amount of structural (membrane) lipid classes after wounding relative to unwounded level for each class. **(C)** Amounts of minor plastidic lipid classes relative to the unwounded level for each class. **(D)** Amounts of minor extraplastidic lipid classes relative to the unwounded level for each class. * indicates significant changes compared to unwounded samples, T-test, $P < 0.001$, $n = 31$. Error bars are standard deviation.



[View larger version](#)

Figure 2. Proposed reactions of membrane lipids in the wounding response in *Arabidopsis* leaves. **(a)** Lipid head-group acylation and lipolytic reactions producing monoacyl lipid molecular species; **(b)** galactosylation of lipids and other reactions producing DAG, PA, and TAG; **(c)** formation of sterol glucosides and acyl sterol glucosides; **(d)** increase of fully desaturated PC and PA fractions during the wounding response; and **(e)** oxidation of lipids in the wounding response. See text for details.

Glycerolipid head-group acylation and lipolytic reactions producing monoacyl lipid molecular species

Fatty acylation of the lipid head group of monogalactosyldiacylglycerol (MGDG) was originally described by Heinz.⁶ MGDG acylation occurs when a fatty acyl chain is transferred from digalactosyldiacylglycerol (DGDG) to the 6-position on the galactose of MGDG,^{7,8} producing an acylated MGDG and digalactosylmonoacylglycerol (DGMG) ([Fig. 2, block a](#)). More recently,

oxidized acylated MGDGs have been identified in Arabidopsis leaves following bacterial infection or wounding; the most common molecular species is an MGDG with oxidized fatty acids in the 1- and 2- positions (oxophytodienoyl/dinor-oxophytodienoyl; OPDA/dnOPDA), acylated on galactose with OPDA. This species was given the common name “Arabidopside E”.^{9,10} It is now apparent that a wide variety of acylated MGDG molecular species and DGMGs are produced upon wounding of Arabidopsis leaves ([Fig. 1C and D](#)).^{5,8,11-13} Additionally phosphatidylglycerol (PG) with an acyl chain on the terminal glycerol and small amounts of head-group acylated DGDGs are also produced in Arabidopsis leaves when they are wounded ([Fig. 1C](#)).^{5,11,13} DGMG may also be produced by the acylhydrolases, such as DAD1 and DAD1-like enzymes, localized in chloroplasts and capable of acting on galactolipids.¹⁴


In addition to DGMG, mono-acyl glycerolipids produced in response to wounding include lysophosphatidylcholines, which increase more than 2 fold at 45 min after wounding, and lysophosphatidylethanolamines ([Fig. 1D](#)). These may be formed from PC and PE by the action of acylhydrolases such as the patatin-like PLA2s (e.g., refs. [15-16](#)).

Galactosylation of galactolipids and other reactions producing DAG, PA, and TAG

In response to freezing, tri- and tetra-galactosyldiacylglycerol (TrGDG and TeGDG) are formed by the processive addition of galactose moieties from available MGDG molecular species, forming first DGDG, and then TrGDG and TeGDG by the enzyme SENSITIVE TO FREEZING 2 (SFR2) ([Fig. 2, block b](#)).^{17,18} The side product, DAG, can be converted to TAG by an

acyltransferase.¹⁷ DAG that has accumulated in the chloroplast envelopes during non-freezing conditions can enter PL metabolism, possibly through action of a DAG kinase.¹⁹⁻²⁰ Following wounding, the tight co-occurrence of PA(34:6) with TrGDG(34:6) and TeGDG(34:6), and parallel reduction of MGDG(34:6),⁵ is consistent with this pathway of processive galactosylation and PA(34:6) as a by-product.

Because previous data suggested that the presence of SFR2 is only critical to plant survival following freezing,²¹⁻²² the role of SFR2 in wounding was confirmed using an Arabidopsis line lacking SFR2, *sfr2-3*.¹⁷ The data are included as Supplemental Table 1. Data extracted from this data set and shown in [Figure 3](#) demonstrate that galactosylation during the wounding response is dependent on the presence of SFR2. In the absence of SFR2, TrGDG and TeGDG are not formed, and PA(34:6) is much reduced. Additionally, TAG(18:3/34:6), a plastid-derived molecular species containing 16:3 in the “34:6” moiety, is not formed during wounding in *sfr2-3* ([Fig. 3D](#)). These data indicate that wounding produces a measurable activation of SFR2 in leaves of wild-type plants, and confirms that PA is correlated with SFR2 activity, adding to the known effects of SFR2. Further investigation of the physical characteristics associated with the wounding suggested that tissue desiccation occurred around the wounding site. If SFR2 is responding to the desiccation, this would be consistent with observed production of oligogalactolipids during desiccation of the resurrection plant.²³



[View larger version](#)

Figure 3. Formation of TrGDG, TeGDG, PA(34:6), and 16:3-containing TAG is abolished in *sfr2-3* (SALK_106253) plants.

Leaves were harvested from unwounded or wounded plants 45 min after wounding as in ref. 5. In the indicated lipids, “34:6” is composed of 18:3 and 16:3 fatty acyl chains. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. ** indicates a significant change at $P < 0.01$, * indicates a significant change at $P < 0.05$, T-test, $n = 6$. Error bars are standard deviation.

Small increases in DAG levels are detectable during the leaf's response to wounding ([Fig. 1D](#)). Like PA, DAG can accumulate after wounding as a result of multiple reactions, including SFR2 acting on MGDG and phospholipase C acting on phospholipids. Production of PA via DAG by a phospholipase C-diacylglycerol kinase pathway has been detected in Arabidopsis in low-temperature stress.²⁴ PA is also formed from DAG by DAG kinase in plant-fungus interaction.²⁵ Formation of PA via DAG and DAG kinase was also shown to play roles in abscisic acid response and plant tolerance to drought and salt stresses.²⁶ Both DAG derived from extraplastidic phospholipids and DAG formed from MGDG are converted to triacylglycerols (TAGs) during the wounding response. This is likely to occur by the action of PDAT1.²⁰ Indeed, TAG levels increase over a prolonged period after wounding ([Fig. 1D](#)). The formation of TAG was recently demonstrated to be an essential intermediate in the delivery of plastidic fatty acids to peroxisomal β -oxidation, to maintain membrane lipid homeostasis in leaves.²⁰

In addition to the SFR2 and phospholipase C-diacylglycerol kinase pathways, PA can be produced by hydrolysis of phospholipids by

phospholipase D (PLD) during wounding stress.⁴ About half of the PA formed in the first 15 min after wounding can be attributed to the action of PLD α 1.⁴ PA species formed by PLD contain DAG moieties characteristic of extraplastidic phospholipids (e.g., 34:3, 34:2, 36:5, and 36:4), distinguishing them from the PA(34:6) derived from MGDG. Additionally, PAs derived from extraplastidic PLs and PA(34:6) from MGDG are formed with different kinetics; specifically, the maximal increase in PA-containing fatty acids from extraplastidic lipids occurs more quickly than maximal formation of PA(34:6) ([Fig. 1C and D](#)).

Formation of sterol glucosides and acyl sterol glucosides

Vu et al.⁵ identified increases in sterol glucosides and acyl sterol glucosides upon wounding ([Fig. 1](#) and [Fig. 2, block c](#)). Sterol glucosides probably are formed by UDP-Glc:sterol glycosyltransferase(s).²⁷ Acyl sterol glucosides are likely to be formed by acylation of sterol glucosides, but acylating enzymes acting on sterol glucosides are not identified.

Increase of fully desaturated PC and PA fractions during the wounding response

One of the more subtle changes detected in response to wounding was an increase in the unsaturation of the PC and PA pools, with fully desaturated 34:6 and 36:6 species particularly increased. “34:6” represents an 18:3/16:3 acyl chain combination, while “36:6” represents an 18:3/18:3 combination. Unsaturation indices and percentage of the PA and PC classes that can be attributed to these fully desaturated species are shown in [Table 1](#). In response to wounding, the unsaturation of PC and PA increases significantly. In particular, the proportion of PC(36:6) and PA(36:6) species as a% of the total head group class increases at 45 min after wounding, and

the proportion of PC(34:6), and PA(34:6), species increases strongly at 6 h after wounding. The increase in PC(36:6) and PA(36:6) may result from an increased amount of desaturase activity per PC molecule due to a decrease in the size of the PC pool ([Fig. 2, block d](#)). However, given that SFR2 and other activities are forming PL precursors from the highly desaturated plastid pool of MGDG, the formation of PC and PA species containing 16:3 likely arises from transfer of 16:3 from the plastids. The identity of the molecule transferred is not clear, but the strong incorporation of 16:3 into PA could suggest that PA or intact DAG is the transferred species.

Table 1.
PA and PC unsaturation is increased in wounded plants. In particular, this is due to increases in 36:6 and 34:6 molecular species. The unsaturation index is a measure of the average unsaturation of the class. The unsaturation index of a lipid species is calculated by multiplying the number of double bonds in the lipid by the fraction of that lipid in the lipid class divided by the number of acyl chains in the lipid (i.e., 2 for diacyl lipids). The unsaturation index of the class is the total of the unsaturation index of all lipid species in that class as percent of the total indicates the percentage of that class composed on the indicated lipid species. One-way ANOVA was performed to compare the unsaturation indices or lipid percentages among the three treatments. If a significant difference ($p < 0.001$; $n = 31$) was detected for a lipid, a Tukey post-hoc test was performed to identify the differing treatments. * indicates that the value in the wounded sample is significantly increased compared to the unwounded sample.

	Unwounded	45 min after wounding	6 h after wounding
PA class	1.00 ± 0.04	1.69* ± 0.01	1.66* ± 0.02
PC class	1.93 ± 0.07	1.86* ± 0.07	1.97* ± 0.01

[See full table](#)

Table 1. PA and PC unsaturation is increased in wounded plants. In particular, this is due to increases in 36:6 and 34:6 molecular species. The unsaturation index is a measure of the average unsaturation of the fatty acyl chains in each class. The unsaturation index of a lipid species is calculated by

multiplying the number of double bonds by the fraction of that lipid in the lipid class divided by the number of acyl chains in the lipid (i.e., 2 for diacyl lipids). The unsaturation index of the class is the total of the unsaturation index of all lipid species in that class. The PA and PC species as percent of the total indicate the percentage of that class composed on the indicated lipid species. One-way ANOVA was performed to compare the unsaturation indices or lipid percentages among the three treatments. If a significant difference ($p < 0.001$; $n = 31$) was detected for a lipid, a Tukey post-hoc test was performed to identify the differing treatments. * indicates that the value in the wounded sample is significantly increased compared to the unwounded sample.


Oxidation of lipids in the wounding response

While oxidation of extraplastidic phospholipids, such as PC and PE, during wounding is minimal ([Fig. 1D](#)),^{5,8,12} plastidic lipids including PG, MGDG, DGDG, and acylated MGDG are converted to a plethora of oxidized molecular species during wounding ([Fig. 2, block e](#)).

In response to wounding, total oxidized DGDG and MGDG of leaves increased 5 to 6-fold at 45 min after wounding.⁵ Oxidized membrane lipids described by Vu et al.,^{5,12} include those potentially produced by established pathways, such as OPDA, and numerous molecular species whose structures and biosynthetic pathways are less clearly delineated. Chemical formulas for detected molecular species include those consistent with the presence of hydroxy fatty acids, such as 18:2-O (hydroxylated 18:2) and 18:3-O (hydroxylated 18:3), ketols (such as 16:3-2O and 18:3-2O), and phytoprostanes (e.g., 18:4-2O and 18:4-3O).^{5,12} The two most abundant oxidized MGDG species are Arabidopside A, also known as MGDG(OPDA/dnOPDA) and MGDG(34:8-2O), and Arabidopside B, also known as MGDG(diOPDA) and MGDG(36:8-2O), which are increased ~100 and ~50 fold at 45 min after wounding, respectively.²⁸ The most abundant oxidized DGDG, Arabidopside D, also known as DGDG(diOPDA) and DGDG(36:8-2O), increased ~45 fold at 45 min compared to its level in unwounded plants.²⁹ Oxidized acylated MGDG species are also formed quickly and in relatively high amounts. Arabidopside E, MGDG(OPDA/dnOPDA) acylated on galactose with OPDA, and also known as acMGDG(18:4-O/34:8-2O), is increased over 200-fold at 45 min after wounding. The formation of OPDA-containing galactolipids seems to be restricted to Arabidopsis and related species.³⁰ However, other plant species produce membrane lipids with oxidatively modified acyl chains, such as colneleic acid-containing PtdIns in potato tubers; etherolenic acid-containing DAGs (linolipin A and B) and etherolenic acid containing DGDGs (linolipin C and D) in flax leaves upon responding to freezing-thawing.³¹⁻³³

In earlier work, a more resolved time-course experiment demonstrated that MGDG(OPDA/dnOPDA), MGDG(OPDA/OPDA), and DGDG(OPDA/OPDA) were formed faster than their counterparts with a single oxylipin chain.³⁴ Nilsson et al.³⁵ demonstrated that oxidation of trienoic fatty acids to OPDA and dnOPDA occurs on intact galactolipid molecules and does not involve hydrolysis of the acyl chains from the backbone. Together, these data suggest that lipoxygenase acts shortly after wounding and is more likely to oxidize both chains on a galactolipid than a single chain.

[Figure 4](#) shows levels of 4 types of acMGDG: acMGDG with 3 normal chains, acMGDG with 1 out of 3 fatty acyl chains oxidized, acMGDG with 2 out of 3 fatty acyl chains oxidized, and acMGDG with 3 oxidized chains. The fully oxidized acMGDG pool is induced most quickly, and to the highest final level: 87 fold at 45 min, with a drop at 6 h. Other acMGDG pools with a mixture of oxidized and normal chains (or all normal chains) were produced more slowly, and levels were higher at 6 h after wounding than at 45 min. This again suggests that oxidation happens quickly after wounding, that there is a preference for multiple oxidation of acyl chains on a single molecule, and that oxidized chains are cleared by replacement with non-oxidized chains at later times.



[View larger version](#)

Figure 4. Levels of 4 types of acMGDG as a function of oxidation. This figure is based on data in Vu et al.⁵ acMGDG with 3 oxidized chains, acMGDG with 2 out of 3 fatty acyl chains oxidized, acMGDG with 1 out of 3 fatty acyl chains oxidized, and acMGDG with 3 normal chains are shown. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. Levels are compared to the level in unwounded plants. The total of the categorized acMGDG accounts for 71% of acMGDG at 45 min and 6 h after wounding; 29% of acMGDG cannot be readily categorized, due to ambiguity in acyl chain identification. * indicates a significant change, T-test, $P < 0.001$, $n = 31$. Error bars are standard deviation.

Conclusion and Perspectives

The data presented here and in Vu et al.⁵ demonstrate that membrane lipids undergo extensive, diverse modifications during wounding. These changes raise important questions concerning the function of the modifications. Wounding of plant tissues can be caused by abiotic stresses, such as wind, rain, and hail or biotic stresses, such as insect attack, grazing, or pathogen infection. Our recent studies indicate that distinctive types of ac-lipids are produced in plants responding to different stresses, such as freezing, wounding, and pathogen infection.¹² In Arabidopsis, the acMGDGs Arabidopsides E and G have antimicrobial functions and are proposed to be reserves for quick release of oxylipins in response to stress. On the other hand, ac-MGDGs acylated with un-oxidized, normal fatty acids (e.g., 16:0) are not oxylipin reserves and likely have other functions. However, the cellular and physiological functions are virtually unknown for most of the wound-induced lipid metabolites and modifications. Potentially, they could be involved in protecting cells from oxidative damage, provide stress pattern recognition and/or cell signaling. The identification of the

compounds and the deduced metabolic pathways for their production will facilitate greatly functional investigations.

It is also noted that our data on whole leaves do not provide informations about potentially different levels of responsiveness to wounding in various leaf cells. Such differences might be due to differences in cell types or physical locations relative to the wounding sites. Toward this end, application of methodology with spatial resolution capacity, such as mass spectrometry imaging (reviewed by Horn & Chapman),³⁶ is likely provide to novel insights that promote functional studies of wound-induced membrane lipids.

Data and metadata from this work and the work of Vu et al.⁵ are available in the metabolomics-focused database, Plant/Eukaryotic and Microbial Systems Resource (PMR; <http://www.metnetdb.org/PMR/>).³⁷ In PMR, the data on lipid changes during leaf wounding in Arabidopsis can be plotted interactively and otherwise explored further. A screenshot from PMR showing an interactive volcano plot of the wounding data is provided in [Figure 5](#). The experiment is publically accessible at <http://www.metnetdb.org/PMR/experiments/?expid=243>.



[View larger version](#)

Figure 5. Screenshot of Arabidopsis wounding data volcano plot at Plant/Eukaryotic and Microbial Systems Resource (<http://www.metnetdb.org/PMR/>). Mouse-over of any point in the plot provides information about the metabolite and its change in response to wounding. Data include levels of fatty acyl chains in complex lipids as well as levels of intact complex lipid species.

Supplementary Material

[Open In Web Browser](#)

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The authors thank the reviewer for excellent suggestions. The authors are grateful to Mary Roth and Pamela Tamura for their technical assistance. The lipid analyses described in this work were performed at the Kansas Lipidomics Research Center (KLRC).

Funding

The work was supported by National Science Foundation MCB 0920663 and MCB 1413036. Instrument acquisition at KLRC was supported by National Science Foundation (EPS 0236913, DBI 0521587, DBI1228622), Kansas Technology Enterprise Corporation, K-IDeA Networks of Biomedical Research Excellence (INBRE) of National Institute of Health (P20GM103418), and Kansas State University. Contribution 15–373-J from the Kansas Agricultural Experiment Station.

Articles from Plant Signaling & Behavior are provided here courtesy of Landes Bioscience

PMC Copyright Notice

The articles available from the PMC site are protected by copyright, even though access is free. Copyright is held by the respective authors or publishers who provide these articles to PMC. Users of PMC are responsible for complying with the terms and conditions defined by the copyright holder.

Users should assume that standard copyright protection applies to articles in PMC, unless an article contains an explicit license statement that gives a user additional reuse or redistribution rights. PMC does not allow automated/bulk downloading of articles that have standard copyright protection.

See the copyright notice on the PMC site, <http://www.ncbi.nlm.nih.gov/pmc/about/copyright/>, for further details and specific exceptions.

References

1. G GlauserE GrataL DubugnonS RudazEE FarmerJ-L Wolfender. Spatial and temporal dynamics of jasmonate synthesis and accumulation in Arabidopsis in response to wounding. *J Biol Chem* 2008; 283:16400-07; PMID:18400744; <http://dx.doi.org/10.1074/jbc.M80176020018400744>
2. AJ KooX GaoAD JonesGA Howe. A rapid wound signal activates the systemic synthesis of bioactive jasmonates in Arabidopsis. *Plant J* 2009; 59:974-86; PMID:19473329; <http://dx.doi.org/10.1111/j.1365-313X.2009.03924.x19473329>
3. A ChauvinD CaldelariJL WolfenderEE Farmer. Four 13-lipoxygenases contribute to rapid jasmonate synthesis in wounded Arabidopsis thaliana leaves: a role for lipoxygenase 6 in responses to long-distance wound signals. *New Phytol* 2013; 197:566-75; PMID:23171345; <http://dx.doi.org/10.1111/nph.1202923171345>
4. CA ZienC WangX WangR Welti. In-vivo substrates and the contribution of the common phospholipase D, PLD α , to wound-induced metabolism of lipids in Arabidopsis. *Biochim Biophys Acta* 2001; 1530:236-48; PMID:11239826; [http://dx.doi.org/10.1016/S1388-1981\(01\)00091-911239826](http://dx.doi.org/10.1016/S1388-1981(01)00091-911239826)
5. HS VuS ShivaMR RothP TamuraL ZhengM LiS SarowarS HoneyD McEllhineyP Hinkes. Lipid changes after leaf wounding in Arabidopsis thaliana: Expanded lipidomic data form the basis for lipid co-occurrence analysis. *Plant J* 2014; 80:728-43; PMID:25200898; <http://dx.doi.org/10.1111/tpj.1265925200898>
6. E Heinz. Acylgalactosyl diglyceride from leaf homogenates. *Biochim Biophys Acta* 1967;


- 144:321-32; PMID:6064609; [http://dx.doi.org/10.1016/0005-2760\(67\)90161-0](http://dx.doi.org/10.1016/0005-2760(67)90161-0)
7. E Heinz. On the enzymatic formation of acylgalactosyl diglyceride. *Biochim Biophys Acta* 1967; 144:333-44; PMID:6064610; [http://dx.doi.org/10.1016/0005-2760\(67\)90162-2](http://dx.doi.org/10.1016/0005-2760(67)90162-2)
 8. HS VuMR RothP TamuraT SamarakoonS ShivaS HoneyK LoweEA SchmelzTD WilliamsR Welti. Head-group acylation of monogalactosyldiacylglycerol is a common stress response, and the acyl-galactose acyl composition varies with the plant species and applied stress. *Physiol Plant* 2014; 150:517-28; PMID:24286212; <http://dx.doi.org/10.1111/ppl.12132>
 9. MX AnderssonM HambergO KourtchenkoA BrunnströmKL McPhailWH GerwickC Göbell FeussnerM Ellerström. Oxylipin profiling of the hypersensitive response in *Arabidopsis thaliana*. Formation of a novel oxo-phytodienoic acid-containing galactolipid *Arabidopside* E. *J Biol Chem* 2006; 281:31528-37; PMID:16923817; <http://dx.doi.org/10.1074/jbc.M60482020016923817>
 10. O KourtchenkoMX AnderssonM HambergA BrunnströmC GöbelKL McPhailWH GerwickI FeussnerM Ellerström. (Oxo-phytodienoic acid-containing galactolipids in *Arabidopsis*: jasmonate signaling dependence. *Plant Physiol* 2007; 145:1658-69; PMID:17951463; <http://dx.doi.org/10.1104/pp.107.10475217951463>
 11. A IbrahimAL SchützJM GalanoC HerrfurthK FeussnerT DurandF BrodhunI Feussner. The alphabet of galactolipids in *Arabidopsis thaliana*. *Front Plant Sci* 2011; 2:95; PMID:22639619; <http://dx.doi.org/10.3389/fpls.2011.0009522639619>
 12. HS VuP TamuraNA GalevaR ChaturvediTD WilliamsX WangJ ShahR Welti. Direct infusion mass spectrometry of oxylipin-containing *Arabidopsis thaliana* membrane lipids reveals varied patterns in different stress responses. *Plant Physiol* 2012; 158:324-39; PMID:22086419; <http://dx.doi.org/10.1104/pp.111.19028022086419>
 13. AK NilssonON JohanssonP FahlbergF SteinhartMB GustavssonM EllerströmMX Andersson. Formation of oxidized phosphatidylinositol and 12-oxo-phytodienoic acid containing acylated phosphatidylglycerol during the hypersensitive response in *Arabidopsis*. *Phytochem* 2014; 101:65-75; PMID:24559746; <http://dx.doi.org/10.1016/j.phytochem.2014.01.02024559746>
 14. YS SeoEY KimJH KimWT Kim. 2009. Enzymatic characterization of class I DAD1-like acylhydrolase members targeted to chloroplast in *Arabidopsis*. *FEBS Lett* 2009; 583:2301-07; PMID:19527719; <http://dx.doi.org/10.1016/j.febslet.2009.06.02119527719>
 15. W YangY ZhengSC BahnX PanM LiHS VuMR RothB ScheuR WeltiX Wang. Patatin-related

- phospholipase A pPLAII α modulates oxylipin formation and water loss in *Arabidopsis thaliana*. *Mol Plant* 2012; 5:452-60; PMID:22259021;
<http://dx.doi.org/10.1093/mp/ssr11822259021>
16. M LiSC BahnL GuoW MusgraveH BergR WeltiX Wang. Patatin-related phospholipase pPLAIII β -induced changes in lipid metabolism alter cellulose content and cell elongation in *Arabidopsis*. *Plant Cell* 2011; 23:1107-23; PMID:2144778821447788
 17. ER MoelleringB MuthanC Benning. Freezing tolerance in plants requires lipid remodeling at the outer chloroplast membrane. *Science* 2010; 330:226-28; PMID:20798281;
<http://dx.doi.org/10.1126/science.119180320798281>
 18. RL RostonK WangLA KuhnC Benning. Structural determinants allowing transferase activity in SENSITIVE TO FREEZING 2, classified as a family I glycosyl hydrolase. *J Biol Chem* 2014; 289:26089-106; PMID:25100720;
<http://dx.doi.org/10.1074/jbc.M114.57669425100720>
 19. B MuthanRL RostonJE Froehlich, Benning C Probing *Arabidopsis* chloroplast diacylglycerol pools by selectively targeting bacterial diacylglycerol kinase to suborganellar membranes. *Plant Physiol* 2013; 163:61-74; PMID:23839866;
<http://dx.doi.org/10.1104/pp.113.22251323839866>
 20. J FanC YanR RostonJ ShanklinC Xu. *Arabidopsis* lipins, PDAT1 acyltransferases, and SDP1 triacylglycerol lipase synergistically direct fatty acids toward β -oxidation, thereby maintaining membrane lipid homeostasis. *Plant Cell* 2014; 26:4119-34; PMID:25293755;
<http://dx.doi.org/10.1105/tpc.114.13037725293755>
 21. G ThorlbyN FourrierG Warren. The SENSITIVE TO FREEZING2 gene, required for freezing tolerance in *Arabidopsis thaliana*, encodes a β -glucosidase. *Plant Cell* 2004; 16:2192-203; PMID:1525826815258268
 22. N FourrierJ BédardeE Lopez-JuezA BarbrookJ BowyerP JarvisG WarrenG Thorlby. A role for SENSITIVE TO FREEZING2 in protecting chloroplasts against freeze-induced damage in *Arabidopsis*. *Plant J* 2008; 55:734-45; PMID:18466306; <http://dx.doi.org/10.1111/j.1365-313X.2008.03549.x18466306>
 23. F GasullaK Vom DorpI DombrinkU ZähringerN GischP DörmannD Bartels. The role of lipid metabolism in the acquisition of desiccation tolerance in *Craterostigma plantagineum*: a comparative approach. *Plant J* 2013; 75:726-41; PMID:23672245;
<http://dx.doi.org/10.1111/tpj.1224123672245>
 24. SA AriszR van WijkW RoelsJK ZhuMA HaringT Munnik. Rapid phosphatidic acid accumulation in response to low temperature stress in *Arabidopsis* is generated through

- diacylglycerol kinase. *Front Plant Sci* 2013; 4:1; PMID:23346092;
<http://dx.doi.org/10.3389/fpls.2013.0000123346092>
25. CF De JongAM LaxaltBOR BargmannPJGM De WitMHAI JoostenT Munnik Phosphatidic acid accumulation is an early response in the Cf-4/Avr4 interaction. *Plant J* 2004; 39:1-1215200638
 26. C PetersM LiR NarashimhanR RothR WeltiX Wang. Nonspecific phospholipase C NPC4 promotes responses to abscisic acid and tolerance to hyperosmotic stress in Arabidopsis. *Plant Cell* 2010; 22:2642-59; PMID:20699393;
<http://dx.doi.org/10.1105/tpc.109.07172020699393>
 27. S DeBoltWR ScheibleK SchrickM AuerF BeissonV BischoffP Bouvier-NavéA CarrollK HematyY Li Mutations in UDP-glucose:sterol glucosyltransferase in Arabidopsis cause transparent testa phenotype and suberization defect in seeds. *Plant Physiol* 2009; 151:78-87; PMID:19641030; <http://dx.doi.org/10.1104/pp.109.14058219641030>
 28. Y HisamatsuN GotoK HasegawaH Shigemori. Arabidopsides A and B, two new oxylipins from Arabidopsis thaliana. *Tetrahedron Lett* 2003; 44:5553-6; PMID:1584495915844959
 29. Y HisamatsuN GotoM SekiguchiK HasegawaH Shigemori Oxylipins arabidopsides C and D from Arabidopsis thaliana. *J Nat Prod* 2005; 68:600-315844959
 30. C BöttcherEW Weiler. Cyclo-oxylipin-galactolipids in plants: occurrence and dynamics. *Planta* 2007; 226:629-37; PMID:17404756; <http://dx.doi.org/10.1007/s00425-007-0511-517404756>
 31. ML FauconnierTD WilliamsM MarlierR. Welti Potato tuber phospholipids containing colneleic acid in the 2-position. *FEBS Lett* 2003; 538:155-158; PMID:12633870;
[http://dx.doi.org/10.1016/S0014-5793\(03\)00171-612633870](http://dx.doi.org/10.1016/S0014-5793(03)00171-612633870)
 32. IR ChechetkinFK MukhitovaAS BlufardAY YarinLL AntsyginaAN Grechkin. Unprecedented pathogen-inducible complex linolipins A and B. *FEBS J* 2009; 276:4463-72; PMID:19645727; <http://dx.doi.org/10.1111/j.1742-4658.2009.07153.x19645727>
 33. IR ChechetkinAS BlufardBI KhairutdinovFK MukhitovaSS GorinaAY YarinLL AntsyginaAN Grechkin. Isolation and structure elucidation of linolipins C and D, complex oxylipins from flax leaves. *Phytochemistry* 2013; 96:110-6; PMID:24042063;
<http://dx.doi.org/10.1016/j.phytochem.2013.08.01024042063>
 34. CM BusemanP TamuraAA SparksEJ BaughmanS MaattaJ ZhaoMR RothSW EschJ ShahTD Williams Wounding stimulates the accumulation of glycerolipids containing oxophytodienoic acid and dinor-oxophytodienoic acid in Arabidopsis leaves. *Plant Physiol* 2006; 142:28-39; PMID:16844834; <http://dx.doi.org/10.1104/pp.106.08211516844834>

35. AK NilssonP FahlbergM EllerströmMX Andersson. Oxo-phytodienoic acid (OPDA) is formed on fatty acids esterified to galactolipids after tissue disruption in *Arabidopsis thaliana*. *FEBS Lett* 2012; 586:2483-87; PMID:22728240;
<http://dx.doi.org/10.1016/j.febslet.2012.06.01022728240>
36. P HornKD Chapman. Lipidomics in situ: insights into plant lipid metabolism from high resolution spatial maps of metabolites. *Prog Lipid Res* 2014; 54:32-52; PMID:24480404;
<http://dx.doi.org/10.1016/j.plipres.2014.01.00324480404>
37. M HurAA CampbellM Almeida-de-MacedoL LiN RansomA JoseM CrispinB NikolauES Wurtele. A global approach to analysis and interpretation of metabolic data for plant natural product discovery. *Nat Prod Rep* 2013; 30:565-83; PMID:23447050;
<http://dx.doi.org/10.1039/c3np20111b23447050>

[\[Back\]](#)







Figure 1.

Changes in leaf lipid classes following pressure wounding on leaves of wild-type *Arabidopsis thaliana* across the mid-vein with a hemostat. Leaves were harvested from unwounded or wounded plants 45 min and 6 h after wounding.⁵ This figure is based on data in ref. 5. **(A)** Normalized mass spectral intensity for lipids in the panels **(B, C, and D)**. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. Values over the bars indicate the percent of the total lipids represented by the bar. **(B)** Amount of structural (membrane) lipid classes after wounding relative to unwounded level for each class. **(C)** Amounts of minor plastidic lipid classes relative to the unwounded level for each class. **(D)** Amounts of minor extraplastidic lipid classes relative to the unwounded level for each class. * indicates significant changes compared to unwounded samples, T-test, $P < 0.001$, $n = 31$. Error bars are standard deviation.

[\[Back\]](#)

[\[Back\]](#)




Figure 2.

Proposed reactions of membrane lipids in the wounding response in Arabidopsis leaves. **(a)** Lipid head-group acylation and lipolytic reactions producing monoacyl lipid molecular species; **(b)** galactosylation of lipids and other reactions producing DAG, PA, and TAG; **(c)** formation of sterol glucosides and acyl sterol glucosides; **(d)** increase of fully desaturated PC and PA fractions during the wounding response; and **e.** oxidation of lipids in the wounding response. See text for details.

[\[Back\]](#)

[\[Back\]](#)




Figure 3.

Formation of TrGDG, TeGDG, PA(34:6), and 16:3-containing TAG is abolished in *sfr2-3* (SALK_106253) plants. Leaves were harvested from unwounded or wounded plants 45 min after wounding as in ref. 5. In the indicated lipids, “34:6” is composed of 18:3 and 16:3 fatty acyl chains. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. ** indicates a significant change at $P < 0.01$, * indicates a significant change at $P < 0.05$, T-test, $n = 6$. Error bars are standard deviation.

[\[Back\]](#)

[\[Back\]](#)

Table 1.

PA and PC unsaturation is increased in wounded plants. In particular, this is due to increases in 36:6 and 34:6 molecular species. The unsaturation index is a measure of the average unsaturation of the fatty acyl chains in each class. The unsaturation index of a lipid species is calculated by multiplying the number of double bonds by the fraction of that lipid in the lipid class divided by the number of acyl chains in the lipid (i.e., 2 for diacyl lipids). The unsaturation index of the class is the total of the unsaturation index of all lipid species in that class. The PA and PC species as percent of the total indicate the percentage of that class composed on the indicated lipid species. One-way ANOVA was performed to compare the unsaturation indices or lipid percentages among the three treatments. If a significant difference ($p < 0.001$; $n = 31$) was detected for a lipid, a Tukey post-hoc test was performed to identify the differing treatments. * indicates that the value in the wounded sample is significantly increased compared to the unwounded sample.

	Unwounded	45 min after wounding	6 h after wounding
Unsaturation index (n = 31)			
PA class	1.60 ± 0.04	1.69* ± 0.01	1.66* ± 0.02
PC class	1.83 ± 0.02	1.86* ± 0.02	1.87* ± 0.01
PA species as percent of total PA			
PA(36:6)	5.1 ± 0.8	6.6* ± 0.5	5.8* ± 0.6
PA(34:6)	0.3 ± 0.3	0.3 ± 0.1	1.6* ± 0.4

PC species as percent of total
PC

PC(36:6)	10.7 ± 0.8	12.0* ± 0.8	11.9* ± 0.9
PC(34:6)	0.2 ± 0.0	0.3* ± 0.0	0.5* ± 0.1

[\[Back\]](#)

[\[Back\]](#)




Figure 4.

Levels of 4 types of acMGDG as a function of oxidation. This figure is based on data in Vu et al.⁵ acMGDG with 3 oxidized chains, acMGDG with 2 out of 3 fatty acyl chains oxidized, acMGDG with 1 out of 3 fatty acyl chains oxidized, and acMGDG with 3 normal chains are shown. 1 unit of normalized mass spectral signal equals the signal obtained from 1 pmol of internal standard. Levels are compared to the level in unwounded plants. The total of the categorized acMGDG accounts for 71% of acMGDG at 45 min and 6 h after wounding; 29% of acMGDG cannot be readily categorized, due to ambiguity in acyl chain identification. * indicates a significant change, T-test, $P < 0.001$, $n = 31$. Error bars are standard deviation.

[\[Back\]](#)

[Back]




Figure 5.

Screenshot of Arabidopsis wounding data volcano plot at Plant/Eukaryotic and Microbial Systems Resource (<http://www.metnetdb.org/PMR/>). Mouse-over of any point in the plot provides information about the metabolite and its change in response to wounding. Data include levels of fatty acyl chains in complex lipids as well as levels of intact complex lipid species.

[Back]

Table of Contents

Modifications of membrane lipids in response to wounding of <i>Arabidopsis thaliana</i> leaves	1
---	---