

## University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications: Department of Entomology

Entomology, Department of

---

2016

# Efficacy of Transport Mikron Against Nuisance Ants When Applied Around Structures, 2015

Sanjay Basnet

*University of Nebraska-Lincoln*, [sbasnet2@unl.edu](mailto:sbasnet2@unl.edu)

Dina Richman

*University of Nebraska-Lincoln*, [dina.richman@fmc.com](mailto:dina.richman@fmc.com)

Ralph B. Narain

*University of Nebraska-Lincoln*, [usralphn@aol.com](mailto:usralphn@aol.com)

Shripat T. Kamble

*University of Nebraska-Lincoln*, [skamble1@unl.edu](mailto:skamble1@unl.edu)

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>

 Part of the [Entomology Commons](#)

---

Basnet, Sanjay; Richman, Dina; Narain, Ralph B.; and Kamble, Shripat T., "Efficacy of Transport Mikron Against Nuisance Ants When Applied Around Structures, 2015" (2016). *Faculty Publications: Department of Entomology*. 468.  
<http://digitalcommons.unl.edu/entomologyfacpub/468>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## URBAN, INDUSTRIAL, AND STRUCTURAL PESTS

# Efficacy of Transport Mikron Against Nuisance Ants When Applied Around Structures, 2015

Sanjay Basnet,<sup>1</sup> Dina Richman, Ralph B. Narain, and Shripat T. Kamble

Department of Entomology, University of Nebraska, Lincoln, NE 68583-0816, Phone: (402) 472-8632, Fax: (402) 472-4687 (sbasnet2@unl.edu; Dina.richman@fmc.com; usralphn@aol.com; skamble1@unl.edu) and <sup>1</sup>Corresponding author, email: sbasnet2@unl.edu

Subject Editor: Eric Natwick

acrobat ant | *Crematogaster* spp.

false honey ant | *Prenolepis imparis*

odorous house ant | *Tapinoma sessile*

pavement ant | *Tetramorium caespitum*

thief ant | *Solenopsis molesta*

Ants are nuisance pests around the homes and other structures. Insecticide baits and sprays are commonly used for controlling ants. The research was conducted to determine efficacy of Transport Mikron and Temprid SC against these nuisance ants. The trial was conducted around two buildings on East Campus, University of Nebraska-Lincoln, Lincoln, NE. A total of 20 experimental plots were established with 3 ft buffer zone between individual plots. The dimension of each experimental plot consists of 20 ft length × 10 ft width + 3 ft high on foundation wall (260 ft<sup>2</sup>). Each treatment was replicated four times yielding 1040 ft<sup>2</sup>. Four plots were randomly assigned to one of the five treatments. The experimental design was a completely randomized design (CRD). Pretreatment ant population was monitored by placing four ant bait-tube traps made of cylindrical plastic tube (17 × 100 mm (VWR, Chicago, IL)) in each experimental unit on 20 Aug. Peanut butter (0.5 inch diam) on paper was used as a bait in each. Foraging nuisance ant populations were pooled across species, counted, and used as a baseline for calculating the percent reduction in ant

populations resulting from treatments. All insecticide treatments were applied using a 1-gallon B&G Compressed Air Sprayer (Jackson, GA) on 25 Aug. Ants were monitored at 1, 7, 14, and 31 days after insecticide treatment (DAT) using the same techniques used for pretreatment assessment. The percent reduction trends were evaluated using pre- and post-treatment nuisance ant population counts. Data were analyzed as repeated measures ANOVA ( $P < 0.05$ ) using PROC GLIMMIX (SAS 9.4. SAS institute, NC). Means were analyzed/sorted by time with significant differences being collection interval specific.

All treatments significantly reduce ant population as compared to untreated through 31 DAT. The highest rate of Transport Mikron reduced ant population consistently through 31 DAT. No significant differences were observed among different rates of Transport Mikron throughout the study. No significant difference was observed between the Temprid SC and Transport Mikron (three rates) at 1 DAT, however, there were significant differences at 7 DAT and 31 DAT (Table 1).

**Table 1**

Product/formulation	Rate (oz/gal/1000 ft <sup>2</sup> )	Percent reduction in ant population			
		1 DAT (8/26/2015)	7 DAT (9/1/2015)	14 DAT (9/8/2015)	31 DAT (9/25/2015)
Transport Mikron (high) DL	1.25	99.62 ± 0.38 a*	97.06 ± 1.88 a	99.81 ± 0.19 a	98.68 ± 1.32 a
Transport Mikron (medium) DL	0.63	99.61 ± 0.39 a	95.61 ± 2.44 a	90.73 ± 7.12 ab	78.94 ± 14.61 a
Transport Mikron (low) DL	0.42	99.57 ± 0.43 a	96.71 ± 2.06 a	88.24 ± 8.93 ab	91 ± 4.27 a
Temprid SC DL	0.27	96.86 ± 2.68 a	75.55 ± 6.87 b	69.37 ± 16.7 b	34.79 ± 28.11 b
Untreated	NA	12 ± 7.05 b	0 ± 0 c	13.9 ± 8.57 c	2.89 ± 2.89 c

\*Means within columns followed by same letter are not statistically different