

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications in Food Science and  
Technology

Food Science and Technology Department

---

2009

# Genomic Regions Conserved in Lineage II *Escherichia coli* O157:H7 Strains

Marina Steele

*Public Health Agency of Canada*

Kim Ziebell

*Public Health Agency of Canada*

Yongxiang Zhang

*Public Health Agency of Canada*

Andrew K. Benson

*University of Nebraska-Lincoln*, [abenson1@unl.edu](mailto:abenson1@unl.edu)

Roger Johnson

*Public Health Agency of Canada*

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/foodsciefacpub>

 Part of the [Food Science Commons](#)

---

Steele, Marina; Ziebell, Kim; Zhang, Yongxiang; Benson, Andrew K.; Johnson, Roger; Laing, Chad; Taboada, Eduardo; and Gannon, Victor, "Genomic Regions Conserved in Lineage II *Escherichia coli* O157:H7 Strains" (2009). *Faculty Publications in Food Science and Technology*. 177.

<http://digitalcommons.unl.edu/foodsciefacpub/177>

This Article is brought to you for free and open access by the Food Science and Technology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in Food Science and Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Marina Steele, Kim Ziebell, Yongxiang Zhang, Andrew K. Benson, Roger Johnson, Chad Laing, Eduardo Taboada, and Victor Gannon

## Genomic Regions Conserved in Lineage II *Escherichia coli* O157:H7 Strains<sup>∇</sup>

Marina Steele,<sup>1</sup> Kim Ziebell,<sup>1</sup> Yongxiang Zhang,<sup>2</sup> Andrew Benson,<sup>3</sup> Roger Johnson,<sup>1</sup>  
Chad Laing,<sup>2</sup> Eduardo Taboada,<sup>2</sup> and Victor Gannon<sup>2\*</sup>

Laboratory for Foodborne Zoonoses, Public Health Agency of Canada, Guelph, Ontario, Canada<sup>1</sup>; Laboratory for Foodborne Zoonoses, Public Health Agency of Canada, Lethbridge, Alberta, Canada<sup>2</sup>; and Department of Food Science and Technology, University of Nebraska, Lincoln, Nebraska<sup>3</sup>

Received 12 September 2008/Accepted 18 March 2009

**Populations of the food- and waterborne pathogen *Escherichia coli* O157:H7 are comprised of two major lineages. Recent studies have shown that specific genotypes within these lineages differ substantially in the frequencies with which they are associated with human clinical disease. While the nucleotide sequences of the genomes of lineage I strains *E. coli* O157 Sakai and EDL9333 have been determined, much less is known about the genomes of lineage II strains. In this study, suppression subtractive hybridization (SSH) was used to identify genomic features that define lineage II populations. Three SSH experiments were performed, yielding 1,085 genomic fragments consisting of 811 contigs. Bacteriophage sequences were identified in 11.3% of the contigs, 9% showed insertions and 2.3% deletions with respect to *E. coli* O157:H7 Sakai, and 23.2% did not have significant identity to annotated sequences in GenBank. In order to test for the presence of these novel loci in lineage I and II strains, 27 PCR primer sets were designed based on sequences from these contigs. All but two of these PCR targets were found in the majority (51.9% to 100%) of 27 lineage II strains but in no more than one (<6%) of the 17 lineage I strains. Several of these lineage II-related fragments contain insertions/deletions that may play an important role in virulence. These lineage II-related loci were also shown to be useful markers for genotyping of *E. coli* O157:H7 strains isolated from human and animal sources.**

Enterohemorrhagic *Escherichia coli* is associated with diarrhea, hemorrhagic colitis, and hemolytic-uremic syndrome in humans (31). *E. coli* serotype O157:H7 predominates in epidemics and sporadic cases of enterohemorrhagic *E. coli*-related infections in the United States, Canada, Japan, and the United Kingdom (12). Cattle are considered the most important reservoir of *E. coli* O157:H7 (10, 24, 37, 41), and foods contaminated with bovine feces are thought to be the most common source of human infection with this pathogen (27, 33). The two most important virulence factors of the organism are the production of one or more Shiga toxins (Stx) (6, 20, 32) and the ability to attach to and efface microvilli of host intestinal cells (AE). Stx genes are encoded by temperate bacteriophage inserted in the bacterial chromosome, and genes responsible for the AE phenotype are located on the locus of enterocyte effacement (LEE) as well as other pathogenicity islands (4, 17). All *E. coli* O157:H7 strains also possess a large plasmid which is thought to play a role in virulence (10, 40, 42).

Octamer-based genome scanning (OBGS) was first used to show that *E. coli* O157 strains from the United States and Australia could be subdivided into two genetically distinct lineages (21, 22, 46). While both *E. coli* O157:H7 lineages are associated with human disease and are isolated from cattle, there is a bias in the host distribution between the two lineages, with a significantly higher proportion of lineage I strains isolated from humans than lineage II strains. Several recent stud-

ies have shown that there are inherent differences in gene content and expression between populations of lineage I and lineage II *E. coli* O157:H7 strains. Lejeune et al. (26) reported that the antiterminator Q gene of the *stx*<sub>2</sub>-converting bacteriophage 933W was found in all nine OBGS lineage I strains examined but in only two of seven lineage II strains, suggesting that there may be lineage-specific differences in toxin production. Dowd and Ishizaki (9) used DNA microarray analysis to examine expression of 610 *E. coli* O157:H7 genes and showed that lineage I and lineage II *E. coli* O157:H7 strains have evolved distinct patterns of gene expression which may alter their virulence and their ability to survive in different microenvironments and colonize the intestines of different hosts (9, 28, 38).

The observations of lineage host bias have been supported and extended by studies using a six-locus-based multiplex PCR termed the lineage-specific polymorphism assay (LSPA-6) (46). However, Ziebell et al. (48) have recently shown that not all LSPA-6 types within lineage II are host biased; e.g., LSPA-6 type 211111 isolation rates from humans and cattle were significantly different from those of other lineage II LSPA-6 types. Therefore, a clearer definition is required of not only the differences between lineages but also the differences among clonal groups within lineages.

The genome sequences of two *E. coli* O157:H7 strains, Sakai and EDL933 (14, 36), have been determined; however, both of these strains are of lineage I, and there are presently no completed and fully annotated genome sequences available for lineage II strains. In our laboratory, comparative studies utilizing suppression subtractive hybridization (SSH) and comparative genomic hybridization revealed numerous potential

\* Corresponding author. Mailing address: Laboratory for Foodborne Zoonoses, Public Health Agency of Canada, 1st floor, C.F.I.A. Building, Lethbridge, Alberta T1J 3Z4, Canada. Phone: (403) 382-5514. Fax: (403) 381-1202. E-mail: gannonv@inspection.gc.ca.

<sup>∇</sup> Published ahead of print on 27 March 2009.

virulence factors that are conserved in lineage I strains and that are rare or absent in lineage II strains (42, 47). In this study, we have used SSH to identify genomic regions present in *E. coli* O157:H7 lineage II strains that are absent from lineage I strains. We wished to examine the distribution of these novel gene segments in *E. coli* O157:H7 strains and gain insight into their origins and functions. We also attempted to identify molecular markers specific to lineage II strains as well as other markers that would be useful in the genetic subtyping or molecular fingerprinting of *E. coli* O157:H7 strains in population and epidemiological studies (25). This information may be helpful in the identification of genotypes of the organism associated with specific phenotypes of both lesser and greater virulence (29).

#### MATERIALS AND METHODS

**Bacterial strains.** OBGS type strains (93-001, FDA 516-520, and FRIK 523-2001) were previously described by Kim et al. (21). Zap strains were obtained from David Gally at the University of Edinburgh (30). *Escherichia coli* O157:H7 strains EDL933 (ATCC 700927) and Sakai (RIMD 0509952) were obtained from the American Type Culture Collection (Manassas, VA). The remaining strains were isolated from human infections or cattle in Canada (Table 1). LSPA-6 genotyping of these strains was performed as previously described (48).

**Preparation of the SSH DNA library.** Bacterial cultures were grown overnight in brain heart infusion broth (Difco, Becton, Dickinson and Company, Franklin Lakes, NJ) in a 37°C shaker-incubator (200 rpm) and genomic DNA was extracted from harvested cells using the DNeasy tissue kit (Qiagen, Valencia, CA). The purity and concentration of the genomic DNA were assessed using a Nanodrop ND-1000 spectrophotometer (Nanodrop Technologies, Rockland, DE).

SSHs were performed using the Clontech PCR-Select bacterial genome subtraction kit (BD Biosciences, Palo Alto, CA). In addition to the RsaI-digested DNA recommended in the Clontech kit, SSHs were also performed on AluI- and HaeIII-digested DNA to increase the diversity of DNA fragments obtained. Advantage polymerase mix (BD Biosciences) was used during the amplification steps. Three sets of SSH experiments were performed. In the first set, *E. coli* FRIK 920 (LSPA-6 222222) was subtracted with *E. coli* Sakai (LSPA-6 111111). In the second set, *E. coli* FRIK 1999 (LSPA-6 222222) was subtracted with *E. coli* Sakai. In the third set, *E. coli* FRIK 2001 (LSPA-6 212232) was subtracted with *E. coli* 93-001 (LSPA-6 111111).

The SSH DNA fragments isolated in these experiments were cloned into the pCR2.1-TOPO plasmid vector, using the TOPO TA cloning kit (Invitrogen, Carlsbad, CA) and plated onto LB agar (Difco) containing 50 µg/ml of ampicillin or kanamycin (Sigma-Aldrich Canada, Oakville, ON, Canada). Prior to cloning, the fragments were incubated for 10 min at 72°C in the presence of AmpliTaq DNA polymerase (Applied Biosystems, Foster City, CA) to ensure that all DNA fragments possessed the necessary "A" overhang.

**DNA sequencing of SSH DNA library.** PCR amplicons were generated directly from the SSH library clones by transferring whole bacterial cells to 20-µl aliquots of PCR master mix containing 1× buffer II (Applied Biosystems), 1.5 mM MgCl<sub>2</sub>, 0.2 mM each deoxynucleoside triphosphates, 1 U AmpliTaq DNA polymerase (Applied Biosystems), and 0.2 µM each M13 Forward (5'-GTAAAAC GACGGCCAG-3') and M13 Reverse (5'-CAGGAAACAGCTATGAC-3') PCR primer sets (Invitrogen). An initial 10-min incubation at 94°C performed to lyse the cells was followed by 30 cycles of 1 min at 94°C, 1 min at 55°C, and 1 min at 72°C and a final 10-min extension step at 72°C. PCR amplicons were purified by passage through superfine Sephadex G-50 (Sigma) packed into a MultiScreen 96-well filtration plate (Millipore, Billerica, MA) and sequenced with the DYEnamic ET terminator cycle sequencing kit (GE Health Care, Piscataway, NJ) using M13 Forward and M13 Reverse primer sets. Sequencing products were purified by passage through superfine Sephadex G-50 packed into a MultiScreen 96-well filtration plate prior to nucleotide sequencing on a MegaBace 500 capillary sequencer (GE Health Care).

**DNA sequence analysis of the SSH DNA library.** Phred base calling of sequence trace data was performed using the Interphase program (CodonCode Corporation, Dedham, MA). Base-called sequences obtained from each set of SSH experiments were analyzed by the SeqMan 5.05 sequence analysis program (DNASTar Inc., Madison, WI). This program removed vector and adaptor sequences and grouped sequences displaying at least 80% sequence homology into contiguous consensus sequences or "contigs." Searches were performed on con-

TABLE 1. *Escherichia coli* O157:H7 strains included in the study (*n* = 44)

Strain	LSPA-6 genotype	Source	Country of origin <sup>a</sup>	Phage type
ECI1375	111111	Bovine	Canada	14
ECI1382	111111	Bovine	Canada	1
ECI563	111111	Bovine	Canada	14
ECI577	111111	Bovine	Canada	4
ECI603	111111	Bovine	Canada	87
ECI653	111111	Bovine	Canada	32
ECI665	111111	Bovine	Canada	1
93001	111111	Human	U.S.A.	14
ECI309	111111	Human	Canada	14
ECI320	111111	Human	Canada	14
ECI485	111111	Human	Canada	14
EDL933	111111	Human	U.S.A.	21
FDA 516	111111	Human	U.S.A.	21
FDA 518	111111	Human	U.S.A.	21
FDA 520	111111	Human	U.S.A.	1
FRIK 523	111111	Human	U.S.A.	34
Sakai	111111	Human	Japan	32
Zap0032	211111	Bovine	Scotland	8
Zap0054	211111	Bovine	Scotland	32
ECI504	211111	Human	Canada	2
ECI882	211111	Human	Canada	1
Zap0058	211111	Human	Scotland	87
ECI241	212222	Human	Canada	74
FRIK 2001	212232	Bovine	U.S.A.	54
ECI240	212232	Human	Canada	54
EC19930200	222221	Bovine	Canada	23
FRIK 1985	222232	Bovine	U.S.A.	45
EC19920005	222222	Bovine	Canada	67
EC19920027	222222	Bovine	Canada	34
EC19920171	222222	Bovine	Canada	23
EC19970520	222222	Bovine	Canada	67
EC20011139	222222	Bovine	Canada	82
EC20030223	222222	Bovine	Canada	74
EC20030289	222222	Bovine	Canada	23
ECI1433	222222	Bovine	Canada	23
ECI564	222222	Bovine	Canada	23
ECI633	222222	Bovine	Canada	23
FRIK 1990	222222	Bovine	U.S.A.	54
FRIK 1999	222222	Bovine	U.S.A.	23
FRIK 920	222222	Bovine	U.S.A.	23
ECI306	222222	Human	Canada	23
ER6554	222222	Human	Canada	23
ER6666	222222	Human	Canada	40
ER6816	222222	Human	Canada	40

<sup>a</sup> U.S.A., United States.

tigs using BLASTN and BLASTX (2) from the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/>).

**PCR screening of putative lineage II conserved markers.** Twenty-seven PCR primer sets were designed to detect frequently occurring sequences within the SSH libraries. These primer sets were used to screen 17 *E. coli* O157:H7 LSPA-6 111111 strains, 17 genotype 222222 strains, five genotype 211111 strains, and five strains of other genotypes (Table 1). Primer Select 5.08 (DNASTar Inc.) was used to design PCR primer sets (Table 2). All PCR assays were performed in duplicate in 20-µl reaction mixture volumes containing 1× buffer II (Applied Biosystems), 1.5 mM MgCl<sub>2</sub>, 0.2 mM deoxynucleoside triphosphates, 1 U AmpliTaq DNA polymerase (Applied Biosystems), 0.2 µM primer sets, and 0.5 ng genomic DNA template. All PCRs included an initial 2-min denaturation step at 94°C, followed by 30 cycles of 1 min denaturation at 94°C, 1 min annealing, and 1 min extension at 72°C, and a final 10-min extension at 72°C. The primer sequences and annealing temperatures for each PCR are listed in Table 2. Positive PCR controls included the SSH tester *E. coli* strains FRIK 920, FRIK 1999, and FRIK 2001 and negative controls included *Campylobacter jejuni* ATCC 33560 and a no-template blank. *E. coli* FRIK 920 amplicons obtained from all primer sets were purified using the QIAquick PCR purification kit (Qiagen) and sequenced

TABLE 2. Primer sets used in PCR screening of lineage II conserved markers among *E. coli* O157:H7 strains

Class	Primer set	Forward primer	Reverse primer	Temp (°C)
B	A02	5'-GCTGGCAGACAATGGCGAGTTT-3'	5'-GATACGGCTTGCGGGCTGTG-3'	59.3
	A03	5'-GCCACGGCATTAAATAACGCTTCTT-3'	5'-CCAGGCATCACTACCCGCAAAAA-3'	57.4
	A04	5'-GGAGACGGGGCGCACTGATTG-3'	5'-CTCCACGATTTTCGGGGCTTATGTA-3'	56.2
	A05	5'-TGCGGCATCTGGTCATTCTGGTT-3'	5'-ACGGTCTGCGCGAGTTTTGTTT-3'	57.3
	A09	5'-AGGGGGCATATTTTCATTGTTGTGT-3'	5'-TAATCTCTTTCGCCTGCCCTTCTT-3'	56.4
	A10	5'-CGAGGCGTGCGGAGAGGA-3'	5'-TTCAGCGCCAGTTCGGTTTTTC-3'	58.8
	A11	5'-AAAACCGTATGATCTATTATTCCT-3'	5'-TTTTTATTATGTTGGCTGAGTTAC-3'	50.3
	A13	5'-ATGCCAGCCACCACCACTCCAC-3'	5'-CTTACCCGCGCCGAACCTATCCTT-3'	58.2
	B01	5'-AAGCCCCACGTAATTCCTGACA-3'	5'-CATTTCCCGCGCCTGACTGAGA-3'	60.7
	B02	5'-AAGCGGCATAAATGGCAGACAGAC-3'	5'-AACGCCACGCTTTACCTTTACCC-3'	57.8
	B03	5'-AATCCGCCCGTTTTACTGA-3'	5'-TGTGCCTGATAGCTCCTTCTTTTT-3'	54.9
	B06	5'-CCCCCTGTTGCGTCTGCTGAAAA-3'	5'-CGGCCATCCATGCGGTGACTT-3'	60.8
	B07	5'-TAAAGCGCTGGTGGGGAAAGGTAG-3'	5'-CCAGAACGCGGGCACAAAAA-3'	54.5
	B08	5'-TGCCGACATCGCCAGTTG-3'	5'-GCCATGCTTCAGGGAGATAAAAT-3'	55.7
	B10	5'-TGTGCGTCTTCAGTTCGTCA-3'	5'-TAGGTGTTCCGCGTCTGTAAAAG-3'	57.8
	B16	5'-CATGCTTCTGCCCTGAGTT-3'	5'-GCGCCGTGCTTATGAAA-3'	54.8
C	A14	5'-TGCCGACAACCTCCCACAGATACT-3'	5'-TCACTTCCGCTACAACACGCACAT-3'	57.8
	A15	5'-CGCCAAAGCAGCACAGCAGGATA-3'	5'-AGAAAAACAGCGAAGGCGATGAT-3'	53.4
D	A06c	5'-CCGCCTGCGATGGTGGTTGC-3'	5'-GGGCGCGGGTGATTTTGCTCTC-3'	60.2
	A07	5'-ATGCGATCGCCTTCTTCAA-3'	5'-TACCATACACGCCACAGTTTTTA-3'	52.3
	B15	5'-CCCGCTGGCAGGCATTGAAG-3'	5'-GGCGCAGCGGACACGAG-3'	60.4
E	A01	5'-ACCAAGGCATCCCCGTGTGAA-3'	5'-ATAATCCGCTGGGGCTGGCTGAC-3'	60.8
	B05	5'-GGTTTTCCGGCACTTTCCACTCCA-3'	5'-ATCCTGCCGGGCGAACATCCTTAT-3'	58.1
	B12	5'-TGAACACCCGAGCAACA-3'	5'-CGCCGCATCTACTCCTATCG-3'	54.2
	B18	5'-AATAACTCGGCTTTTGTCTTTT-3'	5'-AATACTCCGTTTCTGTCTAATCC-3'	52.5
F	A12	5'-GGGCGGACTTTGTTTGGTTGAA-3'	5'-GCCTGGCGGAAATGGACTGTAT-3'	55.9
	B13	5'-CTGGATGCGGCAAAACCTGT-3'	5'-GCCCTTCTCTACGCAAATCAT-3'	55.0

on an ABI Prism 277 DNA sequencer (Applied Biosystems, Foster City, CA) to verify that they corresponded to the sequences from the original contig assembly.

**Analysis of distribution of lineage II conserved regions in *E. coli* O157:H7 strains.** Results of PCR screening assays targeting lineage II conserved markers within the 44 strains tested were converted into presence/absence binary data. A neighbor-joining dendrogram was generated based on a matrix of pairwise distances calculated from the binary data, using the proportion of shared markers as a distance metric. Based on similar BLASTN homologies and strain distributions, some PCR primer sets appeared to recognize the same locus. To avoid biasing the dendrogram results, a single representative PCR assay from each of these 18 loci was included in the final binary data. The data were subsequently bootstrapped, and a neighbor-joining consensus dendrogram was created using the neighbor and consense programs from Phylip v3.16 (11). *E. coli* K-12 (5) was used as an out-group strain in the dendrogram since in silico analysis indicated

that none of the primer sets would have generated a PCR product of the correct size in this strain.

RESULTS AND DISCUSSION

**SSH DNA library sequences.** The three lineage II-minus-lineage I SSH libraries yielded 811 contigs, which were obtained from 1,085 clones. Each contig represented between 1 and 27 of the clones obtained, with an average of  $1.58 \pm 1.90$  clones per contig (Table 3). Sequence identities of the contigs were obtained from BLASTN searches, and sequences were binned into six different classes (A to F), based on sequence

TABLE 3. Frequency of sequence classes within SSH DNA libraries

Sequence class <sup>a</sup>	No. (%) of clones present in:				Range	Avg no. of clones/contig
	All contigs (n = 811)	3 SSH (n = 16)	2 SSH (n = 73)	1 SSH (n = 722)		
A	503 (62.0)	5 (31.3)	30 (41.1)	468 (64.7)	1-7	1.30 ± 0.82
B	92 (11.3)	10 (66.7)	34 (46.6)	48 (6.6)	1-16	2.90 ± 2.64
C	72 (8.9)	3 (18.8)	12 (16.4)	57 (7.9)	1-7	1.60 ± 1.22
D	19 (2.3)	0 (0)	6 (8.2)	13 (1.8)	1-5	1.79 ± 1.13
E	80 (9.9)	1 (13.3)	8 (12.7)	72 (9.7)	1-27	3.01 ± 4.67
F	188 (23.3)	5 (31.3)	14 (19.2)	177 (24.5)	1-11	1.67 ± 1.65
All					1-27	1.58 ± 1.90

<sup>a</sup> Class A, sequences with homology to *E. coli* Sakai (14) genes; class B, sequences with homology to bacteriophage sequences not found in *E. coli* Sakai; class C, putative mosaic sequences with homology to both *E. coli* Sakai and non-Sakai DNA sequences within the same contig; class D, sequences with homology to *E. coli* Sakai genes but lacking synteny with the *E. coli* Sakai chromosome; class E, sequences with homology to previously published nonbacteriophage and non-Sakai DNA sequences; class F, sequences with regions containing no homology to any annotated DNA sequences in GenBank.

TABLE 4. Distribution of putative lineage II conserved markers among *E. coli* O157:H7 strains from various lineages

Class	Primer set	SSH(s) <sup>a</sup>	Predicted amplicon size (bp)	No. of indicated markers/total no. (%)			Sequence homology	
				Lineage I (LSPA-6 111111)	Lineage II (LSPA-6 222222)	Lineage II (LSPA-6 211111)		Other lineage II
B	A10	2, 3	435	0/17 (0)	17/17 (100)	3/5 (60)	4/5 (80)	<i>E. coli</i> UTT189 (7) (96% homology over 1,388 nt) genes with homology to bacteriophage HK97 gp56 and bacteriophage N12p gene <i>dnaB</i> homologue
	A11	1, 2, 3	365	0/17 (0)	17/17 (100)	3/5 (60)	4/5 (80)	Bacteriophage BP-4795 (8) (95% homology over 762 nt), encoding a hypothetical protein
	A13	1, 2, 3	204	0/17 (0)	17/17 (100)	0/5 (0)	5/5 (100)	<i>E. coli</i> K-12 (5) ORFs b1155 to b1157 (81% homology over 1,260 nt), encoding hypothetical proteins and a putative tail fiber assembly protein within the cryptic prophage e14
	B02	1, 2, 3	136	0/17 (0)	13/17 (76.5)	5/5 (100)	3/5 (60)	Bacteriophage P2 (accession no. AF063097) genes J and I (94% homology over 1,019 nt), encoding baseplate assembly proteins
	A05	1, 2, 3	246					<i>E. coli</i> Sakai (14) ORF ECs0284 (88% homology over 338 nt), encoding a DNA invertase, adjacent to 213 nt of DNA sequence, without significant homology to known DNA sequences adjacent to bacteriophage P2 (accession no. AF063097) FI tail sheath protein gene (95% homology over 546 nt)
	B03	1, 2, 3	746	0/17 (0)	17/17 (100)	0/5 (0)	5/5 (100)	<i>E. coli</i> O1091 P4-like bacteriophage ORFs ecO109Im and <i>psu</i> (23) (98% homology over 1,248 nt), encoding methyltransferase and <i>Psu</i> -like proteins
	B06	1, 2, 3	327	0/17 (0)	17/17 (100)	0/5 (0)	5/5 (100)	<i>S. boydii</i> strain Sb227 (45) ORFs SBO4168 to SBO4170 (97% homologous over 1,352 nt) and SBO3220 (98% homology over 170 nt), encoding a putative bacteriophage protein and conserved hypothetical proteins
	A09	2, 3	572					<i>S. boydii</i> strain Sb227 (45) ORFs SBO4164 and SBO4165 (95% homology over 1,366 nt), encoding a prophage P4 integrase and a putative DNA primase
	B08	2, 3	616	0/17 (0)	17/17 (100)	3/5 (60)	5/5 (100)	Bacteriophage HK022 (19) (93% homology over 632 nt, with gaps), intergenic region between genes encoding an excisionase and an Eca22 protein
	B10	1, 2, 3	272	0/17 (0)	17/17 (100)	5/5 (100)	5/5 (100)	<i>S. boydii</i> strain Sb227 (45) ORF SBO2131 (99% homology over 892 nt), encoding conserved hypothetical protein
C	A02	1, 2	582					<i>S. boydii</i> strain Sb227 (45) ORFs SBO2122 to SBO2125 (94% homology over 1,013 nt), encoding conserved hypothetical protein and prophage capsid proteins
	A03	2, 3	483					<i>S. boydii</i> strain Sb227 (45) ORFs SBO2131 to SBO2132 (98% homology over 995 nt), encoding conserved hypothetical proteins
	A04	2, 3	438					<i>S. boydii</i> strain Sb227 (45) ORFs SBO2126 to SBO2127 (97% homology over 684 nt), encoding a conserved hypothetical protein and a hypothetical bacteriophage protein
	B16	1, 2, 3	830	0/17 (0)	17/17 (100)	0/5 (0)	5/5 (100)	<i>S. flexneri</i> bacteriophage V (16) ORFs <i>int</i> , <i>xis</i> , <i>orf28</i> , <i>orf29</i> , <i>orf30</i> , and <i>orf31</i> (97% homologous over 2,453 nt), encoding an integrase, an excisionase, and hypothetical proteins
	B01	1, 2, 3	316					<i>S. flexneri</i> bacteriophage V (16) ORFs <i>orf39</i> , <i>orf40</i> , and <i>dam</i> (96% homology over 823 nt), encoding a replication protein, a hypothetical protein, and a DNA adenine methylase
	B07	1, 3	97					<i>S. flexneri</i> bacteriophage V (16) ORFs <i>orf5</i> and <i>ORF6</i> (98% homology over 803 nt), encoding a capsid protein and a hypothetical protein
	A14	2, 3	474	0/17 (0)	17/17 (100)	4/5 (80)	5/5 (100)	599 nt of DNA sequence without significant homology to known DNA sequences adjacent to <i>E. coli</i> Sakai (14) ORFs ECs1599 and ECs1600 (90 to 91% homologous over 367 nt), encoding hypothetical proteins
	A15	1, 2, 3	1,068	0/17 (0)	17/17 (100)	4/5 (80)	4/5 (80)	247 nt of DNA sequence without significant homology to known DNA sequences adjacent to <i>E. coli</i> Sakai (14) ORFs ECs1557 and ECs1558 (96% homology over 1,968 nt), encoding a putative antirepressor protein and a putative tail assembly protein
	A06c	2	1,212	1/17 (5.9)	17/17 (100)	5/5 (100)	5/5 (100)	<i>E. coli</i> Sakai (14) intergenic region between ORFs ECs1159 and ECs1160 (98% homology over 365 nt), encoding a hypothetical protein and a putative integrase, adjacent to ECs1252 (95% homologous over 929 nt with a single 82-nt gap), encoding a putative transport protein
	D	A07	1, 3	226	0/17 (0)	15/17 (88.2)	5/5 (100)	4/5 (80)

	3	558	0/17 (0)	17/17 (100)	1/5 (20)	5/5 (100)	
B15	3	558	0/17 (0)	17/17 (100)	1/5 (20)	5/5 (100)	<i>E. coli</i> Sakai (14) ORF ECs1688 (100% homology over 378 nt), encoding a murein transglycosylase, adjacent to ORFs ECs1705 and ECs1706 (99% homologous over 672 nt), encoding a putative dihydroxyacetone kinase and a hypothetical protein
E	1	432	0/17 (0)	12/17 (70.6)	0/5 (0)	2/5 (40)	<i>E. coli</i> strain EC93 contact-dependent inhibition ( <i>cdtB</i> ) gene (3) (98% homologous over 523 nt)
B05	1	512	0/17 (0)	1/17 (5.9)	0/5 (0)	0/5 (0)	ColE6-CT14 colicin plasmid (1) colicin E6 gene (99% homologous over 1,605 nt)
B12	2, 3	258	0/17 (0)	17/17 (100)	0/5 (0)	5/5 (100)	<i>E. coli</i> UTI89 (7) ORFs (91% homologous over 710 nt), encoding a regulatory protein and a hypothetical protein
B18	1	290	5/17 (29.4)	8/17 (47.1)	0/5 (0)	2/5 (40)	pColD-157 plasmid (15) colicin D-157 activity protein (99% homology over 921 nt)
F	1, 2, 3	446	0/17 (0)	16/17 (94.1)	4/5 (80)	4/5 (80)	643 nt of DNA sequence without significant homology to known DNA sequences
B13	1, 2	507	0/17 (0)	14/17 (82.4)	4/5 (80)	5/5 (100)	DNA sequence without significant homology to known DNA sequences

\* Sequence found in the following SSH experiments: 1, *E. coli* FRIK 920-*E. coli* Sakai; 2, *E. coli* FRIK 1999-*E. coli* Sakai; 3, *E. coli* FRIK 2001-*E. coli* 93-001.

identity to *E. coli* O157:H7 strain Sakai (A), non-Sakai bacteriophage (B), transitional sequences between lineage-common and lineage II strain-specific DNA (C), sequence deletions in the *E. coli* O157:H7 lineage II strain with respect to strain Sakai (D), nonbacteriophage DNA found in bacteria other than O157 strain Sakai (E), and those bearing little or no sequence identity to sequences in the GenBank NR database (F). Because many contigs contained sequences that could be ascribed to more than one class, the relative proportions of the different classes exceed 100%.

Class A contigs possessed at least 90% identity over at least 90% of their length to sequences from *E. coli* Sakai (14) and likely represent driver DNA contamination or lineage II sequences with only minor deviations from that in the lineage I *E. coli* Sakai genome. These sequences accounted for 503 (62.0%) of the contigs and were not analyzed further. The remaining 308 contigs were reasoned to be enriched for sequences that are unique to lineage II strains. In order to determine the distribution of these DNA sequences in our *E. coli* O157:H7 collection of 44 strains of different LSPA-6 genotypes (Table 1), PCR primers were designed to detect representative target sequences from these contigs (Table 2). The 27 PCR targets tested included 11 of the 16 sequences detected in all three SSH experiments, 11 of the 73 sequences detected in two SSH experiments, and five sequences detected in single SSH experiments (Tables 3 and 4). Class B, C, E, and F sequences represent insertions of DNA segments in lineage II strains with respect to the lineage I *E. coli* Sakai genome, and class D sequences represent deletions in lineage II strains with respect to the position of the element in the *E. coli* Sakai genome. The approximate location of these contigs and their corresponding PCR targets in the *E. coli* O157:H7 genome are shown in Fig. 1.

**Distribution of lineage II SSH sequences among *E. coli* O157:H7 strains.** Class B loci possessed at least 80% sequence identity over at least 50 bp to bacteriophage sequences not found in *E. coli* O157:H7 lineage I strain Sakai (Table 4). These sequences were present in 92 (11.3%) of the 811 contigs, ranged in size from 66 to 3,180 bp, and represented 73,574 of the 565,341 nucleotides (13.0%) from all of the contigs. The 16 primer sets from class B loci amplified DNA from the majority of lineage II strains but not from any of the lineage I strains. Some of these primer sets generated identical strain distribution patterns and were derived from contigs sharing homology with different segments of a single prophage that may be unique to lineage II strains. The class B target sequences share identity with several genes of other lambdoid prophage, including prophage from *E. coli* UTI89 (7), *stx*<sub>1</sub>-converting bacteriophage BP-4795 from *E. coli* O84:H4 4795/97 (8), cryptic prophage e14 from *E. coli* K-12 (5), *E. coli* bacteriophage P2 (accession number AF063097), P4-like bacteriophage from *E. coli* H709c (23), two different *Shigella boydii* Sb227 bacteriophages (45), and *E. coli* bacteriophage HK022 (19). It is well known that bacteriophages are major contributors of genetic diversity in *E. coli* O157 and other enteric bacteria (21, 22, 34, 39, 40). Further characterization of these lineage II conserved bacteriophages will be required to determine their effect on the phenotype of this group of *E. coli* O157:H7 strains.

Class C loci likely represent transitional sequences between lineage-common and strain- or lineage-specific DNA. They





in size from 438 to 1,427 bp, and represented 17,213 (3.0%) of all contig nucleotides. Two of the class D primer sets, A06c and A07, amplify sequences flanking the insertion sites of the Sakai Stx2-encoding Sp5 and the NLE-encoding Sp10, respectively. In contrast to all *E. coli* O157:H7 lineage I strains, most lineage II strains produced amplicons with these primers, showing that Sp5 and Sp10 are not present in the respective lineage I chromosomal prophage insertion sites. This finding is supported by other studies that have shown that the chromosomal location of the *stx*<sub>2</sub>-transducing prophage Sp5 varies among *E. coli* O157:H7 strains (35, 40). The third primer set was based on a contig with identity to S-loop 83. In lineage II strains, the *E. coli* Sakai open reading frame (ORF) ECs1688 is juxtaposed next to ECs1705 due to an apparent insertion/deletion in the intervening ORFs. This region encodes a putative iron transport gene cluster with identity to the *prxA-modD-yc73-fepC* region of the pyonephritis and cystitis pathogenicity island of *E. coli* CFT073 (13). All three of the class D loci were also identified in a previous study of lineage I-specific genomic segments (42). In the latter study, a segment of DNA in the tellurite resistance and adherence-conferring genomic island carrying the *perC*-like regulatory protein gene, *pchD*, was also found to be missing in lineage II strains. The fact that the genes contained within these regions include *stx*<sub>2</sub> in Sp5, a putative iron transport gene cluster in S-loop 83, and a gene with similarity to the *perC*-like LEE1 regulator in the tellurite resistance and adherence-conferring genomic island provides strong evidence of key differences in virulence-associated traits between these two lineages.

Class E loci possessed at least 80% sequence identity over at least 50 bp to nonbacteriophage DNA sequences found in bacteria other than *E. coli* O157:H7 strain Sakai, ranged in size from 51 to 3,110 bp, and represented 9.9% (80 of 811) of the contigs or 54,989 (9.7%) of all contig nucleotides. Primer set A01 amplified a target with identity to the *E. coli* strain EC93 *cdiB* gene (3). The *cdiB* gene is part of a cell contact-dependent growth inhibition system in *E. coli* EC93 and uropathogenic *E. coli* strains, which has been shown to inhibit in vitro growth of *E. coli* K-12 (3). While the *cdiB* gene was highly prevalent among lineage II strains of LSPA-6 222222 (Table 4), lineage I strains were missing this operon. The lineage II conserved region amplified by primer set B12 has identity to a putative regulatory gene and two hypothetical genes found in the uropathogenic *E. coli* strain UTI89 (7) and the avian pathogenic *E. coli* serotype O1 strain (18). This sequence was absent from lineage I strains but present in all lineage II strains except for genotype LSPA-6 211111 strains.

The other two class E PCR primer sets, B05 and B18, represent colicin genes identified at high frequency within the *E. coli* FRIK 920-*E. coli* Sakai SSH library but not in either of the two remaining SSH experiments. These targets were not lineage specific. The colicin E gene was amplified by the B05 primers and was found only in the strain FRIK 920, while the colicin D gene targeted by primer set B18 was found among both lineages but more commonly among genotype LSPA-6 222222 strains (8/17 or 47.1%) than lineage I genotype LSPA-6 111111 strains (5/17 or 29.4%) (Table 4).

Class F contigs had very little or no identity to sequences in the GenBank NR database. These were >50 bp in length and contained <80% sequence identity over any 50-bp stretch

within the contig or contained no region of >50 bp in length with >80% sequence identity (criteria used for inclusion in classes B, C, and E). Class F represented 23.2% (188/811) of the contigs, ranged in size from 51 bp to 3,485 bp, and included 150,034 (26.5%) of all contig nucleotides. PCR analysis with the A12 and B13 primer sets confirmed that these segments are present in a large number of lineage II strains and are therefore not due to artifacts of the SSH methodology. Although the function of these genes is unknown, the fact that they are conserved in lineage II strains, coupled with their prevalence within the SSH libraries (23.3%), would argue that they play an important functional role in the survival of the organism.

PCR amplicons generated from tester strains were sequenced and compared with the original contigs to confirm correct assembly of the SSH sequences. The sequences obtained were 89 to 100% identical to the original contigs, suggesting that the original assemblies were correct and representative of the actual genomic sequences. The high proportion of primer sets (25/27) that amplified these sequences in the *E. coli* O157:H7 lineage II strains tested illustrates that SSH is a robust method for isolating strain-specific sequences.

**Genotyping of *E. coli* O157:H7 strains based on lineage II conserved loci.** Sequences identified in this study suggest that genomic content and organization vary within lineage I and lineage II populations. Even within a lineage, diversity can be observed, as few loci are conserved in every isolate tested within a lineage. Nonetheless, when heritability of the SSH markers is examined as a whole, the extent of genomic diversity distinguishing the two lineages is impressive. As illustrated in Fig. 2, neighbor-joining analysis of the PCR data (converted to binary strings) shows that the major LSPA-6 genotypes behave as a group with respect to genomic diversity. With the exception of a single LSPA-6 111111 strain, human phage type 14 strain ECI-485, *E. coli* O157:H7 strains from lineage I (LSPA-6 111111) and from lineage II (other LSPA-6 genotypes) clustered separately (ECI-485 clustered with the lineage II strains, not with the other lineage I strains). While the LSPA-6 markers are within backbone regions of the genome, many of the lineage-specific loci found in this study are located within mobile genetic elements such as phage; this congruence implies that selection is uniquely shaping the genomes of these two lineages.

Within lineage II, the LSPA-6 genotype 211111 strains showed significant strain-strain diversity (Fig. 2). We have previously shown that LSPA-6 genotype 211111 strains group separately from other *E. coli* O157:H7 strains and share genomic characteristics of lineage I or lineage II populations at independent loci (42), leading us to categorize them as lineage I/II strains. This finding, coupled with the fact that the lineage II SSH markers are not uniformly distributed within LSPA-6 genotype 211111, suggests that there is significant diversity among lineage I/II strains and that they are distinct from lineage I and lineage II strains (47).

Statistical analysis of the distribution of lineage I and lineage II strains has previously shown biases in the frequency with which some subtypes are found in bovine or human clinical samples (46). In this report and our previous studies, we have identified genomic features that differentiate populations of these lineages. As observed in other studies, bacteriophages are important contributors to genomic diversity in this organ-




FIG. 2. Phylogenetic relatedness of *E. coli* O157:H7 strains based on the presence of 18 lineage II conserved molecular markers. A neighbor-joining dendrogram was generated based on a matrix of pairwise distances calculated from the binary data by using the proportion of shared lineage II markers as a distance metric. Marker distribution is shown on the right, with positive PCR results indicated by black boxes and negative results indicated by white boxes. The level of support as a proportion of 500 bootstrap replicates is indicated on each branch on the dendrogram.

ism. Whether these phages carry unique combinations of virulence genes and regulatory genes or otherwise influence physiological traits of the two lineages remains to be determined.

A large number of studies now have confirmed that genomic alterations associated with bacteriophage differentiate subpopulations of *E. coli* O157:H7 (34, 39, 40). Unfortunately, many of these studies have not been done within a phylogenetic context, so it is difficult to determine how those results relate to lineage I and II groupings. Nonetheless, collectively,

the studies show that the phage-mediated events are common and that they are also the common events discriminating the two lineages. Since both lineages coexist temporally and spatially, we would expect that they would be exposed to similar phage pools in nature. However, it would appear that these lineages have different host distributions and that they may have different selective pressures or differ in the strategies used to deal with these selective forces. It is also possible that they are uniquely susceptible to lysogeny by certain bacteriophages.

Recently we have shown phage type to be lineage-related in *E. coli* O157:H7 strains (48). The mode of acquisition of these phage-related elements and their effect on phenotype are at present unknown but will become clearer as these lineage-specific loci begin to be studied.

A number of the differentially distributed elements have the potential to affect the ecology and pathogenicity of *E. coli* O157:H7. Within lineage I strains, these include genomic regions containing the *stx*<sub>2</sub>-transducing Sakai prophage Sp5 and a putative iron transport gene cluster with homology to the *prfA-modD-yc73-fepC* gene cluster of *E. coli* CFT073 (13). Within lineage II strains, these include genes for a putative contact-dependent inhibition system (3) and a putative regulatory gene found within the uropathogenic *E. coli* strain UTI89. In addition, this study identified the existence of other bacteriophage-related elements that appear to be conserved among lineage II *E. coli* O157:H7 strains. These may possess factors which influence the virulence and host distribution of *E. coli* O157:H7 lineages.

#### ACKNOWLEDGMENTS

We thank Shelley Frost for technical assistance and the Canadian Food Inspection Agency (CFIA) for allowing part of this research to be conducted at the Animal Diseases Research Institute, Lethbridge, Alberta, Canada.

This research was supported by grants from Health Canada's Office of Biotechnology and Science.

#### REFERENCES

- Akutsu, A., H. Masaki, and T. Ohta. 1989. Molecular structure and immunity specificity of colicin E6, an evolutionary intermediate between E-group colicins and cloacin DF13. *J. Bacteriol.* **171**:6430–6436.
- Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Aoki, S. K., R. Pamma, A. D. Hernday, J. E. Bickham, B. A. Braaten, and D. A. Low. 2005. Contact-dependent inhibition of growth in *Escherichia coli*. *Science* **309**:1245–1248.
- Blanco, M., N. L. Padola, A. Kruger, M. E. Sanz, J. E. Blanco, E. A. Gonzalez, G. Dahbi, A. Mora, M. I. Bernardes, A. I. Etcheverria, G. H. Arroyo, P. M. Lucchesi, A. E. Parma, and J. Blanco. 2004. Virulence genes and intimin types of Shiga-toxin-producing *Escherichia coli* isolated from cattle and beef products in Argentina. *Int. Microbiol.* **7**:269–276.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453–1474.
- Boerlin, P., S. A. McEwen, F. Boerlin-Petzold, J. B. Wilson, R. P. Johnson, and C. L. Gyles. 1999. Associations between virulence factors of Shiga toxin-producing *Escherichia coli* and disease in humans. *J. Clin. Microbiol.* **37**:497–503.
- Chen, S. L., C. S. Hung, J. Xu, C. S. Reigstad, V. Magrini, A. Sabo, D. Blasiar, T. Bieri, R. R. Meyer, P. Ozersky, J. R. Armstrong, R. S. Fulton, J. P. Latreille, J. Spieth, T. M. Hooton, E. R. Mardis, S. J. Hultgren, and J. I. Gordon. 2006. Identification of genes subject to positive selection in uropathogenic strains of *Escherichia coli*: a comparative genomics approach. *Proc. Natl. Acad. Sci. USA* **103**:5977–5982.
- Creuzburg, K., J. Recktenwald, V. Kuhle, S. Herold, M. Hensel, and H. Schmidt. 2005. The Shiga toxin 1-converting bacteriophage BP-4795 encodes an NleA-like type III effector protein. *J. Bacteriol.* **187**:8494–8498.
- Dowd, S. E., and H. Ishizaki. 2006. Microarray based comparison of two *Escherichia coli* O157:H7 lineages. *BMC Microbiol.* **6**:30.
- Elder, R. O., J. E. Keen, G. R. Siragusa, G. A. Barkocy-Gallagher, M. Koohmaraie, and W. W. Laegreid. 2000. Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proc. Natl. Acad. Sci. USA* **97**:2999–3003.
- Felsenstein, J. 1989. PHYLIP-phylogeny inference package (version 3.2). *Cladistics* **5**:164–166.
- Griffin, P. M. 1998. Epidemiology of Shiga toxin-producing *Escherichia coli* infections in humans in the United States, p. 15–22. In J. B. Kaper and A. D. O'Brien (ed.), *Escherichia coli* O157:H7 and other Shiga toxin-producing *E. coli* strains. American Society for Microbiology, Washington, DC.
- Guyer, D. M., J. S. Kao, and H. L. Mobley. 1998. Genomic analysis of a pathogenicity island in uropathogenic *Escherichia coli* CFT073: distribution of homologous sequences among isolates from patients with pyelonephritis, cystitis, and catheter-associated bacteriuria and from fecal samples. *Infect. Immun.* **66**:4411–4417.
- Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C. G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**:11–22.
- Hofinger, C., H. Karch, and H. Schmidt. 1998. Structure and function of plasmid pColD157 of enterohemorrhagic *Escherichia coli* O157 and its distribution among strains from patients with diarrhea and hemolytic-uremic syndrome. *J. Clin. Microbiol.* **36**:24–29.
- Huan, P. T., B. L. Whittle, D. A. Bastin, A. A. Lindberg, and N. K. Verma. 1997. *Shigella flexneri* type-specific antigen V: cloning, sequencing and characterization of the glucosyl transferase gene of temperate bacteriophage SfV. *Gene* **195**:207–216.
- Jarvis, K. G., and J. B. Kaper. 1996. Secretion of extracellular proteins by enterohemorrhagic *Escherichia coli* via a putative type III secretion system. *Infect. Immun.* **64**:4826–4829.
- Johnson, T. J., S. Kariyawasam, Y. Wannemuehler, P. Mangiamale, S. J. Johnson, C. Doetkott, J. A. Skyberg, A. M. Lynne, J. R. Johnson, and L. K. Nolan. 2007. The genome sequence of avian pathogenic *Escherichia coli* strain O1:K1:H7 shares strong similarities with human extraintestinal pathogenic *E. coli* genomes. *J. Bacteriol.* **189**:3228–3236.
- Juhala, R. J., M. E. Ford, R. L. Duda, A. Youtton, G. F. Hatfull, and R. W. Hendrix. 2000. Genomic sequences of bacteriophage HK97 and HK022: pervasive genetic mosaicism in the lambdaoid bacteriophage. *J. Mol. Biol.* **299**:27–51.
- Karmali, M. A., B. T. Steele, M. Petric, and C. Lim. 1983. Sporadic cases of haemolytic-uraemic syndrome associated with faecal cytotoxin and cytotoxin-producing *Escherichia coli* in stools. *Lancet* **i**:619–620.
- Kim, J., J. Niefeldt, and A. K. Benson. 1999. Octamer-based genome scanning distinguishes a unique subpopulation of *Escherichia coli* O157:H7 strains in cattle. *Proc. Natl. Acad. Sci. USA* **96**:13288–13293.
- Kim, J., J. Niefeldt, J. Ju, J. Wise, N. Fegan, P. Desmarchelier, and A. K. Benson. 2001. Ancestral divergence, genome diversification, and phylogeographic variation in subpopulations of sorbitol-negative, β-glucuronidase-negative enterohemorrhagic *Escherichia coli* O157. *J. Bacteriol.* **183**:6885–6897.
- Kita, K., J. Tsuda, T. Kato, K. Okamoto, H. Yanase, and M. Tanaka. 1999. Evidence of horizontal transfer of the EcoO1091 restriction-modification gene to *Escherichia coli* chromosomal DNA. *J. Bacteriol.* **181**:6822–6827.
- Laegreid, W. W., R. O. Elder, and J. E. Keen. 1999. Prevalence of *Escherichia coli* O157:H7 in range beef calves at weaning. *Epidemiol. Infect.* **123**:291–298.
- Laing, C., C. Pegg, D. Yawney, K. Ziebell, M. Steele, R. Johnson, J. E. Thomas, E. N. Taboada, Y. Zhang, and V. P. J. Gannon. 2008. Rapid determination of *Escherichia coli* O157:H7 lineage types and molecular subtypes by using comparative genomic fingerprinting. *Appl. Environ. Microbiol.* **74**:6606–6615.
- Lejeune, J. T., S. T. Abedon, K. Takemura, N. P. Christie, and S. Sreevatsan. 2004. Human *Escherichia coli* O157:H7 genetic marker in isolates of bovine origin. *Emerg. Infect. Dis.* **10**:1482–1485.
- Locking, M. E., S. J. O'Brien, W. J. Reilly, E. M. Wright, D. M. Campbell, J. E. Coia, L. M. Browning, and C. N. Ramsay. 2001. Risk factors for sporadic cases of *Escherichia coli* O157 infection: the importance of contact with animal excreta. *Epidemiol. Infect.* **127**:215–220.
- Malone, A. S., A. E. Yousef, and J. T. LeJeune. 2007. Association of prophage antiterminator Q alleles and susceptibility to food-processing treatments applied to *Escherichia coli* O157 in laboratory media. *J. Food Prot.* **70**:2617–2619.
- Manning, S. D., A. S. Motiwala, A. C. Springman, W. Qi, D. W. Lacher, L. M. Ouellette, J. M. Mladonicky, P. Somsel, J. T. Rudrik, S. E. Dietrich, W. Zhang, B. Swaminathan, D. Alland, and T. S. Whitam. 2008. Variation in virulence among clades of *Escherichia coli* O157:H7 associated with disease outbreaks. *Proc. Natl. Acad. Sci. USA* **105**:4868–4873.
- McNally, A., A. J. Roe, S. Simpson, F. M. Thomson-Carter, D. E. Hoey, C. Currie, T. Chakraborty, D. G. Smith, and D. L. Gally. 2001. Differences in levels of secreted locus of enterocyte effacement proteins between human disease-associated and bovine *Escherichia coli* O157. *Infect. Immun.* **69**:5107–5114.
- Nataro, J. P., and J. B. Kaper. 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**:142–201.
- O'Brien, A. D., V. L. Tesh, A. Donohue-Rolfe, M. P. Jackson, S. Olsnes, K. Sandvig, A. Lindberg, and G. T. Keusch. 1992. Shiga toxin: biochemistry, genetics, mode of action, and role in pathogenesis. *Curr. Top. Microbiol. Immunol.* **180**:65–94.
- O'Brien, S. J., G. K. Adak, and C. Gilham. 2001. Contact with farming environment as a major risk factor for Shiga toxin (Vero cytotoxin)-produc-

- ing *Escherichia coli* O157 infection in humans. *Emerg. Infect. Dis.* **7**:1049–1051.
34. Ohnishi, M., K. Kurokawa, and T. Hayashi. 2001. Diversification of *Escherichia coli* genomes: are bacteriophage the major contributors? *Trends Microbiol.* **9**:481–485.
  35. Ohnishi, M., J. Terajima, K. Kurokawa, K. Nakayama, T. Murata, K. Tamura, Y. Ogura, H. Watanabe, and T. Hayashi. 2002. Genomic diversity of enterohemorrhagic *Escherichia coli* O157 revealed by whole genome PCR scanning. *Proc. Natl. Acad. Sci. USA* **99**:17043–17048.
  36. Perna, N. T., G. Plunkett III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grotbeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamouisis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
  37. Sargeant, J. M., M. W. Sanderson, R. A. Smith, and D. D. Griffin. 2003. *Escherichia coli* O157 in feedlot cattle feces and water in four major feeder-cattle states in the USA. *Prev. Vet. Med.* **61**:127–135.
  38. Saridakis, C. E., R. P. Johnson, A. Benson, K. Ziebell, and C. L. Gyles. 2004. Influence of animal origin and lineage on survival of *Escherichia coli* O157:H7 strains in strong and weak acid challenges. *J. Food Prot.* **67**:1591–1596.
  39. Sato, T., T. Shimizu, M. Watarai, M. Kobayashi, S. Kano, T. Hamabata, Y. Takeda, and S. Yamasaki. 2003. Distinctiveness of the genomic sequence of Shiga toxin 2-converting phage isolated from *Escherichia coli* O157:H7 Okayama strain as compared to other Shiga toxin 2-converting phages. *Gene* **309**:35–48.
  40. Shaikh, N., and P. I. Tarr. 2003. *Escherichia coli* O157:H7 Shiga toxin-encoding bacteriophage: integrations, excisions, truncations, and evolutionary implications. *J. Bacteriol.* **185**:3596–3605.
  41. Smith, D., M. Blackford, S. Younts, R. Moxley, J. Gray, L. Hungerford, T. Milton, and T. Klopfenstein. 2001. Ecological relationships between the prevalence of cattle shedding *Escherichia coli* O157:H7 and characteristics of the cattle or conditions of the feedlot pen. *J. Food Prot.* **64**:1899–1903.
  42. Steele, M., K. Ziebell, Y. Zhang, A. Benson, P. Konczy, R. Johnson, and V. Gannon. 2007. Identification of *Escherichia coli* O157:H7 genomic regions conserved in strains with a genotype associated with human infection. *Appl. Environ. Microbiol.* **73**:22–31.
  43. Stothard, P., and D. S. Wishart. 2005. Circular genome visualization and exploration using CGView. *Bioinformatics* **21**:537–539.
  44. Tarr, P. I., S. S. Bilge, J. C. J. Vary, S. Jelacic, R. L. Habeeb, T. R. Ward, M. R. Baylor, and T. E. Besser. 2000. Iha: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.* **68**:1400–1407.
  45. Yang, F., J. Yang, X. Zhang, L. Chen, Y. Jiang, Y. Yan, X. Tang, J. Wang, Z. Xiong, J. Dong, Y. Xue, Y. Zhu, X. Xu, L. Sun, S. Chen, H. Nie, J. Peng, J. Xu, Y. Wang, Z. Yuan, Y. Wen, Z. Yao, Y. Shen, B. Qiang, Y. Hou, J. Yu, and Q. Jin. 2005. Genome dynamics and diversity of *Shigella* species, the etiologic agents of bacillary dysentery. *Nucleic Acids Res.* **33**:6445–6458.
  46. Yang, Z., J. Kovar, J. Kim, J. Nietfeldt, D. R. Smith, R. A. Moxley, M. E. Olson, P. D. Fey, and A. K. Benson. 2004. Identification of common subpopulations of non-sorbitol-fermenting,  $\beta$ -glucuronidase-negative *Escherichia coli* O157:H7 from bovine production environments and human clinical samples. *Appl. Environ. Microbiol.* **70**:6846–6854.
  47. Zhang, Y., C. Laing, M. Steele, K. Ziebell, R. Johnson, A. K. Benson, E. Taboada, and V. P. Gannon. 2007. Genome evolution in major *Escherichia coli* O157:H7 lineages. *BMC Genomics* **8**:121.
  48. Ziebell, K., M. Steele, Y. Zhang, A. Benson, E. N. Taboada, C. Laing, S. McEwen, B. Ciebin, R. Johnson, and V. Gannon. 2008. Genotypic characterization and prevalence of virulence factors among Canadian *Escherichia coli* O157:H7 strains. *Appl. Environ. Microbiol.* **74**:4314–4323.