

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Faculty Publications in the Biological Sciences

Papers in the Biological Sciences

2012

Multiple exposures to drought 'train' transcriptional responses in *Arabidopsis*

Yong Ding

University of Nebraska-Lincoln

Michael Fromm

University of Nebraska-Lincoln, mfromm2@unl.edu

Zoya Avramova

University of Nebraska-Lincoln, zavramova2@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscifacpub>

 Part of the [Biology Commons](http://digitalcommons.unl.edu/bioscifacpub)

Ding, Yong; Fromm, Michael; and Avramova, Zoya, "Multiple exposures to drought 'train' transcriptional responses in *Arabidopsis*" (2012). *Faculty Publications in the Biological Sciences*. 535.

<http://digitalcommons.unl.edu/bioscifacpub/535>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in the Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

ARTICLE

Received 27 Oct 2011 | Accepted 6 Feb 2012 | Published 13 Mar 2012

DOI: 10.1038/ncomms1732

Multiple exposures to drought 'train' transcriptional responses in *Arabidopsis*

Yong Ding¹, Michael Fromm² & Zoya Avramova¹

Pre-exposure to stress may alter plants' subsequent responses by producing faster and/or stronger reactions implying that plants exercise a form of 'stress memory'. The mechanisms of plants' stress memory responses are poorly understood leaving this fundamental biological question unanswered. Here we show that during recurring dehydration stresses *Arabidopsis* plants display transcriptional stress memory demonstrated by an increase in the rate of transcription and elevated transcript levels of a subset of the stress-response genes (trainable genes). During recovery (watered) states, trainable genes produce transcripts at basal (preinduced) levels, but remain associated with atypically high H3K4me3 and Ser5P polymerase II levels, indicating that RNA polymerase II is stalled. This is the first example of a stalled RNA polymerase II and its involvement in transcriptional memory in plants. These newly discovered phenomena might be a general feature of plant stress-response systems and could lead to novel approaches for increasing the flexibility of a plant's ability to respond to the environment.

¹ University of Nebraska School of Biological Sciences, 1901 Vine Street, Lincoln, 68588, USA. ² University of Nebraska Center for Biotechnology and Center for Plant Science Innovation, 1901 Vine Street, Lincoln, 68588, USA. Correspondence and requests for materials should be addressed to Z.A. (email: zavramova2@unlnotes.unl.edu).

Plants' responses to abiotic stresses involve coordinated actions to change cellular physiology and gene expression patterns¹. A previous exposure to a stress may alter a plant's subsequent stress response by producing faster and/or stronger reactions that may provide the benefits of enhanced protection^{2,3}. Altered responses to consecutive stresses imply that plants exercise a form of 'stress memory'⁴. For example, tobacco plants pre-exposed to methyl jasmonate increased nicotine pools 2 days earlier when exposed again as compared with plants without previous exposure⁵; Pretreatment of plants with salicylic acid or its synthetic analogue, benzothiadiazole *S*-methyl ester, resulted in increased transcription from a subset of genes upon a subsequent stress^{6,7}. Some stress memory effects could be perpetuated to the next generation, as observed with flagellin or ultraviolet radiation⁸ treatments. The molecular mechanisms associated with stress memory are largely unknown. Accumulation of dormant signalling molecules that are recruited only on exposure to stress, or proteins involved in their synthesis, have been proposed².

Epigenetic control systems offer an alternative mechanism for responses that are dynamic and persistent: a gene, or a network, is activated in response to a stimulus, but once the stimulus is removed, a mark could be left to facilitate altered, quicker or more potent responses to subsequent stresses. Therefore, epigenetic systems may act as the conduit for environmental cues initiating both short- and long-term changes in gene expression in response to a stress. The formal definition of epigenetics includes a heritable (mitotic or meiotic) component. However, the finding that the functional states of neurons, which almost never divide, involve epigenetic phenomena⁹ has prompted a refined definition suggesting a broader meaning that excludes the strict requirement for heritability¹⁰.

In this paper, we use the term 'epigenetics' within this broader context as we study responses in leaf cells that have predominantly ceased division. It is important to emphasize also that we distinguish between a chromatin mark (reflecting a modification that is dynamically associated with a process, but is removed at the conclusion of that process) and an epigenetic mark to imply persistence longer than the initial stimulus that caused the chromatin mark and that affects subsequent performances of the gene. We define 'transcriptional memory' to mean that a type of information persists after the plant has recovered from the initial stress and that the 'memory' influences subsequent transcriptional responses. We found that during recurring dehydration stresses *Arabidopsis* plants demonstrate transcriptional memory of the stress as displayed by increased rates of transcription and elevated transcript levels from a subset of the stress-response genes (trainable genes). The transcriptional memory is associated with two distinct marks found only at the trainable genes during their recovery from stress-induced transcription: high levels of trimethylated histone H3 Lys4 (H3K4me3) nucleosomes and stalled Ser5P Pol II. In contrast, H3K4me3 and Ser5P dynamically increase when non-trainable genes are induced and then decrease to basal levels during the recovery. At the trainable genes, H3K4me3 and Ser5P polymerase II (Pol II) persist as 'memory marks' for as long as the transcriptional memory responses last. The low transcription rates and high Ser5P Pol II levels during the watered recovery conditions provide the first evidence of a stalled RNA Pol II in plants.

Results

Dehydration stress-trained *Arabidopsis* plants. To determine whether plants retain a memory of a previous dehydration stress (trained plants), we developed a repetitive dehydration/rehydration system of air-drying for 2 h and rehydration for 22 h (Methods). Non-trained plants wilted faster than trained plants (Fig. 1a) and their leaves lost water at a faster rate than trained plants (Fig. 1b). The leaf relative water content (RWC) at the beginning of each stress treatment (measured at the end of the R1–R3 recovery periods) was




Figure 1 | Response to dehydration stress of trained or non-trained plants. (a) Appearance of 3-week-old *Arabidopsis* plants to 1-h exposure to air if not previously stressed (non-trained) or previously stressed with three consecutive cycles of stress/recovery treatments (trained). (b) Fresh weights of leaves, after air drying for the indicated times, for plants experiencing their first stress (dark blue line) or for trained plants exposed to a second (red line), third (brown line) or fourth (light blue line) stress cycles. Experiments in a were repeated at least 5 times. Three independent experiments were performed for b, each with three replicates, each containing 8–10 leaves. The representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates. (c) Relative water content (RWC) measured during the initial watered (W) state and on recovery (R1–R3) from one or more stresses, before a subsequent dehydration stress (second, third or fourth stress) as described in b.

similar to the RWC of initial preinduced watered (W) state (Fig. 1c). The results showed that despite losing water during air exposure, rehydration brought the RWC to the initial prestress levels, indicating plants have fully recovered and start each dehydration stress test at similar initial RWCs. These results are consistent with drought hardening observed in horticultural plants¹¹, supporting the idea that trained plants alter their responses after an initial dehydration stress.

Trainable and non-trainable *Arabidopsis* genes. Trained plants also responded differently at the transcriptional level. We analysed four genes widely used as markers for dehydration-inducible genes (*RD29A*, *RD29B*, *RAB18* and *COR15A* (refs 12–15)) and observed two distinct types of expression patterns during repeated stresses (Fig. 2a,b). The *RD29A* and *COR15A* genes repetitiously increased their transcripts to about the same level during each stress




Figure 2 | Transcript levels of non-trainable and trainable genes in plants before and after single or multiple dehydration stresses. RNA was isolated from watered plants (initial state defined as watered, W) and from plants subjected to one to four dehydration stresses of 2 h of air drying (S1–S4) separated by 22-h intervals of watered recovery treatments (R1–R3). Transcript levels for (a) *RD29A*; (b) *COR15A*; (c) *RD29B*; (d) *RAB18* genes were measured by reverse transcription and real-time quantitative PCR. *UBQ10* was used as an internal control. Genes are considered trained if the transcript level in S1 is considerably less than in subsequent stresses, as occurring for *RD29B* and *RAB18*. Experiments were repeated at least three times, each with three replicates, and the representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates.

(non-trainable genes). In contrast, *RD29B* and *RAB18* produced considerably higher transcript levels during one or more subsequent stresses relative to the initial stress (the trainable genes). A critical result is that during the recovery (watered) states both the non-trainable and the trainable genes return to their initial (non-stressed) transcript levels. Restoration of each gene's activity to its prestressed level when stress signalling was alleviated indicates lack of a persistent induction. Return to baseline transcript levels from the trainable genes during recovery, together with a higher induction level on a following stress, illustrates the concept of 'transcriptional memory'.

To establish whether transcriptional memory was carried also by other dehydration stress–response genes, we searched for genes displaying increased transcript levels in multiply stressed plants. Five additional non-trainable genes (Supplementary Fig. S1a,b) and five additional trainable genes were identified (Supplementary Fig. S1a,c).

Collectively, our results uncovered a hitherto unknown existence of two distinct subsets within the dehydration stress–response gene fraction: the repetitively responding non-trainable genes and the trainable genes, which increased the magnitude of their subsequent transcriptional response, relative to their initial stress response.

Transcription rates of trainable and non-trainable genes. Restoration of transcript levels to the basal non-induced levels during the watered recoveries indicated that response genes' mRNAs were not persisting from one stress to the next. Whether elevated transcript levels were due to increased rates of transcription was determined by nuclear run on assays. The non-trainable *RD29A* and *COR15A* genes had similar transcription rates during each stress treatment (Fig. 3a,b). In contrast, the trainable *RD29B* and *RAB18* genes displayed threefold higher transcription rates during the repeated stress (S4) relative to their rates during the first (S1) stress (Fig. 3c,d). Similar transcription rates during the initial (prestressed) and the recovery (watered) phases for both the trainable and non-trainable genes indicated that in the absence of dehydration stress,




Figure 3 | Nuclear run on assays of non-trained or trained plants. Nuclei were isolated from non-stressed watered (W), singly (S1) or repetitively stressed plants. The repetitively stressed plants were dehydration stressed 3 times and then nuclei isolated before (recovery R3) or during a fourth dehydration stress (S4). Isolated nuclei were used in run on transcription reactions containing biotin-UTP to label newly synthesized transcripts. The nascent biotin-labelled transcripts were purified by binding to streptavidin beads and quantitated by quantitative PCR (qPCR) for (a) *RD29A*; (b) *COR15A*; (c) *RD29B*; (d) *RAB18* genes. Labelling experiments were repeated at three times, each with three reverse transcription–qPCR measurements, and the representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates. *UBQ10* was used as internal control.

transcription was restored to the initial levels and that the increased transcript production by the trainable genes was regulated, at least in part, at the level of transcription.

Distribution of the elongating (Ser2P) Pol II. Next we investigated the possible roles of RNA Pol II and the chromatin histone H3K4me3 modification in the different transcriptional behaviour of the trainable and the non-trainable genes. The distribution of the actively elongating form of Pol II, phosphorylated at serine 2 (Ser2P) of the consensus $(Y_1S_2P_3T_4S_5P_6S_7)_{34}$ CTD repeat, was analysed during the induced and recovery phases of the dehydration–stress cycle. Active transcription elongation is associated with high Ser2P levels accumulating towards the 3'-ends of actively transcribed genes¹⁶. In agreement, the Ser2P distribution profiles showed a peak towards the 3'-ends of both the non-trainable and the trainable genes when induced by stress (Fig. 4a,b).

The Ser2P distribution during both the non-induced and the stress-induced phases of the cycle provided independent support that the transcript levels measured in recovery or during stress were regulated at the level of transcription. First, transcription of both trainable and non-trainable genes during watered recovery returns to their initial levels in watered plants, as Ser2P levels during R3 are similar to those in watered. Second, during stress-induced transcription, the trainable (*RD29B* and *RAB18*) genes accumulate higher Ser2P in S4 than in S1, consistent with their enhanced transcription. Contrasting this pattern, the non-trainable (*RD29A* and *COR15A*) genes accumulate similar Ser2P amounts during S1 or S4, consistent with their transcriptional responses.

The distribution profiles for the elongating Ser2P were also established for the additional five non-trainable and five trainable genes (Supplementary Fig. S1a–c). The results from these non-trainable and trainable groups were similar to the patterns displayed by the non-trainable *RD29A* and *COR15A*, and trainable *RD29B* and *RAB18*, respectively. Thereby, Ser2P levels correlated with transcription rates and transcript levels. As the operational definition of memory marks is that they must last longer than the stimulus, we conclude that Ser2P does not function as a memory mark.




Figure 4 | Profiles of Pol II and of H3K4me3 at the non-trainable and trainable genes. (a) Schematic diagram of the genes, with the promoter region (to left), the 5' untranslated region (grey box), exons (dark box), introns (thin lines between exons) and indicating the regions (1 to 6) analysed by chromatin immunoprecipitation and real-time qPCR (ChIP-qPCR). (b-d) ChIP-qPCR analysis of the amounts of specific modifications present at regions 1-6 for the non-stressed watered (W, dark blue), singly stressed (S1, red) and trained plants (three stress cycles) before recovery (R3, brown) or during a fourth dehydration stress (S4, light blue). The results with *ACT7* are shown as an internal control; (b) distribution profiles of Ser2P Pol II; (c) distribution profiles of Ser5P Pol II; (d) distribution profiles of H3K4me3. Experiments were repeated at least four times, each with three replicates, and the representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates.

Stalled Pol II and H3K4me3 as memory marks. The distribution of the phosphorylated serine 5 form of Pol II (Ser5P), associated with transcription initiation/early elongation or pausing/stalling^{16,17}, was analysed next. Ser5P levels follow the transcription rates and transcript abundances from the non-trainable *RD29A* and *COR15A* genes during all treatments: the initial (S1) and repetitively stressed states (S4) show similar Ser5P levels; similar levels were also found in the prestressed (W) and recovery (R3) states (Fig. 4c). These patterns were confirmed in the five additional non-trainable genes as well (Supplementary Fig. S1b).

For the trainable genes, the Ser5P patterns were different: higher for the repetitively stressed (S4) than for singly stressed (S1) plants (Fig. 4c, Supplementary Fig. S1c), consistent with their transcript levels (Fig. 2) and transcription rates (Fig. 3). The most remarkable difference, however, was the atypically high Ser5P level in the watered conditions, despite the low transcription of *RD29B* and *RAB18*. During recovery (R3), Ser5P was retained at levels similar to the levels generated during the first stress (S1) at *RD29B* and *RAB18* (Fig. 4c). Elevated Ser5P was also displayed by the five additional trainable genes during recovery (Supplementary Fig. S1c). Retention of high Ser5P presence at trained genes during the watered recovery when transcription rates were low suggested that Ser5P Pol II was stalled. Stalled Pol II was found only at the trainable dehydration stress-response genes during their recovery from an induced transcription. Thereby, elevated Ser5P Pol II during recovery behaves as a memory mark from a previously active state.

To determine whether transcriptional memory was also associated with a histone modification, we analysed the H3K4me3 profiles.

For all non-trainable genes, the H3K4me3 modification increased to similar levels on each stress induction and decreased to baseline levels in watered plants (Fig. 4d and Supplementary Fig. S1b). For the trainable genes, H3K4me3 also increased on gene induction, but repetitively stressed (S4) plants displayed higher H3K4me3 than singly stressed (S1) plants (Fig. 4d; Supplementary Fig. S1c). This pattern is in agreement with their increased transcript abundances and transcription rates (Figs 2b,c, and 3b,c). Importantly, however, during recovery (R3) the trainable genes in repetitively stressed plants had higher H3K4me3 levels than in the non-stressed state (F); the elevated H3K4me3 levels were similar to those of the singly stressed (S1) plants (Fig. 4d), indicating that H3K4me3 modification was retained on the nucleosomes of the trainable genes during the recovery states. Retention of high H3K4me3 levels in R3, comparable with S1 was also displayed by the other five trainable genes analysed (Supplementary Fig. S1c). Furthermore, analyses of histone H3 distribution at representative trainable, or non-trainable, genes revealed similar nucleosome levels during the prestressed (W) and the last recovery (R3) states (Fig. 5). The higher H3K4me3 levels in R3 relative to those in watered, thereby, reflect H3K4me3 modifications retained from the preceding transcription.

Therefore, at the non-trainable genes, H3K4me3 acts as a reversible chromatin mark dynamically changing with transcript levels. At the trainable genes, H3K4me3 acts as a persistent epigenetic mark associated with the transcriptional memory displayed by these genes.

Length of the transcriptional memory persistence. To establish how long transcriptional memory can persist in the absence of




Figure 5 | Nucleosomal distribution at the non-trainable and trainable genes. The columns represent the amounts of Histone H3 measured by ChIP-PCR with antibodies specific for histone H3. Numbers under the columns indicate tested regions along the gene sequences as indicated in Fig. 4a. Data from fresh non-stressed tissues (blue columns) and stressed plants in the R3 recovery interval (brown columns) are shown for the non-trainable *RD29A*, *COR15A* and the trainable *RD29B*, *RAB18* genes. Data are from three biological sample replicas.

inducing signals, trained and untrained plants were watered for 3, 5 or 7 days in soil, and then their transcript levels were measured before and after a subsequent dehydration stress. In trained plants, after 3 or 5 days of recovery, the *RD29B* and *RAB18* genes displayed super-induced transcript levels when exposed to dehydration stress. After 7 days, however, when plants were stressed, the *RD29B* and *RAB18* transcripts were induced at non-trained levels (Fig. 6a,b). The non-trainable (*RD29A* and *COR15A*) genes showed little variation in their responses on any of these days (Fig. 6c,d). We conclude that dehydration stress-induced transcriptional memory persists for 5 days, but is lost after 7 days, under watered conditions.

Next, we analysed how long the memory marks persisted in the absence of induced transcription. We found that at the trainable genes, the high Ser5P Pol II and H3K4me3 levels were also retained for 3 and 5 days, but not for 7 days (Fig. 6e–h). Thereby, the similar duration of transcriptional memory and of elevated Ser5P Pol II and H3K4me3 levels strengthens their association and our conclusion that Ser-5P Pol II and H3K4me3 levels function as memory marks at the genes displaying transcriptional memory.

Abscisic acid and transcriptional memory. The phytohormone ABA induces transcription from *Arabidopsis* dehydration stress-response genes¹⁵ and was examined as a potential factor in stress memory responses. Endogenous ABA levels repetitiously increased upon each dehydration stress, but returned to low levels during each recovery treatment (Fig. 6i). This pattern also occurred after the extended recovery periods, as similar levels of endogenous ABA were produced when plants were stressed after 3, 5 or 7 days recovery in watered soil (Fig. 6i). The high ABA levels were similar in each stress cycle and rule out a model wherein higher levels of ABA in subsequent stresses were responsible for the super-induced transcript levels of the trainable genes. The residual amount of ABA after dehydration stress is slightly higher than the initial level in non-stressed plants but, apparently, insufficient to induce transcription in R1–R3. Further, this residual ABA level is present after 3, 5 and 7 days of watering (Fig. 6i), even though transcriptional memory is lost after 7 days of watering (Fig. 6a,b). The results indicate that induced ABA levels were not sufficient to cause super-induced transcript levels and residual ABA levels were not sufficient to maintain a memory response.

Role of ABA-regulated ABRE-binding transcription factors. Recent studies have identified ABA-responsive elements (ABREs) AREB1, AREB2 and ARF3 as basic domain leucine zipper transcription factors (TFs) that act as master regulators to cooperatively regulate gene expression from a large number of plant dehydration stress-response genes¹⁸. These TFs bind to ABREs found in the promoters of ABA-inducible genes and require ABA for their activity. AREB1, AREB2 and ARF3 have largely overlapping functions, therefore we used a plant genotype with mutations in all three genes (*areb1/areb2/arf3* triple mutant)¹⁸ to assess the potential effect of the three TFs on the transcriptional memory response. This

analysis revealed that transcript levels from both the non-trainable (*RD29A* and *COR15A*) and trainable (*RD29B* and *RAB18*) genes were strongly diminished in the triple mutant plants (Fig. 7a–d). However, despite the strong reduction in transcription, both *RD29B* and *RAB18* produced much higher levels of transcripts in S4 than in S1 (Fig. 7c,d), indicating that transcriptional memory was still functional. Thereby, although critical for inducing maximal transcription in response to dehydration stress, AREB1, AREB2 and ARF3 are not essential for the transcriptional memory response.

Role of ATX1 in transcriptional memory. As *Arabidopsis* Trithorax-like 1 (*ATX1*) regulates a large number of dehydration stress-response genes^{19–21}, its potential role in the memory responses was investigated here. Dehydration-induced transcript levels were strongly diminished in *atx1* plants, consistent with its activating role at these genes (Fig. 8a–d). However, the trainable *RD29B* and *RAB18* genes still produced increased transcripts in trained, relative to untrained, *atx1* plants (Fig. 8c,d) and retained H3K4me3 levels at elevated levels during the watered recovery states (Fig. 8e). Trained or untrained *atx1* plants also showed differences in the rates of leaf water loss, albeit smaller than the differences between trained and untrained wild-type plants (Fig. 8f). Together these results indicate that transcriptional memory responses were attenuated in *atx1* mutants but stress memory was still occurring.

Discussion

An altered ability of *Arabidopsis* leaf cells to retain water during repetitive exposures to dehydration stress (Fig. 1) as well as the increased transcriptional responses from a subset of genes (Fig. 2, Supplementary Fig. S1) illustrate the concept of a memory from an earlier stress. Our method for applying multiple dehydration stress/recovery treatments in a relatively short time, as opposed to prolonged soil-water withdrawal/recovery treatments, allowed us to observe this phenomenon and may explain why it has evaded detection in *Arabidopsis* so far. It is particularly relevant to point out that our experimental system is similar to the day/night oscillations in the leaf water potential that plants experience in the field during low soil water potential conditions^{22–24}. It is tempting to suggest that under natural conditions stress memory is activated by the previous day's dehydration stress, persists through the recovery period at night (when transpiration is less and leaf water potential recovers) and then facilitates the plant's response to dehydration stress encountered during the next day, when adequate soil water is unavailable to maintain leaf water potentials at the higher transpiration rates occurring during the day. The duration of dehydration stress memory, persisting for up to 5 days in trained *Arabidopsis* plants recovering in watered soil (Fig. 6), is sufficiently long and may be a mechanism used by plants during soil-water deficit conditions.

A well-studied example of epigenetic memory in plants occurring under natural conditions is the regulation of flowering time in *Arabidopsis* upon vernalization, which involves repression of the




Figure 6 | Length of the transcriptional memory in trained plants. The relative levels of mRNA of trained plants after a recovery period under watered conditions for 3 or more days, before and after a subsequent dehydration stress are shown in **a–d**. Plants were not stressed (Control) or were dehydration stressed 4 times (Trained), with 22-h recovery periods before receiving water in soil for 3, 5 or 7 days (first, second and third columns, respectively). After the indicated days of watering, plants received water (W) or were dehydration stressed (D) for 2 h before the indicated analyses. **(a,b)** Transcript levels of the trainable (*RD29B* and *RAB18*) genes. **(c,d)** Transcript levels of the non-trainable (*RD29A* and *COR15A*) genes measured by reverse transcription–quantitative PCR (qPCR). *UBQ10* was used as an internal control. **(e)** H3K4me3 levels at the *RD29B* gene measured by ChIP–qPCR, relative to *ACT7*, shown in **f** at the regions showing the peak accumulation of the respective marks at each gene (see Fig. 4 for location of the peak regions). **(g)** Ser5P levels at the *RD29B* gene measured by ChIP–qPCR, relative to *ACT7* shown in **h**, at the regions showing the peak accumulation of the respective marks at each gene (see Fig. 4 for location of the peak regions). Results from measurements during days 3, 5 or 7 after recovery in soil are shown in the first, second and third columns, respectively. **(i)** ABA levels before stress (W), after one to four stresses (S1–S4) or one to three recoveries (R1–R3) and before (W) or after a dehydration stress (D) that was initiated after 5 or 7 days of watering of control or trained plants. Experiments in **a–c** were repeated at least three times, each with three replicates, and the representative experiments shown indicates the mean \pm s.e.m., $n=3$ replicates. Experiments in **d** were performed twice, with two replicates, and the representative experiment shown indicates the mean \pm range, $n=2$ replicates.

homeotic *FLC* gene. Prolonged cold initiates pathways that are maintained for the remainder of the plant's life cycle^{25,26}. We emphasize that the epigenetic memory at the *FLC* locus involves repressive polycomb group and small RNA-mediated silencing mechanisms²⁷. In contrast, the vernalization-regulated transcription of the *VRN1* gene is positively regulated. Its induced transcription is maintained throughout the plant development^{28,29} but it does not show intervening periods of low transcription. The inducible transcriptional memory described here is different from the persistent expression of *VRN1*, as the transcription of the trainable gene cycles in correlation with the stress or recovery conditions, indicating that memory is preserved during the low-expression recovery periods.

A signature feature of the trained genes is the preservation of elevated H3K4me3 during their transcriptionally less active recovery periods. Changes in histone modifications, including H3K4me3, in response to environmental stresses have been reported for cold treatments, high salinity, hypoxia and drought-induced transcrip-

tion from response genes in a variety of plant species and tissue cultures^{30–32}. However, maintenance of these stress-induced chromatin changes after the removal of the initial signal has not been reported, except for some plant defence–response genes. H3K4me3 accumulates on the nucleosomes of defence–response genes before their induction by a pathogen attack³³ or upon chemical priming^{6,7}. Priming leads to an increased systemic immunity response on subsequent treatments and stronger transcriptional induction of response-related genes relative to non-exposed plants^{6,7,34,35}. Elevated H3K4me3 levels in the primed state and the subsequent stronger transcriptional response from primed defence genes resembles the transcriptional memory described here. A major difference, however, is that priming establishes H3K4me3 before activation of transcription from defence genes^{6,7}. As such, the H3K4me3 does not occur as a mark retained from a previously active state (as with trainable genes) but, rather, appears as a 'preparatory' mark for future inductions³³.

Transcriptional memory, lasting 1–12 h and involving the H3K4me3 modification, has been reported for *Saccharomyces cerevisiae* genes^{36–39}. A major distinction, however, is that the memory mechanism in yeast does not involve stalled Pol II. Stalled Ser5P Pol II during recovery from previous transcriptionally active periods is another signature feature of the *Arabidopsis* trained genes.




Figure 7 | The role of three ABA-dependent transcription factors in the transcriptional memory. The effects of three ABA-dependent transcription factors, AREB1, AREB2 and ABF3, were examined in the wild-type (dark blue columns) and *areb1/areb2/abf3* triple mutant backgrounds (light blue columns) before and after multiple dehydration stresses. Transcript levels were determined in plants during their prestressed (W), first stressed (S1) and fourth stressed (S4) periods. Transcript levels of the non-trainable (a) *RD29A* and (b) *COR15A* genes. Transcript levels of the trainable (c) *RD29B* and (d) *RAB18* genes. Note the scale for *RD29B* is logarithmic. *UBQ10* was used as an internal control. Experiments were repeated two times, each with three replicates, and the representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates.

This study provides the first example of a stalled Pol II at plant genes. Found at the 5'-ends of inactive metazoan developmental, or stress regulated genes, stalled (paused) Pol II has been proposed to enhance rapid transcriptional responses⁴⁰ or to allow for a synchronous induction of the same gene sets in distinct cell populations at a particular developmental stage⁴¹. Although the molecular mechanism is unclear, stalled Pol II is associated with the massively increased transcription during later inductions¹⁷. A major difference from the trainable *Arabidopsis* genes is that Ser5P Pol II is present at the developmental animal genes before their transcriptional activation¹⁷. In contrast, the Ser5P Pol II levels at the plant genes are low before induction, but remain at elevated levels across a broader region of the gene during the recovery that follows active transcription (Fig. 4, Supplementary Fig. S1).

The most important remaining question is what factors cause transcriptional memory and the Ser5P and H3K4me3 association with the trainable genes. To address this question we analysed the possible involvement of ABA, a key signalling molecule in dehydration stress, of three master TFs regulated by ABA, and of the histone H3K4me3 methyltransferase, ATX1, regulating the expression of the examined genes.

ABA signalling has a critical role in the dehydration stress response and regulates transcription from a large number of *Arabidopsis* genes⁴², including those studied here. Earlier, in genome-wide analyses, we found that the ABA-inducible genes displayed a broader H3K4me3 distribution profiles along the genes' sequences that was distinct from the typical H3K4me3 accumulation at +300 bp downstream of the transcription start sites⁴³. Here we establish that this broader H3K4me3 profile is a feature of both non-trainable and trainable response genes.

During each dehydration stress, endogenous ABA increased to about the same level (Fig. 6i), thus precluding a model wherein higher induction of ABA levels can account for the super-induction of trainable genes observed in S4 versus S1. Moreover, after 7 days of recovery in watered soil, exposure of trained plants to a dehydration stress results in a non-trained transcriptional response, despite comparable levels of ABA in plants that were dehydration stressed




Figure 8 | Role of ATX1 in the dehydration stress memory. Transcript levels in wild-type (WS, shown in dark blue) or *atx1* (shown in light blue) plants measured before stress (W), after one to four stresses (S1–S4), or one to three recoveries (R1–R3). Results relative to *UBQ10* are shown for the non-trainable (a) *RD29A*, (b) *COR15A* and the trainable (c) *RD29B* and (d) *RAB18* genes; (e) H3K4me3 levels measured by ChIP-quantitative PCR during the watered or dehydration stress treatments for WS (dark blue) and *atx1* (light blue) plants for the trainable *RD29B* gene; the profiles of the control *ACT7* gene is shown in the panel to the right. (f) Rates of water loss from detached leaves of wild-type (WS) or *atx1* plants during a first or fourth dehydration stresses: dark blue and red lines illustrate data for WS, first and fourth stresses, respectively; brown and light blue lines illustrate data for *atx1* during the first and fourth stresses, respectively. Experiments were repeated at least twice, each with three replicates, and the representative experiment shown indicates the mean \pm s.e.m., $n = 3$ replicates.

after recovery for 5 or 7 days (Fig. 6a,b). The loss of transcriptional memory by day 7 coincided with the loss of the elevated levels of Ser5P and H3K4me3 at the trainable genes. Thereby, despite the presence of high ABA levels induced by the stress, a trained transcriptional response is not displayed when the memory and signature marks are lost.

We also note that during recovery, ABA levels are slightly higher than in the initial watered stage (Fig. 6i). Importantly, higher baseline ABA levels were present 5 days after watering when transcriptional memory was still active and were also present 7 days after watering (Fig. 6i) when transcriptional memory was no longer active (Fig. 6a,b). This result demonstrates that the higher ABA basal levels during recovery periods are not sufficient for maintaining the transcriptional memory. Therefore, we conclude that although ABA is critically required for inducing the transcriptional responses from both non-trainable and trainable genes¹⁵, ABA levels do not explain the super-induction of trained genes or the maintenance of transcriptional memory in the recovery state.

Persistence of higher levels of key TFs after the first stress event is another potential mechanism for transcriptional memory. Our results here indicate that accumulation of active TFs is unlikely as transcript levels and transcription rates return to basal levels during the recovery (watered) phases. To assess the role of three master TFs regulated by ABA and known to induce *RD29B* and *RAB18* during dehydration stress, we analysed dehydration stress responses in a triple mutant deficient in the *AREB1*, *AREB2* and *ABF3* genes¹⁸. Despite the strong reduction in transcription in *RD29B* and *RAB18* transcript levels, the transcriptional memory response was still preserved in triple mutant plants (Fig. 7c,d). Thereby, although *AREB1*, *AREB2* and *ABF3* are critical for the robust response of the *RD29B* and *RAB18* trained genes during both the initial and repetitive stress responses, they are not essential for the transcriptional memory.

Finally, we emphasize that the results in the *atx1* background provide independent evidence that the trainable behaviour is not directly linked to the level of transcriptional activity, as trainable genes in *atx1* mutants produce reduced transcript levels and display lower levels of retained H3K4me3, while maintaining trainability (Fig. 8).

Thereby, the key TFs, in addition to ABA and the chromatin modifier ATX1, are not the factors causing the transcriptional memory response, although they strongly affect the magnitude of the transcriptional responses. Trainability is displayed by a distinct gene subset within the larger dehydration stress–response gene set and is associated with high levels of Ser5P Pol II and H3K4me3 during the recovery periods. The different behaviour of the trainable and non-trainable genes allows plants to modify and adjust their responses during repeated episodes of dehydration stress. Further studies would be needed to elucidate the signalling pathways and mechanisms that cause the dehydration-induced transcriptional memory in plants.

Methods

Plant growth and treatments. Wild-type and *atx1 Arabidopsis* plants were grown in potting soil in growth rooms at 22 °C with a 12-h light photoperiod and light intensity of 180 μmol m⁻² s⁻¹. Dehydration stress was initiated by removing the plants from soil, washing any remaining soil from their roots, and then air-drying for 2 h. This exposure resulted in a decrease in fresh tissue mass to ~65% of untreated (control) leaves, and is annotated as Stress1 (S1). In preliminary experiments, we have established that on reaching ~65% of RWC, whole plants (or detached tissues) could recover fully on rehydration, indicating that this level of water loss is reversible and non-lethal. Additional advantage of this method is that developmental differences between the plants experiencing different numbers of stress cycles are minimized. Recovery (R1) was achieved by placing plants in humid chambers for 22 h with their roots in a few drops of water. For a subsequent stress treatment, R1 plants were gently blotted onto filter paper to remove water and air-dried for 2 h (S2) followed by a recovery (R2). The same procedures were repeated for S3, R3 and S4. Prolonged watering of 3, 5 or 7 days was accomplished by replanting the plants in soil until harvest or a terminal dehydration stress.

To exclude the possibility that potential wounding of roots during extraction from soil affected the transcriptional response of the tested genes, we analysed plants removed from soil but kept in humid chambers throughout the stress–recovery cycle as controls. In the absence of dehydration stress, these control plants did not induce the dehydration stress–response genes (Supplementary Fig. S2). Further, when these control plants were exposed to their first dehydration stress, they responded with induced transcript levels typical for a S1 response (Supplementary Fig. S2). We concluded that potentially occurring root breakage does not have an observable effect on the dehydration stress responses we are measuring.

Water loss and RWC measurements. Relative water content was measured to follow the plant's water status. This value was calculated using the formula: RWC (%) = [(FW – DW)/(TW – DW)] × 100% (ref. 44), where FW is fresh weight, DW is dry weight and TW is turgid weight. Leaves were detached from plants and immediately weighed to determine their FW. The same leaves were submerged in deionized water for 24 h, blotted dry and weighed to determine their TW. Measurements of DW were taken after the leaves were oven-dried (65 °C) in brown paper bags for 24 h.

Nuclear run on assays. Nuclei were isolated from the tissue ground in a mortar and pestle, following the method of Meng and Lemaux⁴⁵, except that fresh, non-frozen tissues were used and the nuclei were purified on a 2 M sucrose pad⁴⁶. Nuclear run on assays were performed with 0.25 mM biotin-16-UTP and 0.75 mM of ATP, CTP and GTP in the transcription reaction. The Trizol (Invitrogen, Carlsbad, CA, USA) isolated RNA was treated with DNase I, and then purified on an RNeasy column (Qiagen, Valencia, CA, USA), and bound to streptavidin magnetic beads (Invitrogen). RNA on the washed beads was reversed transcribed with random primers, the complementary DNA eluted at 98 °C, and specific cDNA abundances were measured by quantitative PCR. The relative transcription rate of specific genes was quantitated with the 2^{-ΔΔC_t} calculation, according to the manufacturer's software (Bio-Rad, Hercules, CA, USA), where ΔΔC_t is the difference in the threshold cycles and the reference to *UBQ10*. Primers used are shown in Supplementary Table S1.

Chromatin immunoprecipitation assay. The ChIP assay was performed according to the described method^{20,47}. The specific antibodies (1:150 dilution) used for Ser2P Pol II (ab5095, Abcam, Cambridge, MA, USA, Lot: 703307); Ser5P Pol II (ab5131, Abcam, Lot: 806890); trimethyl-H3K4 (ab1012, Abcam, Lot: GR561731-1) or H3 (ab1791, Abcam, Lot: 517990) were added for an overnight incubation at 4 °C. The antibody–protein complexes were isolated by binding to protein A or protein G beads. The washed beads were heated at 65 °C for 8 h with proteinase K to reverse the formaldehyde crosslinking and digest proteins. The sample was then extracted with phenol/chloroform and the DNA precipitated in ethanol and resuspended in water. Purified DNA was analysed by real-time PCR with gene-specific primers shown in Supplementary Table S1.

Reverse transcription and real-time PCR. Total RNA isolation and reverse transcription with oligo(dT) (18418-012, Invitrogen) were performed as described previously²⁰. The amounts of individual genes were measured with gene-specific primers by real-time PCR analysis with a CyclerIQ real-time PCR Instrument (Bio-Rad) and SYBR Green mixture (Bio-Rad). The relative expression or amount of specific genes was quantitated with the 2^{-ΔΔC_t} calculation⁴⁸, according to the manufacturer's software (Bio-Rad), where ΔΔC_t is the difference in the threshold cycles and the reference housekeeping gene, which was ubiquitin for expression analyses or relative to input DNA for ChIP assays. The specific primers used are shown in Supplementary Table S1.

References

1. Arnholdt-Schmitt, B. Stress-induced cell reprogramming. A role for global genome regulation? *Plant Physiol.* **136**, 2579–2586 (2004).
2. Conrath, U. *et al.* Priming: getting ready for battle. *Mol. Plant Microbe Interact.* **19**, 1062–1071 (2006).
3. Ton, J. *et al.* Priming by airborne signals boosts direct and indirect resistance in maize. *Plant J.* **49**, 16–26 (2007).
4. Bruce, T. J., Matthes, M. C., Napier, J. A. & Pickett, J. A. Stressful 'memories' of plants: evidence for possible mechanisms. *Plant Sci.* **173**, 603–608 (2007).
5. Baldwin, I. T. & Schmelz, E. Immunological 'memory' in the induced accumulation of nicotine in wild tobacco. *Ecology* **77**, 236–246 (1996).
6. Conrath, U. Molecular aspects of defence priming. *Trends Plant Sci.* **16**, 524–531 (2011).
7. Jaskiewicz, M., Conrath, U. & Peterhansel, C. Chromatin modification acts as a memory for systemic acquired resistance in the plant stress response. *EMBO Rep.* **12**, 50–55 (2011).
8. Molinier, J., Ries, G., Zipfel, C. & Hohn, B. Transgenerational memory of stress in plants. *Nature* **442**, 1046–1049 (2006).
9. Hong, E. J., West, A. E. & Greenberg, M. E. Transcriptional control of cognitive development. *Curr. Opin. Neurobiol.* **15**, 21–28 (2005).
10. Bird, A. Perceptions of epigenetics. *Nature* **447**, 396–398 (2007).

11. Kozlowski, T. & Pallardy, S. Acclimation and adaptive responses of woody plants to environmental stresses **68**, 270–334 (2002).
12. Baker, S. S., Wilhelm, K. S. & Thomashow, M. F. The 5′-region of *Arabidopsis thaliana* cor15a has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol. Biol.* **24**, 701–713 (1994).
13. Lang, V. & Palva, E. T. The expression of a rab-related gene, rab18, is induced by abscisic acid during the cold acclimation process of *Arabidopsis thaliana* (L.) Heynh. *Plant Mol. Biol.* **20**, 951–962 (1992).
14. Yamaguchi-Shinozaki, K. & Shinozaki, K. A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* **6**, 251–264 (1994).
15. Cutler, S. R., Rodriguez, P. L., Finkelstein, R. R. & Abrams, S. R. Abscisic acid: emergence of a core signaling network. *Annu. Rev. Plant Biol.* **61**, 651–679 (2010).
16. Nechaev, S. & Adelman, K. Pol II waiting in the starting gates: regulating the transition from transcription initiation into productive elongation. *Biochim. Biophys. Acta* **1809**, 34–45 (2011).
17. Levine, M. Paused RNA polymerase II as a developmental checkpoint. *Cell* **145**, 502–511 (2011).
18. Yoshida, T. *et al.* AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation. *Plant J.* **61**, 672–685 (2010).
19. Ding, Y., Avramova, Z. & Fromm, M. The *Arabidopsis* trithorax-like factor ATX1 functions in dehydration stress responses via ABA-dependent and ABA-independent pathways. *Plant J.* **66**, 735–744 (2011).
20. Ding, Y., Avramova, Z. & Fromm, M. Two distinct roles of ARABIDOPSIS HOMOLOG OF TRITHORAX1 (ATX1) at promoters and within transcribed regions of ATX1-regulated genes. *Plant Cell* **23**, 350–363 (2011).
21. Ding, Y. *et al.* The *Arabidopsis* chromatin modifier ATX1, the myotubularin-like AtMTM and the response to drought. *Plant Signal Behav.* **4**, 1049–1058 (2009).
22. Bahrn, A., Jensen, C. R., Asch, F. & Mogensen, V. O. Drought-induced changes in xylem pH, ionic composition, and ABA concentration act as early signals in field-grown maize (*Zea mays* L.). *J. Exp. Bot.* **53**, 251–263 (2002).
23. Jongdee, B., Fukao, S. & Cooper, M. Leaf water potential and osmotic adjustment as physiological traits to improve drought tolerance in rice. *Field Crop Res.* **76**, 153–163 (2002).
24. Liu, G.-L. *et al.* Panicle water potential, a physiological trait to identify drought tolerance in rice. *J. Integr. Plant Biol.* **49**, 1464–1469 (2007).
25. Sung, S. & Amasino, R. M. Vernalization and epigenetics: how plants remember winter. *Curr. Opin. Plant Biol.* **7**, 4–10 (2004).
26. Sheldon, C. C. *et al.* Resetting of FLOWERING LOCUS C expression after epigenetic repression by vernalization. *Proc. Natl Acad. Sci. USA* **105**, 2214–2219 (2008).
27. Swiezewski, S. *et al.* Small RNA-mediated chromatin silencing directed to the 3′ region of the *Arabidopsis* gene encoding the developmental regulator, FLC. *Proc. Natl Acad. Sci. USA* **104**, 3633–3638 (2007).
28. Distelfeld, A., Li, C. & Dubcovsky, J. Regulation of flowering in temperate cereals. *Curr. Opin. Plant Biol.* **12**, 178–184 (2009).
29. Greenup, A., Peacock, W. J., Dennis, E. S. & Trevaskis, B. The molecular biology of seasonal flowering-responses in *Arabidopsis* and the cereals. *Ann. Bot.* **103**, 1165–1172 (2009).
30. Tsuji, H., Saika, H., Tsutsumi, N., Hirai, A. & Nakazono, M. Dynamic and reversible changes in histone H3-Lys4 methylation and H3 acetylation occurring at submergence-inducible genes in rice. *Plant Cell Physiol* **47**, 995–1003 (2006).
31. Sokol, A., Kwiatkowska, A., Jerzmanowski, A. & Prymakowska-Bosak, M. Up-regulation of stress-inducible genes in tobacco and *Arabidopsis* cells in response to abiotic stresses and ABA treatment correlates with dynamic changes in histone H3 and H4 modifications. *Planta* **227**, 245–254 (2007).
32. Kim, J. M. *et al.* Alterations of lysine modifications on the histone H3 N-tail under drought stress conditions in *Arabidopsis thaliana*. *Plant Cell Physiol.* **49**, 1580–1588 (2008).
33. Alvarez-Venegas, R., Abdallat, A. A., Guo, M., Alfano, J. R. & Avramova, Z. Epigenetic control of a transcription factor at the cross section of two antagonistic pathways. *Epigenetics* **2**, 106–113 (2007).
34. Jung, H. W., Tschaplinski, T. J., Wang, L., Glazebrook, J. & Greenberg, J. T. Priming in systemic plant immunity. *Science* **324**, 89–91 (2009).
35. van den Burg, H. A. & Takken, F. L. Does chromatin remodeling mark systemic acquired resistance? *Trends Plant Sci.* **14**, 286–294 (2009).
36. Ng, H. H., Robert, F., Young, R. A. & Struhl, K. Targeted recruitment of Set1 histone methylase by elongating Pol II provides a localized mark and memory of recent transcriptional activity. *Mol. Cell* **11**, 709–719 (2003).
37. Kundu, S. & Peterson, C. L. Dominant role for signal transduction in the transcriptional memory of yeast GAL genes. *Mol. Cell Biol.* **30**, 2330–2340 (2010).
38. Light, W. H., Brickner, D. G., Brand, V. R. & Brickner, J. H. Interaction of a DNA zip code with the nuclear pore complex promotes H2A.Z incorporation and INO1 transcriptional memory. *Mol. Cell* **40**, 112–125 (2010).
39. Tan-Wong, S. M., Wijayatilake, H. D. & Proudfoot, N. J. Gene loops function to maintain transcriptional memory through interaction with the nuclear pore complex. *Genes Dev.* **23**, 2610–2624 (2009).
40. Nechaev, S. & Adelman, K. Promoter-proximal Pol II: when stalling speeds things up. *Cell Cycle* **7**, 1539–1544 (2008).
41. Boettiger, A. N. & Levine, M. Synchronous and stochastic patterns of gene activation in the *Drosophila* embryo. *Science* **325**, 471–473 (2009).
42. Fujita, Y., Fujita, M., Shinozaki, K. & Yamaguchi-Shinozaki, K. ABA-mediated transcriptional regulation in response to osmotic stress in plants. *J. Plant Res.* **124**, 509–525 (2011).
43. van Dijk, K. *et al.* Dynamic changes in genome-wide histone H3 lysine 4 methylation patterns in response to dehydration stress in *Arabidopsis thaliana*. *BMC Plant Biol.* **10**, 238 (2010).
44. Hewlett, J. & Kramer, P. The measurement of water deficits in broadleaf plants. *Protoplasma* **57**, 381–391 (1962).
45. Meng, L. & Lemaux, P. G. A simple and rapid method for nuclear run-on transcription assays in plants. *Plant Mol. Biol. Rep.* **21**, 65–71 (2003).
46. Kanazawa, A., O’Dell, M., Hellens, R. P., Hitchin, E. & Metzloff, M. Mini-scale method for nuclear run-on transcription assay in plants. *Plant Mol. Biol. Rep.* **18**, 377–383 (2000).
47. Ding, Y. *et al.* SDG714, a histone H3K9 methyltransferase, is involved in Tos17 DNA methylation and transposition in rice. *Plant Cell* **19**, 9–22 (2007).
48. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).

Acknowledgements

We are grateful to Dr. Yamaguchi-Shinozaki for kindly providing the *areb1/areb2/abf3* triple mutant seeds and to Dr. Karin van Dijk for critically reading the manuscript. Partial support was provided by NSF award MCB-1121898 to Z.A. and M.F.

Author contributions

Y.D., M.F. and Z.A. conceived the study and designed experiments; Y.D. performed the experiments with input from M.F. All authors took part in the interpretation of results and preparation of the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Ding, Y. *et al.* Multiple exposures to drought ‘train’ transcriptional responses in *Arabidopsis*. *Nat. Commun.* **3**:740 doi: 10.1038/ncomms1732 (2012).