

1 Functional screening and *in vitro* analysis reveals thioesterases with enhanced substrate
2 specificity profiles that improve short-chain fatty acid production in *Escherichia coli*

3

4 Matthew D. McMahon^a and Kristala L.J. Prather^{a,b,#}

5

6 ^aDepartment of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA,
7 USA; ^bSynthetic Biology Engineering Research Center (SynBERC), Massachusetts Institute of
8 Technology, Cambridge, MA, USA.

9

10 Running Head: Thioesterases for short-chain fatty acid biosynthesis

11

12 #Address Correspondance to Kristala L.J. Prather, kljp@mit.edu

13 **Abstract**

14 Short-chain fatty acid (SCFAs) biosynthesis is pertinent to production of biofuels, industrial
15 compounds, and pharmaceuticals from renewable resources. To expand on *Escherichia coli*
16 SCFA products, we previously implemented a coenzyme A (CoA)-dependent pathway that
17 condenses acetyl-CoA to a diverse group of short-chain fatty acyl-CoAs. To increase product
18 titers and reduce premature pathway termination products, we describe *in vivo* and *in vitro*
19 analyses to understand and improve the specificity of the acyl-CoA thioesterase enzyme, which
20 releases fatty acids from CoA. A total of 62 putative bacterial thioesterases, including 23 from
21 the cow rumen microbiome, were inserted into a pathway that condenses acetyl-CoA to an acyl-
22 CoA molecule derived from exogenously provided propionic or isobutyric acid. Functional
23 screening revealed thioesterases that increase production of saturated (valerate), unsaturated
24 (trans-2-pentenoate) and branched (4-methylvalerate) SCFAs compared to overexpression of *E.*
25 *coli* thioesterase *tesB* or native expression of endogenous thioesterases. To determine if altered
26 thioesterase acyl-CoA substrate specificity caused the increase in product titers, six of the most
27 promising enzymes were analyzed *in vitro*. Biochemical assays revealed that the most productive
28 thioesterases rely on promiscuous activity, but have greater specificity for product-associated
29 acyl-CoAs than for precursor acyl-CoAs. Here we introduce novel thioesterases with improved
30 specificity for saturated, branched and unsaturated short-chain acyl-CoAs, thereby expanding the
31 diversity of potential fatty acid products while increasing titers of current products. The growing
32 uncertainty associated with protein database annotations denotes this study as a model for
33 isolating functional biochemical pathway enzymes in situations where experimental evidence of
34 enzyme function is absent.

35

36 **1. Introduction**

37 The potential for producing biofuels, industrial compounds, and pharmaceuticals from
38 renewable resources has led to an increased interest in short-chain (C2 – C7) fatty acid (SCFA)
39 biosynthesis. Developing recombinant strains to produce these molecules could lead to
40 production of polymers such as polyhydroxyalkanoates and pharmaceuticals such as statins from
41 fossil fuel alternatives (1, 2). Supporting the development of microbially-synthesized products is
42 the reduced cost and increased prevalence of genomic sequencing. The resulting profusion of
43 metabolic diversity provides a wealth of potential enzymes with known genetic sequences for
44 improving biosynthetic pathways. For example, recent genomic sequencing has unveiled the
45 metabolic diversity of important members from the cow rumen microbiome, an environment rich
46 in SCFAs (3, 4). These genome sequences provide an opportunity to find enzymes that improve
47 production and specificity in SCFA biosynthesis pathways.

48 One such pathway that would benefit from improved enzyme specificity is Coenzyme A
49 (CoA)-dependent biosynthesis of SCFAs. In the CoA-dependent pathway, a thiolase enzyme
50 condenses an acyl-CoA molecule with acetyl-CoA (Figure 1). The resulting 3-ketoacyl-CoA
51 molecule is then sequentially reduced by reductase, crotonase, and enoyl-reductase enzymes
52 before the 3-hydroxy, unsaturated, and saturated fatty acids, respectively, are cleaved from CoA
53 using a thioesterase enzyme. Previous work from our lab has used this pathway with acetyl-CoA
54 and glycolyl, propionyl, or isobutyryl-CoA as the condensed substrates to produce a variety of
55 SCFAs using *E. coli* TesB including 3-hydroxyvalerate (3-hydroxypentanoate) (1),
56 dihydroxybutyrate (5), 3-hydroxy-4-methylvalerate (5), and a variety of alcohols (6). One benefit
57 of CoA-dependent biosynthesis of SCFAs is the diversity of potential products (7); however,
58 with this diversity comes a need for selective enzymes that increase final product titers by

59 minimizing substrate flux to undesired byproducts. The selectivity of the final enzyme in the
60 pathway, the thioesterase, is of particular importance because it influences the product profile by
61 catalyzing fatty acid release from CoA at each step of the pathway (Figure 1) and is important
62 for secretion of fatty acid products (8). While many acyl-ACP thioesterases have been
63 investigated for improved fatty acid production, acyl-CoA thioesterases are not as well explored
64 (8-10). Despite its preference for acyl-CoAs in the C14-C18 range (11), the *E. coli* acyl-CoA
65 thioesterase TesB produces diverse SCFAs (12). However, locating more selective thioesterases
66 may reduce byproduct formation and increase final product titers.

67 Selecting individual thioesterase enzymes for functional screening against short-chain
68 fatty acyl-CoAs is challenging because much of their vast phylogenetic and functional diversity
69 is poorly understood (13). While many thioesterases have been explored for long-chain fatty acid
70 production (14-17), few studies have focused on those that prefer short-chain acyl-CoAs. Several
71 broad specificity acyl-CoA thioesterases including *E. coli* TesB and *Saccharomyces cerevisiae*
72 Pte1p can be used for SCFA production, but lack the specificity necessary for optimizing
73 biosynthetic pathways (11, 18). One approach to selecting thioesterases for functional screening
74 that improves the likelihood of finding enzymes with the desired specificity is to investigate
75 those proteins with similarity to commonly used and effective enzymes. However, the absence of
76 known selective short-chain acyl-CoA thioesterases restricts this approach. Sampling candidates
77 more broadly will increase opportunities for finding enzymes with new substrate specificities,
78 but will also increase the number of thioesterases with undesired activities. Some combination of
79 these routes can be used to screen sufficient phylogenetic breadth while also increasing the
80 sample size of TesB-like thioesterases to reveal enzymes with greater specificity for short-chain
81 acyl-CoAs.

82 To address this need, we functionally screened 62 putative thioesterases in a pathway for
83 production of the SCFAs 3-hydroxyvalerate, trans-2-pentenoate, valerate, and 4-methylvalerate
84 in *E. coli*. Six enzymes were homologous to variants active on short-chain acyl-CoAs, while the
85 remaining 56 represent all of the annotated thioesterases from five bacterial strains of interest.
86 Based upon *in vivo* fatty acid titers of those recombinant strains, six enzymes were chosen for *in*
87 *vitro* analysis on a broad range of acyl-CoAs to determine their substrate preferences. The
88 combination of *in vivo* and *in vitro* data indicate that we have uncovered thioesterases with
89 greater specificity for and production of unsaturated, saturated, and branched SCFAs in *E. coli*
90 relative to TesB and endogenous thioesterase activity.

91

92 **2. Methods**

93 2.1. Bacterial Strains

94 Rumen isolate *Prevotella ruminicola* 23 was obtained from Dr. Roderick Mackie of the
95 University of Illinois, Urbana-Champaign (USA). Genomic DNA from rumen isolates
96 *Fibrobacter succinogenes* S85 and *Ruminococcus albus* 7 were obtained from Dr. Paul Weimer
97 of the US Dairy Forage Research Center, USDA-Agricultural Research Service, Madison (USA).
98 *Alcanivorax borkumensis* SK2 (ATCC# 700651) and genomic DNA from *Pseudomonas*
99 *aeruginosa* PAO1 (ATCC# 47085) were purchased from the ATCC. Genomic DNA of
100 *Rhodopseudomonas palustris* CGA009 was obtained from Dr. Caroline Harwood of the
101 University of Washington, Seattle (USA). *Pseudomonas syringae* pv. *maculicola* ES4326 was
102 obtained from Dr. Fred Ausubel at Massachusetts General Hospital (Boston, MA, USA). *E. coli*
103 MG1655 (DE3) $\Delta endA \Delta recA$ was previously constructed in our lab (1).

104

105 2.2. Plasmid and Strain Construction

106 Plasmid pET/*ter/bktB/pct* was constructed previously (6) and plasmid pCDF/*phaB/phaJ4*
107 was constructed by subcloning *phaB* from pET/*bktB/phaB* (5) into multiple cloning site I by
108 NdeI/XhoI restriction digest and *phaJ4* into multiple cloning site II of pCDFDuet-1
109 (EMDmillipore) using primers listed in Table S1. Plasmid pET/*bktB/pct* was constructed from
110 pET/*ter/bktB/pct* by BamHI/NotI restriction digest followed by treatment with Mung Bean
111 Nuclease (New England Biolabs) and blunt ligation of the 8 kb fragment.

112 Genomic DNA was isolated from bacterial strains using the Wizard Genomic DNA
113 Purification Kit (Promega). Custom oligonucleotides were purchased for the PCR amplification
114 of all individual thioesterases and CoA ligases from purified genomic DNA (Integrated DNA
115 Technologies). Primers used for amplification are listed in Table S1. Following amplification,
116 individual genes were inserted into the expression vector pACYCDuet-1 (EMDmillipore) using
117 polymerase incomplete primer extension (PIPE)-based cloning (19). The genes encoding *R.*
118 *palustris* CoA ligases FcsA (Rpa4267) and VcsA (Rpa3299) (20) and the genes encoding the six
119 thioesterases chosen for further analysis were inserted into vector pTEV5 for protein purification
120 using PIPE-based methods (21). The pTEV5 construct produced an enzyme with an N-terminal
121 hexahistidine tag removable by TEV protease. Due to solubility problems in pTEV5, the genes
122 encoding thioesterases Pr1687 and Fs2108 were amplified from genomic DNA and cloned using
123 restriction enzymes NdeI and EcoRI into pTYB22 (New England Biolabs) which produced an
124 enzyme with an N-terminal chitin-binding domain removable by intein self-cleavage. Plasmid
125 sequences were confirmed using PCR amplification and DNA sequencing (GENEWIZ).

126 Gene deletions of *yciA*, *yigI*, and *tesB* in *E. coli* MG1655 (DE3) $\Delta endA \Delta recA$ were made
127 using P1 transduction with strains JW1245-1, JW5588-1, and JW0442-1, respectively, from the

128 Keio collection as donor cells (22). The kanamycin resistance gene was removed using FLP-
129 mediated recombination as previously described (23).

130

131 2.3. Culture Conditions

132 Recombinant strains of *E. coli* MG1655 (DE3) $\Delta endA \Delta recA$ were grown at 30°C in
133 Luria-Bertani (LB) medium overnight in a shaking incubator at 250 rpm. 50 μ L of the overnight
134 culture was used to inoculate a 50 mL LB culture supplemented with 10 g/L glucose containing
135 100 mg/L ampicillin, 50 mg/L streptomycin, and, when pACYCDuet-1 was present, 16 mg/L
136 chloramphenicol. Cultures were grown at 30°C until an optical density at 600 nm (OD_{600}) of 0.8
137 was reached, at which point isopropyl- β -D-1-thiogalactopyranoside (IPTG) was added to a final
138 concentration of 1 mM with either propionate or isobutyrate to a final concentration of 15 mM.
139 Cultures were incubated at 30°C for 48 h post-induction prior to fatty acid titer determination.

140

141 2.4. Fatty Acid Analysis

142 Culture samples were centrifuged to pellet cells and culture supernatant was removed for
143 HPLC analysis. A 5 μ l sample of culture supernatant was injected into an Agilent 1100 series
144 instrument equipped with refractive index detection (RID). Samples were processed through an
145 Aminex[®] HPX-87H anion-exchange column (Bio-Rad Laboratories) with isocratic flow of 5 mM
146 H₂SO₄ mobile phase at a rate of 0.6 mL/min and column and detector temperatures set to 35°C.
147 Concentrations of valerate, 3-hydroxyvalerate, trans-2-pentenoate, and 4-methylvalerate were
148 determined using linear regression of external standards.

149

150 2.5. Protein Purification

151 Acyl-CoA ligases FcsA and VcsA and thioesterases EcTesB, PpTesB, EcYdiI and Pr655
152 were overproduced using pTEV5 constructs in *E. coli* BL21Star (DE3) (Invitrogen). One liter of
153 cells were grown at 30°C in LB medium containing 100 mg/L ampicillin until an OD₆₀₀ of 0.5
154 was reached, at which point IPTG was added to the cultures at a concentration of 100 mg/L. Post
155 induction, the cells were grown for 15 h at 30°C, then harvested by centrifugation and
156 resuspended in 2.5x v/w Buffer A (50 mM Tris-HCl pH 8.0, 300 mM NaCl, 10% v/v glycerol).
157 Protein purification followed previously described protocols for nickel chelate chromatography
158 followed by cleavage with TEV protease (24). Proteins were flash frozen in liquid nitrogen and
159 the concentration was determined using the Bradford assay with bovine serum albumin as a
160 standard (25) (Bio-Rad).

161 Pr1687 and Fs2108 protein was purified using the Intein-Mediated Purification with an
162 Affinity Chitin-binding Tag (IMPACT) expression vector pTYB22 (New England Biolabs).
163 Cells were grown and induced using the same conditions as described for pTEV5 constructs and
164 proteins were purified with chitin-affinity chromatography followed by intein cleavage mediated
165 by dithiothreitol (DTT) using previously described conditions (26). Proteins were stored and
166 quantified as described for nickel chelate chromatography purified proteins.

167

168 2.6. Enzymatic Synthesis of Acyl-CoAs

169 Acyl-CoA synthesis reactions were carried out in 50 mM HEPES pH 7.5, 1 mM DTT, 5
170 mM MgCl₂, 5 mM ATP, and 2 mM CoA. Fatty acid substrate was added at 7.5 mM for butyrate,
171 3-hydroxyvalerate, trans-2-pentenoate, valerate, 4-methylvalerate, hexanoate, octanoate,
172 decanoate, and dodecanoate while 2 mM fatty acid substrate was used for tetradecanoate. To
173 increase solubility, 1% and 3% w/v triton-X 100 was added to reactions with dodecanoate and

174 tetradecanoate respectively. CoA ligase enzymes were added at 500 nM for all reactions. VcsA
175 was added to butyrate, 3-hydroxyvalerate, valerate, 4-methylvalerate, and hexanoate reactions,
176 while FcsA was added to octanoate, decanoate, dodecanoate, and tetradecanoate reactions.
177 Reactions were run overnight at room temperature for all substrates except trans-2-pentenoate,
178 which was run at 30°C for 6 h because these conditions reduced the appearance of degradation
179 products. Acyl-CoA ligases were precipitated from reactions at 95°C for 5 min, and then
180 removed by centrifugation.

181 Acyl-CoA products were purified from substrates using an Agilent 1200 series HPLC
182 with diode array detection (DAD). A 100 µL reaction volume was injected onto an Agilent
183 Eclipse XDB-C18 column and separation achieved using a mobile phase of 50 mM ammonium
184 acetate, 0.1% m/v acetic acid (Solvent A) - 50 mM ammonium acetate, 0.1% m/v acetic acid,
185 70% v/v acetonitrile (Solvent B) gradient. The method began at 100% Solvent A from 0-5 min,
186 followed by a 0-100% gradient of solvent B from 5-50 minutes, followed by an isocratic step of
187 100% solvent B from 50-55 min. The gradient was run at a flow rate of 1 mL/min and CoA was
188 monitored by measuring absorbance at 258 nM. Fractions containing peaks corresponding to
189 acyl-CoAs were collected, flash frozen in liquid nitrogen, and lyophilized. Dried acyl-CoAs were
190 then resuspended in water and the concentration was determined by the absorbance at 258 nM
191 using the molar extinction coefficient of CoA ($14,328 \text{ M}^{-1} \text{ cm}^{-1}$) within the linear range of
192 detection (27).

193

194 2.7. Thioesterase Activity Assays

195 Thioesterase activity was measured using 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB)
196 which reacts with free CoA after it is released through thioesterase-mediated bond cleavage.

197 Reactions were carried out in 1 mM DTNB, 100 mM HEPES pH 8.0, 20 μ M acyl-CoA and were
198 run in the linear range of each thioesterase. Substrate concentrations matched those used
199 previously for EcTesB analysis (11). Reaction progress was monitored through the change in
200 absorbance at 412 nm using the molar extinction coefficient of 5-thio-2-nitrobenzoate (14,150
201 $M^{-1} \text{ cm}^{-1}$), which is formed when DTNB reacts with free CoA (28).

202

203 **3. Results**

204 3.1 *In Vivo* Functional Screening for Thioesterases that Alter Fatty Acid Production

205 Acyl-CoA thioesterase substrate specificity plays an important role in deciding the SCFA
206 profile produced using a CoA-based pathway. For example, if the target product is a saturated
207 fatty acid, a thioesterase with broad substrate specificity will release fatty acids at each step in
208 the biosynthetic pathway, depleting substrate pools and reducing the final product titer (Figure
209 1). To find thioesterases that increase specificity for and production of 3-hydroxy, unsaturated,
210 saturated and branched fatty acids in the pathway, 62 putative thioesterases were chosen for
211 screening. A group of six thioesterases, three TesB and three TesB2 enzymes, were chosen that
212 had 38-50% amino acid identity to *E. coli* MG1655 TesB (EcTesB), which makes 3-hydroxy and
213 saturated fatty acids *in vivo* (1, 6, 29), and 40-41% amino acid identity to *A. borkumensis* TesB2
214 enzyme, which was previously described as having specificity for 3-hydroxy acyl-CoAs (30)
215 (Table 1).

216 To incorporate greater phylogenetic and functional diversity of thioesterases, the
217 remaining 56 proteins encompassed all of the annotated thioesterases from five bacterial strains
218 (Table 1). *E. coli* MG1655 (Accession: NC_000913.2) thioesterases were chosen with the dual
219 purpose of identifying those that increase product titers of desired SCFAs for use in heterologous

220 pathways and those that reduce product titers and should be targeted for deletion from our host
221 strain. The second source of thioesterases, *Pseudomonas putida* KT2440 (Accession:
222 NC_002947.3), was chosen for its phylogenetic similarity to *E. coli* and because it is a known
223 producer of polyhydroxyalkanoates, which may indicate the presence of thioesterases with
224 specificity to 3-hydroxy acyl-CoAs (31). The remaining three organisms, *Prevotella ruminicola*
225 23 (Accession: NC_014033.1), *Fibrobacter succinogenes* S85 (Accession: NC_017448.1), and
226 *Ruminococcus albus* 7 (Accession: NC_014833.1), were chosen because they are prevalent in the
227 cow rumen microbiome and contribute to the high concentrations of SCFAs found there (3, 4,
228 32).

229 Each of the 62 putative thioesterases was individually overexpressed in *E. coli* containing
230 all the necessary genes for CoA-dependent biosynthesis of valerate (Figure 1). Previous work
231 from our lab identified *Treponema denticola* Ter, *Megaphaera elsdenii* Pct, and *Cupriavidus*
232 *necator* (formerly *R. eutropha* H16) BktB, PhaB, and PhaJ4 as suitable upstream pathway
233 enzymes for valerate production (6, unpublished data). Cells were grown in LB containing
234 glucose until log phase growth, then pathway genes were induced and cultures were
235 supplemented with either propionate or isobutyrate for straight or branched SCFA biosynthesis,
236 respectively (Figure 1). LC-based analysis of the culture supernatant was used to screen for
237 enzymes that altered the fatty acid product profiles. Specifically, the resulting chromatograms
238 were examined for increases and decreases of 3-hydroxyvalerate, trans-2-pentenoate, valerate,
239 and 4-methylvalerate. Profiles produced by recombinant strains were compared to controls with
240 either no recombinant thioesterase or with overproduced EcTesB, a thioesterase used previously
241 in our group (1, 5, 6).

242 Overproduction of greater than 20% of the thioesterases functionally screened in this
243 study resulted in observable changes in substrate, intermediate, and product titers relative to the
244 control strains (Table S2). Four thioesterases, EcYciA, Pp1466, Pp3807, and Pp4975, were
245 associated with significant reduction in valerate and 4-methylvalerate production combined with
246 increased acetate and propionate titers (Table S2). Two thioesterases, Pr1510 and Fs368, were
247 associated with a production phenotype involving reduced titers of measured substrates,
248 intermediates, and final products combined with increased glucose consumption (Table S2). The
249 most logical explanation for this phenotype is channeling of substrates to long-chain fatty acid
250 biosynthesis. Because long-chain fatty acids could not be quantified with our HPLC system,
251 additional experiments outside the scope of this work are required to determine the activity
252 profile of these enzymes.

253

254 3.2 Host Strain Development and Identification of Thioesterases for *In Vitro* Analysis

255 From the 12 annotated *E. coli* thioesterases screened during this study, three were chosen
256 for deletion from the host strain. The *tesB* gene was deleted because its overexpression resulted
257 in increased titers of 3-hydroxyvalerate, which could be a final product, but also acts as a shunt
258 product in the formation of trans-2-pentenoate or valerate; *yciA* was deleted because its
259 overexpression increased final titers of the precursor-derived and exogenously supplied acids
260 acetate and propionate, which correlates with previous work (33), while decreasing final product
261 titers of valerate and 4-methylvalerate; and *yigI* was deleted because its overexpression resulted
262 in decreased production of 4-methylvalerate and increased final acetate and isobutyrate titers.
263 The resulting triple thioesterase deletion strain *E. coli* MG1655 (DE3) $\Delta endA \Delta recA \Delta tesB$
264 $\Delta yciA \Delta yigI$ was used for further *in vivo* analysis of several active thioesterases.

265 From the full set of 62 functionally screened thioesterases, EcTesB, PpTesB, EcYdiI,
266 Fs2108, Pr655, and Pr1687 were chosen for more detailed *in vivo* and *in vitro* analysis because
267 their overproduction resulted in increased titers of 3-hydroxyvalerate, trans-2-pentenoate,
268 valerate, or 4-methylvalerate. Propionate and isobutyrate feeding experiments were performed in
269 triplicate with these six thioesterases in both *E. coli* MG1655 (DE3) $\Delta endA \Delta recA$ and the triple
270 thioesterase deletion strain containing pET/*ter/bktB/pct* and pCDF/*phaB/phaJ4* in an effort to
271 distinguish recombinant thioesterase activity from background and potentially increase final
272 product titers (Figure 2a, b). The greatest 3-hydroxyvalerate titers were produced in recombinant
273 strains overproducing TesB enzymes from *E. coli* and *P. putida* when compared with the other
274 60 putative thioesterases (Figure 2). The maximum titer (2.163 g/L) resulted from
275 overproduction of EcTesB (Table 2), and is similar to previously published titers from our lab
276 (1). 3-hydroxyvalerate titers were 20% lower in the triple deletion strain overproducing EcTesB
277 than in *E. coli* with native thioesterases.

278 Observable titers of unsaturated SCFAs were uncommon among the recombinant strains
279 with only Pr655 overproduction resulting in detectable trans-2-pentenoate during initial
280 screening (Figure 2a, b). To determine whether strains containing the other five thioesterases
281 selected for further analysis could produce detectable unsaturated SCFAs *in vivo*, each was
282 introduced into a strain lacking the downstream enoyl-reductase gene *ter*, which does not
283 produce the favored substrate valeryl-CoA (Figure 2c, d). In these recombinant strains, product
284 titers of trans-2-pentenoate were greatest for EcYdiI, Pr655, and Pr1687, with EcYdiI
285 overproduction resulting in the greatest trans-2-pentenoate titer, 695 mg/L (Table 2). To our
286 knowledge, this is the largest published titer of an unsaturated SCFA from an engineered
287 pathway in *E. coli*.

288 Three thioesterases, Fs2108, Pr1687, and EcYdiI, were chosen for further analysis
289 because their overproduction increased titers of the saturated acids valerate and 4-methylvalerate
290 while decreasing shunt product titers (Figure 2a, b). Overproduction of thioesterase Fs2108
291 achieves the greatest final titers for both the straight SCFA valerate and the branched SCFA 4-
292 methylvalerate (Table 2) with 3-hydroxy and unsaturated fatty acid intermediate titers below the
293 limit of detection. While overproduction of Fs2108 resulted in a modest 15% improvement in
294 valerate titer over the EcTesB control, an improvement of greater than 200% was observed for 4-
295 methylvalerate titer over the same control. This 200% improvement was possible in part because
296 4-methylvalerate titers were significantly lower than valerate titers for all 62 thioesterases tested,
297 leaving more room for improvement. Overproduction of Fs2108 in the deletion strain resulted in
298 approximately 10% more valerate and 30% more 4-methylvalerate than in the host containing all
299 native thioesterases, which likely results from reduced hydrolysis of 3-hydroxyacyl-CoA and
300 trans-2-acyl-CoA precursors. No 4-methylvalerate production was observed from the triple
301 deletion strain when isobutyrate was supplied without thioesterase overexpression, which
302 indicates that the remaining native thioesterases have poor activity on 4-methylvaleryl-CoA
303 (Figure 2a, b).

304

305 3.3 Determination of *In Vitro* SCFA Substrate Specificity for Active Thioesterases

306 To determine the substrate preferences for the selected thioesterases, all six were
307 overexpressed in *E. coli* BL21Star (DE3) and purified. Overexpression of *Pr1687* and *Fs2108*
308 required the use of a vector that inserted an N-terminal chitin binding domain to obtain soluble
309 protein. EcTesB, PpTesB, EcYdiI, and Pr655 were soluble with an N-terminal hexahistidine tag.
310 Thioesterases were then tested for *in vitro* acyl-CoA hydrolysis activity on 20 μ M acetyl-CoA,

311 butyryl-CoA, valeryl-CoA, hexanoyl-CoA, octanoyl-CoA, decanoyl-CoA, dodecanoyl-CoA and
312 tetradecanoyl-CoA using an Ellman's reagent-based assay described previously for determining
313 specific activity (11) (Figure 3a). EcTesB, Fs2108 and Pr1687 showed a preference for longer-
314 chain acyl-CoAs. PpTesB also showed a preference for longer acyl-CoAs, but the relationship
315 was less linear than for EcTesB, Fs2108 and Pr1687. Pr655 showed a strong preference for
316 butyryl and valeryl-CoAs and activity on acyl-CoAs longer than hexanoyl-CoA was below the
317 limit of detection for this assay. EcYdiI had low activity for all substrates tested. A recent
318 publication showing that EcYdiI has strong activity on the aromatic compound 1,4-dihydroxy-2-
319 naphthoyl-CoA provides justification for the weak activity of this enzyme on the substrates
320 provided in this study (34).

321 Specific activities were also measured for all six thioesterases on 3-hydroxy, unsaturated,
322 saturated, and branched valeryl-CoAs (Figure 3b). For 3-hydroxyvaleryl-CoA, EcTesB and
323 PpTesB had 6-fold greater specific activity than the next best thioesterase. Both EcYdiI and
324 Pr655 showed low activity for 3-hydroxyvaleryl-CoA, while specific activity of Fs2108 and
325 Pr1687 for the same substrate was below the limit of detection. No thioesterase had specific
326 activity greater than $2 \mu\text{M CoA min}^{-1} \text{ mg protein}^{-1}$ on trans-2-pentenoyl-CoA. Pr655, EcTesB,
327 PpTesB, and EcYdiI displayed similar specific activities for this substrate; however, the small
328 differences in specific activity translated into significant deviations in trans-2-pentenoate
329 product titers when the thioesterases were overproduced in recombinant strains lacking the
330 downstream enzyme Ter (Figure 3b). Fs2108 and Pr1687 specific activity on trans-2-pentenoyl-
331 CoA was below the limit of detection, but *in vivo* product profiles of strains lacking *ter* reflect a
332 slight preference for 3-hydroxyvaleryl-CoA over trans-2-pentenoyl-CoA (Figure 2d).

333 All six enzymes selected for further analysis had detectable activity on valeryl-CoA;
334 however, EcTesB, PpTesB, Pr655 and Fs2108 had much greater activity relative to EcYdiI and
335 Pr1687 (Figure 3b). Activity against the 4-methylvaleryl-CoA (branched) substrate was greater
336 than or equal to activity on straight valeryl-CoA for EcTesB, PpTesB, Fs2108, and Pr1687.
337 EcYdiI and Pr655 both showed weak activity against 4-methylvaleryl-CoA. *In vitro* analysis of
338 the enzymes associated with the greatest titers of valerate revealed that Fs2108 had greater
339 specific activity for both valeryl-CoA and 4-methylvaleryl-CoA than either Pr1687 or EcYdiI
340 (Figure 3b). Further, Fs2108 specific activity for both 3-hydroxyvaleryl-CoA and trans-2-
341 pentenyl-CoA precursors was below the limit of detection for *in vitro* assays. The greater
342 specific activity of Fs2108 on 4-methylvaleryl-CoA and valeryl-CoA combined with reduced
343 specific activity on precursor acyl-CoAs likely accounts for the increased 4-methylvalerate and
344 valerate titers found for Fs2108.

345

346 **4. Discussion**

347 In this study, 62 putative thioesterases were screened for increased product titers of 3-
348 hydroxy, unsaturated, saturated and branched products of interest. From these, six were chosen
349 for more thorough *in vivo* and *in vitro* analysis. Relating the specific activity profiles of EcTesB,
350 PpTesB, EcYdiI, Fs2108, Pr655 and Pr1687 with their *in vivo* product profiles of 3-
351 hydroxyvalerate, trans-2-pentenoate, valerate, or 4-methylvalerate leads to several important
352 conclusions. First, the enzymes with the greatest *in vivo* product titers always had lower specific
353 activity for the associated acyl-CoA than for alternative CoA substrates. For example, Fs2108
354 has a six-fold higher specific activity for tetradecanoyl-CoA than for valeryl-CoA indicating that
355 valeryl-CoA is not the enzyme's preferred substrate. The disparity between the specific activities

356 of Fs2108 for these two substrates indicates that our pathway relies on the promiscuous
357 thioesterase activity of Fs2108. This is particularly evident for two thioesterases, EcYdiI and
358 Pr1697, that were associated with increased valerate production despite having much higher
359 specific activity for long-chain acyl-CoAs in the case of Pr1687, or aromatic acyl-CoAs for
360 EcYdiI (34). The fact that all six thioesterases that were investigated *in vitro* prefer alternate
361 substrates over those provided in our pathways suggests that both protein engineering and future
362 bioprospecting efforts could further improve on the short-chain fatty acyl-CoA thioesterases
363 discovered here.

364 Comparing the *in vivo* product titers with *in vitro* substrate specificities of the six chosen
365 thioesterases also suggests that after a specific activity level of 1-5 $\mu\text{M CoA min}^{-1} \text{ mg protein}^{-1}$ is
366 reached for a given acyl-CoA, the enzyme's activity for the pathway precursor acyl-CoAs
367 becomes an important factor influencing final product titer, illustrating the importance of
368 selecting pathway thioesterases with reduced activity on precursor acyl-CoAs. For example,
369 strains overproducing thioesterases Pr1687 and EcYdiI produce more valerate than those
370 overproducing EcTesB despite EcTesB having 30-fold greater specific activity for valeryl-CoA
371 than Pr1687 and EcYdiI (Figure 2b). Accounting for this difference is the observation that
372 EcTesB also has strong specific activity on the precursor 3-hydroxyvaleryl-CoA that translates
373 into 3-hydroxyvalerate production while Pr1687 does not have detectable activity on 3-
374 hydroxyvaleryl-CoA. Further supporting the importance of reduced specific activity on pathway
375 precursors is the observation that recombinant strains overproducing Pr655 produce much less
376 valerate than strains overproducing EcYdiI and Pr1687 (Figure 2a, b) even though Pr655
377 maintains greater specific activity for valeryl-CoA than thioesterases EcYdiI and Pr1687 (Figure
378 3b). The increased specific activity of Pr655 on 3-hydroxyvaleryl-CoA and trans-2-pentenoyl-

379 CoA precursors relative to EcYdiI and Pr1687 suggests that the reduced valerate titer results
380 from increased precursor acyl-CoAs hydrolysis.

381 Confounding our conclusion that thioesterases with low specific activity on precursor
382 acyl-CoAs have improved final product titers is the observation that recombinant strains
383 overexpressing *ydiI* produce high valerate titers with trans-2-pentenoate titers below the limit of
384 detection despite EcYdiI exhibiting higher *in vitro* specific activity for 20 μ M trans-2-pentenoyl-
385 CoA than for 20 μ M valeryl-CoA (Figures 2a, 3b). In our recombinant strains, EcYdiI and the
386 other screened thioesterases are competing with the downstream pathway enzyme Ter for the
387 substrate trans-2-pentenoyl-CoA. One plausible explanation for the absence of detectable trans-
388 2-pentenoate from this recombinant strain is that EcYdiI has a K_m , a measure of substrate
389 affinity, for both trans-2-pentenoyl-CoA and valeryl-CoA greater than the 20 μ M concentration
390 used in our *in vitro* assay, which is reasonable to assume because EcYdiI has evolved for
391 specificity towards aromatic acids (34). If Ter has a lower K_m for trans-2-pentenoyl-CoA than
392 EcYdiI, then it could reduce the intracellular concentration of trans-2-pentenoyl-CoA below 20
393 μ M, causing EcYdiI activity for this compound to become physiologically irrelevant. In this
394 situation, the intracellular valeryl-CoA concentration could increase to a point where low-level
395 specific thioesterase activity would result in significant valerate production. Pr655, on the other
396 hand, could have a lower K_m for both trans-2-pentenoyl-CoA and valeryl-CoA, which would
397 compete more effectively for the trans-2-pentenoyl-CoA with downstream enzyme Ter and allow
398 continued low level production of trans-2-pentenoate (Figure 2a). Determination of kinetic
399 parameters for these non-cognate substrates is limited by the low concentrations of acyl-CoAs
400 obtained from enzymatic synthesis and LC-based purification.

401 The thioesterase substrate preferences found by our *in vitro* experiments indicate the
402 breadth of thioesterase functional diversity screened in this study and underline the importance
403 of sampling in the selection of pathway enzymes (Figure 3a and b). The TesB enzymes of *E. coli*
404 and *P. putida* represent the most phylogenetically (50% amino acid identity) and functionally
405 (Figure 3a and b) similar enzymes, while the remaining four thioesterases appear both
406 phylogenetically and functionally disparate. The TesB thioesterases showed both similar *in vivo*
407 product profiles and *in vitro* specific activity profiles, which suggests that future screens aimed at
408 acquiring more diverse TesB function should sample enzymes with lower similarity at the amino
409 acid level than the *E. coli* and *P. putida* homologs. The diversity of the remaining 60 thioesterase
410 phenotypes is an indication of the functional diversity inherent to bacterial thioesterases and
411 suggests that many opportunities remain for isolating enzymes with improved specificity over
412 those currently used in CoA-dependent biosynthetic pathways.

413 Our study also highlights the challenges associated with enzyme selection for metabolic
414 pathways. A common route to selecting pathway enzymes is to rely on proposed function of
415 known enzymes in databases and the literature or choose enzymes homologous to those with
416 proposed functions. Unfortunately, enzyme annotation in public databases has degraded as
417 functional analysis has not kept up with the rate of sequence deposition (35). The absence of
418 credible studies on short-chain acyl-CoA thioesterases prompted us to implement a broad
419 functional screen of diverse candidates. Our results demonstrate the power of a well-designed
420 screen for isolating uncharacterized or poorly characterized enzymes that improve product titers.
421 Our methods for screening and *in vitro* characterization of substrate specificity serve as a
422 template for investigating other poorly characterized enzyme functions for pathway
423 development. The *E. coli* thioesterase EcYdiI serves as an example of an enzyme whose

424 documented function may eliminate it from contention (34), but once included in our functional
425 screen, maintains the appropriate levels of promiscuous activity in intracellular conditions to
426 provide the highest specificity for unsaturated SCFAs discovered so far. It is unlikely that
427 selecting enzymes based solely upon annotated function would have identified this level of
428 activity because short-chain acyl-coA thioesterase activity remains poorly characterized.

429 During this study we isolated acyl-CoA thioesterases from diverse bacterial sources that
430 increase production of saturated, unsaturated, and branched SCFAs through improvements in
431 acyl-CoA substrate specificity. By comparing *in vivo* product profiles with *in vitro* specific
432 activities of thioesterases that produced the greatest 3-hydroxyvalerate, trans-2-pentenoate,
433 valerate, and 4-methylvalerate titers, we discovered that the most productive thioesterases found
434 during functional screening: 1) use promiscuous activity to produce the SCFAs monitored in this
435 study and 2) maintain low specific activity for pathway precursors relative to the preferred acyl-
436 CoA. These findings indicate this study as a model for isolating enzymes for biochemical
437 pathway functions that are poorly characterized in the literature. Further, the thioesterases we
438 identified provide opportunities for increasing titers of desirable products as well as developing
439 new pathways for the production of unsaturated SCFAs. We have also provided important *in*
440 *vivo* and *in vitro* data on the production phenotypes and substrate specificities of poorly
441 characterized acyl-CoA thioesterases for short-chain acyl-CoAs that are valuable for future
442 bioprospecting and engineering studies.

443

444 **Acknowledgements**

445 We would like to thank the U.S. Department of Agriculture Agricultural Research Service for
446 providing genomic DNA from cow rumen strains. This work was supported by the Institute for

447 Collaborative Biotechnologies through grant W911NF-09-0001 from the U.S. Army Research
448 Office. The content of the information does not necessarily reflect the position or the policy of
449 the Government, and no official endorsement should be inferred.

450 **References**

- 451 1. **Tseng HC, Harwell CL, Martin CH, Prather KL.** 2010. Biosynthesis of chiral 3-
452 hydroxyvalerate from single propionate-unrelated carbon sources in metabolically
453 engineered *E. coli*. *Microb Cell Fact* **9**:96.
- 454 2. **Lee SH, Park OJ.** 2009. Uses and production of chiral 3-hydroxy-gamma-
455 butyrolactones and structurally related chemicals. *Appl Microbiol Biotechnol* **84**:817-
456 828.
- 457 3. **Suen G, Weimer PJ, Stevenson DM, Aylward FO, Boyum J, Deneke J, Drinkwater
458 C, Ivanova NN, Mikhailova N, Chertkov O, Goodwin LA, Currie CR, Mead D,
459 Brumm PJ.** 2011. The complete genome sequence of *Fibrobacter succinogenes* S85
460 reveals a cellulolytic and metabolic specialist. *PLoS One* **6**:e18814.
- 461 4. **Purushe J, Fouts DE, Morrison M, White BA, Mackie RI, Coutinho PM, Henrissat
462 B, Nelson KE.** 2010. Comparative genome analysis of *Prevotella ruminicola* and
463 *Prevotella bryantii*: insights into their environmental niche. *Microb Ecol* **60**:721-729.
- 464 5. **Martin CH, Dhamankar H, Tseng HC, Sheppard MJ, Reisch CR, Prather KL.** 2013.
465 A platform pathway for production of 3-hydroxyacids provides a biosynthetic route to 3-
466 hydroxy-gamma-butyrolactone. *Nat Commun* **4**:1414.
- 467 6. **Tseng HC, Prather KL.** 2012. Controlled biosynthesis of odd-chain fuels and chemicals
468 via engineered modular metabolic pathways. *Proc Natl Acad Sci U S A* **109**:17925-
469 17930.
- 470 7. **Torella JP, Ford TJ, Kim SN, Chen AM, Way JC, Silver PA.** 2013. Tailored fatty
471 acid synthesis via dynamic control of fatty acid elongation. *Proc Natl Acad Sci U S A*.
472 **110**:11290-11295.

- 473 8. **Handke P, Lynch SA, Gill RT.** 2011. Application and engineering of fatty acid
474 biosynthesis in *Escherichia coli* for advanced fuels and chemicals. *Metab Eng* **13**:28-37.
- 475 9. **Jing F, Cantu DC, Tvaruzkova J, Chipman JP, Nikolau BJ, Yandeau-Nelson MD,**
476 **Reilly PJ.** 2011. Phylogenetic and experimental characterization of an acyl-ACP
477 thioesterase family reveals significant diversity in enzymatic specificity and activity.
478 *BMC Biochem* **12**:44.
- 479 10. **Lu X, Vora H, Khosla C.** 2008. Overproduction of free fatty acids in *E. coli*:
480 implications for biodiesel production. *Metab Eng* **10**:333-339.
- 481 11. **Nie L, Ren Y, Janakiraman A, Smith S, Schulz H.** 2008. A novel paradigm of fatty
482 acid beta-oxidation exemplified by the thioesterase-dependent partial degradation of
483 conjugated linoleic acid that fully supports growth of *Escherichia coli*. *Biochemistry*
484 **47**:9618-9626.
- 485 12. **Naggert J, Narasimhan ML, DeVeaux L, Cho H, Randhawa ZI, Cronan JE, Jr.,**
486 **Green BN, Smith S.** 1991. Cloning, sequencing, and characterization of *Escherichia coli*
487 thioesterase II. *J Biol Chem* **266**:11044-11050.
- 488 13. **Cantu DC, Chen Y, Reilly PJ.** 2010. Thioesterases: a new perspective based on their
489 primary and tertiary structures. *Protein Sci* **19**:1281-1295.
- 490 14. **Voelker TA, Davies HM.** 1994. Alteration of the specificity and regulation of fatty acid
491 synthesis of *Escherichia coli* by expression of a plant medium-chain acyl-acyl carrier
492 protein thioesterase. *J Bacteriol* **176**:7320-7327.
- 493 15. **Liu A, Tan X, Yao L, Lu X.** 2013. Fatty alcohol production in engineered *E. coli*
494 expressing *Marinobacter* fatty acyl-CoA reductases. *Appl Microbiol Biotechnol*.

- 495 16. **Liu T, Vora H, Khosla C.** 2010. Quantitative analysis and engineering of fatty acid
496 biosynthesis in *E. coli*. *Metab Eng* **12**:378-386.
- 497 17. **Gronenberg LS, Marcheschi RJ, Liao JC.** 2013. Next generation biofuel engineering
498 in prokaryotes. *Curr Opin Chem Biol* **17**:462-471.
- 499 18. **Maeda I, Delessert S, Hasegawa S, Seto Y, Zuber S, Poirier Y.** 2006. The peroxisomal
500 Acyl-CoA thioesterase Pte1p from *Saccharomyces cerevisiae* is required for efficient
501 degradation of short straight chain and branched chain fatty acids. *J Biol Chem*
502 **281**:11729-11735.
- 503 19. **Klock HE, Koesema EJ, Knuth MW, Lesley SA.** 2008. Combining the polymerase
504 incomplete primer extension method for cloning and mutagenesis with microscreening to
505 accelerate structural genomics efforts. *Proteins* **71**:982-994.
- 506 20. **Crosby HA, Pelletier DA, Hurst GB, Escalante-Semerena JC.** 2012. System-wide
507 studies of N-lysine acetylation in *Rhodospseudomonas palustris* reveal substrate
508 specificity of protein acetyltransferases. *J Biol Chem* **287**:15590-15601.
- 509 21. **Rocco CJ, Dennison KL, Klenchin VA, Rayment I, Escalante-Semerena JC.** 2008.
510 Construction and use of new cloning vectors for the rapid isolation of recombinant
511 proteins from *Escherichia coli*. *Plasmid* **59**:231-237.
- 512 22. **Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita**
513 **M, Wanner BL, Mori H.** 2006. Construction of *Escherichia coli* K-12 in-frame, single-
514 gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**. doi:10.1038/msb4100050
515 (2006).
- 516 23. **Datsenko KA, Wanner BL.** 2000. One-step inactivation of chromosomal genes in
517 *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**:6640-6645.

- 518 24. **McMahon MD, Rush JS, Thomas MG.** 2012. Analyses of MbtB, MbtE, and MbtF
519 suggest revisions to the mycobactin biosynthesis pathway in *Mycobacterium*
520 *tuberculosis*. J Bacteriol **194**:2809-2818.
- 521 25. **Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram
522 quantities of protein utilizing the principle of protein-dye binding. Anal Biochem **72**:248-
523 254.
- 524 26. **Felnagle EA, Barkei JJ, Park H, Podevels AM, McMahon MD, Drott DW, Thomas**
525 **MG.** 2010. MbtH-like proteins as integral components of bacterial nonribosomal peptide
526 synthetases. Biochemistry **49**:8815-8817.
- 527 27. **Giljanovic J, Prkic A.** 2010. Determination of coenzyme A (CoASH) in the presence of
528 different thiols by using flow-injection with a UV/Vis spectrophotometric detector and
529 potentiometric determination of CoASH using an iodide ISE. Molecules **15**:100-113.
- 530 28. **Riener CK, Kada G, Gruber HJ.** 2002. Quick measurement of protein sulfhydryls with
531 Ellman's reagent and with 4,4'-dithiodipyridine. Anal Bioanal Chem **373**:266-276.
- 532 29. **Zheng Z, Gong Q, Liu T, Deng Y, Chen JC, Chen GQ.** 2004. Thioesterase II of
533 *Escherichia coli* plays an important role in 3-hydroxydecanoic acid production. Appl
534 Environ Microbiol **70**:3807-3813.
- 535 30. **Sabirova JS, Ferrer M, Lunsdorf H, Wray V, Kalscheuer R, Steinbuchel A, Timmis**
536 **KN, Golyshin PN.** 2006. Mutation in a "tesB-like" hydroxyacyl-coenzyme A-specific
537 thioesterase gene causes hyperproduction of extracellular polyhydroxyalkanoates by
538 *Alcanivorax borkumensis* SK2. J Bacteriol **188**:8452-8459.

- 539 31. **Huijberts GN, Eggink G, de Waard P, Huisman GW, Witholt B.** 1992. *Pseudomonas*
540 *putida* KT2442 cultivated on glucose accumulates poly(3-hydroxyalkanoates) consisting
541 of saturated and unsaturated monomers. *Appl Environ Microbiol* **58**:536-544.
- 542 32. **Bergman EN.** 1990. Energy contributions of volatile fatty acids from the gastrointestinal
543 tract in various species. *Physiol Rev* **70**:567-590.
- 544 33. **Clomburg JM, Vick JE, Blankschien MD, Rodriguez-Moya M, Gonzalez R.** 2012. A
545 synthetic biology approach to engineer a functional reversal of the beta-oxidation cycle.
546 *ACS Synth Biol* **1**:541-554.
- 547 34. **Chen M, Ma X, Chen X, Jiang M, Song H, Guo Z.** 2013. Identification of a Hotdog
548 Fold Thioesterase Involved in the Biosynthesis of Menaquinone in *Escherichia coli*. *J*
549 *Bacteriol* **195**:2768-2775.
- 550 35. **Schnoes AM, Brown SD, Dodevski I, Babbitt PC.** 2009. Annotation error in public
551 databases: misannotation of molecular function in enzyme superfamilies. *PLoS Comput*
552 *Biol* **5**:e1000605.

553

554 **Figures.**

555

556 Figure 1. Biochemical pathway and LC-based *in vivo* screen for thioesterases with improved
557 substrate specificity. The pathway in *E. coli* for the biosynthesis of SCFAs contains enzymes Pct
558 (*Megaphaera elsdenii*), BktB (*Cupriavidus necator*), PhaB (*Cupriavidus necator*), PhaJ4
559 (*Cupriavidus necator*), and Ter (*Treponema denticola*) with varied thioesterases. The dashed
560 bond indicates the additional carbon incorporated into the fatty acids with feeding of isobutyrate
561 in place of propionate.

562
563 Figure 2. *In vivo* fatty acid production. Titers of fatty acids produced from (a,c) *E. coli* MG1655
564 (DE3) $\Delta endA \Delta recA$ and (b,d) *E. coli* MG1655 (DE3) $\Delta endA \Delta recA \Delta tesB \Delta yciA \Delta yigI$
565 expressing CoA-based pathways for the biosynthesis of (a,b) valerate or 4-methylvalerate, (c,d)
566 trans-2-pentenoate, with six different thioesterases. The product titers shown were obtained from
567 three separate shake flasks for each thioesterase with error bars representing standard deviations.
568 Each strain contained plasmids pCDF/*phaB/phaJ4* with either (a,b) pET/*ter/bktB/pct* or (c,d)
569 pET/*bktB/pct*. Each strain also contained plasmid pACYCDuet-1 with the thioesterases displayed
570 on the X-axis. Absence of product bars for a thioesterase indicates that titers were below the
571 limit of detection for the HPLC RID detector.


572 *4-methylvalerate titers were determined using three separate cultures for each thioesterase fed
573 15 mM isobutyrate in the place of propionate.

574
575 Figure 3. Acyl-CoA specificity of six active thioesterases. (a) *In vitro* specific activity of six
576 thioesterases on C2-C14 acyl-CoAs. Specific activities represent the average from three
577 enzymatic reactions with error bars representing standard deviations. They were determined in
578 the linear range for each thioesterase with acyl-CoA concentrations of 20 μ M for C2, acetyl-
579 CoA; C4, butyryl-CoA; C5, valeryl-CoA; C6, hexanoyl-CoA; C8, octanoyl-CoA; C10, decanoyl-
580 CoA; C12, dodecanoyl-CoA; C14, tetradecanoyl-CoA. (b) *In vitro* specific activity of six
581 thioesterases on saturated, valeryl-CoA; branched, 4-methylvaleryl-CoA; 3-hydroxy, 3-
582 hydroxyvaleryl-CoA; unsaturated, trans-2-pentenoyl-CoA. Inset is enlarged activities of EcYdiI
583 and Pr655 for comparison of low level activities. Specific activities were determined using the


584 same conditions described in (a). The absence of a bar for a given substrate/thioesterase specific
585 activity indicates that the activity was below the limit of detection at absorbance of 412 nm.

586




Figure 1.




a.



b.



a.



b.




Table 1. Thioesterases screened for activity in a biosynthetic pathway for the production of short-chain fatty acids

TesB & TesB2 enzymes	<i>E. coli</i> MG1655	<i>P. putida</i> KT2440	<i>P. ruminicola</i> 23	<i>R. albus</i> 7	<i>F. succinogenes</i> S85
<i>E. coli</i> TesB ^a	EcTesA	Pp244	Pr655	Ral865	Fsu266
<i>P. putida</i> TesB	EcPaaI	Pp254	Pr1075	Ral874	Fsu270
<i>R. opacus</i> TesB	EcEntH	Pp262	Pr1498	Ral880	Fsu368
<i>P. syringae</i> TesB2	EcFadM	Pp301	Pr1510	Ral1843	Fsu803
<i>A. borkumensis</i> TesB2 ^b	EcYbfF	Pp580	Pr1668	Ral1929	Fsu944
<i>P. aeruginosa</i> PAOI TesB2	EcYbgC	Pp1218	Pr1687	Ral1938	Fsu1747
	EcYbhC	Pp1466	Pr2385	Ral2059	Fsu2108
	EcYciA	Pp1980		Ral2801	
	EcYiiD	Pp2050		Ral3109	
	EcYigl	Pp2308			
	EcYdil	Pp2318			
		Pp3281			
		Pp3807			
		Pp4105			
		Pp4180			
		Pp4181			
	Pp4975				
	Pp5198				
	Pp5331				
	Pp5356				
	PpPhaJ1				

^a(11)

^b(30)

Table 2. Maximum fatty acid titers observed for each product of the CoA-dependent pathway

Strains	Maximum Titters (g/L)			
	3-Hydroxyvalerate ^a	Trans-2-pentenoate ^b	Valerate ^c	4-Methylvalerate ^c
MG1655 $\Delta endA$ $\Delta recA$	2.163±0.078	0.695±0.002	1.425±0.011	0.428±0.019
MG1655 $\Delta endA$ $\Delta recA$ $\Delta tesB$ $\Delta yciA$ $\Delta yigI$	1.940±0.099	0.609±0.015	1.540±0.034	0.570±0.041

^aThioesterase EcTesB

^bThioesterase EcYdiI

^cThioesterase Fs2108