

# Bosonic anomalies, induced fractional quantum numbers, and degenerate zero modes: The anomalous edge physics of symmetry-protected topological states

Juven C. Wang,<sup>1,2,\*</sup> Luiz H. Santos,<sup>2,†</sup> and Xiao-Gang Wen<sup>1,2,3,‡</sup>

<sup>1</sup>*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>2</sup>*Perimeter Institute for Theoretical Physics, Waterloo, ON, Canada N2L 2Y5*

<sup>3</sup>*Institute for Advanced Study, Tsinghua University, Beijing, 100084, P. R. China*

(Received 21 July 2014; revised manuscript received 7 April 2015; published 22 May 2015)

The boundary of symmetry-protected topological states (SPTs) can harbor new quantum anomaly phenomena. In this work, we characterize the bosonic anomalies introduced by the 1+1D non-onsite-symmetric gapless edge modes of (2+1)D bulk bosonic SPTs with a generic finite Abelian group symmetry (isomorphic to  $G = \prod_i Z_{N_i} = Z_{N_1} \times Z_{N_2} \times Z_{N_3} \times \dots$ ). We demonstrate that some classes of SPTs (termed “Type II”) trap fractional quantum numbers (such as fractional  $Z_N$  charges) at the 0D kink of the symmetry-breaking domain walls, while some classes of SPTs (termed “Type III”) have degenerate zero energy modes (carrying the projective representation protected by the unbroken part of the symmetry), either near the 0D kink of a symmetry-breaking domain wall, or on a symmetry-preserving 1D system dimensionally reduced from a thin 2D tube with a monodromy defect 1D line embedded. More generally, the energy spectrum and conformal dimensions of gapless edge modes under an external gauge flux insertion (or twisted by a branch cut, i.e., a monodromy defect line) through the 1D ring can distinguish many SPT classes. We provide a manifest correspondence from the physical phenomena, the induced fractional quantum number, and the zero energy mode degeneracy to the mathematical concept of cocycles that appears in the group cohomology classification of SPTs, thus achieving a concrete physical materialization of the cocycles. The aforementioned edge properties are formulated in terms of a long wavelength continuum field theory involving scalar chiral bosons, as well as in terms of matrix product operators and discrete quantum lattice models. Our lattice approach yields a regularization with anomalous non-onsite symmetry for the field theory description. We also formulate some bosonic anomalies in terms of the Goldstone-Wilczek formula.

DOI: [10.1103/PhysRevB.91.195134](https://doi.org/10.1103/PhysRevB.91.195134)

PACS number(s): 71.27.+a, 11.10.-z, 03.65.Aa, 73.43.-f

## I. INTRODUCTION

Symmetry dictates the conservation law and the corresponding conserved current on classical actions in classical physics, such as by Noether’s theorem [1]. However, as it is now well known, there is a potential obstruction of some classical symmetry to be promoted to a consistent symmetry in the quantum level. This is the paradigm of “quantum anomalies” [2].

Quantum anomalies occur in our real-world physics, such as pion decaying to two photons via Adler-Bell-Jackiw chiral anomaly [3–5]. Anomalies also constrain beautifully on the Standard Model of particle physics, in particular to the Glashow-Weinberg-Salam theory, via anomaly cancellations of gauge and gravitational couplings. The above two familiar examples of anomalies concern chiral fermions and continuous symmetry [e.g. U(1), SU(2), SU(3)]. Out of curiosity, we ask the following: “Are there concrete examples of quantum anomalies for bosons instead? And anomalies for discrete symmetries? Are they potentially testable experimentally in the laboratory in the near future?”

In this work, we address the question affirmatively and demonstrate that “bosonic anomalies for discrete symmetries” can be expected on the boundary of some interacting bosonic symmetry-protected topological states (SPTs) in condensed

matter systems [6,7]. [Such interacting bosonic SPTs may be realized in the future by applying the ultracold bosonic gas controlled by optical lattice [8] (see a recent proposal and reference therein [9]).] Our work thus will address some of the interplays between “symmetry,” “quantum anomaly,” and “topology.”

There has been rapid progress on exploring the entangled quantum states with gapless edge modes protected by some global symmetry. The classic example is the one dimensional (1D, one dimensional space and one dimensional time, or 1+1D) Haldane spin-1 chain with SO(3) spin rotational symmetry [10,11]. Another renown example is topological insulators, which are protected by fermion number conservation U(1) symmetry and time-reversal symmetry  $Z_2^T$  [12–17]. A topological insulator may be realized in a noninteracting free fermion system, while there are so-called bosonic SPTs, which can only happen in an interacting bosonic system.

In attempting to understand various phases of interacting bosonic systems, it is important to try to characterize them in terms of unique physical properties. The goal of this paper is to address this question for bosonic SPTs in 2D. Let us motivate our question in the simplest scenario of the 1D SPTs given by the spin-1 Haldane chain. The Hamiltonian conserves spin rotation and time-reversal symmetries and the ground state is formed by singlets in the bulk. Bulk excitations are formed by breaking singlets, a process that requires an energy gap. Its nontrivial property resides on the edges, both of which contain an effective spin- $\frac{1}{2}$  transforming projectively under rotation or time-reversal symmetry. Since the edge spin is

\*juven@mit.edu

†lsantos@perimeterinstitute.ca

‡xwen@perimeterinstitute.ca




FIG. 1. (Color online) The boundary of 2D SPT state harbors 1D gapless edge modes if the global symmetry is preserved (not broken spontaneously or explicitly). The global symmetry transformation  $S$  of 1D edge mode acts in a non-onsite manner, where  $S$  cannot be written as a tensor product form on each site (i.e., the symmetry operator  $S$  acts on more than a single site for its tensor operators, where we show schematically  $S$  acts on two neighbor sites.)

effectively “free,” it renders a twofold degeneracy (per edge) in the spectrum. Hence, here the mathematical concept of projective representations is directly connected to spectral zero energy mode degeneracy.

In this work, we will show that edge modes of bosonic SPTs in 2D can also provide physical signatures of the bulk state. We will study the 2D bosonic SPTs with 1D edge modes on the boundary (see Fig. 1), protected by a global symmetry  $G$  of a generic cyclic group  $G = \prod_i Z_{N_i} = Z_{N_1} \times Z_{N_2} \times Z_{N_3} \times \dots$  (to which any finite Abelian group is isomorphic). Our basic result is that point defects on the 1D edge are associated to induced  $Z_N$  charge (referred as Type II bosonic anomaly in Sec. IV) or protected degeneracies (referred as Type III bosonic anomaly in Sec. V) for some classes of SPTs.

The edge modes of our focus have the property that they can only be gapped out if the symmetry is broken. In a description around a gapless 1+1D Luttinger-liquid-like fixed point, this means that putative interacting energy-gap-opening terms (sine-Gordon cosine terms) violate the symmetry and are therefore forbidden (which does not rule out the possibility that a gap may open by symmetry breaking). The suppression of all these gap-opening terms is a manifestation that counterpropagating modes carry different global charges, which, consequentially implies that backscattering processes violate the symmetry. Thus an important step in capturing the edge properties of SPTs is to construct the symmetry transformation that endows counterpropagating modes with this anomalous property. We will study this *anomalous non-onsite symmetry* explicitly.

Recently, several theoretical approaches have been developed to understand bosonic SPTs, such as using group cohomology [6,7,18], lattice models [19–23], matrix product states [20,22], field theory techniques [22–27], or projective construction [28–32]. One of the goals of this paper is to address the connections among miscellaneous approaches by working out a few specific examples. To this end, we specifically highlight three learned aspects about SPTs:

1. *Non-onsite symmetry on the edge.* An important feature of SPT is that the *global symmetry* acting on a local Hamiltonian of edge modes is realized *non-onsite* [20–22]. For a given symmetry group  $G$ , the non-onsite symmetry means that its symmetry transformation *cannot* be written as a tensor product form on each site [6,20]

$$U(g)_{\text{non-onsite}} \neq \otimes_i U_i(g), \quad (1)$$

for  $g \in G$  of the symmetry group. On the other hand, the onsite symmetry transformation  $U(g)$  can be written in a tensor product form acting on each site  $i$  [6,20], i.e.,  $U(g)_{\text{onsite}} = \otimes_i U_i(g)$ , for  $g \in G$ . [The symmetry transformation acts as an operator  $U(g)$  with  $g \in G$ , transforming the state  $|v\rangle$  globally by  $U(g)|v\rangle$ .] Therefore, to study the SPT edge mode, one should realize how the non-onsite symmetry acts on the boundary as in Fig. 1.

2. *Group cohomology construction.* It has been proposed that  $d + 1$  dimensional ( $d + 1$ D) SPTs of symmetry-group- $G$  interacting boson system can be constructed by the number of distinct cocycles in the  $(d + 1)$ th cohomology group  $\mathcal{H}^{d+1}(G, U(1))$ , with  $U(1)$  coefficient [6,33]. (See also the first use of cocycle in the high energy context by Jackiw in Refs. [34,35].) While another general framework of cobordism theory is subsequently proposed [36] to account for subtleties when symmetry  $G$  involves time reversal [25], in our work we will focus on a finite Abelian symmetry group  $G = \prod_i Z_{N_i}$ , where the group cohomology is a complete classification.

3. *Surface anomalies.* It has been proposed that the edge modes of SPTs are the source of gauge anomalies, while that of intrinsic topological orders are the source of gravitational anomalies [37]. SPT boundary states are known to show at least one of three properties: (1) symmetry-preserving gapless edge modes, (2) symmetry-breaking gapped edge modes, (3) symmetry-preserving gapped edge modes with surface topological order [25,38–41].

### Bosonic anomalies realized on the SPT edge

The three aspects 1, 2, 3 above had hinted at the bosonic anomalies harbored on the boundary of interacting bosonic SPTs. In this work, we focus on *characterizing the bosonic anomalies as precisely as possible*, and attempt to connect our *bosonic anomalies to the notion defined in the high energy physics context*. In short, we aim to *make connections between the meanings of boundary bosonic anomalies studied in both high energy physics and condensed matter theory*.

We will examine a generic finite Abelian  $G = \prod_i Z_{N_i}$  bosonic SPTs, and study *what is truly anomalous* about the edge under the case of 1 and 2 above. (Since it is forbidden to have any intrinsic topological order in a 1D edge, we do not have scenario 3.) We focus on addressing the properties of its 1+1D edge modes, their anomalous non-onsite symmetry and bosonic anomalies from three different perspectives, (i) quantum lattice models, (ii) matrix product states, and (iii) quantum field theory, while connecting them to cocycles of group cohomology.

We shall now define the meaning of quantum anomaly in a language appreciable by both high energy physics and




FIG. 2. (Color online) The intuitive way to view the bulk-boundary correspondence for edge modes of SPTs (or intrinsic topological order) under the flux insertion, or equivalently the monodromy defect/branch cut (blue dashed line) modifying the bulk and the edge Hamiltonians. SPTs located on a large sphere with two holes with flux-in and flux-out is analogous to a Laughlin-type flux insertion through a cylinder, inducing anomalous edge modes (red arrows) moving along the opposite directions on two edges.

condensed matter communities:

The quantum anomaly is *an obstruction of a symmetry of a theory to be fully regularized for a full quantum theory as an onsite symmetry on the UV-cut-off lattice in the same spacetime dimension.*

According to this definition, to characterize our bosonic anomalies, we will find several possible obstructions to regulate the symmetry at the quantum level:

*Obstruction of onsite symmetries.* Consistently we will find throughout our examples to fully regularize our SPTs 1D edge theory on the 1D lattice Hamiltonian requires the *non-onsite symmetry*, namely, *realizing the symmetry anomalously*. The non-onsite symmetry on the edge cannot be “dynamically gauged” on its own spacetime dimension [19–22,37], thus this also implies the following obstruction.

*Obstruction of the same spacetime dimension.* We will show that the physical observables for gapless edge modes (the case 1) are their energy spectral shifts [22] under symmetry-preserving external flux insertion through a compact 1D ring.

TABLE I. A summary of bosonic anomalies as 1D edge physical observables to detect the 2+1D SPT of  $G = Z_{N_1} \times Z_{N_2} \times Z_{N_3}$  symmetry, here we use  $p_i, p_{ij}, p_{ijk}$  to label the SPT class index in the third cohomology group  $\mathcal{H}^3(G, U(1))$ . For Type II class  $p_{12} \in \mathbb{Z}_{N_{12}}$ , we can use a unit of  $Z_{N_1}$ -symmetry-breaking domain wall to induce a  $\frac{p_{12}}{N_{12}}$  fractional  $Z_{N_2}$  charge (see Sec. IV). For Type III class  $p_{123} \in \mathbb{Z}_{N_{123}}$ , we can either use  $Z_{N_1}$ -symmetry-breaking domain wall or use  $Z_{N_1}$ -symmetry-preserving flux insertion (effectively a monodromy defect) through 1D ring to trap  $N_{123}$  multiple degenerate zero energy modes (see Sec. V). For Type I class  $p_1 \in \mathbb{Z}_{N_1}$ , our proposed physical observable is the energy spectrum [or conformal dimension  $\tilde{\Delta}(\mathcal{P})$  as a function of momentum  $\mathcal{P}$ , see Ref. [22]] shift under the flux insertion. This energy spectral shift also works for all other (Type II and III) classes (see Sec. VI). This table serves as topological invariants for Type I, II, and III bosonic SPTs in the context of Ref. [44].

Group cohomology 3-cocycle	$p$ in $\mathcal{H}^3(G, U(1))$	Bosonic anomalies and physical observables		
		induced fractional charge	degenerate zero energy modes	$\tilde{\Delta}(\mathcal{P})$ under flux/monodromy
Type I $p_1$ : Eq. (8)	$\mathbb{Z}_{N_1}$	No	No	Yes
Type II $p_{12}$ : Eq. (9)	$\mathbb{Z}_{N_{12}}$	Yes, $\frac{p_{12}}{N_{12}}$ of $Z_{N_2}$ charge	No	Yes
Type III $p_{123}$ : Eq. (10)	$\mathbb{Z}_{N_{123}}$	No	Yes, $N_{123}$ degeneracy	Yes

The energy spectral shift is caused by the Laughlin-type flux insertion of Fig. 2. The *flux insertion* can be equivalently regarded as an effective *branch cut* modifying the Hamiltonian (blue dashed line in Fig. 2) connecting from the edge to an extra dimensional bulk. Thus the spectral shifts also indicate the transportation of quantum numbers from one edge to the other edge. This can be regarded as the anomaly requiring an *extra dimensional bulk*.

*Nonperturbative effects.* We know that the familiar Adler-Bell-Jackiw anomaly of *chiral fermions* [3,4], observed in the pion decay in particle physics, can be captured by the perturbative one-loop Feynman diagram. However, importantly, the result is nonperturbative, being exact from low energy IR to high energy UV. This effect can be further confirmed via Fujikawa’s path integral method [42] nonperturbatively. Instead of the well-known *chiral fermionic anomalies*, do we have *bosonic anomalies* with these nonperturbative effects?

Indeed, yes, we will show two other kinds of bosonic anomalies with nonperturbative effects with symmetry-breaking gapped edges [the case (2)]: One kind of consequent anomalies for Type II SPTs under  $Z_{N_1}$  symmetry-breaking domain walls is the induced fractional  $Z_{N_2}$  charge trapped near 0D kink of gapped domain walls. Amazingly, through a fermionization/bosonization procedure, we can apply the field theoretic Goldstone-Wilczek method to capture the one-loop Feynman diagram effect *nonperturbatively*, as this fractional charge is known to be robust without higher-loop diagrammatic corrections [43]. We will term this a Type II bosonic anomaly.

The second kind of anomalies for symmetry-breaking gapped edge [the case (2)] is that the edge is gapped under  $Z_{N_1}$  symmetry-breaking domain walls, with consequent degenerate zero energy ground states due to the projective representation of other symmetries  $Z_{N_2} \times Z_{N_3}$ . The zero mode degeneracy is found to be  $\text{gcd}(N_1, N_2, N_3)$ -fold. We will term this a Type III bosonic anomaly.

The paper is organized as follows. In Sec. II, we start with some basic results in group cohomology and its  $n$ -cocycles. The readers who are not familiar with group cohomology may either take the chance to learn the basics or skip it and proceed to Sec. III. We set up Type I [22], II, and III SPT lattice constructions in Sec. III, their matrix product operators, and low energy field theory. Remarkably, the Type III non-onsite symmetry transformation is distinct from the Types I and



TABLE II. Given a generic finite Abelian global symmetry group (isomorphic to a cyclic group  $G = \prod_{i=1}^m \mathbb{Z}_{N_i} = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{N_3} \times \dots$ ), here we provide the data of group cohomology and their corresponding realization as symmetry-protected topological (SPT) states by using (i) quantum lattice models, (ii) matrix product operators (MPO), and (iii) quantum field theory approach. The classification labels  $p_1, p_{12}, p_{ijk}$  belong to the Type I  $\mathbb{Z}_N$  class, Type II  $\mathbb{Z}_{N_{12}}$  ( $\equiv \mathbb{Z}_{\text{gcd}(N_1, N_2)}$ ) class, Type III  $\mathbb{Z}_{N_{123}}$  ( $\equiv \mathbb{Z}_{\text{gcd}(N_1, N_2, N_3)}$ ) class (all labeled in blue color in the table), respectively.

3-cocycle	Min. symm. group $G$	$\mathcal{H}^3(G, \text{U}(1))$	Lattice models $S; H$	MPOs $S$	Field theories $S$
Type I $p_1$ : Eq. (8)	$\mathbb{Z}_N$	$\mathbb{Z}_N$	Eq. (27); Eq. (37)	Eq. (19)	Eq. (42)
Type II $p_{12}$ : Eq. (9)	$\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}$	$\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{N_{12}}$	Eq. (28); Eq. (37)	Eq. (19)	Eq. (42)
Type III $p_{123}$ : Eq. (10)	$\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{N_3}$	$\prod_{1 \leq i < j \leq 3} \mathbb{Z}_{N_i} \times \mathbb{Z}_{N_{ij}} \times \mathbb{Z}_{N_{123}}$	Eq. (31); Eq. (37)	Eq. (20)	Eq. (47)

II; it introduces a new quantum number, a different charge vector coupling  $Q$  for the conserved current term. *Although the Type III symmetry  $G$  is Abelian, its symmetry transformation operator has a noncommutative non-Abelian feature thus yielding degenerate zero energy modes.* In Secs. IV and V, we study the physical observables for bosonic anomalies of these SPT: induced fractional quantum numbers and degenerate zero energy modes. In Sec. VI, we work on the *twisted sector*: the effect of gauge flux insertion through a 1D ring effectively captured by using a branch cut or so-called monodromy defect [44] modifying the original Hamiltonian [22]. The *twisted non-onsite symmetry transformation* and *twisted lattice Hamiltonians* are studied, which spectral shift response under flux insertion provides physical observables to distinguish different SPTs [22,45] applicable for all Type I, II, and III SPTs. Our main results are summarized in Tables I, II, and III.

[Note: Our notation for finite cyclic group is either  $\mathbb{Z}_N$  or  $\mathbb{Z}_N$ , although mathematically they are the same. We denote  $\mathbb{Z}_N$  for the symmetry group  $G$ , the discrete gauge  $\mathbb{Z}_N$  flux, or the  $\mathbb{Z}_N$  variables. We denote  $\mathbb{Z}_N$  only for the classes of SPT classification. In addition, we denote  $n+1$ D as  $n$  dimensional space and one dimensional time, and denote  $n$ D as  $n$  dimensional space. We also denote  $\text{gcd}(N_i, N_j) \equiv N_{ij}$  and  $\text{gcd}(N_i, N_j, N_l) \equiv N_{ijl}$  with gcd standing for the greatest common divisor.]

## II. GROUP COHOMOLOGY AND COCYCLES

In this section, we will gather the information known and predicted by the group cohomology approach [6]. First, it has been predicted that the  $d+1$ D bosonic SPTs can be constructed by a mathematical object: the  $(d+1)$ th Borel cohomology group  $\mathcal{H}^{d+1}(G, \text{U}(1))$  of  $G$  over  $G$ -module  $\text{U}(1)$  [6,33]. (It is almost complete classification for bosons,

if without considering time-reversal symmetry.) The SPT classification itself as  $\mathcal{H}^{d+1}(G, \text{U}(1))$  also forms a group structure. Throughout the paper, we study a generic cyclic group  $G = \prod_{i=1}^m \mathbb{Z}_{N_i} = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{N_3} \times \dots$ . It is generic enough in the sense that any finite Abelian group is isomorphic to such a finite cyclic group  $G$ . We can thus compute its third cohomology group (see also Refs. [6,46]),

$$\mathcal{H}^3(G, \text{U}(1)) = \prod_{1 \leq i < j < l \leq m} \mathbb{Z}_{N_i} \times \mathbb{Z}_{\text{gcd}(N_i, N_j)} \times \mathbb{Z}_{\text{gcd}(N_i, N_j, N_l)}. \quad (2)$$

Here  $\text{gcd}(N_i, N_j, \dots)$  stands for the greatest common divisor among the numbers  $(N_i, N_j, \dots)$ . For simplicity, we denote  $\text{gcd}(N_i, N_j) \equiv N_{ij}$  and  $\text{gcd}(N_i, N_j, N_l) \equiv N_{ijl}$ . This cohomology group predicts that there are  $\mathbb{Z}_{N_i} \times \mathbb{Z}_{N_{ij}} \times \mathbb{Z}_{N_{ijl}}$  distinct classes for SPTs. One can find explicit 3-cocycles, such that each distinct 3-cocycle labels the distinct classes in SPTs. [More generally,  $(d+1)$ -cocycles for  $(d+1)$ th cohomology group  $\mathcal{H}^{d+1}(G, \text{U}(1))$ .] The  $n$ -cochain is a mapping  $\omega(A_1, A_2, \dots, A_n): G^n \rightarrow \text{U}(1)$  [which inputs  $A_i \in G$ ,  $i = 1, \dots, n$ , and outputs a  $\text{U}(1)$  phase]. The  $n$ -cochains satisfy the group multiplication rules:

$$(\omega_1 \omega_2)(A_1, \dots, A_n) = \omega_1(A_1, \dots, A_n) \omega_2(A_1, \dots, A_n), \quad (3)$$

thus forming an Abelian group. The  $n$ -cocycle is a  $n$ -cochain additionally satisfying the  $n$ -cocycle conditions  $\delta\omega = 1$ . The 3-cocycle condition (a pentagon relation) is

$$\delta\omega(A, B, C, D) = \frac{\omega(B, C, D)\omega(A, BC, D)\omega(A, B, C)}{\omega(AB, C, D)\omega(A, B, CD)} = 1 \quad (4)$$

with  $A, B, C, D \in G$ . One should check that the distinct 3-cocycles are not equivalent by 3-coboundaries, i.e., any  $\omega_1(A, B, C)$  is equivalent to  $\omega_2(A, B, C)$  if they are identified

TABLE III. The phase  $e^{i\Theta_L}$  on a domain wall  $D_u$  acted by  $\mathbb{Z}_{N_v}$  symmetry  $S_v$ . This phase is computed at the left kink (the site  $r_1$ ). The first column shows SPT class labels  $p$ . The second and the third columns show the computation of phases. The last column interprets whether the phase indicates a nontrivial induced  $\mathbb{Z}_N$  charge. Only Type II SPT class with  $p_{12} \neq 0$  contains nontrivial induced  $\mathbb{Z}_{N_2}$  charge with a unit of  $\mathbf{p}_{12}/N_{12}$  trapped at the kink of  $\mathbb{Z}_{N_1}$ -symmetry-breaking domain walls. Here  $n_3$  is the exponent inside the  $W^{\text{III}}$  matrix,  $n_3 = 0, 1, \dots, N_3 - 1$ , for each subblock within the total  $N_3$  subblocks [5].  $N_{12} \equiv \text{gcd}(N_1, N_2)$  and  $N_{123} \equiv \text{gcd}(N_1, N_2, N_3)$ .

SPT class	$e^{i\Theta_L}$ of $D_{N_u} \psi\rangle$ acted by $\mathbb{Z}_{N_v}$ symmetry $S_v$	$e^{i\Theta_L}$ of $D_{N_u} \psi\rangle$ under a soliton $\int_0^L dx \partial_x \phi_u = 2\pi$	Fractional charge
Type I $p_1$	$S_{N_1}^{(p_1)} D_{N_1}^{(p_1)} S_{N_1}^{(p_1)\dagger} \rightarrow e^{i\Theta_L} = e^{i \frac{2\pi p_1}{N_1}}$	$S_{N_1}^{(p_1)} (D_{N_1}^{(p_1)})^{N_1} S_{N_1}^{(p_1)\dagger} \rightarrow e^{i\Theta_L} = e^{i \frac{2\pi p_1}{N_1}}$	No
Type II $p_{12}$	$S_{N_2}^{(p_{12})} D_{N_1}^{(p_{12})} S_{N_2}^{(p_{12})\dagger} \rightarrow e^{i\Theta_L} = e^{i \frac{2\pi}{N_2} \frac{\mathbf{p}_{12}}{N_{12}}}$	$S_{N_2}^{(p_{12})} (D_{N_1}^{(p_{12})})^{N_{12}} S_{N_2}^{(p_{12})\dagger} \rightarrow e^{i\Theta_L} = e^{i \frac{2\pi p_{12}}{N_2}}$	Yes [Eqs. (54), (58)]
Type III $p_{123}$	$S_{N_2}^{(p_{123})} D_{N_1}^{(p_{123})} S_{N_2}^{(p_{123})\dagger} \rightarrow e^{i\Theta_L} = e^{i \frac{2\pi p_{123} n_3}{N_{123}}}$	$S_{N_2}^{(p_{123})} (D_{N_1}^{(p_{123})})^{N_{123}} S_{N_2}^{(p_{123})\dagger} \rightarrow e^{i\Theta_L} = 1$	No




FIG. 3. The illustration of 1D lattice model with  $M$  sites on a compact ring.

by 3-coboundaries  $\delta\Omega(A, B, C)$ :

$$\frac{\omega_1(A, B, C)}{\omega_2(A, B, C)} = \delta\Omega(A, B, C) = \frac{\Omega(B, C)\Omega(A, BC)}{\Omega(AB, C)\Omega(A, B)} \quad (5)$$

with some 2-cochain  $\Omega(B, C)$ . The 3-cochains form a group  $C^3$ , the 3-cochains satisfying the 3-cocycle conditions Eq. (4) further form a subgroup  $Z^3$ , and the 3-coboundaries satisfying Eq. (5) further form a subgroup  $B^3$  [since  $\delta^2\Omega(A, B, C) = 1$ ]. Overall

$$B^3 \subset Z^3 \subset C^3. \quad (6)$$

The third cohomology group is exactly a kernel  $Z^3$  (the group of 3-cocycles) mod out image  $B^3$  (the group of 3-coboundary) relation

$$\mathcal{H}^3(G, \text{U}(1)) = Z^3 / B^3. \quad (7)$$

For any finite Abelian group  $G$ , we can derive the distinct 3-cocycles satisfying Eq. (4) (but not identified as 3-coboundary by Eq. (5)):

$$\omega_{\text{I}}^{(i)}(A, B, C) = \exp\left(\frac{2\pi i p_i}{N_i^2} a_i(b_i + c_i - [b_i + c_i])\right), \quad (8)$$

$$\omega_{\text{II}}^{(ij)}(A, B, C) = \exp\left(\frac{2\pi i p_{ij}}{N_i N_j} a_i(b_j + c_j - [b_j + c_j])\right), \quad (9)$$

$$\omega_{\text{III}}^{(ijl)}(A, B, C) = \exp\left(\frac{2\pi i p_{ijl}}{\text{gcd}(N_i, N_j, N_l)} a_i b_j c_l\right), \quad (10)$$

so-called Type I, II, and III 3-cocycles [46], respectively. Here  $A, B, C \in G$ . We denote that  $A = (a_1, a_2, a_3, \dots)$ , where  $a_i \in Z_{N_i}$ , and similarly for  $B, C$ . And  $[b_i + c_i]$  are defined as the  $(b_i + c_i) \bmod N_i$ , the module elements in  $Z_{N_i}$ . In Table II, we summarize some data of group cohomology and their corresponding realization as SPT by using (i) quantum lattice model, (ii) matrix product states, and (iii) quantum field theory approach. In Sec. III, we will demonstrate their explicit construction for Type I, II, and III 3-cocycles and their corresponding Type I, II, and III SPTs.

### III. SPTS WITH $Z_{N_1} \times Z_{N_2} \times Z_{N_3}$ SYMMETRY

We will now go further to consider the edge modes of lattice Hamiltonian with  $G = Z_{N_1} \times Z_{N_2} \times Z_{N_3}$  symmetry on a compact ring with  $M$  sites (Fig. 3). Since there are at most three finite Abelian subgroup indices shown in Eqs. (8), (9), and (10), such a finite group with three Abelian discrete subgroups is the minimal example containing necessary and

sufficient information to explore finite Abelian SPTs. Such a symmetry group  $G$  may have nontrivial SPT class of Type I, II, and III SPTs. Apparently, the Type I SPTs studied in our previous work happen [22], which are the class of  $p_u \in \mathbb{Z}_{N_u}$  in  $\mathcal{H}^3(Z_{N_1} \times Z_{N_2} \times Z_{N_3}, \text{U}(1))$  of Eq. (2). Here and below we denote  $u, v, w \in \{1, 2, 3\}$  and  $u, v, w$  are distinct. We will also introduce the new class where  $Z_{N_u}$  and  $Z_{N_v}$  rotor models “talk to each other.” This will be the mixed Type II class  $p_{uv} \in \mathbb{Z}_{N_{uv}}$ , where symmetry transformation of  $Z_{N_1}$  global symmetry will affect the  $Z_{N_2}$  rotor models, while similarly  $Z_{N_2}$  global symmetry will affect the  $Z_{N_1}$  rotor models. There is a new class where three  $Z_{N_1}$ ,  $Z_{N_2}$ ,  $Z_{N_3}$  rotor models *directly talk to each other*. This will be the exotic Type III class  $p_{123} \in \mathbb{Z}_{N_{123}}$ , where the symmetry transformation of  $Z_{N_u}$  global symmetry will affect the mixed  $Z_{N_v}, Z_{N_w}$  rotor models *in a mutual way*.

To verify that our model construction corresponding to the Type I, II, and III 3-cocycles in Eqs. (8), (9), and (10), we will implement a technique called “matrix product operators” in Sec. III A. We would like to realize a discrete lattice model in Sec. III B and a continuum field theory in Sec. III C to capture the essence of these classes of SPTs.

#### A. Matrix product operators and cocycles

There are various advantages to put a quantum system on a discretized lattice, better than viewing it as a continuum field theory. For example, one advantage is that the symmetry transformation can be regularized so to understand its property such as onsite or non-onsite. Another advantage is that we can simulate our model by considering a discretized finite system with a finite dimensional Hilbert space. For our purpose, to regularize a quantum system on a discrete lattice, we will first use the matrix product operators (MPO) formalism (see Refs. [21,47] and references therein) to formulate our symmetry transformations corresponding to nontrivial 3-cocycles in the third cohomology group in  $\mathcal{H}^3(Z_{N_1} \times Z_{N_2}, \text{U}(1)) = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{N_{12}}$ .

First we formulate the unitary operator  $S$  as the MPO:

$$S = \sum_{\{j, j'\}} \text{tr} [T_{\alpha_1 \alpha_2}^{j_1 j'_1} T_{\alpha_2 \alpha_3}^{j_2 j'_2} \dots T_{\alpha_M \alpha_1}^{j_M j'_M}] |j'_1, \dots, j'_M\rangle \langle j_1, \dots, j_M| \quad (11)$$

with its coefficient taking the trace (tr) of a series of onsite tensor  $T(g)$  on a lattice, and input a state  $|j_1, \dots, j_M\rangle$  and output another state  $|j'_1, \dots, j'_M\rangle$ .  $T = T(g)$  is a tensor with multi-indices and with dependency on a group element  $g \in G$  for a symmetry group. This is the operator formalism of matrix product states (MPS). Here *physical indices*  $j_1, j_2, \dots, j_M$  and  $j'_1, j'_2, \dots, j'_M$  are labeled by input/output physical eigenvalues (here  $Z_N$  rotor angle), the subindices  $1, 2, \dots, M$  are the physical site indices. There are also *virtual indices*  $\alpha_1, \alpha_2, \dots, \alpha_M$  which are traced in the end. Summing over all the operations from  $\{j, j'\}$  indices, we shall reproduce the symmetry transformation operator  $S$ .

What MPO really helps us is that *by contracting MPO tensors  $T(g)$  of  $G$ -symmetry transformation  $S$  (here  $g \in G$ ) in different sequence on the effective 1D lattice of SPT edge modes, it can reveal the nontrivial projective phase*

corresponds to the nontrivial 3-cocycles of the cohomology group.

To find out the projective phase  $e^{i\theta(g_a, g_b, g_c)}$ , following we use the facts of tensors  $T(g_a), T(g_b), T(g_c)$  acting on the same site with group elements  $g_a, g_b, g_c$ . We know a generic projective relation:

$$T(g_a g_b) = P_{g_a, g_b}^\dagger T(g_a) T(g_b) P_{g_a, g_b}. \quad (12)$$

Here  $P_{g_a, g_b}$  is the projection operator. We contract three tensors in two different orders

$$(P_{g_a, g_b} \otimes I_3) P_{g_a, g_b, g_c} \simeq e^{i\theta(g_a, g_b, g_c)} (I_1 \otimes P_{g_b, g_c}) P_{g_a, g_b, g_c}. \quad (13)$$

$$\begin{aligned} & (T^{\phi_{in}^{(1)}, \phi_{out}^{(1)}, \phi_{in}^{(2)}, \phi_{out}^{(2)}})^{(p_1, p_{12})}_{\varphi_\alpha^{(1)}, \varphi_\beta^{(1)}, \varphi_\alpha^{(2)}, \varphi_\beta^{(2)}, N_1} \left( \frac{2\pi k_1}{N_1} \right) \\ &= \delta \left( \phi_{out}^{(1)} - \phi_{in}^{(1)} - \frac{2\pi k_1}{N_1} \right) \delta(\phi_{out}^{(2)} - \phi_{in}^{(2)}) \int d\varphi_\alpha^{(1)} d\varphi_\beta^{(1)} |\varphi_\beta^{(1)}\rangle \langle \varphi_\alpha^{(1)}| \delta(\varphi_\beta^{(1)} - \phi_{in}^{(1)}) e^{ip_1 k_1 (\varphi_\alpha^{(1)} - \phi_{in}^{(1)}) / N_1} \\ & \quad \times \int d\tilde{\varphi}_\alpha^{(2)} d\tilde{\varphi}_\beta^{(2)} |\tilde{\varphi}_\beta^{(2)}\rangle \langle \tilde{\varphi}_\alpha^{(2)}| \delta(\tilde{\varphi}_\beta^{(2)} - \tilde{\varphi}_{in}^{(2)}) e^{ip_{12} k_1 (\tilde{\varphi}_\alpha^{(2)} - \tilde{\varphi}_{in}^{(2)}) / N_1}. \end{aligned} \quad (14)$$

We propose the Type III  $T(g)$  tensor with  $p_{123} \in \mathbb{Z}_{N_{123}}$  as

$$\begin{aligned} & (T^{\phi_{in}^{(1)}, \phi_{out}^{(1)}, \phi_{in}^{(2)}, \phi_{out}^{(2)}, \phi_{in}^{(3)}, \phi_{out}^{(3)}})^{(p_{123})}_{\varphi_\alpha^{(1)}, \varphi_\beta^{(1)}, \varphi_\alpha^{(2)}, \varphi_\beta^{(2)}, \varphi_\alpha^{(3)}, \varphi_\beta^{(3)}, N_1, N_2, N_3} \left( \frac{2\pi k_1}{N_1}, \frac{2\pi k_2}{N_2}, \frac{2\pi k_3}{N_3} \right) \\ &= \prod_{u, v, w \in \{1, 2, 3\}} \int d\varphi_\alpha^{(u)} |\phi_{in}^{(u)}\rangle \langle \varphi_\alpha^{(u)}| \exp \left[ i p_{123} \epsilon^{uvw} k_u \frac{(\varphi_\alpha^{(v)} \phi_{in}^{(w)})_r}{N_u} \frac{N_1 N_2 N_3}{2\pi \text{gcd}(N_1, N_2, N_3)} \right] |\phi_{out}^{(u)}\rangle \langle \phi_{in}^{(u)}|. \end{aligned} \quad (15)$$

Here we consider a lattice with both  $\phi^{(u)}, \varphi^{(u)}$  as  $Z_{N_u}$  rotor angles. The tilde notation  $\tilde{\phi}^{(u)}, \tilde{\varphi}^{(u)}$ , for example on  $\tilde{\phi}^{(2)}$ , means that the variables are in units of  $\frac{2\pi}{N_{12}}$ , but not in  $\frac{2\pi}{N_2}$  units (the reason will become explicit later when we regularize the Hamiltonian on a lattice in Sec. III B).

Taking Eq. (14), by computing the projection operator  $P_{g_a, g_b}$  via Eq. (12), we derive the projective phase from Eq. (13):

$$e^{i\theta(g_a, g_b, g_c)} = e^{ip_1 \frac{2\pi}{N} m_c \frac{m_a + m_b - [m_a + m_b]_N}{N}} = \omega_1^{(i)}(m_c, m_a, m_b), \quad (16)$$

which the complex projective phase indeed induces the Type I 3-cocycle  $\omega_1^{(i)}(m_c, m_a, m_b)$  of Eq. (8) in the third cohomology group  $\mathcal{H}^3(Z_N, U(1)) = \mathbb{Z}_N$ . (Up to the index redefinition  $p_1 \rightarrow -p_1$ .) We further derive the projective phase as Type II 3-cocycle of Eq. (9):

$$\begin{aligned} e^{i\theta(g_a, g_b, g_c)} &= e^{ip_{12} \left( \frac{2\pi m_c^{(1)}}{N_1} \right) (m_a^{(2)} + m_b^{(2)}) - [m_a^{(2)} + m_b^{(2)}]_{N_2} / N_2} \\ &= \omega_{II}^{(ij)}(m_3, m_1, m_2) \end{aligned} \quad (17)$$

up to the index redefinition  $p_{12} \rightarrow -p_{12}$ . Here  $[m_a + m_b]_N$  with subindex  $N$  means taking the value module  $N$ .

Taking Eq. (15), we can also derive the projective phase  $e^{i\theta(g_a, g_b, g_c)}$  of Type III  $T(g)$  tensor as

$$\begin{aligned} e^{i\theta(g_a, g_b, g_c)} &= e^{i2\pi p_{123} \epsilon^{uvw} \left( \frac{m_c^{(u)}}{N_u} \frac{m_a^{(v)}}{N_v} \frac{m_b^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{N_{123}}} \\ &\simeq \omega_{III}^{(uvw)}(m_c, m_a, m_b). \end{aligned} \quad (18)$$

The left-hand side contracts the  $a, b$  first then with the  $c$ , while the right-hand side contracts the  $b, c$  first then with the  $a$ . Here  $\simeq$  means the equivalence is up to a projection out of unparallel states. We can derive  $P_{g_a, g_b}$  by observing that  $P_{g_a, g_b}$  inputs one state and outputs two states [5].

For Type I SPT class, this MPO formalism has been done quite carefully in Refs. [21, 22]. Here we generalize it to other SPTs, below we input a group element with  $g = (k_1, k_2, k_3)$  and  $k_1 \in \mathbb{Z}_{N_1}, k_2 \in \mathbb{Z}_{N_2}, k_3 \in \mathbb{Z}_{N_3}$ . Without losing generality, we focus on the symmetry Type I index  $p_1 \in \mathbb{Z}_{N_1}$ , Type II index  $p_{12} \in \mathbb{Z}_{N_{12}}$ , Type III index  $p_{123} \in \mathbb{Z}_{N_{123}}$ . By index relabeling, we can fulfill all SPT symmetries within the classification in Eq. (2).

We propose our  $T(g)$  tensor for Type I [21, 22] and II symmetries with  $p_1 \in \mathbb{Z}_{N_1}, p_{12} \in \mathbb{Z}_{N_{12}}$  as

Adjust  $p_{123}$  index (i.e., setting only the  $p_{123}$  index in  $m_c^{(1)} m_a^{(2)} m_b^{(3)}$  to be nonzero, while others  $p_{213} = p_{312} = 0$ ), and compute Eq. (13) with only  $p_{123}$  index, we can recover the projective phase reveals Type III 3-cocycle in Eq. (10).

By Eq. (11), we verify that  $T(g)$  of Types I and II in Eq. (14) renders the symmetry transformation operator  $S_{N_1}^{(p_1, p_{12})}$ :

$$\begin{aligned} S_{N_1}^{(p_1, p_{12})} &= \prod_{j=1}^M e^{i2\pi L_j^{(1)} / N_1} \exp \left[ i \frac{p_1}{N_1} (\phi_{j+1}^{(1)} - \phi_j^{(1)})_r \right] \\ & \quad \times \exp \left[ i \frac{p_{12}}{N_1} (\tilde{\phi}_{j+1}^{(2)} - \tilde{\phi}_j^{(2)})_r \right], \end{aligned} \quad (19)$$

where  $j$  are the site indices from 1 to  $M$  shown in Fig. 3.

By Eq. (11), we verify that  $T(g)$  of Type III in Eq. (15) renders the symmetry transformation operator  $S_{N_1, N_2, N_3}^{(p_{123})}$ :

$$S_{N_1, N_2, N_3}^{(p_{123})} = \prod_{j=1}^M \left( \prod_{u, v, w \in \{1, 2, 3\}} e^{i2\pi L_j^{(u)} / N_u} W_{j, j+1}^{III} \right) \quad (20)$$

with

$$W_{j, j+1}^{III} \equiv \prod_{u, v, w \in \{1, 2, 3\}} e^{i \frac{N_1 N_2 N_3}{2\pi N_{123}} \epsilon^{uvw} \frac{p_{123}}{N_u} (\phi_{j+1}^{(v)} \phi_j^{(w)})}. \quad (21)$$

For both Eqs. (19) and (20), there is an onsite piece  $\langle \phi_j^{(u)} | e^{i2\pi L_j^{(u)}/N_u} | \phi_j^{(u)} \rangle$  and also extra non-onsite symmetry transformation parts: namely,  $\exp[i\frac{p_1}{N_1}(\phi_{j+1}^{(1)} - \phi_j^{(1)})_r]$ ,  $\exp[i\frac{p_{12}}{N_1}(\tilde{\phi}_{j+1}^{(2)} - \tilde{\phi}_j^{(2)})_r]$ , and  $W_{j,j+1}^{\text{III}}$ . We introduce an angular momentum operator  $L_j^{(u)}$  conjugate to  $\phi_j^{(u)}$ , such that the  $e^{i2\pi L_j^{(u)}/N_u}$  shifts the rotor angle by  $\frac{2\pi}{N_u}$  unit, from  $|\phi_j^{(u)}\rangle$  to  $|\phi_j^{(u)} + \frac{2\pi}{N_u}\rangle$ . The subindex  $r$  means that we further *regularize the variable to a discrete compact rotor angle*.

Meanwhile  $p_1 = p_1 \bmod N_1$ ,  $p_{12} = p_{12} \bmod N_{12}$ , and  $p_{123} = p_{123} \bmod N_{123}$ ; these demonstrate that our MPO construction fulfills all classes in Eq. (2) as we desire. So far we have achieved the SPT symmetry transformation operators (19) and (20) via MPO. Other technical derivations on MPO formalism are preserved in [5].

### B. Lattice model

To construct a lattice model, we require the minimal ingredients: (i)  $Z_{N_u}$  operators (with  $Z_{N_u}$  variables), (ii) Hilbert space (the state space where  $Z_{N_u}$  operators act on) consists with  $Z_{N_u}$  variables state. Again we denote  $u = 1, 2, 3$  for  $Z_{N_1}, Z_{N_2}, Z_{N_3}$  symmetry. We can naturally choose the  $Z_{N_u}$  variable  $\omega_u \equiv e^{i2\pi/N_u}$ , such that  $\omega_u^{N_u} = 1$ . Here and following we will redefine the quantum state and operators from the MPO basis in Sec. III A to a lattice basis via

$$\phi_j^{(u)} \rightarrow \phi_{u,j}, \quad L_l^{(u)} \rightarrow L_{u,l}. \quad (22)$$

The natural physical states on a single site are the  $Z_{N_u}$  rotor angle state  $|\phi_u = 0\rangle, |\phi_u = 2\pi/N_u\rangle, \dots, |\phi_u = 2\pi(N_u - 1)/N_u\rangle$ .

One can find a dual state of rotor angle state  $|\phi_u\rangle$ , the angular momentum  $|L_u\rangle$ , such that the basis from  $|\phi_u\rangle$  can transform to  $|L_u\rangle$  via the Fourier transformation  $|\phi_u\rangle = \sum_{L_u=0}^{N_u-1} \frac{1}{\sqrt{N_u}} e^{iL_u\phi_u} |L_u\rangle$ . One can find two proper operators  $\sigma^{(u)}, \tau^{(u)}$  which make  $|\phi_u\rangle$  and  $|L_u\rangle$  their own eigenstates, respectively. With a site index  $j$  ( $j = 1, \dots, M$ ), we can project  $\sigma_j^{(u)}, \tau_j^{(u)}$  operators into the rotor angle  $|\phi_{u,j}\rangle$  basis, so we can derive  $\sigma_j^{(u)}, \tau_j^{(u)}$  operators as  $N_u \times N_u$  matrices. Their forms are

$$\sigma_j^{(u)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega_u & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \omega_u^{N_u-1} \end{pmatrix}_j = \langle \phi_{u,j} | e^{i\hat{\phi}_j^{(u)}} | \phi_{u,j} \rangle, \quad (23)$$

$$\tau_j^{(u)} = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & 0 & 0 & \dots & 1 & 0 \end{pmatrix}_j = \langle \phi_{u,j} | e^{i2\pi\hat{L}_j^{(u)}/N} | \phi_{u,j} \rangle. \quad (24)$$

Operators and variables satisfy the analog property mentioned in Ref. [22], such as  $(\tau^{(u)})_j^{N_u} = (\sigma^{(u)})_j^{N_u} = \mathbb{1}$ ,  $\tau_j^{(u)\dagger} \sigma_j^{(u)} \tau_j^{(u)} = \omega_u \sigma_j^{(u)}$ . It also enforces the canonical conjugation relation on  $\hat{\phi}^{(u)}$  and  $\hat{L}^{(u)}$  operators, i.e.,  $[\hat{\phi}_j^{(u)}, \hat{L}_l^{(v)}] = i \delta_{(j,l)} \delta_{(u,v)}$  with the

symmetry group index  $u, v$  and the site indices  $j, l$ . Here  $|\phi\rangle$  and  $|L\rangle$  are eigenstates of  $\hat{\phi}$  and  $\hat{L}$  operators, respectively.

The linear combination of all  $|\phi_1\rangle |\phi_2\rangle |\phi_3\rangle$  states form a complete  $N_1 \times N_2 \times N_3$  dimensional Hilbert space on a single site.

### 1. Symmetry transformations

*Type I and II  $Z_{N_1} \times Z_{N_2}$  symmetry transformations.* First, we warm up with a generic  $Z_N$  lattice model realizing the SPT edge modes on a 1D ring with  $M$  sites (Fig. 3). It has been emphasized in Refs. [6,21] that the SPT edge modes have a special non-onsite symmetry transformation, which means that its symmetry transformation cannot be written as a tensor product form on each site, thus  $U(g)_{\text{non-onsite}} \neq \otimes_i U_i(g)$ . In general, the symmetry transformation contains an onsite part and another non-onsite part. The trivial class of SPT (trivial bulk insulator) with unprotected gapped edge modes can be achieved by a simple Hamiltonian as  $-\lambda \sum_{j=1}^M (\tau_j + \tau_j^\dagger)$ . [Notice that for the simplest  $Z_2$  symmetry, the  $\tau_j$  operator reduces to a spin operator  $(\sigma_z)_j$ .] The simple way to find an onsite operator which this Hamiltonian respects and which acts at each site is the  $\prod_{j=1}^M \tau_j$ , a series of  $\tau_j$ . On the other hand, to capture the *non-onsite* symmetry transformation, we can use a *domain wall* variable in Ref. [22], where the symmetry transformation contains information stored nonlocally between different sites (here we will use the minimum construction: symmetry stored nonlocally between *two nearest neighbored sites*). Based on the understanding of previous work [19,21,22], we propose this non-onsite symmetry transformation  $U_{j,j+1}$  with a domain wall  $(N_{\text{dw}})_{j,j+1}$  operator acting nonlocally on sites  $j$  and  $j+1$  as

$$U_{j,j+1} \equiv \exp\left(i\frac{p}{N}\frac{2\pi}{N}(\delta N_{\text{dw}})_{j,j+1}\right) \equiv \exp\left[i\frac{p}{N}(\phi_{1,j+1} - \phi_{1,j})_r\right]. \quad (25)$$

The justification of non-onsite symmetry operator (25) realizing SPT edge symmetry is based on MPO formalism already done in Sec. III A. The domain wall operator  $(\delta N_{\text{dw}})_{j,j+1}$  counts the number of units of  $Z_N$  angle between sites  $j$  and  $j+1$ , so indeed  $(2\pi/N)(\delta N_{\text{dw}})_{j,j+1} = (\phi_{1,j+1} - \phi_{1,j})_r$ . The subindex  $r$  means that we need to further *regularize the variable to a discrete  $Z_N$  angle*. Here we insert a  $p$  index, which is just an available free index with  $p = p \bmod N$ . From Sec. III A,  $p$  is indeed the classification index for the  $p$ th of  $Z_N$  class in the third cohomology group  $\mathcal{H}^3(Z_N, \text{U}(1)) = \mathbb{Z}_N$ .

Now the question is how should we fully regularize this  $U_{j,j+1}$  operator into terms of  $Z_N$  operators  $\sigma_j^\dagger$  and  $\sigma_{j+1}$ . We see the fact that the  $N$ th power of  $U_{j,j+1}$  renders a constraint

$$U_{j,j+1}^N = (\exp[i\phi_{1,j}]^\dagger \exp[i\phi_{1,j+1}])^p = (\sigma_j^\dagger \sigma_{j+1})^p \quad (26)$$

(since  $\exp[i\phi_{1,j}]_{ab} = \langle \phi_a | e^{i\phi_j} | \phi_b \rangle = \sigma_{ab,j}$ ). More explicitly, we can write it as a polynomial ansatz  $U_{j,j+1} = \exp[\frac{i}{N} \sum_{a=0}^{N-1} q_a (\sigma_j^\dagger \sigma_{j+1})^a]$ . The non-onsite symmetry operator  $U_{j,j+1}$  reduces to a problem of solving polynomial coefficients  $q_a$  by the constraint (26). Indeed we can solve the constraint explicitly, thus the non-onsite symmetry transformation operator acting on a  $M$ -site ring from  $j = 1, \dots, M$



is derived:

$$U_{j,j+1} = e^{-i\frac{2\pi}{N_2} p \left\{ \left( \frac{N-1}{2} \right) \mathbb{1} + \sum_{a=1}^{N-1} \frac{(\sigma_j^\dagger \sigma_{j+1})^a}{(\omega^a)^{N-1}} \right\}}. \quad (27)$$

For a lattice SPT model with  $G = Z_{N_1} \times Z_{N_2}$ , we can convert MPO's symmetry transformation Eq. (19) to a lattice variable via Eq. (27). We obtain the  $Z_{N_u}$  symmetry transformation (here and below  $u, v \in \{1, 2\}, u \neq v$ ):

$$\begin{aligned} S_{N_u}^{(p_u, p_{uv})} &\equiv \prod_{j=1}^M e^{i2\pi L_{u,j}/N_u} \exp \left[ i \frac{p_u}{N_u} (\phi_{u,j+1} - \phi_{u,j})_r \right] \\ &\quad \times \exp \left[ i \frac{p_{uv}}{N_u} (\tilde{\phi}_{v,j+2} - \tilde{\phi}_{v,j})_r \right] \\ &= \prod_{j=1}^M \tau_j^{(u)} U_{j,j+1}^{(N_u, p_u)} U_{j,j+2}^{(N_u, p_{uv})} \\ &= \prod_{j=1}^M \tau_j^{(u)} e^{(-i\frac{2\pi}{N_u} p_u \left\{ \left( \frac{N_u-1}{2} \right) \mathbb{1} + \sum_{a=1}^{N_u-1} \frac{(\sigma_j^\dagger \sigma_{j+1})^a}{(\omega^a)^{N_u-1}} \right\})} \\ &\quad \times e^{(-i\frac{2\pi}{N_u N_u} p_{uv} \left\{ \left( \frac{N_u-1}{2} \right) \mathbb{1} + \sum_{a=1}^{N_u-1} \frac{(\tilde{\sigma}_j^\dagger \tilde{\sigma}_{j+2})^a}{(\omega^a)^{N_u-1}} \right\})}. \end{aligned} \quad (28)$$

The operator is unitary, i.e.,  $S_{N_u}^{(p_u, p_{uv})} S_{N_u}^{(p_u, p_{uv})\dagger} = 1$ . Here  $\sigma_{M+j} \equiv \sigma_j$ .

The intervals of rotor angles are

$$\begin{aligned} \phi_{1,j} &\in \left\{ n \frac{2\pi}{N_1} \mid n \in \mathbb{Z} \right\}, \quad \phi_{2,j} \in \left\{ n \frac{2\pi}{N_2} \mid n \in \mathbb{Z} \right\}, \\ \tilde{\phi}_{1,j}, \tilde{\phi}_{2,j} &\in \left\{ n \frac{2\pi}{N_{12}} \mid n \in \mathbb{Z} \right\}, \end{aligned} \quad (29)$$

where  $\phi_{1,j}$  is  $Z_{N_1}$  angle,  $\phi_{2,j}$  is  $Z_{N_2}$  angle,  $\tilde{\phi}_{1,j}$  and  $\tilde{\phi}_{2,j}$  are  $Z_{N_{12}}$  angles [recall  $\gcd(N_1, N_2) \equiv N_{12}$ ]. There are some remarks on our above formalism: (i) First, the  $Z_{N_1}, Z_{N_2}$  symmetry transformation (28) including both the Type I indices  $p_1, p_2$  and also Type II indices  $p_{12}$  and  $p_{21}$ . Although  $p_1, p_2$  are distinct indices, but  $p_{12}$  and  $p_{21}$  indices are the same index,  $p_{12} + p_{21} \rightarrow p_{12}$ . The invariance  $p_{12} + p_{21}$  describes the same SPT symmetry class. (ii) For Type I, non-onsite symmetry transformation (with  $p_1$  and  $p_2$ ) is chosen to act on the nearest-neighbor sites (NN: site  $j$  and site- $j+1$ ), but the Type II non-onsite symmetry transformation (with  $p_{12}$  and  $p_{21}$ ) is chosen to be the next-nearest-neighbor sites (NNN: sites  $j$  and  $j+2$ ). The reason is that we have to avoid the nontrivial Type I and II symmetry transformations canceling or interfering with each other. Although in Sec. III C, we will reveal that the low energy field theory description of non-onsite symmetry transformations for both NN and NNN having the same form in the continuum limit. In the absence of Type I index, we can have Type II non-onsite symmetry transformation act on nearest-neighbor sites. (iii) The domain wall picture mentioned in Eq. (25) of Sec. III for Type II  $p_{12}$  class still holds. But here the lattice regularization is different for terms with  $p_{12}, p_{21}$  indices. In order to have distinct  $Z_{\gcd(N_1, N_2)}$  class with the identification  $p_{12} = p_{12} \bmod N_{12}$ . We will expect that performing the  $N_u$  times  $Z_{N_u}$  symmetry transformation on the Type II  $p_{uv}$  non-onsite piece renders a

constraint

$$(U_{j,j+2}^{(N_u, p_{uv})})^{N_u} = (\tilde{\sigma}_j^{(v)\dagger} \tilde{\sigma}_{j+2}^{(v)})^{p_{uv}}. \quad (30)$$

To impose the identification  $p_{12} = p_{12} \bmod N_{12}$  and  $p_{21} = p_{21} \bmod N_{12}$  so that we have distinct  $Z_{\gcd(N_1, N_2)}$  classes for the Type II symmetry class [which leads to impose the constraint  $(\tilde{\sigma}_j^{(1)})^{N_{12}} = (\tilde{\sigma}_j^{(2)})^{N_{12}} = \mathbb{1}$ ], we can regularize the  $\tilde{\sigma}_j^{(1)}, \tilde{\sigma}_j^{(2)}$  operators in terms of  $Z_{\gcd(N_1, N_2)}$  variables.

With  $\omega_{12} \equiv \omega_{21} \equiv e^{i\frac{2\pi}{N_{12}}}$ , we have  $\omega_{12}^{N_{12}} = 1$ . The  $\tilde{\sigma}_j^{(u)}$  matrix has  $N_u \times N_u$  components, for  $u = 1, 2$ . It is block diagonalizable with  $\frac{N_u}{N_{12}}$  subblocks, and each subblock with  $N_{12} \times N_{12}$  components. Our regularization provides the nice property:  $\tau_j^{(1)\dagger} \tilde{\sigma}_j^{(1)} \tau_j^{(1)} = \omega_{12} \sigma_j^{(1)}$  and  $\tau_j^{(2)\dagger} \tilde{\sigma}_j^{(2)} \tau_j^{(2)} = \omega_{12} \sigma_j^{(2)}$ . Using the above procedure to regularize Eq. (19) on a discretized lattice and solve the constraint (30), we obtain an explicit form of lattice-regularized symmetry transformations (28).

For more details on our lattice regularization, see Supplemental Material [5].

*Type III symmetry transformations.* To construct a Type III SPT with a Type III 3-cocycle [Eq. (10)], the key observation is that the 3-cocycle inputs, for example,  $a_1 \in Z_{N_1}, b_2 \in Z_{N_2}, c_3 \in Z_{N_3}$  and outputs a U(1) phase. This implies that the  $Z_{N_1}$  symmetry transformation will affect the mixed  $Z_{N_2}, Z_{N_3}$  rotor models, etc. This observation guides us to write the tensor  $T(g)$  in Eq. (15) and we obtain the symmetry transformation  $S_N^{(p)} = S_{N_1, N_2, N_3}^{(p_{123})}$  as Eq. (20):

$$S_{N_1, N_2, N_3}^{(p_{123})} = \prod_{j=1}^M \left( \prod_{u, v, w \in \{1, 2, 3\}} \tau_j^{(u)} W_{j, j+1}^{\text{III}} \right). \quad (31)$$

There is an onsite piece  $\tau_j \equiv \langle \phi_j | e^{i2\pi L_j^{(u)}/N} | \phi_j \rangle$  and also an extra non-onsite symmetry transformation part  $W_{j, j+1}^{\text{III}}$ . This non-onsite symmetry transformation  $W_{j, j+1}^{\text{III}}$ , acting on the sites  $j$  and  $j+1$ , is defined by the following, and can be further regularized on the lattice:

$$W_{j, j+1}^{\text{III}} = \prod_{u, v, w \in \{1, 2, 3\}} (\sigma_j^{(v)\dagger} \sigma_{j+1}^{(v)})^{e^{uvw} p_{123} - \frac{\ln(\sigma_j^{(w)})_{N_u N_v N_w}}{2\pi N_{123}}}. \quad (32)$$

Here we separate  $Z_{N_1}, Z_{N_2}, Z_{N_3}$  non-onsite symmetry transformations to  $W_{j, j+1; N_1}^{\text{III}}, W_{j, j+1; N_2}^{\text{III}}, W_{j, j+1; N_3}^{\text{III}}$ , respectively. Equations (31) and (32) are fully regularized in terms of  $Z_N$  variables on a lattice, although they contain *anomalous non-onsite symmetry operators* [5].

## 2. Lattice Hamiltonians

We had mentioned the trivial class of SPT Hamiltonian (the class of  $p = 0$ ) for 1D gapped edge:

$$H_N^{(0)} = -\lambda \sum_{j=1}^M (\tau_j + \tau_j^\dagger). \quad (33)$$

Apparently, the Hamiltonian is symmetry preserving with respect to  $S_N^{(0)} \equiv \prod_{j=1}^M \tau_j$ , i.e.,  $S_N^{(0)} H_N^{(0)} (S_N^{(0)})^{-1} = H_N^{(0)}$ . In addition, this Hamiltonian has a symmetry-preserving gapped ground state.



To extend our lattice Hamiltonian construction to  $p \neq 0$  class, intuitively we can view the nontrivial SPT Hamiltonians as close relatives of the trivial Hamiltonian (which preserves the onsite part of the symmetry transformation with  $p = 0$ ), which satisfies the symmetry-preserving constraint, i.e.,

$$S_N^{(p)} H_N^{(p)} (S_N^{(p)})^{-1} = H_N^{(p)}. \quad (34)$$

More explicitly, to construct a SPT Hamiltonian of  $Z_{N_1} \times Z_{N_2} \times Z_{N_3}$  symmetry obeying translation and symmetry transformation invariant (here and following  $u, v, w \in \{1, 2, 3\}$  and  $u, v, w$  are distinct):

$$[H_{N_1, N_2, N_3}^{(p_u, p_{uv}, p_{uvw})}, T] = 0, \quad (35)$$

$$[H_{N_1, N_2, N_3}^{(p_u, p_{uv}, p_{uvw})}, S_N^{(p)}] = 0. \quad (36)$$

Here  $T$  is a translation operator by one lattice site, satisfying  $T^\dagger X_j T = X_{j+1}$ ,  $j = 1, \dots, M$ , for any operator  $X_j$  on the ring such that  $X_{M+1} \equiv X_1$ . Also,  $T$  satisfies  $T^M = \mathbb{1}$ . We can immediately derive the following SPT Hamiltonian satisfying the rules

$$H_{N_1, N_2, N_3}^{(p_u, p_{uv}, p_{uvw})} \equiv -\lambda \sum_{j=1}^M \sum_{\ell=0}^{M-N-1} (S_N^{(p)})^{-\ell} (\tau_j + \tau_j^\dagger) (S_N^{(p)})^\ell + \dots, \quad (37)$$

where we define our notations  $S_N^{(p)} \equiv \prod_{u,v,w \in \{1,2,3\}} S_{N_u}^{(p_u, p_{uv}, p_{uvw})}$  and  $\tau_j \equiv \tau_j^{(1)} \otimes \mathbb{1}_{N_2 \times N_2} \otimes \mathbb{1}_{N_3 \times N_3} + \mathbb{1}_{N_1 \times N_1} \otimes \tau_j^{(2)} \otimes \mathbb{1}_{N_3 \times N_3} + \mathbb{1}_{N_1 \times N_1} \otimes \mathbb{1}_{N_2 \times N_2} \otimes \tau_j^{(3)}$ . Here  $\tau_j$  is a matrix of  $(N_1 \times N_2 \times N_3) \times (N_1 \times N_2 \times N_3)$  components. The tower series of sum over power of  $(S_N^{(p)})$  over  $(\tau_j + \tau_j^\dagger)$  will be shifted upon  $S_N^{(p)} H_N^{(p)} (S_N^{(p)})^{-1}$ , but the overall sum of this Hamiltonian is a symmetry-preserving invariant.

### C. Field theory

From a fully regularized lattice model in the previous section, we attempt to take the low energy limit to realize its corresponding field theory, by identifying the commutation relation  $[\hat{\phi}_j^{(u)}, \hat{L}_l^{(v)}] = i \delta_{(j,l)} \delta_{(u,v)}$  (here  $j, l$  are the site indices,  $u, v \in \{1, 2, 3\}$  are the  $Z_{N_1}, Z_{N_2}, Z_{N_3}$  rotor model indices) in the continuum as

$$\left[ \phi_u(x_1), \frac{1}{2\pi} \partial_x \phi'_v(x_2) \right] = i \delta(x_1 - x_2) \delta_{(u,v)}, \quad (38)$$

which means the  $Z_{N_1}, Z_{N_2}, Z_{N_3}$  lattice operators  $\hat{\phi}_j^{(1)}, \hat{L}_l^{(1)}, \hat{\phi}_j^{(2)}, \hat{L}_l^{(2)}, \hat{\phi}_j^{(3)}, \hat{L}_l^{(3)}$  and field operators  $\phi_1, \phi'_1, \phi_2, \phi'_2, \phi_3, \phi'_3$  are identified by

$$\hat{\phi}_j^{(u)} \rightarrow \phi_u(x_j), \quad \hat{L}_l^{(u)} \rightarrow \frac{1}{2\pi} \partial_x \phi'_u(x_l). \quad (39)$$

We view  $\phi_u$  and  $\phi'_u$  as the dual rotor angles as before, the relation follows as Sec. III C. We have no difficulty to formulate a  $K$  matrix multiplet chiral boson field theory

(nonchiral “doubled” version of Ref. [48]’s action) as

$$\mathbf{S}_{\text{SPT}, \partial \mathcal{M}^2} = \frac{1}{4\pi} \int dt dx (K_{IJ} \partial_t \phi_I \partial_x \phi_J - V_{IJ} \partial_x \phi_I \partial_x \phi_J) + \dots \quad (40)$$

requiring a rank-6 symmetric  $K$  matrix

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (41)$$

with a chiral boson multiplet  $\phi_I(x) = (\phi_1(x), \phi'_1(x), \phi_2(x), \phi'_2(x), \phi_3(x), \phi'_3(x))$ . The commutation relation (38) becomes  $[\phi_I(x_1), K_{I'J} \partial_x \phi_J(x_2)] = 2\pi i \delta_{II'} \delta(x_1 - x_2)$ . The continuum limit of Eq. (28) becomes [49]

$$S_{N_u}^{(p_u, p_{uv})} = \exp \left[ \frac{i}{N_u} \left( \int_0^L dx \partial_x \phi'_u + p_u \int_0^L dx \partial_x \phi_u + 0 \int_0^L dx \partial_x \phi'_v + p_{uv} \int_0^L dx \partial_x \tilde{\phi}_v \right) \right]. \quad (42)$$

Notice that we carefully input a tilde on some  $\tilde{\phi}_v$  fields. We stress the lattice regularization of  $\tilde{\phi}_v$  is different from  $\phi_v$  [see Eq. (29)], which is analogous to  $\tilde{\sigma}^{(1)}, \tilde{\sigma}^{(2)}$  and  $\sigma^{(1)}, \sigma^{(2)}$  in Sec. IIIB1. We should mention two remarks: First, there are higher order terms beyond  $\mathbf{S}_{\text{SPT}, \partial \mathcal{M}^2}$ ’s quadratic terms when taking continuum limit of lattice. At the low energy limit, it shall be reasonable to drop higher order terms. Second, in the nontrivial SPT class (some topological terms  $p_i \neq 0, p_{ij} \neq 0$ ), the  $\det(V) \neq 0$  and all eigenvalues are nonzeros, so the edge modes are gapless. In the trivial insulating class (all topological terms  $p = 0$ ), the  $\det(V) = 0$ , so the edge modes may be gapped (consistent with Sec. IIIB2). Using Eq. (38), we derive the 1D edge global symmetry transformation  $S_{N_u}^{(p_u, p_{uv})}$ , for example,  $S_{N_1}^{(p_1, p_{12})}$  and  $S_{N_2}^{(p_2, p_{21})}$  [49]:

$$S_{N_1}^{(p_1, p_{12})} \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} (S_{N_1}^{(p_1, p_{12})})^{-1} = \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} + \frac{2\pi}{N_1} \begin{pmatrix} 1 \\ p_1 \\ 0 \\ p_{12} \end{pmatrix}, \quad (43)$$

$$S_{N_2}^{(p_2, p_{21})} \begin{pmatrix} \tilde{\phi}_1(x) \\ \tilde{\phi}'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} (S_{N_2}^{(p_2, p_{21})})^{-1} = \begin{pmatrix} \tilde{\phi}_1(x) \\ \tilde{\phi}'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} + \frac{2\pi}{N_2} \begin{pmatrix} 0 \\ p_{21} \\ 1 \\ p_2 \end{pmatrix}. \quad (44)$$

We can see how  $p_{12}, p_{21}$  identify the same index by doing a  $M$  matrix with  $M \in \text{SL}(4, \mathbb{Z})$  transformation on the  $K$  matrix Chern-Simons theory, which redefines the  $\phi$  field, but still describes the same theory. That means  $K \rightarrow K' = M^T K M$  and  $\phi \rightarrow \phi' = M^{-1} \phi$ , and so the symmetry charge vector  $q \rightarrow q' = M^{-1} q$ . By choosing

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 \\ p_1 & 1 & p_{21} & 0 \\ 0 & 0 & 1 & 0 \\ p_{12} & 0 & p_2 & 1 \end{pmatrix}, \quad \text{then the basis is changed to}$$

$$K' = \begin{pmatrix} 2p_1 & 1 & p_{12} + p_{21} & 0 \\ 1 & 0 & 0 & 0 \\ p_{12} + p_{21} & 0 & 2p_2 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

$$q'_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad q'_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}.$$

The theory labeled by  $K_{\text{SPT}}, q_1, q_2$  is equivalent to the one labeled by  $K', q'_1, q'_2$ . Thus we show that  $p_{12} + p_{21} \rightarrow p_{12}$  identifies the same index. There are other ways of using the gauged or probed-field version of topological gauge theory (either on the edge or in the bulk) to identify the gauge theory's symmetry transformation [26] or the bulk braiding statistics [50] to determine this Type II classification  $p_{12} \bmod[\text{gcd}(N_1, N_2)]$ .

The nontrivial fact that when  $p_{12} = N_{12}$  is a trivial class, the symmetry transformation in Eq. (43) may not go back to the trivial symmetry under the condition  $\int_0^L dx \partial_x \tilde{\phi}_1 = \int_0^L dx \partial_x \tilde{\phi}_2 = 2\pi$ , implying a soliton can induce fractional charge (for details see Sec. IV).

Our next goal is deriving Type III symmetry transformation (20). By taking the continuum limit of

$$\epsilon^{(u=1,2,3)(v)(w)} \phi_{in}^{j+1,(v)} \phi_{in}^{j,(w)} \quad (45)$$

$$= [(\phi_{in}^{j+1,(v)} - \phi_{in}^{j,(v)}) \phi_{in}^{j,(w)} - (\phi_{in}^{j+1,(w)} - \phi_{in}^{j,(w)}) \phi_{in}^{j,(v)}]$$

$$\rightarrow [\partial_x \phi_{in}^{(v)}(x) \phi_{in}^{(w)}(x) - \partial_x \phi_{in}^{(w)}(x) \phi_{in}^{(v)}(x)], \quad (46)$$

we can massage the continuum limit of Type III symmetry transformation (20) to  $[\text{gcd}(N_1, N_2, N_3) \equiv N_{123}]$

$$S_{N_1, N_2, N_3}^{(p_{123})} = \prod_{u,v,w \in \{1,2,3\}} \exp \left[ \frac{i}{N_u} \left( \int_0^L dx \partial_x \phi'_u \right) \right]$$

$$\times \exp \left[ i \frac{N_1 N_2 N_3}{2\pi N_{123}} \frac{p_{123}}{N_u} \right]$$

$$\times \int_0^L dx \epsilon^{uvw} \partial_x \phi_v(x) \phi_w(x). \quad (47)$$

Here  $u, v, w \in \{1, 2, 3\}$  is the label of the symmetry group  $Z_{N_1}, Z_{N_2}, Z_{N_3}$ 's indices. Although this Type III class is already known in the group cohomology sense, this Type III field theory symmetry transformation result is exotic, unfamiliar and not yet been well explored in the literature, especially not yet studied in the field theory in the SPT context. Our result is an extension along the work of Refs. [24, 26].

The commutation relation leads to

$$[\phi_I(x_i), K_{I'J} \phi_J(x_j)] = -2\pi i \delta_{II'} \tilde{h}(x_i - x_j). \quad (48)$$

Here  $\tilde{h}(x_i - x_j) \equiv h(x_i - x_j) - 1/2$ , where  $h(x)$  is the Heaviside step function, with  $h(x) = 1$  for  $x \geq 0$  and  $h(x) = 0$  for  $x < 0$ . And  $\tilde{h}(x)$  is  $h(x)$  shifted by  $\frac{1}{2}$ , i.e.,  $\tilde{h}(x) = 1/2$  for  $x \geq 0$  and  $h(x) = -1/2$  for  $x < 0$ . The shifted  $\frac{1}{2}$  value is for the consistency condition for the integration-by-parts and the commutation relation. Using these relations, we derive the

global symmetry transformation  $S_{N_1, N_2, N_3}^{(p_{123})}$  acting on the rotor fields  $\phi_u(x), \phi'_u(x)$  (here  $u \in \{1, 2, 3\}$ ) on the 1D edge by

$$(S_{N_1, N_2, N_3}^{(p_{123})}) \phi_u(x) (S_{N_1, N_2, N_3}^{(p_{123})})^{-1} = \phi_u(x) + \frac{2\pi}{N_u}, \quad (49)$$

$$(S_{N_1, N_2, N_3}^{(p_{123})}) \phi'_u(x) (S_{N_1, N_2, N_3}^{(p_{123})})^{-1}$$

$$= \phi'_u(x) - \epsilon^{uvw} Q \frac{2\pi}{N_v} \left( 2\phi_w(x) - \frac{[\phi_w(L) + \phi_w(0)]}{2} \right), \quad (50)$$

where one can define a Type III symmetry charge  $Q \equiv p_{123} \frac{N_1 N_2 N_3}{2\pi N_{123}}$ . Here the 1D edge is on a compact circle with the length  $L$ , here  $\phi_w(L)$  are  $\phi_w(0)$  taking value at the position  $x = 0$  (also  $x = L$ ). [In the case of infinite 1D line, we shall replace  $\phi_w(L)$  by  $\phi_w(\infty)$  and replace  $\phi_w(0)$  by  $\phi_w(-\infty)$ .] But,  $\phi_w(L)$  may differ from  $\phi_w(0)$  by  $2\pi n$  with some number  $n$  if there is a nontrivial winding, i.e.,

$$\phi_w(L) = \phi_w(0) + 2\pi n = 2\pi \frac{n_w}{N_w} + 2\pi n, \quad (51)$$

where we apply the fact that  $\phi_w(0)$  is a  $Z_{N_w}$  rotor angle. So Eq. (50) effectively results in a shift  $+\epsilon^{uvw} p_{123} \frac{N_u}{N_{123}} (2\pi n_w + \pi N_w n)$  and a rotation  $\epsilon^{uvw} Q \frac{2\pi}{N_v} [2\phi_w(x)]$ . Since  $\frac{N_u}{N_{123}}$  is necessarily an integer, the symmetry transformation  $(S_{N_1, N_2, N_3}^{(p_{123})}) \phi'_u(x) (S_{N_1, N_2, N_3}^{(p_{123})})^{-1}$  will shift by a  $2\pi$  multiple if  $p_{123} \frac{N_u}{N_{123}} N_w n$  is an even integer.

By realizing the field theory symmetry transformation, we have obtained all classes of SPT edge field theory within the group cohomology  $\mathcal{H}^3(Z_{N_1} \times Z_{N_2} \times Z_{N_3}, U(1))$  with  $p_u \in \mathbb{Z}_{N_u}, p_{uv} \in \mathbb{Z}_{N_{uv}}, p_{123} \in \mathbb{Z}_{N_{123}}$ .

#### IV. TYPE II BOSONIC ANOMALY: FRACTIONAL QUANTUM NUMBERS TRAPPED AT THE DOMAIN WALLS

We now apply the tools we develop in Sec. III to capture physical observables for these SPTs. We propose the experimental/numerical signatures for certain SPT with Type II class  $p_{12} \neq 0$  with (at least) two symmetry groups  $Z_{N_1} \times Z_{N_2}$ , also as a way to study the physical measurements for Type II bosonic anomaly. We show that when the  $Z_{N_1}$  symmetry is broken by  $Z_{N_1}$  domain wall created on a ring, there will be some fractionalized  $Z_{N_2}$  charges induced near the kink. We will demonstrate our field theory method can easily capture this effect.

##### A. Field theory approach: Fractional $Z_N$ charge trapped at the kink of $Z_N$ symmetry-breaking domain walls

Consider the  $Z_{N_1}$  domain wall is created on a ring (the  $Z_{N_1}$  symmetry is broken), then the  $Z_{N_1}$  domain wall can be captured by  $\phi_1(x)$  for  $x \in [0, L)$  takes some constant value  $\phi_0$  while  $\phi_1(L)$  shifted by  $2\pi \frac{n_1}{N_1}$  away from  $\phi_0$ . This means that  $\phi_1(x)$  has the fractional winding number

$$\int_0^L dx \partial_x \phi_1 = \phi_1(L) - \phi_1(0) = 2\pi \frac{n_1}{N_1}. \quad (52)$$




FIG. 4. We expect some fractional charge trapped near a single kink around  $x = 0$  (i.e.,  $x = 0 + \epsilon$ ) and  $x = L$  (i.e.,  $x = 0 - \epsilon$ ) in the domain walls. For  $Z_{N_1}$ -symmetry-breaking domain wall with a kink jump  $\Delta\phi_1 = 2\pi \frac{n_{12}}{N_{12}}$ , we predict that the fractionalized  $(n_{12}/N_{12})p_{21}$  units of  $Z_{N_2}$  charge are induced.

Also, recall Eq. (42) that the Type II  $p_{21} \neq 0$  (and  $p_1 = 0, p_2 = 0$ )  $Z_{N_2}$  symmetry transformation

$$S_{N_2}^{(p_2, p_{21})} = \exp \left[ \frac{i}{N_2} \left( p_{21} \int_0^L dx \partial_x \tilde{\phi}_1 + \int_0^L dx \partial_x \phi'_2 \right) \right] \quad (53)$$

can measure the induced  $Z_{N_2}$  charge on a state  $|\Psi_{\text{domain}}\rangle$  with this domain wall feature as

$$\begin{aligned} S_{N_2}^{(p_2, p_{21})} |\Psi_{\text{domain}}\rangle &= \exp \left[ \frac{i p_{21}}{N_2} [\tilde{\phi}_1(L) - \tilde{\phi}_1(0)] \right] |\Psi_{\text{domain}}\rangle \\ &= \exp \left[ \left( 2\pi i \frac{n_{12} p_{21}}{N_{12} N_2} \right) \right] |\Psi_{\text{domain}}\rangle. \end{aligned} \quad (54)$$

We also adopt two facts: First,  $\int_0^L dx \partial_x \tilde{\phi}_1 = 2\pi \frac{n_{12}}{N_{12}}$  with some integer  $n_{12}$ , where the  $\tilde{\phi}_1$  is regularized in a unit of  $2\pi/N_{12}$ . Second, as  $Z_{N_2}$  symmetry is not broken, both  $\phi_2$  and  $\phi'_2$  have no domain walls, then the above evaluation takes into account that  $\int_0^L dx \partial_x \phi'_2 = 0$ . This implies that induced charge is fractionalized  $(n_{12}/N_{12})p_{21}$  (recall  $p_{12}, p_{21} \in \mathbb{Z}_{N_{12}}$ )  $Z_{N_2}$  charge. This is the fractional charge trapped at the configuration of a single kink in Fig. 4.

On the other hand, one can imagine a series of  $N_{12}$  number of  $Z_{N_1}$ -symmetry-breaking domain wall each breaks to different vacuum expectation value (v.e.v.) where the domain wall in the region  $[0, x_1], [x_1, x_2], \dots, [x_{N_{12}-1}, x_{N_{12}} = L]$  with their symmetry-breaking  $\phi_1$  value at  $0, 2\pi \frac{1}{N_{12}}, 2\pi \frac{2}{N_{12}}, \dots, 2\pi \frac{N_{12}-1}{N_{12}}$ . This means a nontrivial winding number, like a soliton effect (see Fig. 5),  $\int_0^L dx \partial_x \tilde{\phi}_1 = 2\pi$  and  $S_{N_2}^{(p_2, p_{21})} |\Psi_{\text{domain wall}}\rangle = \exp[2\pi i \frac{p_{21}}{N_2}] |\Psi_{\text{domain wall}}\rangle$  capturing  $p_{21}$  integer units of  $Z_{N_2}$  charge at  $N_{12}$  kinks for totally  $N_{12}$  domain walls, in the configuration of Fig. 5. In average, each kink captures the  $p_{21}/N_{12}$  fractional units of  $Z_{N_2}$  charge.




FIG. 5. A nontrivial winding  $\int_0^L dx \partial_x \phi(x) = 2\pi$ . This is like a soliton (or particle) insertion. For  $N_{12}$  number of  $Z_{N_1}$ -symmetry-breaking domain walls, we predict that the integer  $p_{21}$  units of total induced  $Z_{N_2}$  charge on a 1D ring. On average, each kink captures a  $p_{21}/N_{12}$  fractional units of  $Z_{N_2}$  charge.

Similarly, we can consider the  $Z_{N_2}$  domain wall is created on a ring (the  $Z_{N_2}$  symmetry is broken), then the  $Z_{N_2}$  domain wall can be captured by  $\phi_2(x)$  soliton profile for  $x \in [0, L]$ . We consider a series of  $N_{12}$  number of  $Z_{N_2}$ -symmetry-breaking domain walls, each breaks to different v.e.v. (with an overall profile of  $\int_0^L dx \partial_x \tilde{\phi}_2 = 2\pi$ ). By  $S_{N_1}^{(p_1, p_{12})} |\Psi_{\text{domain wall}}\rangle = \exp[2\pi i \frac{p_{12}}{N_1}] |\Psi_{\text{domain wall}}\rangle$ , the  $N_{12}$  kinks of domain wall captures  $p_{12}$  integer units of  $Z_{N_1}$  charge for totally  $N_{12}$  domain wall, as in Fig. 5. On average, each domain wall captures  $p_{12}/N_{12}$  fractional units of  $Z_{N_1}$  charge.

## B. Goldstone-Wilczek formula and fractional quantum number

It is interesting to view our result above in light of the Goldstone-Wilczek (GW) approach [43]. We warm up by computing  $\frac{1}{2}$ -fermion charge found by Jackiw-Rebbi [51] using the GW method. We will then do a more general case for SPT. The construction, valid for 1D systems, works as follows.

*Jackiw-Rebbi model.* Consider a Lagrangian describing spinless fermions  $\psi(x)$  coupled to a classical background profile  $\lambda(x)$  via a term  $\lambda \psi^\dagger \sigma_3 \psi$ . In the high temperature phase, the v.e.v. of  $\lambda$  is zero and no mass is generated for the fermions. In the low temperature phase, the  $\lambda$  acquires two degenerate vacuum values  $\pm(\lambda)$  that are related by a  $Z_2$  symmetry. Generically we have

$$\langle \lambda \rangle \cos[\phi(x) - \theta(x)], \quad (55)$$

where we use the bosonization dictionary  $\psi^\dagger \sigma_3 \psi \rightarrow \cos[\phi(x)]$  and a phase change  $\Delta\theta = \pi$  captures the existence of a domain wall separating regions with opposite values of the v.e.v. of  $\lambda$ . From the fact that the fermion density  $\rho(x) = \psi^\dagger(x)\psi(x) = \frac{1}{2\pi} \partial_x \phi(x)$  (and the current  $J^\mu = \psi^\dagger \gamma^\mu \psi = \frac{1}{2\pi} \epsilon^{\mu\nu} \partial_\nu \phi$ ), it follows that the induced charge  $Q_{\text{dw}}$  on the kink by a domain wall is

$$Q_{\text{dw}} = \int_{x_0-\epsilon}^{x_0+\epsilon} dx \rho(x) = \int_{x_0-\epsilon}^{x_0+\epsilon} dx \frac{1}{2\pi} \partial_x \phi(x) = \frac{1}{2}, \quad (56)$$

where  $x_0$  denotes the center of the domain wall.

*Type II bosonic anomalies.* We now consider the case where the  $Z_{N_1}$  symmetry is spontaneously broken into different “vacuum” regions. This can be captured by an effective term in the Hamiltonian of the form

$$H_{sb} = -\lambda \cos[\phi_1(x) - \theta(x)], \quad \lambda > 0 \quad (57)$$

and the ground state is obtained, in the large  $\lambda$  limit, by phase locking  $\phi_1 = \theta$ , which opens a gap in the spectrum.

Different domain wall regions are described by different choices of the profile  $\theta(x)$ , as discussed in Sec. IV A. In particular, if we have  $\theta(x) = \theta_k(x)$  and  $\theta_k(x) = (k-1)2\pi/N_{12}$ , for  $x \in [(k-1)L/N_{12}, kL/N_{12}]$ ,  $k = 1, \dots, N_{12}$ . Then, we see that a domain wall separating regions  $k$  and  $k+1$  (where the phase difference is  $2\pi/N_{12}$ ) induces a  $Z_{N_2}$  charge given by

$$\begin{aligned} \delta Q_{k, k+1} &= \int_{kL/N_{12}-\epsilon}^{kL/N_{12}+\epsilon} dx \delta \rho_2(x) \\ &= \frac{1}{2\pi} \int_{kL/N_{12}-\epsilon}^{kL/N_{12}+\epsilon} dx \frac{p_{12}}{N_2} \partial_x \phi_1 = \frac{p_{12}}{N_2 N_{12}}. \end{aligned} \quad (58)$$




FIG. 6. (Color online) A profile of several domain walls, each with kinks and antikinks (in blue color). For  $Z_{N_1}$ -symmetry-breaking domain wall, each single kink can trap fractionalized  $Z_{N_2}$  charge. However, overall there is no nontrivial winding  $\int_0^L dx \partial_x \phi_1(x) = 0$  (i.e., no net soliton insertion), so there is no net induced charge on the whole 1D ring.

This implies a fractional of  $p_{12}/N_{12}$  induced  $Z_{N_2}$  charge on a single kink of  $Z_{N_1}$ -symmetry-breaking domain walls, consistent with Eq. (54).

Some remarks follow: If the system is placed on a ring, note the following: (i) First, with net soliton (or particle) insertions, then the total charge induced is nonzero (see Fig. 5). (ii) Second, without net soliton (or particle) insertions, then the total charge induced is obviously zero, as domain walls necessarily come in pairs with opposite charges on the kink and the antikink (see Fig. 6). (iii) One can also capture this bosonic anomaly in the fermionized language using the one-loop diagram under soliton background [43] shown in Fig. 7. (iv) A related phenomena has also been examined recently where fractionalized boundary excitations cause that the symmetry-broken boundary cannot be proliferated to restore the symmetry [52].

### C. Lattice approach: Projective phase observed at domain walls

Now we would like to formulate a fully regularized lattice approach to derive the induced fractional charge, and compare to the complementary field theory done in Sec. IV A and Goldstone-Wilczek approach in Sec. IV B. Below our notation follows Sec. III. Recall that in the case of a system with onsite symmetry, such as  $Z_N$  rotor model on a 1D ring with a simple Hamiltonian of  $\sum_j (\sigma_j + \sigma_j^\dagger)$ , there is an onsite symmetry transformation  $S = \prod_j \tau_j$  acting on the full ring. We can simply take a segment (from the site  $r_1$  to  $r_2$ )




FIG. 7. In the fermionized language, one can capture the anomaly effect on induced (fractional) charge/current under soliton background by the one-loop diagram [43]. The solid line represents fermions, the wavy line represents the external (gauge) field coupling to the induced current  $J^\mu$  (or charge  $J^0$ ), and the dashed line represents the scalar soliton (domain walls) background. Here in Sec. IV B, instead of fermionizing the theory, we directly address in the bosonized language to capture the bosonic anomaly.

of the symmetry transformation defined as a  $D$  operator  $D(r_1, r_2) \equiv \prod_{j=r_1}^{r_2} \tau_j$ . The  $D$  operator does the job to flip the measurement on  $\langle \sigma_\ell \rangle$ . What we mean is that  $\langle \psi | \sigma_\ell | \psi \rangle$  and  $\langle \psi' | \sigma_\ell | \psi' \rangle \equiv \langle \psi | D^\dagger \sigma_\ell D | \psi \rangle = e^{i2\pi/N} \langle \psi | \sigma_\ell | \psi \rangle$  are distinct by a phase  $e^{i2\pi/N}$  as long as  $\ell \in [r_1, r_2]$ . Thus, the  $D$  operator creates domain wall profile.

For our case of SPT edge modes with non-onsite symmetry studied here, we are ready to generalize the above and take a line segment of non-onsite symmetry transformation with symmetry  $Z_{N_u}$  (from the site  $r_1$  to  $r_2$ ) and define it as a  $D_{N_u}$  operator  $D_{N_u}(r_1, r_2) \equiv \prod_{j=r_1}^{r_2} \tau_j^{(u)} \prod_{j=r_1}^{r_2} U_{j,j+1} W_{j,j+1}^{III}$  [from the expression of  $S_{N_u}$ , with the onsite piece  $\tau_j^{(u)}$  and the non-onsite piece  $U_{j,j+1}$  in Eq. (28) and  $W_{j,j+1}$  in Eq. (32)]. This  $D$  operator effectively creates domain wall on the state with a kink (at the  $r_1$ ) and antikink (at the  $r_2$ ) feature, such as in Fig. 6. The total net charge on this type of domain wall (with equal numbers of kink and antikinks) is zero, due to no net soliton insertion (i.e., no net winding, so  $\int_0^L \partial_x \phi dx = 0$ ). However, by well separating kinks and antikinks, we can still compute the phase gained at each single kink [5]. We consider the induced charge measurement by  $S(D|\psi)$ , which is  $(SDS^\dagger)S|\psi\rangle = e^{i(\Theta_0 + \Theta)}D|\psi\rangle$ , where  $\Theta_0$  is from the initial charge (i.e.,  $S|\psi\rangle \equiv e^{i\Theta_0}|\psi\rangle$ ) and  $\Theta$  is from the charge gained on the kink. The measurement of symmetry  $S$  producing a phase  $e^{i\Theta}$  implies a nontrivial induced charge trapped at the kink of domain walls. We compute the phase at the left kink on a domain wall for all Type I, II, and III SPT classes, and summarize them in Table III.

In Table III, although we obtain  $e^{i\Theta_L}$  for each type, but there are some words of caution for interpreting it.

(i) For Type I class, with the  $Z_{N_1}$ -symmetry-breaking domain wall, there is no notion of induced  $Z_{N_1}$  charge since there is no  $Z_{N_1}$  symmetry (already broken) to respect.

(ii)  $(D_N^{(p)})^n$  captures  $n$  units of  $Z_N$ -symmetry-breaking domain wall. The calculation  $S_N^{(p)}(D_N^{(p)})^n S_N^{(p)\dagger}$  renders a  $e^{i\Theta_L}$  phase for the left kink and a  $e^{i\Theta_R} = e^{-i\Theta_L}$  phase for the right antikink. Our formalism is analogous to Ref. [52], where we choose the domain operator as a segment of symmetry transformation. For Type II class, if we have operators  $(D_{N_1}^{(p_{12})})^0$  acting on the interval  $[0, x_1)$ , while  $(D_{N_1}^{(p_{12})})^1$  acting on the interval  $[x_1, x_2)$ , ..., and  $(D_{N_1}^{(p_{12})})^{N_{12}}$  acting on the interval  $[x_{N_{12}-1}, x_{N_{12}} = L)$ , then we create the domain wall profile shown in Fig. 6. It is easy to see that due to charge cancellation on each kink/antikink, the  $S_{N_2}^{(p_{12})}(D_{N_1}^{(p_{12})})^{N_{12}} S_{N_2}^{(p_{12})\dagger}$  measurement on a left kink captures the same amount of charge trapped by a nontrivial soliton:  $\int_0^L dx \partial_x \phi_u = 2\pi$  [5].

(iii) For Type II class, we consider  $Z_{N_1}$ -symmetry-breaking domain wall (broken to a unit of  $\Delta\phi_1 = 2\pi/N_{12}$ ), and find that there is induced  $Z_{N_2}$  charge with a unit of  $\mathbf{p}_{12}/\mathbf{N}_{12}$ , consistent with field theory approach in Eqs. (54) and (58). For a total winding is  $\int_0^L dx \partial_x \phi_1 = 2\pi$ , there is also a nontrivial induced  $\mathbf{p}_{12}$  units of  $Z_{N_2}$  charge. Suppose a soliton generates this winding number 1 domain wall profile, even if  $p_{12} = N_{12}$  is identified as the trivial class as  $p_{12} = 0$ , we can observe  $\mathbf{N}_{12}$  units of  $Z_{N_2}$  charge, which is in general still not  $N_2$  units of  $Z_{N_2}$  charge. These phenomena have no analogs in Type I, and can be traced back to the discussion in Sec. III C.



(iv) For Type III class, with a  $Z_{N_1}$ -symmetry-breaking domain wall: On one hand, the  $\Theta_L$  phase written in terms of  $Z_{N_2}$  or  $Z_{N_3}$  charge unit is nonfractionalized but integer. On the other hand, we will find in Sec. VB that the  $Z_{N_2}$ ,  $Z_{N_3}$  symmetry transformation surprisingly no longer commutes. So, there is no proper notion of induced  $Z_{N_2}$ ,  $Z_{N_3}$  charge at all in the Type III class.

### V. TYPE III BOSONIC ANOMALY: DEGENERATE ZERO ENERGY MODES (PROJECTIVE REPRESENTATION)

We apply the tools we develop in Secs. II and III to study the physical measurements for Type III bosonic anomaly.

#### A. Field theory approach: Degenerate zero energy modes trapped at the kink of $Z_N$ symmetry-breaking domain walls

We propose the experimental/numerical signature for certain SPT with Type III symmetric class  $p_{123} \neq 0$  under the case of (at least) three symmetry group  $Z_{N_1} \times Z_{N_2} \times Z_{N_3}$ . Under the presence of a  $Z_{N_1}$  symmetry-breaking domain wall (without losing generality, we can also assume it to be any  $Z_{N_u}$ ), we can detect that the remained unbroken symmetry  $Z_{N_2} \times Z_{N_3}$  carries projective representation. More precisely, under the  $Z_{N_1}$  domain-wall profile,

$$\int_0^L dx \partial_x \phi_1 = \phi_1(L) - \phi_1(0) = 2\pi \frac{n_1}{N_1}, \quad (59)$$

we compute the commutator between two unbroken symmetry operators (47):

$$S_{N_2}^{(p_{231})} S_{N_3}^{(p_{312})} = S_{N_3}^{(p_{312})} S_{N_2}^{(p_{231})} e^{i \frac{2\pi n_1}{N_{123}} p_{123}}, \quad (60)$$

$$[\ln S_{N_2}^{(p_{231})}, \ln S_{N_3}^{(p_{312})}] = i \frac{2\pi n_1}{N_{123}} p_{123}, \quad (61)$$

where we identify the index  $(p_{231} + p_{312}) \rightarrow p_{123}$  as the same one. This noncommutative relation (60) indicates that  $S_{N_2}^{(p_{231})}$  and  $S_{N_3}^{(p_{312})}$  are not in a linear representation, but in a projective representation of  $Z_{N_2}$ ,  $Z_{N_3}$  symmetry. This is analogous to the commutator  $[T_x, T_y]$  of magnetic translations  $T_x, T_y$  along the  $x, y$  direction on a  $\mathbb{T}^2$  torus for a filling fraction  $1/k$  fractional quantum Hall state [described by  $U(1)_k$  level- $k$  Chern-Simons theory] [53]:

$$e^{iT_x} e^{iT_y} = e^{iT_y} e^{iT_x} e^{i2\pi/k}, \quad (62)$$

$$[T_x, T_y] = -i2\pi/k, \quad (63)$$

where one studies its ground states on a  $\mathbb{T}^2$  torus with a compactified  $x$  and  $y$  direction gives  $k$ -fold degeneracy. The  $k$  degenerate ground states are  $|\psi_m\rangle$  with  $m = 0, 1, \dots, k-1$ , while  $|\psi_m\rangle = |\psi_{m+k}\rangle$ . The ground states are chosen to satisfy  $e^{iT_x} |\psi_m\rangle = e^{i\frac{2\pi m}{k}} |\psi_m\rangle$ ,  $e^{iT_y} |\psi_m\rangle = |\psi_{m+1}\rangle$ . Similarly, for Eq. (60) we have a  $\mathbb{T}^2$  torus compactified in  $\phi_2$  and  $\phi_3$  directions, such that the following: (i) There is a  $N_{123}$ -fold degeneracy for zero energy modes at the domain wall. We can count the degeneracy by constructing the orthogonal ground states: consider the eigenstate  $|\psi_m\rangle$  of unitary operator  $S_{N_2}^{(p_{231})}$ , it implies that  $S_{N_2}^{(p_{231})} |\psi_m\rangle = e^{i\frac{2\pi n_1}{N_{123}} p_{123} m} |\psi_m\rangle$ .  $S_{N_3}^{(p_{312})} |\psi_m\rangle = |\psi_{m+1}\rangle$ . As long as  $\gcd(n_1 p_{123}, N_{123}) = 1$ , we




FIG. 8. (Color online) (a) The induced 2-cocycle from a 2+1D  $M^3 = M^2 \times I^1$  topology with a symmetry-preserving  $Z_{N_u}$  flux  $A$  insertion. (b) Here  $M^2 = S^1 \times I^1$  is a 2D spatial cylinder, composed by  $A$  and  $B$ , with another extra time dimension  $I^1$ . Along the  $B$  line we insert a monodromy defect of  $Z_{N_1}$ , such that  $A$  has a nontrivial group element value  $A = g_1' g_1^{-1} = g_2' g_2^{-1} = g_3' g_3^{-1} \in Z_{N_1}$ . The induced 2-cocycle  $\beta_A(B, C)$  is a nontrivial element in  $\mathcal{H}^2(Z_{N_u} \times Z_{N_w}, U(1)) = \mathbb{Z}_{N_{vw}}$  (here  $u, v, w$  cyclic as  $e^{uvw} = 1$ ), thus which carries a projective representation. (c) A monodromy defect can be viewed as a branch cut induced by a  $\Phi_B$  flux insertion (both modifying the Hamiltonians). (d) This means that when we do dimensional reduction on the compact ring  $S^1$  and view the reduced system as a 1D line segment, there are  $N_{123}$  degenerate zero energy modes (due to the nontrivial projective representation).

have  $N_{123}$ -fold degeneracy of  $|\psi_m\rangle$  with  $m = 0, \dots, N_{123} - 1$ . (ii) Equation (60) means the symmetry is realized projectively for the trapped zero energy modes at the domain wall.

We observe these are the signatures of Type III bosonic anomaly. This Type III anomaly in principle can be also captured by the perspective of *decorated  $\mathbb{Z}_{N_1}$  domain walls* of Ref. [23] with projective  $\mathbb{Z}_{N_2} \times \mathbb{Z}_{N_3}$  symmetry.

#### B. Cocycle approach: Degenerate zero energy modes from $Z_N$ -symmetry-preserving monodromy defect (branch cut) dimensional reduction from 2D to 1D

In Sec. VB, we had shown the symmetry-breaking domain wall would induce degenerate zero energy modes for Type III SPT. In this section, we will further show that a symmetry-preserving  $Z_{N_1}$  flux insertion (or a monodromy defect or branch cut modifying the Hamiltonian as in Refs. [22,44]) can also have degenerate zero energy modes. This is the case (see Fig. 8) when we put the system on a 2D cylinder and dimensionally reduce it to a 1D line along the monodromy defect. In this case, there is no domain wall, and the  $Z_{N_1}$  symmetry is not broken (but only translational symmetry is broken near the monodromy defect/branch cut).

In the following discussion, we will directly use 3-cocycles  $\omega_3$  from cohomology group  $\mathcal{H}^3(G, U(1))$  to detect the Type III bosonic anomaly. For convenience we use the nonhomogeneous cocycles (the lattice gauge theory cocycles), although there is no difficulty to convert it to homogeneous cocycles (SPT cocycles). The definition of the lattice gauge theory  $n$ -cocycles is indeed related to SPT  $n$ -cocycles [6,44,54–56]:

$$\begin{aligned} \omega_n(A_1, A_2, \dots, A_n) &= v_n(A_1 A_2 \dots A_n, A_2 \dots A_n, \dots, A_n, 1) \\ &= v_n(\tilde{A}_1, \tilde{A}_2, \dots, \tilde{A}_n, 1), \end{aligned} \quad (64)$$




FIG. 9. The triangulation of a  $M^3 = M^2 \times I^1$  topology (here  $M^2$  is a spatial cylinder composed by the  $A$  and  $B$  directions, with a  $I^1$  time) into three tetrahedra with branched structures.

where  $\tilde{A}_j \equiv A_j A_{j+1} \dots A_n$ . For 3-cocycles

$$\begin{aligned} \omega_3(A, B, C) &= v_3(ABC, BC, C, 1) \\ &\Rightarrow \omega_3(g_{01}, g_{12}, g_{23}) = \omega_3(g_0 g_1^{-1}, g_1 g_2^{-1}, g_2 g_3^{-1}) \\ &= v_3(g_0 g_3^{-1}, g_1 g_3^{-1}, g_2 g_3^{-1}, 1) = v_3(g_0, g_1, g_2, g_3). \end{aligned} \quad (65)$$

Here  $A = g_{01}$ ,  $B = g_{12}$ ,  $C = g_{23}$ , with  $g_{ab} \equiv g_a g_b^{-1}$ . We use the fact that SPT  $n$ -cocycle  $v_n$  belongs to the  $G$ -module, such that for  $r$  are group elements of  $G$ , it obeys  $r v_n(r_0, r_1, \dots, r_{n-1}, 1) = v_n(r r_0, r r_1, \dots, r r_{n-1}, r)$  (here we consider only Abelian group  $G = \prod_i Z_{N_i}$ ). In our case, we do not have time-reversal symmetry, so group action  $g$  on the  $G$ -module is trivial.


In short, there is no obstacle so that we can simply use the lattice gauge theory 3-cocycle  $\omega(A, B, C)$  to study the SPT 3-cocycle  $v(ABC, BC, C, 1)$ . Our goal is to design a geometry of 3-manifold  $M^3 = M^2 \times I^1$  with  $M^2$  the 2D cylinder with flux insertion (or monodromy defect) and with the  $I^1$  time direction [see Fig. 8(a)] with a set of 3-cocycles as tetrahedra filling this geometry (Fig. 9). All we need to do is compute the 2+1D SPT path integral  $\mathbf{Z}_{\text{SPT}}$  (i.e., partition function) using 3-cocycles  $\omega_3$  [44]:

$$\mathbf{Z}_{\text{SPT}} = |G|^{-N_v} \sum_{\{g_v\}} \prod_i [\omega_3^{s_i}(\{g_{v_a} g_{v_b}^{-1}\})]. \quad (66)$$


Here  $|G|$  is the order of the symmetry group,  $N_v$  is the number of vertices,  $\omega_3$  is 3-cocycle, and  $s_i$  is the exponent 1 or  $-1$  (i.e.,  $\dagger$ ) depending on the orientation of each tetrahedron (3-simplex). The summing over group elements  $g_v$  on the vertex produces a symmetry-preserving ground state. We consider a specific  $M^3$ , a 3-complex of Fig. 8(a), which can be decomposed into tetrahedra (each as a 3-simplex) shown in Fig. 9. There the three-dimensional spacetime manifold is under triangulation (or cellularization) into three tetrahedra.

We now go back to remark that the 3-cocycle condition in Eq. (4) indeed means that the path integral  $\mathbf{Z}_{\text{SPT}}$  on the 3-sphere  $S^3$  (as the surface the 4-ball  $B^4$ ) will be trivial as 1. The 3-coboundary condition in Eq. (5) means to identify the same topological terms (i.e., 3-cocycle) up to total derivative terms. There is a specific way (called the *branching structure*) to determine the orientation of tetrahedron, thus to determine the sign of  $s$  for 3-cocycles  $\omega_3^s$  by the determinant of volume  $s \equiv \det(\vec{v}_{32}, \vec{v}_{31}, \vec{v}_{30})$ . Two examples of the orientation

$s = +1, -1$  are



$$= \omega_3(g_0 g_1^{-1}, g_1 g_2^{-1}, g_2 g_3^{-1}) \quad (67)$$



$$= \omega_3^{-1}(g_0 g_1^{-1}, g_1 g_2^{-1}, g_2 g_3^{-1}). \quad (70)$$

Here we define the numeric ordering  $g_1' < g_2' < g_3' < g_4' < g_1 < g_2 < g_3 < g_4$ , and our arrows connect from the higher to lower ordering.

Now we can compute the induced 2-cocycle (the dimensional reduced 1+1D path integral) with a given inserted flux  $A$ , determined from three tetrahedra of 3-cocycles [see Fig. 9 and Eq. (71)].

$$\beta_A(B, C) \equiv \begin{array}{c} g_3 \\ \diagup \quad \diagdown \\ g_1 \quad g_2 \\ \diagdown \quad \diagup \\ g_1' \quad g_2' \end{array} \cdot \begin{array}{c} g_3 \\ \diagup \quad \diagdown \\ g_2 \quad g_2' \\ \diagdown \quad \diagup \\ g_1' \quad g_2' \end{array} \cdot \begin{array}{c} g_3 \\ \diagup \quad \diagdown \\ g_2' \quad g_3' \\ \diagdown \quad \diagup \\ g_1' \quad g_2' \end{array} \quad (71)$$

$$\begin{aligned} &= \frac{\omega(A, B, C)^{-1} \omega(ABA^{-1}, A, C)}{\omega(ABA^{-1}, ACA^{-1}, A)} = \frac{\omega(B, A, C)}{\omega(A, B, C) \omega(B, C, A)} \\ &= \frac{\omega(g_1 g_2^{-1}, g_1' g_1^{-1}, g_2 g_3^{-1})}{\omega(g_1' g_1^{-1}, g_1 g_2^{-1}, g_2 g_3^{-1}) \omega(g_1 g_2^{-1}, g_2 g_3^{-1}, g_1' g_1^{-1})}. \end{aligned} \quad (72)$$

We show that among the Type I, II, and III 3-cocycles discussed in Sec. II, only when  $\omega_3$  is the Type III 3-cocycle  $\omega_{\text{III}}$  [of Eq. (10)], this induced 2-cochain is nontrivial (i.e., a 2-cocycle but not a 2-coboundary). In that case,

$$\beta_A(B, C) = \exp \left[ i \frac{2\pi}{N_{123}} (b_1 a_2 c_3 - a_1 b_2 c_3 - b_1 c_2 a_3) \right]. \quad (73)$$

If we insert  $Z_{N_1}$  flux  $A = (a_1, 0, 0)$ , then we shall compare Eq. (73) with the nontrivial 2-cocycle  $\omega_2(B, C)$  in  $\mathcal{H}^2(Z_{N_2} \times Z_{N_3}, \text{U}(1)) = Z_{N_{23}}$ ,

$$\omega_2(B, C) = \exp \left[ i \frac{2\pi}{N_{23}} (b_2 c_3) \right]. \quad (74)$$

The  $\beta_A(B, C)$  is indeed nontrivial 2-cocycle as  $\omega_2(B, C)$  in the second cohomology group  $\mathcal{H}^2(Z_{N_2} \times Z_{N_3}, \text{U}(1))$ . In the following, we like to argue that Eq. (74) implies the projective representation of the symmetry group  $Z_{N_2} \times Z_{N_3}$ . Our argument is based on two facts. First, the dimensionally reduced SPTs in

terms of spacetime partition function (74) is a nontrivial 1+1D SPT [57]. We can physically understand it from the symmetry twist as a branch cut modifying the Hamiltonian [55,57] (see also Sec. VI). Second, from Ref. [6]'s Sec. VI, we know that the 1+1D SPT symmetry transformation  $\otimes_x U^x(g)$  along the 1D's  $x$  site is dictated by 2-cocycle. The onsite tensor  $S(g) \equiv \otimes_x U^x(g)$  acting on a chain of 1D SPT renders

$$S(g)|\alpha_L, \dots, \alpha_R\rangle = \frac{\omega_2(\alpha_L^{-1}g^{-1}, g)}{\omega_2(\alpha_R^{-1}g^{-1}, g)}|g\alpha_L, \dots, g\alpha_R\rangle, \quad (75)$$

where  $\alpha_L$  and  $\alpha_R$  are the two ends of the chain, with  $g, \alpha_L, \alpha_R, \dots \in G$  all in the symmetry group. We can derive that the effective degree of freedom on the 0D edge  $|\alpha_L\rangle$  forms a projective representation of symmetry, we find

$$\begin{aligned} S(B)S(C)|\alpha_L\rangle &= \frac{\omega_2(\alpha_L^{-1}C^{-1}B^{-1}, B)\omega_2(\alpha_L^{-1}C^{-1}, B)}{\omega_2(\alpha_L^{-1}C^{-1}B^{-1}, BC)}S(BC)|\alpha_L\rangle \\ &= \omega_2(B, C)S(BC)|\alpha_L\rangle. \end{aligned} \quad (76)$$

In the last line, we implement the 2-cocycle condition of  $\omega_2$ :  $\delta\omega_2(a, b, c) = \frac{\omega_2(b, c)\omega_2(a, bc)}{\omega_2(ab, c)\omega_2(a, b)} = 1$ . The projective representation of symmetry transformation  $S(B)S(C) = \omega_2(B, C)S(BC)$  is explicitly derived, and the projective phase is the 2-cocycle  $\omega_2(B, C)$  classified by  $\mathcal{H}^2(G, U(1))$ . Interestingly, the symmetry transformations on two ends together will form a linear representation, namely,  $S(B)S(C)|\alpha_L, \dots, \alpha_R\rangle = S(BC)|\alpha_L, \dots, \alpha_R\rangle$  [6].

The same argument holds when  $A$  is  $Z_{N_2}$  flux or  $Z_{N_3}$  flux. From Sec. V, the projective representation of symmetry implies the nontrivial ground state degeneracy if we view the system as a dimensionally reduced 1D line segment as in Fig. 8(d). From the  $N_{123}$  factor in Eq. (73), we conclude there is  $N_{123}$ -fold degenerated zero energy modes.

We should make two more remarks: (i) The precise 1+1D path integral is actually summing over  $g_v$  with a fixed flux  $A$  as  $\mathbf{Z}_{\text{SPT}} = |G|^{-N_v} \sum_{\{g_v\}; \text{fixed } A} \beta_A(B, C)$ , but overall our discussion above still holds. (ii) We have used 3-cocycle to construct a symmetry-preserving SPT ground state under  $Z_{N_1}$  flux insertion. We can see that indeed a  $Z_{N_1}$ -symmetry-breaking domain wall of Fig. 10 can be done in almost the same calculation, using 3-cocycles filling a 2+1D spacetime complex [Fig. 10(a)]. Although in Fig. 10(a), we need to fix the group elements  $g_1 = g_2$  on one side (in the time independent domain wall profile, we need to fix  $g_1 = g_2 = g_3$ ) and/or fix  $g'_1 = g'_2$  on the other side. Remarkably, we conclude that both the  $Z_{N_1}$ -symmetry-preserving flux insertion and  $Z_{N_1}$ -symmetry-breaking domain wall both provide  $N_{123}$ -fold degenerate ground states (from the nontrivial projective representation for the  $Z_{N_2}, Z_{N_3}$  symmetry). The symmetry-breaking case is consistent with Sec. V B.

## VI. TYPE I, II, AND III CLASS OBSERVABLES: FLUX INSERTION AND NONDYNAMICALLY ‘‘GAUGING’’ THE NON-ONSITE SYMMETRY

With the Type I, II, and III SPT lattice models built in Sec. III, in principle we can perform numerical simulations to measure their physical observables, such as (i) the energy




FIG. 10. (Color online) The  $Z_{N_1}$ -symmetry-breaking domain wall along the red  $\times$  mark and/or orange  $+$  mark, which induces  $N_{123}$ -fold degenerate zero energy modes. The situation is very similar to Fig. 8 (however, there was  $Z_{N_1}$ -symmetry-preserving flux insertion). We show that in both cases the induced 2-cochain from calculating path integral  $\mathbf{Z}_{\text{SPT}}$  renders a nontrivial 2-cocycle of  $\mathcal{H}^2(Z_{N_2} \times Z_{N_3}, U(1)) = \mathbb{Z}_{N_{23}}$ , thus carrying nontrivial projective representation of symmetry.

spectrum, (ii) the entanglement entropy, and (iii) the central charge of the edge modes. Those are the physical observables for the ‘‘untwisted sectors,’’ and we would like to further achieve more physical observables on the lattice, by applying the parallel discussion in Ref. [22], using  $Z_N$  gauge flux insertions through the 1D ring. The similar idea can be applied to detect SPTs numerically [45]. The gauge flux insertion on the SPT edge modes (lattice Hamiltonian) is like *gauging its non-onsite symmetry in a nondynamical way*. We emphasize *gauging in a nondynamical way* because the gauge flux is not a local degree of freedom on each site, but a global effect. The Hamiltonian affected by gauge flux insertions can be realized as the Hamiltonian with twisted boundary conditions (see an analogy made in Fig. 11). Another way to phrase the flux insertion is that it creates a monodromy defect [44] (or a branch cut) which modify both the bulk and the edge Hamiltonian. Namely, our flux insertion acts effectively as the *symmetry twist* [55,57] modifying the Hamiltonian. Here we outline the twisted boundary conditions on the Type I, II, and III SPT lattice models of Sec. III.

We first review the work done in Ref. [22] of Type I SPT class and then extend it to Type II and III classes. (We leave some tedious calculation to Appendix D.) We aim to build a lattice model with twisted boundary conditions to capture the edge modes physics in the presence of a unit of  $Z_N$  flux




FIG. 11. (Color online) (a) Thread a gauge flux  $\Phi_B$  through a 1D ring (the boundary of 2D SPT). (b) The gauge flux is effectively captured by a branch cut (the dashed line in the blue color). Twisted boundary condition is applied on the branch cut. The (a) and (b) are equivalent in the sense that both cases capture the equivalent physical observables, such as the energy spectrum.




FIG. 12. (Color online) The illustration of an effective 1D lattice model with  $M$  sites on a compact ring under a discrete  $Z_N$  flux insertion. Effectively the gauge flux insertion is captured by a branch cut located between the site  $M$  and the site 1. This results in a  $Z_N$  variable  $\omega$  insertion as a twist effect modifying the lattice Hamiltonian around the site  $M$  and the site 1.

insertion. Since the gauge flux effectively introduces a branch cut breaking the translational symmetry of  $T$  (as shown in Fig. 11), the gauged (or twisted) Hamiltonian, say  $\tilde{H}_N^{(p)}$ , is *not* invariant with respect to translational operator  $T$ , say  $[\tilde{H}_N^{(p)}, T] \neq 0$ . The challenge of constructing  $\tilde{H}_N^{(p)}$  is to first find a new (so-called *magnetic or twisted*) translation operator  $\tilde{T}^{(p)}$  incorporating the gauge flux effect at the branch cut, in Fig. 11(b), and in Fig. 12, say, the branch cut is between the sites  $M$  and the 1. We propose two principles to construct the twisted lattice model. The first general principle is that a string of  $M$  units of *twisted translation operator*  $\tilde{T}^{(p)}$  renders a *twisted symmetry transformation*  $\tilde{S}_N^{(p)}$  incorporating a  $Z_N$  unit flux:

$$\tilde{S}_N^{(p)} \equiv (\tilde{T}^{(p)})^M = \tilde{S}_N^{(p)} (U_{M,1}^{(N,p)} [\sigma_M^\dagger \sigma_1])^{-1} U_{M,1}^{(N,p)} [\omega \sigma_M^\dagger \sigma_1], \quad (77)$$

with the unitary operator  $(\tilde{T}^{(p)})$ , i.e.,  $(\tilde{T}^{(p)})^\dagger \tilde{T}^{(p)} = \mathbb{1}$ . We clarify that  $U_{M,1}^{(N,p)}$  is from Eq. (27), where  $U_{M,1}^{(N,p)}[\dots] \equiv U_{M,1}^{(N,p)} \circ [\dots]$  means  $U_{M,1}^{(N,p)}$  is a function of  $\dots$  variables. For example,  $U_{M,1}^{(N,p)}[\omega \sigma_M^\dagger \sigma_1]$  means that the variable  $\sigma_M^\dagger \sigma_1$  in Eq. (27) is replaced by  $\omega \sigma_M^\dagger \sigma_1$  with an extra  $\omega$  insertion. The second principle is that the twisted Hamiltonian is invariant in respect of the twisted translation operator, thus also invariant in respect of twisted symmetry transformation, i.e.,

$$[\tilde{H}_N^{(p)}, \tilde{T}^{(p)}] = 0, \quad [\tilde{H}_N^{(p)}, \tilde{S}_N^{(p)}] = 0. \quad (78)$$

We solve Eq. (77) by finding the twisted lattice translation operator

$$\tilde{T}^{(p)} = T [U_{M,1}^{(N,p)} (\sigma_M^\dagger \sigma_1)] \tau_1, \quad (79)$$

for each  $p \in \mathbb{Z}_N$  class. For the  $s$  units of  $Z_N$  flux, we have the generalization of  $\tilde{T}^{(p)}$  from a unit  $Z_N$  flux as

$$\tilde{T}^{(p)}|_s = T (U_{M,1}^{(N,p)} [\sigma_M^\dagger \sigma_1])^s \tau_1^s. \quad (80)$$

Indeed, there is no difficulty to extend this construction to Type II and III classes. For Type II SPT classes [with nonzero indices  $p_{12}$  and  $p_{21}$  of Eq. (28), while  $p_1 = p_2 = 0$ ] the non-onsite symmetry transformation can be reduced from NNN to NN coupling term  $U_{j,j+2}^{(N_1,p_{12})} \rightarrow U_{j,j+1}^{(N_1,p_{12})}$ , also from  $U_{j,j+2}^{(N_2,p_{21})} \rightarrow U_{j,j+1}^{(N_2,p_{21})}$ . The Type II twisted symmetry

transformation has exactly the same form as Eq. (77) except replacing the  $U$ . For Type III SPT classes, the Type III twisted symmetry transformation also has the same form as Eq. (77) except replacing the  $U$  to  $W$  in Eq. (32). The second principle in Eq. (78) also follows.

### Twisted Hamiltonian

The twisted Hamiltonian  $\tilde{H}_{N_1, N_2}^{(p_1, p_2, p_{12})}$  can be readily constructed from  $H_{N_1, N_2}^{(p_1, p_2, p_{12})}$  of Eq. (37), with the condition (78). (An explicit example for Type I SPT 1D lattice Hamiltonian with a gauge flux insertion has been derived in Ref. [22], which we shall not repeat here.)

Notice that the twisted nontrivial Hamiltonian breaks the SPT global symmetry [i.e., if  $p \neq 0 \pmod{N}$ , then  $[\tilde{H}_N^{(p)}, \mathcal{S}_N^{(p)}] \neq 0$ ], which can be regarded as the sign of  $Z_N$  anomaly [37]. On the other hand, in the trivial state  $p = 0$ , Eq. (77) yields  $\tilde{S}_N^{(p=0)} = \mathcal{S}_N^{(p=0)} = \prod_{j=1}^M \tau_j$ , where the twisted trivial Hamiltonian still *commutes* with the global  $Z_N$  onsite symmetry, and the twisted boundary effect is nothing but the usual toroidal boundary conditions [58]. (See also a discussion along the context of SPT and the orbifolds [59].)

The twisted Hamiltonian provides distinct low energy spectrum due to the gauge flux insertion (or the symmetry twist). The energy spectrum thus can be a physical observable to distinguish SPTs. Analytically we can use the field theoretic mode expansion for multiplet scalar chiral bosons  $\Phi_I(x) = \phi_{0I} + K_{IJ}^{-1} P_{\phi_j} \frac{2\pi}{L} x + i \sum_{n \neq 0} \frac{1}{n} \alpha_{I,n} e^{-inx \frac{2\pi}{L}}$ , with zero modes  $\phi_{0I}$  and winding modes  $P_{\phi_j}$  satisfying the commutator  $[\phi_{0I}, P_{\phi_j}] = i \delta_{IJ}$ . The Fourier modes satisfy a generalized Kac-Moody algebra:  $[\alpha_{I,n}, \alpha_{J,m}] = n K_{IJ}^{-1} \delta_{n,-m}$ . The low energy Hamiltonian, in terms of various quadratic mode expansions, becomes

$$H = \frac{(2\pi)^2}{4\pi L} \left[ V_{IJ} K_{IJ}^{-1} K_{JL}^{-1} P_{\phi_{I1}} P_{\phi_{L2}} + \sum_{n \neq 0} V_{IJ} \alpha_{I,n} \alpha_{J,-n} \right] + \dots \quad (81)$$

Following the procedure outlined in Ref. [22] with gauge flux (compared to the ungauged case in Ref. [23]), taking into account the twisted boundary conditions, we expect the conformal dimension of gapless edge modes of central charge  $c = 1$  free bosons labeled by the primary states  $|n_1, m_1, n_2, m_2\rangle$  (all parameters are integers) with the same compactification radius  $R$  for Type I and II SPTs (for simplicity, we assume  $N_1 = N_2 \equiv N$ ):

$$\begin{aligned} \tilde{\Delta}_N^{(p_1, p_2, p_{12})}(n_1, m_1, n_2, m_2; R) &= \frac{1}{R^2} \left( n_1 + \frac{p_1}{N} + \frac{p_{12}}{N} \right)^2 + \frac{R^2}{4} \left( m_1 + \frac{1}{N} \right)^2 \\ &+ \frac{1}{R^2} \left( n_2 + \frac{p_2}{N} + \frac{p_{21}}{N} \right)^2 + \frac{R^2}{4} \left( m_2 + \frac{1}{N} \right)^2, \end{aligned} \quad (82)$$

which is directly proportional to the energy of twisted Hamiltonian ( $p_{12}$  or  $p_{21}$  can be used interchangeably). The conformal dimension  $\tilde{\Delta}_N^{(p_1, p_2, p_{12})}(\mathcal{P}_u, \mathcal{P}_{uv})$  is intrinsically related to the SPT class labels:  $p_1, p_2, p_{12}$ , and is a function of momentum  $\mathcal{P}_u \equiv (n_u + \frac{p_u}{N} + \frac{p_{uv}}{N})(m_u + \frac{1}{N})$  and



$\mathcal{P}_{uv} \equiv (n_u + \frac{p_u}{N} + \frac{p_{uv}}{N})(m_v + \frac{1}{N})$ . Remarkably, for Type III SPTs, the nature of *noncommutative symmetry* generators will play the key rule, as if the gauged conformal field theory (CFT) and its corresponding gauged dynamical bulk theory has *non-Abelian* features, we will leave this survey for future works. The bottom line is that different classes of SPT’s CFT spectra respond to the flux insertion distinctly, thus we can in principle distinguish Type I, II, and III SPTs.

VII. CONCLUSION

Quantum anomalies have recently been emphasized to be intimately related to classifying and characterizing symmetry-protected topological states (SPTs) and topologically ordered states [37]. While fermionic anomalies are more familiar to the high energy particle physics communities (such as Adler-Bell-Jackiw anomaly [3,4], see Ref. [5]), the bosonic anomalies in our work are less discussed in the literature. For particle physicists, one may attempt to compute the anomaly through (i) a one-loop Feynman diagram of chiral fermions [3,4] or (ii) Fujikawa path integral method [42] by a Jacobi integral measure variation under the symmetry transformation. However, here, in our work, we instead seek another route, a fully bosonic language, to capture bosonic anomalies. We ask *what are the anomalous signals for these bosonic anomalies*. The result is summarized in Table I.

Since some recent papers also discuss the issues of anomalies in the context of SPTs or condensed matter setting [36,60–63,65–67] we shall stress the meaning of *quantum anomaly* more clearly. We shall also ask the following:

“How does the *bosonic anomaly* of our study relate to the context of the known quantum anomaly in the language of *high energy physics*?”

To answer this question, we have defined the following:

“The quantum anomaly is *the obstruction of a symmetry of a theory to be fully regularized for the full quantum theory as an onsite symmetry on the UV-cutoff lattice in the same spacetime dimension*.”

First, this understanding is consistent with the cases of Adler-Bell-Jackiw (ABJ) anomaly, where the symmetry of a classical action *cannot* be a symmetry of any regularization of the full quantum theory. For example, in chiral U(1) anomaly at quantum level, the axial U(1)<sub>A</sub> symmetry is in conflict with the vector U(1)<sub>V</sub> symmetry conservation [3,4,42].

Second, one can further ask, “how can we fully regularize the edge theory with bosonic anomalies on the same spacetime dimension (1+1D) if it has quantum anomalies?” The answer is that “because the (anomalous) symmetry is realized as a *non-onsite* symmetry instead of as an *onsite* symmetry, we can still realize the edge theory on the lattice *anomalously*.” Again, this agrees with our result and the known previous work [6,20–22,68] This regularization with *non-onsite* symmetry indeed is analogous to the Ginsparg-Wilson fermion approach [69] dealing with the fermion doubling problem for chiral fermions using *non-onsite* symmetry [68]. The *non-onsite symmetry* is an *anomalous symmetry*; thus that is why it is difficult to *gauge the non-onsite symmetry locally and dynamically* (see Ref. [68] for a connection between Ginsparg-Wilson fermions and SPTs).

Furthermore, another way to understand the anomaly is that one can regularize the quantum theory with onsite symmetry, if the regularization is done with an *extra dimensional bulk* [6] (thus not in the same spacetime dimension as the boundary). Again, this realization agrees with the quantum anomaly picture leaking quantum numbers through an extra dimensional bulk, shown in Fig. 2.

Let us now summarize the Type I, II, and III bosonic anomalies using the above understanding. To detect Type II bosonic SPTs, we find that the classic model studied by Jackiw-Rebbi [51] or Goldstone-Wilczek [43] offers a similar prototype observable. More precisely, the induced fractional quantum number is found in  $p_{12}$  class in  $G = Z_{N_1} \times Z_{N_2}$  symmetry. For Type II SPTs, the  $Z_{N_1}$ -symmetry-breaking domain wall will gap the edge and then induce a  $\frac{p_{12}}{N_{12}}$  fractional unit of  $Z_{N_2}$  charge (Fig. 4). The fermionized language shown in Fig. 7 can capture the one-loop effect analogous to ABJ anomaly’s one-loop calculation [3,4].

Type III SPTs’ bosonic anomaly provides different phenomena. The  $N_{123}$ -fold degenerate ground states are induced from either the symmetry-breaking domain wall on the 1D edges (Fig. 10) or the symmetry-preserving monodromy defect connecting edges through the bulk of a cylinder (which can be viewed as a dimensional-reduced 1D line system in Fig. 8). We show that the induced projective representation of symmetry under the above two circumstances implies the  $N_{123}$ -fold degenerate zero energy modes [70]. We shall stress that the Type III edge’s symmetry transformation provides a new kind of symmetry charge  $Q$  coupling as  $Q \int \epsilon^{uvw} \partial_x \phi_v(x) \phi_w(x) dx$  in the current term (47), which is rather distinct from the conventional symmetry charge  $q$  coupling as  $q \int \partial_x \phi_u(x) dx$ . While the work done in Refs. [24,26] cannot accommodate Type III class ( $p_{123} \neq 0$ ) SPTs, our approach with a new charge vector  $Q$  goes beyond previous work; thus we expect to obtain the new refined classification for the field theory also for other finite symmetry groups using Eq. (47) and its generalization.

For Type II and III SPT classes, we can characterize them by dimensional reduction to a lower dimensional boundary, and look for its induced quantum number or topological defects (similar effects happen in Majorana zero modes for free-fermion SPT cases [71]). For Type I class  $p_1 \in \mathbb{Z}_{N_1}$ , however, the physical observables we found so far are a bulk probe, instead of having a dimensional reduction to a lower dimensional system trapped with nontrivial quantum number. For Type I SPT probe, either the flux insertion goes through the bulk cylinder or the branch cut/monodromy defects connect from the edges to the bulk (Fig. 2). One can calculate the conformal dimension  $\tilde{\Delta}(\mathcal{P})$  (both analytically and numerically) as a function of momentum  $\mathcal{P}$  [72] in the twisted sector under monodromy defects, and one can show that each SPT class has distinct spectral shift [22].

Meanwhile, this type of probe such as flux insertion/monodromy defect which connects from the boundary to the bulk is essentially a signal of edge anomalous physics. In a sense, we develop an *effective 1D lattice Hamiltonian* with non-onsite symmetry which signals the existence of higher dimensional bulk, just like the edge chiral boson theory signals the bulk Chern-Simons theory. Only through a “nondynamically” gauge-flux insertion are we able to achieve *gauging the non-onsite symmetry effectively with a monodromy*

defect branch cut, shown in Figs. 10 and 2. This provides yet another way to interpret the edge anomaly: the 1D edge modified twisted Hamiltonian incorporating a branch cut does *not* preserve the original symmetry  $G$  (i.e.,  $[\tilde{H}_N^{(p)}, S_N^{(p)}] \neq 0$  in Sec. VI). However, one can readily check the full bulk-edge Hamiltonian description  $\tilde{H}_{N,\text{cylinder}}^{(p)}$  such as a cylinder with two edges in Fig. 2 will preserve the symmetry  $G$  (i.e.,  $[\tilde{H}_{N,\text{cylinder}}^{(p)}, S_N^{(p)}] = 0$ ).

We emphasize that, thanks to realizing the symmetry as a *non-onsite symmetry* on the lattice, all our SPT edge lattice constructions are successfully regularized on discrete space lattice with finite dimensional Hilbert space on the 1D ring. All our lattice models are ready for performing numerical simulations. For future directions, it will be interesting to numerically study physical observables to detect the distinct SPT classes, and also to study the charge transport with two edges on the cylinder *talking to each other* by quantum number pumping process in Fig. 2. This may require a full construction of the extra dimensional 2D bulk lattice, which can address what we mean by *quantum anomalies as some lower dimensional theory leaks certain quantum numbers to an extra dimensional bulk*.

#### ACKNOWLEDGMENTS

J.W. is grateful to F. Wilczek and Y. Nishida for inspiring discussions some years ago about Goldstone-Wilczek method, also to R. Jackiw for pointing out the first use of 3-cocycle in physics in Refs. [34,35]. J.W. thanks L.-Y. Hung for very helpful feedback on the manuscript. This research is supported by NSF Grants No. DMR-1005541, No. NSFC 11074140, and No. NSFC 11274192. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development and Innovation. The research is also supported by the BMO Financial Group and the John Templeton Foundation.

#### APPENDIX A: CHIRAL FERMIONIC ADLER-BELL-JACKIW ANOMALIES AND TOPOLOGICAL PHASES

In contrast to the *bosonic anomalies of discrete symmetries* studied in our main text, here we present a *chiral fermionic anomaly* (ABJ anomalies [3,4]) of a *continuous U(1) symmetry* realized in topological phases in condensed matter.

Specifically we consider a 1+1D U(1) quantum anomaly realization through 1D edge of U(1) quantum Hall state, such as in Fig. 13. We can formulate a Chern-Simons action  $S = \int (\frac{K}{4\pi} a \wedge da + \frac{q}{2\pi} A \wedge da)$  with an internal statistical gauge field  $a$  and an external U(1) electromagnetic gauge field  $A$ . Its 1+1D boundary is described by a (singlet or multiplet) chiral boson theory of a field  $\Phi$  ( $\Phi_L$  on the left edge,  $\Phi_R$  on the right edge). Here the field strength  $F = dA$  is equivalent to the external U(1) flux in the flux-insertion thought experiment threading through the cylinder (see a precise derivation in the Appendix of Ref. [22]). Without losing generality, let us first focus on the boundary with only one edge mode. We derive its




FIG. 13. (Color online) For topological phases, the anomalous current  $J_b$  of the boundary theory along the  $x$  direction leaks to  $J_y$  along the  $y$  direction in the extended bulk system.  $\Phi_B$ -flux insertion  $d\Phi_B/dt = -\oint E dL$  induces the electric  $E_x$  field along the  $x$  direction. The effective Hall effect dictates that  $J_y = \sigma_{xy} E_x = \sigma_{xy} \epsilon^{\mu\nu} \partial_\mu A_\nu$ , with the effective Hall conductance  $\sigma_{xy}$  probed by an external U(1) gauge field  $A$ .

equations of motion as

$$\partial_\mu j_b^\mu = \frac{\sigma_{xy}}{2} \epsilon^{\mu\nu} F_{\mu\nu} = \sigma_{xy} \epsilon^{\mu\nu} \partial_\mu A_\nu = J_y, \quad (\text{A1})$$

$$\partial_\mu j_L = \partial_\mu \left( \frac{q}{2\pi} \epsilon^{\mu\nu} \partial_\nu \Phi_L \right) = \partial_\mu (q \bar{\psi} \gamma^\mu P_L \psi) = +J_y, \quad (\text{A2})$$

$$\partial_\mu j_R = -\partial_\mu \left( \frac{q}{2\pi} \epsilon^{\mu\nu} \partial_\nu \Phi_R \right) = \partial_\mu (q \bar{\psi} \gamma^\mu P_R \psi) = -J_y. \quad (\text{A3})$$

We show the Hall conductance from its definition  $J_y = \sigma_{xy} E_x$  in Eq. (A1), as  $\sigma_{xy} = qK^{-1}q/(2\pi)$ .

Here  $j_b$  stands for the edge current. A left-moving current  $j_L = j_b$  is on one edge, and a right-moving current  $j_R = -j_b$  is on the other edge, shown in Fig. 13. By bosonization, we convert a compact bosonic phase  $\Phi$  to the fermion field  $\psi$ . The vector current is  $j_L + j_R \equiv j_V$ , and the U(1)<sub>V</sub> current is conserved. The axial current is  $j_L - j_R \equiv j_A$ , and we derive the famous ABJ U(1)<sub>A</sub> anomalous current in 1+1D (or Schwinger's 1+1D quantum electrodynamic [QED] anomaly [73]):

$$\partial_\mu j_V^\mu = \partial_\mu (j_L^\mu + j_R^\mu) = 0, \quad (\text{A4})$$

$$\partial_\mu j_A^\mu = \partial_\mu (j_L^\mu - j_R^\mu) = \sigma_{xy} \epsilon^{\mu\nu} F_{\mu\nu}. \quad (\text{A5})$$

This simple bulk-edge derivation is consistent with field theory one-loop calculation through Fig. 14. It shows that the combined boundary theory on the left and right edges [living on the edges of a 2+1D U(1) Chern-Simons theory] can be viewed as a 1+1D anomalous world of Schwinger's 1+1D QED [73]. This is an example of chiral fermionic anomaly of a continuous U(1) symmetry when  $K$  is an odd integer. [When




FIG. 14. In the fermionic language, the 1+1D chiral fermions (represented by the solid line) and the external U(1) gauge field (represented by the wavy curve) contribute to a one-loop Feynman diagram correction to the axial current  $j_A^\mu$ . This leads to the nonconservation of  $j_A^\mu$  as the anomalous current  $\partial_\mu j_A^\mu = \epsilon^{\mu\nu} (qK^{-1}q/2\pi) F_{\mu\nu}$ .

$K$  is an even integer, it becomes a chiral bosonic anomaly of a continuous  $U(1)$  symmetry.]

## APPENDIX B: MATRIX PRODUCT OPERATORS AND LATTICE REGULARIZATION

In this Appendix, we provide detailed calculations about the matrix product operators (MPO) formalism. Contracting three neighbored sites tensor  $T(g_a), T(g_b), T(g_c)$  of  $G$ -symmetry transformation  $S$  (with  $g \in G$ ) in different order will render a relative projective phase. Importantly, if this phase is a nontrivial 3-cocycle, then it readily verifies that our lattice construction maps to the nontrivial class of cohomology group. We also show the details of lattice regularizations in Sec. III.

We now formulate the unitary operator  $S_N^{(p)}$  as the MPO with the form

$$S_N^{(p)} = \sum_{\{j, j'\}} \text{tr} [T_{\alpha_1 \alpha_2}^{j_1 j'_1} T_{\alpha_2 \alpha_3}^{j_2 j'_2} \dots T_{\alpha_M \alpha_1}^{j_M j'_M}] |j'_1, \dots, j'_M\rangle \langle j_1, \dots, j_M|. \quad (\text{B1})$$

This is the operator formalism of matrix product states (MPS). Here *physical indices*  $j_1, j_2, \dots, j_M$  and  $j'_1, j'_2, \dots, j'_M$  are

labeled by input/output physical eigenvalues (here  $Z_N$  rotor angle), the subindices  $1, 2, \dots, M$  are the physical site indices. There are also *virtual indices*  $\alpha_1, \alpha_2, \dots, \alpha_M$  which are traced in the end. Summing over all the operation from  $\{j, j'\}$  indices, we shall reproduce the symmetry transformation operator  $S_N^{(p)}$ .

To find out the projective phase  $e^{i\theta(g_a, g_b, g_c)}$ , we use the facts of tensors  $T(g_a), T(g_b), T(g_c)$  acting on the same site with group elements  $g_a, g_b, g_c$ . There is a generic projective relation

$$T(g_a g_b) = P_{g_a, g_b}^\dagger T(g_a) T(g_b) P_{g_a, g_b}. \quad (\text{B2})$$

Here  $P_{g_a, g_b}$  is the projection operator. We contract three tensors in two different orders:

$$(P_{g_a, g_b} \otimes I_3) P_{g_a g_b, g_c} \simeq e^{i\theta(g_a, g_b, g_c)} (I_1 \otimes P_{g_b, g_c}) P_{g_a, g_b g_c}. \quad (\text{B3})$$

The left-hand side contracts the  $a, b$  first then with the  $c$ , while the right-hand side contracts the  $b, c$  first then with the  $a$ . Here  $\simeq$  means the equivalence is up to a projection out of unparallel states. If the projective phase  $e^{i\theta(g_a, g_b, g_c)}$  happens to be the nontrivial 3-cocycle in a cohomology group, then we reach our goal; this verifies that our SPT lattice constructions (thus also the low energy field theory) map to the nontrivial class of the cohomology group  $\mathcal{H}^3(G, U(1))$ . This is the emphasis of this Appendix.

### 1. Type I and II classes

We first write the  $\tilde{\sigma}_j^{(1)}, \tilde{\sigma}_j^{(2)}$  operators in the lattice regularization for Type II symmetry transformation in Sec. III B:

$$\tilde{\sigma}_j^{(u)} = \begin{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega_{uv} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \omega_{uv}^{\text{gcd}(N_u, N_v)-1} \end{pmatrix} & 0 & 0 & 0 \\ 0 & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega_{uv} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \omega_{uv}^{\text{gcd}(N_u, N_v)-1} \end{pmatrix} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega_{uv} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \omega_{uv}^{\text{gcd}(N_u, N_v)-1} \end{pmatrix} \end{pmatrix}_j \\ = \langle \phi_{u, j} | e^{i\tilde{\phi}_{u, j}} | \phi_{u, j} \rangle. \quad (\text{B4})$$

The  $\tilde{\sigma}_j^{(u)}$  matrix has  $N_u \times N_u$  components. It is block diagonalizable with  $\frac{N_u}{N_{12}}$  subblocks, and each subblock with  $N_{12} \times N_{12}$  components. We now verify that our symmetry transformations (28) [thus the lattice Hamiltonian (37)] correspond to nontrivial 3-cocycles in the third cohomology group in  $\mathcal{H}^3(Z_{N_1} \times Z_{N_2}, U(1)) = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{\text{gcd}(N_1, N_2)}$ . In this section we will focus on  $p_1, p_2$  Type I and  $p_{12}$  Type II class. The test below will verify that we indeed construct a lattice model of the nontrivial SPT of the  $p_{12}$  class with  $p_u \in \mathbb{Z}_{N_u}, p_{12} \in \mathbb{Z}_{\text{gcd}(N_1, N_2)}$ .

The tensor  $T(g)$  and the unitary operator  $S_{N_1}^{(p_1, p_{12})} S_{N_2}^{(p_2, p_{21})}$  as the matrix product operators (MPO) already appeared in the main text, we should not repeat it here.

To find out the projective phase  $e^{i\theta(g_a, g_b, g_c)}$  of three tensors  $T(g_a), T(g_b), T(g_c)$  acting on three neighbored sites, we follow the fact in Eq. (12), and derive the Type I and II projection operators

$$P_{N_1, N_2}^{(p)} \equiv P_{N_1, N_2, (m_a^{(1)}, m_b^{(1)}), (m_a^{(2)}, m_b^{(2)})}^{(p)} = \int d\phi'_{in} d\phi''_{in} \left( \left| \phi'_{in} + \frac{2\pi m_b^{(1)}}{N_1} \right\rangle \langle \phi'_{in} | \right) \left( \left| \phi''_{in} + \frac{2\pi m_b^{(2)}}{N_2} \right\rangle \langle \phi''_{in} | \right) \\ \times e^{ip_1 \phi'_{in} (lm_a^{(1)} + m_b^{(1)})_{N_1} - (m_a^{(1)} + m_b^{(1)})_{N_1}} e^{ip_2 \phi''_{in} (lm_a^{(2)} + m_b^{(2)})_{N_2} - (m_a^{(2)} + m_b^{(2)})_{N_2}} \\ \times e^{ip_{21} (\tilde{\phi}''_{in})_r (lm_a^{(2)} + m_b^{(2)})_{N_2} - (m_a^{(2)} + m_b^{(2)})_{N_2}} e^{ip_{12} (\tilde{\phi}'_{in})_r (lm_a^{(1)} + m_b^{(1)})_{N_1} - (m_a^{(1)} + m_b^{(1)})_{N_1}}, \quad (\text{B5})$$

where  $[m_a + m_b]_N$  with subindex  $N$  means taking the value module  $N$ .  $P_{g_1, g_2}$  inputs one state  $\langle \phi_{in}^{(1)} | \langle \phi_{in}^{(2)} |$  and outputs two states  $(|\phi_{in}^{(1)} + \frac{2\pi m_b^{(1)}}{N_1} \rangle | \phi_{in}^{(1)} \rangle)(|\phi_{in}^{(2)} + \frac{2\pi m_c^{(2)}}{N_2} \rangle | \phi_{in}^{(2)} \rangle)$ . To derive the projective phase  $e^{i\theta(g_a, g_b, g_c)}$ , we start by contracting  $T(g_b)$  and  $T(g_c)$  first, and then the combined tensor contracts with  $T(g_a)$  gives

$$\begin{aligned} (I_1 \otimes P_{g_b, g_c}) P_{g_a, g_b, g_c} &= \int d\phi_{in}^{\prime(1)} d\phi_{in}^{\prime(2)} \left( \left| \phi_{in}^{\prime(1)} + \frac{2\pi(m_b^{(1)} + m_c^{(1)})}{N_1} \right\rangle_a \left| \phi_{in}^{\prime(1)} + \frac{2\pi m_c^{(1)}}{N_1} \right\rangle_b \left| \phi_{in}^{\prime(1)} \right\rangle_c \left| \phi_{in}^{\prime(1)} \right\rangle_{abc} \right) \\ &\times \left( \left| \phi_{in}^{\prime(2)} + \frac{2\pi(m_b^{(2)} + m_c^{(2)})}{N_2} \right\rangle_a \left| \phi_{in}^{\prime(2)} + \frac{2\pi m_c^{(2)}}{N_2} \right\rangle_b \left| \phi_{in}^{\prime(2)} \right\rangle_c \left| \phi_{in}^{\prime(2)} \right\rangle_{abc} \right) \\ &\times e^{ip_1 \phi_{in}^{\prime(1)} (m_a^{(1)} + m_b^{(1)} + m_c^{(1)})_{N_1} - m_a^{(1)} - m_b^{(1)} - m_c^{(1)})/N_1} e^{ip_2 \phi_{in}^{\prime(2)} (m_a^{(2)} + m_b^{(2)} + m_c^{(2)})_{N_2} - m_a^{(2)} - m_b^{(2)} - m_c^{(2)})/N_2} \\ &\times e^{ip_{21} (\bar{\phi}_{in}^{\prime(1)})_r (m_a^{(2)} + m_b^{(2)} + m_c^{(2)})_{N_2} - m_a^{(2)} - m_b^{(2)} - m_c^{(2)})/N_2} e^{ip_{12} (\bar{\phi}_{in}^{\prime(2)})_r (m_a^{(1)} + m_b^{(1)} + m_c^{(1)})_{N_1} - m_a^{(1)} - m_b^{(1)} - m_c^{(1)})/N_1}, \end{aligned} \quad (B6)$$

which inputs one state  $\langle \phi_{in} |$  and outputs three states  $|\phi_{in} + \frac{2\pi}{N}(m_b + m_c)\rangle$ ,  $|\phi_{in} + \frac{2\pi}{N}m_c\rangle$ , and  $|\phi_{in}\rangle$ . Similarly we can derive  $(P_{g_a, g_b} \otimes I_3) P_{g_a, g_b, g_c}$  by contracting  $T(g_a)$  and  $T(g_b)$  first, and then the combined tensor contracts with  $T(g_c)$ . By computing Eq. (B3), with only  $p_1$  index (i.e., setting  $p_2 = p_{12} = 0$ ), we can derive Type I 3-cocycle

$$e^{i\theta(g_a, g_b, g_c)} = e^{ip_1 \left( \frac{2\pi m_c^{(1)}}{N_1} \right) \frac{(m_a^{(1)} + m_b^{(1)})_{N_2} - (m_a^{(1)} + m_b^{(1)})}{N_1}} = \omega_I^{(i)}(m_c, m_a, m_b). \quad (B7)$$

By computing Eq. (B3) with only  $p_{21}$  index (i.e., setting  $p_1 = p_2 = p_{12} = 0$ ), we can recover Type II 3-cocycle

$$e^{i\theta(g_a, g_b, g_c)} = e^{ip_{21} \left( \frac{2\pi m_c^{(1)}}{N_1} \right) ((m_a^{(2)} + m_b^{(2)})_{N_2} - (m_a^{(2)} + m_b^{(2)})) / N_2} = \omega_{II}^{(ij)}(m_c, m_a, m_b), \quad (B8)$$

up to the index redefinition  $p_{21} \rightarrow -p_{12}$ . We thus derive that the projective phase  $e^{i\theta(g_a, g_b, g_c)}$  from MPS tensors corresponds to the group cohomology approach [6]. From here we learn that the inserted  $p_{12}$  and  $p_{21}$  are indeed the same indices because  $e^{ip_{21} \left( \frac{2\pi m_c^{(1)}}{N_1} \right) ((m_a^{(2)} + m_b^{(2)})_{N_2} - (m_a^{(2)} + m_b^{(2)})) / N_2}$  and  $e^{ip_{12} \left( \frac{2\pi m_c^{(2)}}{N_2} \right) ((m_a^{(1)} + m_b^{(1)})_{N_1} - (m_a^{(1)} + m_b^{(1)})) / N_1}$  are equivalent 3-cocycles up to 3-coboundaries [46], meanwhile  $p_{12} = p_{12} \bmod \gcd(N_1, N_2)$ . This demonstrates that our lattice construction fulfills all  $\mathbb{Z}_{\gcd(N_1, N_2)}$  Type II classes of SPT with  $Z_{N_1} \times Z_{N_2}$  symmetry, and also Type I  $\mathbb{Z}_{N_1}, \mathbb{Z}_{N_2}$  classes as we desired.

## 2. Type III class

We first motivate our construction of matrix product operators by observing that Type III 3-cocycle in Eq. (10) inputs, for example,  $a_1 \in Z_{N_1}$ ,  $b_2 \in Z_{N_2}$ ,  $c_3 \in Z_{N_3}$  and outputs a U(1) phase. This implies that the  $Z_{N_1}$  symmetry transformation will affect the mixed  $Z_{N_2}, Z_{N_3}$  rotor models, while similarly  $Z_{N_2}, Z_{N_3}$  global symmetry will cause the same effect. This observation guides us to write the tensor  $T(g)$  and the symmetry transformation  $S_N^{(p)} = S_{N_1, N_2, N_3}^{(p_{123})}$  defined in Sec. III A. We propose the tensor  $T(g)$  and  $S_{N_1, N_2, N_3}^{(p_{123})}$  already in the main text, which we shall not repeat. Let us first understand how to regularize the symmetry operator on the lattice.

## 3. Lattice regularization

We derive the non-onsite symmetry transformation  $W_{j, j+1}^{III}$ , acting on the sites  $j$  and  $j+1$  as

$$W_{j, j+1}^{III} = \prod_{u, v, w \in \{1, 2, 3\}} \exp \left( i \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)} \epsilon^{uvw} \frac{p_{123}}{N_u} (\phi_{in}^{j+1, (v)} \phi_{in}^{j, (w)}) \right) \quad (B9)$$

$$= \prod_{(v, w) = (2, 3), (3, 1), (1, 2)} e^{ip_{123} ((\phi_{in}^{j+1, (v)} - \phi_{in}^{j, (v)}) \phi_{in}^{j, (w)} - (\phi_{in}^{j+1, (w)} - \phi_{in}^{j, (w)}) \phi_{in}^{j, (v)}) \frac{N_1 N_w}{2\pi \gcd(N_1, N_2, N_3)}} \quad (B10)$$

$$= \prod_{(v, w) = (2, 3), (3, 1), (1, 2)} \left( (\sigma_j^{(v)\dagger} \sigma_{j+1}^{(v)}) \phi_{in}^{j, (w)} \left( (\sigma_j^{(w)} \sigma_{j+1}^{(w)\dagger}) \phi_{in}^{j, (v)} \right) \right)^{p_{123} \frac{N_1 N_w}{2\pi \gcd(N_1, N_2, N_3)}} \quad (B11)$$

$$= \prod_{u, v, w \in \{1, 2, 3\}} (\sigma_j^{(v)\dagger} \sigma_{j+1}^{(v)}) e^{uvw p_{123} \frac{\ln(\sigma_j^{(w)}) N_v N_w}{2\pi i \gcd(N_1, N_2, N_3)}} \quad (B12)$$

$$\equiv W_{j, j+1; N_1}^{III} W_{j, j+1; N_2}^{III} W_{j, j+1; N_3}^{III}, \quad (B13)$$



where we separate  $Z_{N_1}$ ,  $Z_{N_2}$ ,  $Z_{N_3}$  non-onsite symmetry transformation to  $W_{j,j+1;N_1}^{\text{III}}$ ,  $W_{j,j+1;N_2}^{\text{III}}$ ,  $W_{j,j+1;N_3}^{\text{III}}$ , respectively. More explicitly, we have  $Z_{N_1}$  non-onsite symmetry transformation

$$W_{j,j+1;N_1}^{\text{III}} = e^{ip_{123}((\phi_{in}^{j+1,(2)} - \phi_{in}^{j,(2)})\phi_{in}^{j,(3)} - (\phi_{in}^{j+1,(3)} - \phi_{in}^{j,(3)})\phi_{in}^{j,(2)}) \frac{N_2 N_3}{2\pi \text{gcd}(N_1, N_2, N_3)}} \quad (\text{B14})$$

$$= ((\sigma_{2,j}^\dagger \sigma_{2,j+1})^{\ln(\sigma_{3,j})} [(\sigma_{3,j} \sigma_{3,j+1}^\dagger)^{\ln(\sigma_{2,j})})]^{p_{123} \frac{N_2 N_3}{2\pi i \text{gcd}(N_1, N_2, N_3)}}, \quad (\text{B15})$$

and  $W_{j,j+1;N_2}^{\text{III}}$ ,  $W_{j,j+1;N_3}^{\text{III}}$  have the analogous forms. We first attempt to regularize this  $W_{j,j+1}^{\text{III}}$  operator by defining

$$\phi_{in}^{j,(u)} \equiv i^{-1} \ln(\sigma_{u,j}) = i^{-1} \begin{pmatrix} \ln[1] & 0 & 0 & 0 \\ 0 & \ln[\omega_u] & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \ln[\omega_u^{N_u-1}] \end{pmatrix}_j = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{2\pi}{N_u} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \frac{2\pi(N_u-1)}{N_u} \end{pmatrix}_j, \quad (\text{B16})$$

here  $u \in \{1, 2, 3\}$ . The challenge of the lattice regularization is to understand what exactly does this operator  $(\sigma_{v,j}^\dagger \sigma_{v,j+1})^{p_{123} \frac{\ln(\sigma_{w,j}) N_v N_w}{2\pi i \text{gcd}(N_1, N_2, N_3)}}$  in Eq. (32) mean on the lattice. Without losing generality, let us take  $(\sigma_{v,j}^\dagger \sigma_{v,j+1})^{\ln(\sigma_{3,j})}$  in  $W_{j,j+1;N_1}^{\text{III}}$  of Eq. (B14) as an example. The answer to this question is that we should view how this operator acts on the combined  $Z_{N_2} \times Z_{N_3}$  states  $|\phi^{(2)}\rangle \otimes |\phi^{(3)}\rangle$ . The  $W_{j,j+1;N_1}^{\text{III}}$  operator is a  $[(N_2)^2 \times (N_3)^2] \times [(N_2)^2 \times (N_3)^2]$ -component matrix acting on the  $(N_2)^2 \times (N_3)^2$  dimensional Hilbert space spanned by the all  $|\phi_j^{(2)}\rangle \otimes |\phi_{j+1}^{(2)}\rangle \otimes |\phi_j^{(3)}\rangle \otimes |\phi_{j+1}^{(3)}\rangle$  states at the sites  $j$  and  $j+1$ . The key is regularizing this operator  $W_{j,j+1;N_1}^{\text{III}}$  explicitly, using Eq. (B16) as

$$(\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\frac{p_{123} \ln(\sigma_{3,j}) N_2 N_3}{2\pi i \text{gcd}(N_1, N_2, N_3)}} = \begin{pmatrix} (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\ln[1]} & 0 & 0 & 0 \\ 0 & (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\ln[\omega_3]} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\ln[\omega_3^{N_3-1}]} \end{pmatrix}_j^{\frac{p_{123} N_2 N_3}{2\pi i \text{gcd}(N_1, N_2, N_3)}} \quad (\text{B17})$$

$$= \begin{pmatrix} (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^0 & 0 & 0 & 0 \\ 0 & (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\frac{p_{123} N_2}{\text{gcd}(N_1, N_2, N_3)}} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & (\sigma_j^{(2)\dagger} \sigma_{j+1}^{(2)})^{\frac{p_{123} N_2 (N_3-1)}{\text{gcd}(N_1, N_2, N_3)}} \end{pmatrix}_j. \quad (\text{B18})$$

We emphasize that each subblock involving  $(\sigma_{2,j}^\dagger \sigma_{2,j+1})$  is a  $(N_2)^2 \times (N_2)^2$ -component matrix. (Here  $\sigma_{2,j+1}$  is a  $N_2 \times N_2$ -component matrix.) There are totally  $N_3 \times N_3$  subblocks. We recall that  $\sigma_2$  are operators defined in this manner in Eq. (23), i.e.,  $\sigma_2 \sim e^{i\phi^{(2)}}$ , with  $\phi^{(2)}$  a  $Z_{N_2}$  variable. Thus, the operator in each subblock has the form

$$(W_{j,j+1;N_1}^{\text{III}}) = ((\sigma_{2,j}^\dagger \sigma_{2,j+1})^{n_3 N_2} [(\sigma_{3,j} \sigma_{3,j+1}^\dagger)^{n_2 N_3}])^{\frac{p_{123}}{\text{gcd}(N_1, N_2, N_3)}}. \quad (\text{B19})$$

The notation  $n_u$  (above  $u = 2$  or  $3$ ) denotes an integer which corresponds to the  $Z_{N_u}$  values for  $|\phi^{(u)} = n_u(2\pi/N_u)\rangle$  state in different subblocks. First, we notice that  $p_{123}$  is identified by  $p_{123} = p_{123} \bmod \text{gcd}(N_1, N_2, N_3)$ . In addition, when  $p_{123}$  is a multiple of  $\text{gcd}(N_1, N_2, N_3)$ , we have  $(W_{j,j+1;N_1}^{\text{III}}) = 1$  (here 1 really means  $\mathbb{1}_{N_2 \times N_2, j} \otimes \mathbb{1}_{N_2 \times N_2, j+1} \otimes \mathbb{1}_{N_3 \times N_3, j} \otimes \mathbb{1}_{N_3 \times N_3, j+1}$ , the identity operator of  $Z_{N_2}$ ,  $Z_{N_3}$  states on sites  $j, j+1$ ). When  $p_{123}$  is not a multiple of  $\text{gcd}(N_1, N_2, N_3)$ , our lattice construction represents a nontrivial non-onsite symmetry transformation ( $W_{j,j+1}^{\text{III}} \neq 1$ ), thus producing a nontrivial SPT labeled by  $p_{123} \in \mathbb{Z}_{\text{gcd}(N_1, N_2, N_3)}$ . One may expect to fully regularize Eq. (B18), we need to solve a constraint  $(W_{j,j+1;N_1}^{\text{III}})^{N_1}$  analogous to Eqs. (26) and (30). But we do not have to: the exponent in Eq. (B18) is already an integer, e.g.,  $\frac{p_{123} N_2 N_3}{\text{gcd}(N_1, N_2, N_3)}$  is necessarily an integer. We note that, as we expected, when  $p_{123} = \text{gcd}(N_1, N_2, N_3)$ , we have  $(W_{j,j+1;N_1}^{\text{III}})^{N_1} = 1$ ; when  $p_{123} \neq \text{gcd}(N_1, N_2, N_3)$ , we have  $(W_{j,j+1;N_1}^{\text{III}})^{N_1} \neq 1$ . Therefore, we have shown Eq. (B18) as the fully regularized  $Z_{N_2}$  operator acting on the  $Z_{N_2} \times Z_{N_3}$  states.

It is straightforward to apply the above  $W_{j,j+1;N_1}^{\text{III}}$  discussion to  $S_{N_1, N_2, N_3}^{(p_{123})}$ ,  $W_{j,j+1}^{\text{III}}$ . We should just regard  $S_{N_1, N_2, N_3}^{(p_{123})}$ ,  $W_{j,j+1}^{\text{III}}$  as operators acting on the Hilbert space with  $Z_{N_1} \times Z_{N_2} \times Z_{N_3}$  states. We can show that all terms in  $W_{j,j+1;N_1}^{\text{III}}$ ,  $W_{j,j+1;N_2}^{\text{III}}$ ,  $W_{j,j+1;N_3}^{\text{III}}$  can be regularized in the same way.

#### 4. Matrix product operators and cocycles

Following, we calculate in details on Type III analog of Eq. (B3) to derive the nontrivial projective phase in MPO formalism, equivalent to the Type III 3-cocycles [Eq. (10)]. We use the fact Eq. (B2) to derive the projection tensor  $P_{g_a, g_b}$ :

$$\begin{aligned} P_{N_1, N_2, N_3}^{(p)} &\equiv P_{N_1, N_2, N_3, (m_a^{(1)}, m_b^{(1)}), (m_a^{(2)}, m_b^{(2)}), (m_a^{(3)}, m_b^{(3)})}^{(p)} \\ &= \prod_{u, v, w \in \{1, 2, 3\}} \int d\phi_{in}^{(u)} \left( \left| \phi_{in}^{(u)} + \frac{2\pi m_b^{(u)}}{N_1} \right| \phi_{in}^{(u)} \right) \left| \phi_{in}^{(u)} \right\rangle \left\langle \phi_{in}^{(u)} \right| e^{i2\pi p_{123} \epsilon^{uvw} \phi_{in}^{(u)} \left( \frac{m_a^{(v)}}{N_v} - \frac{m_b^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)}}}. \end{aligned} \quad (\text{B20})$$

Similar to Eq. (B5),  $P_{g_1, g_2}$  inputs one state  $\langle \phi_{in}^{(1)} | \langle \phi_{in}^{(2)} | \langle \phi_{in}^{(3)} |$  and outputs two states  $(|\phi_{in}^{(1)} + \frac{2\pi m_b^{(1)}}{N_1} \rangle | \phi_{in}^{(1)} \rangle) (|\phi_{in}^{(2)} + \frac{2\pi m_b^{(2)}}{N_2} \rangle | \phi_{in}^{(2)} \rangle) (|\phi_{in}^{(3)} + \frac{2\pi m_b^{(3)}}{N_3} \rangle | \phi_{in}^{(3)} \rangle)$ . For  $(I_1 \otimes P_{g_b, g_c}) P_{g_a, g_b, g_c}$ , we start by contracting  $T(g_b)$  and  $T(g_c)$  first, and then the combined tensor contracts with  $T(g_a)$  give

$$\begin{aligned} (I_1 \otimes P_{g_b, g_c}) P_{g_a, g_b, g_c} &= \prod_{u, v, w \in \{1, 2, 3\}} \int d\phi_{in}^{\prime\prime(u)} \left( \left| \phi_{in}^{\prime\prime(u)} + \frac{2\pi m_b^{(u)}}{N_u} + \frac{2\pi m_c^{(u)}}{N_u} \right| \phi_{in}^{\prime\prime(u)} + \frac{2\pi m_c^{(u)}}{N_u} \right) \left| \phi_{in}^{\prime\prime(u)} \right\rangle \left\langle \phi_{in}^{\prime\prime(u)} \right|_{abc} \\ &\times e^{i2\pi p_{123} \epsilon^{uvw} \phi_{in}^{\prime\prime(u)} \left( \frac{m_b^{(v)}}{N_v} - \frac{m_c^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)}}}. \end{aligned} \quad (\text{B21})$$

In Eq. (B21), we have dropped an extra factor  $e^{i2\pi p_{123} \epsilon^{uvw} \phi_{in}^{\prime\prime(u)} \left( \frac{m_b^{(v)}}{N_v} + \frac{m_c^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)}} = 1$ , because we are dealing with  $Z_N$  variables so the module relation renders the factor to be always trivial as 1.

On the other hand, to derive  $(P_{a,b} \otimes I_3) P_{ab,c}$ , we start by contracting  $T(g_a)$  and  $T(g_b)$  first, and then the combined tensor contracts with  $T(g_c)$ :

$$\begin{aligned} (P_{a,b} \otimes I_3) P_{ab,c} &= \prod_{u, v, w \in \{1, 2, 3\}} \int d\phi_{in}^{\prime\prime(u)} \left( \left| \phi_{in}^{\prime\prime(u)} + \frac{2\pi m_b^{(u)}}{N_u} + \frac{2\pi m_c^{(u)}}{N_u} \right| \phi_{in}^{\prime\prime(u)} + \frac{2\pi m_c^{(u)}}{N_u} \right) \left| \phi_{in}^{\prime\prime(u)} \right\rangle \left\langle \phi_{in}^{\prime\prime(u)} \right|_{abc} \\ &\times e^{i2\pi p_{123} \epsilon^{uvw} \left( \frac{2\pi m_c^{(u)}}{N_u} \right) \left( \frac{m_a^{(v)}}{N_v} - \frac{m_b^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)}}} e^{i2\pi p_{123} \epsilon^{uvw} \phi_{in}^{\prime\prime(u)} \left( \frac{m_a^{(v)}}{N_v} - \frac{m_b^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{2\pi \gcd(N_1, N_2, N_3)}}}. \end{aligned} \quad (\text{B22})$$

Comparing to Eq. (B3), we can derive  $e^{i\theta(g_a, g_b, g_c)}$  in Eq. (B23).

Adjusting the  $p_{123}$  index [i.e., setting Eq. (15)'s  $p_{123} \rightarrow p_{123}/2$ ,  $p_{213} = p_{312} = 0$ ], and computing Eq. (B3) with only the  $p_{123}$  index, we can recover the projective phase revealing Type III 3-cocycle:

$$e^{i\theta(g_1, g_2, g_3)} = e^{i2\pi p_{123} \epsilon^{uvw} \left( \frac{m_c^{(u)}}{N_u} - \frac{m_a^{(v)}}{N_v} - \frac{m_b^{(w)}}{N_w} \right) \frac{N_1 N_2 N_3}{\gcd(N_1, N_2, N_3)}} \simeq \omega_{\text{III}}^{(uvw)}(m_c, m_a, m_b). \quad (\text{B23})$$

#### APPENDIX C: INDUCED FRACTIONALIZED CHARGES AND DOMAIN WALL OPERATORS

Here we fill in more details on computing induced fractionalized charges (Type II bosonic anomaly) via lattice domain wall operators, outlined in Sec. III.C. The symmetry operator is  $S = \prod_j \tau_j \prod_j U_{j, j+1}$  acting on all sites on a 1D compact ring. We define a chain of domain wall operator from the site  $j = r_1$  to the site  $j = r_2$  as  $D(r_1, r_2) \equiv \prod_{j=r_1}^{r_2} \tau_j \prod_{j=r_1}^{r_2} U_{j, j+1}$  which creates a kink at the site  $r_1$  and an antikink at the site  $r_2$ . In the main text, we prescribe a method to capture the fractionalized charge at the kink/antikink based on

$$\begin{aligned} S D(r_1, r_2)^m S^\dagger &= [U(\omega^{-1} \sigma_{r_1-1}^\dagger \sigma_{r_1}) U^\dagger(\sigma_{r_1-1}^\dagger \sigma_{r_1})]^m \\ &\times [U(\omega \sigma_{r_2}^\dagger \sigma_{r_2+1}) U^\dagger(\sigma_{r_2}^\dagger \sigma_{r_2+1})]^m D(r_1, r_2)^m. \end{aligned} \quad (\text{C1})$$

Above we express a generic onsite symmetry operator  $\tau_j$  capturing  $\tau_j^{(u)}$  for  $\prod_u Z_{N_u}$  symmetry. We also express a generic non-onsite symmetry operator in terms of  $U_{j, j+1}$ . An explicit

calculation for Type I's  $U_{j, j+1}$  shows

$$[U(\omega \sigma_r^\dagger \sigma_{r+1}) U^\dagger(\sigma_r^\dagger \sigma_{r+1})]^m = e^{-i \frac{2\pi pm}{N^2} \sum_{a=1}^{N-1} (\sigma_r^\dagger \sigma_{r+1})^a} = e^{i \frac{2\pi pm}{N^2}}, \quad (\text{C2})$$

$$\begin{aligned} [U(\omega^{-1} \sigma_{r-1}^\dagger \sigma_r) U^\dagger(\sigma_{r-1}^\dagger \sigma_r)]^m &= e^{i \frac{2\pi pm}{N^2} \sum_{a=1}^{N-1} (\sigma_{r-1}^\dagger \sigma_r)^a} \\ &= e^{-i \frac{2\pi pm}{N^2}}. \end{aligned} \quad (\text{C3})$$

We can define  $[U(\omega^{-1} \sigma_{r_1-1}^\dagger \sigma_{r_1}) U^\dagger(\sigma_{r_1-1}^\dagger \sigma_{r_1})]^m \equiv e^{i\Theta_L}$  as the fractionalized charge phase measurement on the left kink at  $r_1$  since this operator contributes the phase gained exactly at the kink  $r_1$ . And we can define  $[U(\omega \sigma_{r_2}^\dagger \sigma_{r_2+1}) U^\dagger(\sigma_{r_2}^\dagger \sigma_{r_2+1})]^m \equiv e^{i\Theta_R}$  as the fractionalized charge phase measurement on the right kink at  $r_2$  since this operator contributes the phase gained exactly at the antikink  $r_2$ . Following, we explicitly express a generic non-onsite symmetry operator  $U_{j, j+1}$  in terms of non-onsite symmetry operators of Type I's  $U_{j, j+1}^{(N_u, p_u)}$ , Type II's  $U_{j, j+1}^{(N_u, p_{uv})}$ , Type III's  $W_{j, j+1}^{\text{III}}$ , with  $u, v \in \{1, 2, 3\}$ . The phases gained at the kink can be computed via the quantities  $S D(r_1, r_2)^m S^\dagger$  as follows:

(i) Type I:  $S_{N_1}^{(p_1)}(D_{N_1}^{(p_1)})^m S_{N_1}^{(p_1)\dagger}$  with  $e^{i\Theta_L} = e^{-i\Theta_R} = e^{i\frac{2\pi p_1}{N_1^2}m}$ .

(ii) Type II:  $S_{N_2}^{(p_{12})}(D_{N_1}^{(p_{12})})^m S_{N_2}^{(p_{12})\dagger}$  with  $e^{i\Theta_L} = e^{-i\Theta_R} = e^{i\frac{2\pi p_{12}}{N_2 N_1}m}$ .

(iii) Type III:  $S_{N_2}^{(p_{123})}(D_{N_1}^{(p_{123})})^m S_{N_2}^{(p_{123})\dagger}$  with  $e^{i\Theta_L} = e^{-i\Theta_R} = e^{i\frac{2\pi p_{123} n_3}{N_{123} m}}$ . Here  $n_3 = 0, 1, \dots, N_3 - 1$  is the exponent for each subblock of total  $N_3$  subblocks inside the  $W^{\text{III}}$  matrix (B18).

The systematic interpretation of fractionalized charge is organized in Table III in the main text.

## APPENDIX D: TWISTED SECTORS: TWISTED HAMILTONIAN AND TWISTED NON-ONSITE SYMMETRY TRANSFORMATION

### 1. Type II

We can adopt the discussion in Sec. VI on the twisted translation operator  $\tilde{T}^{(p)}$  and the twisted symmetry transformation  $S_N^{(p)}$  to Type II symmetry class. What we will focus on are the indices  $p_{12}$  and  $p_{21}$  of Eq. (28). We will set  $p_1 = p_2 = 0$  for the sake of simplicity. With this assumption, we can adjust the non-onsite symmetry transformation  $U_{j,j+2}^{(N_1, p_{12})} \rightarrow U_{j,j+1}^{(N_1, p_{12})}$  (from NNN to NN), also from  $U_{j,j+2}^{(N_2, p_{21})} \rightarrow U_{j,j+1}^{(N_2, p_{21})}$ . Here we explicitly indicate that  $U_{j,j+1}^{(N_1, p_{12})}$ ,  $U_{j,j+1}^{(N_2, p_{21})}$  are polynomial functions of  $(\tilde{\sigma}_j^{(2)\dagger} \tilde{\sigma}_{j+1}^{(2)})$ ,  $(\tilde{\sigma}_j^{(1)\dagger} \tilde{\sigma}_{j+1}^{(1)})$  respectively, with  $\tilde{\sigma}^{(1)}$ ,  $\tilde{\sigma}^{(2)}$  carefully being defined in Eq. (B4). The two principles addressed in Sec. VI for Type I are still valid. The first principle becomes defining the twisted symmetry transformation

$$\begin{aligned} \tilde{S}_{N_1}^{(p_{12})} &\equiv (\tilde{T}_{N_1}^{(p_{12})})^M = S_{N_1}^{(p_{12})}(U_{M,1}^{(N_1, p_{12})}[\tilde{\sigma}_M^{(2)\dagger} \tilde{\sigma}_1^{(2)}])^{-1} \\ &\times U_{M,1}^{(N_1, p_{12})}[\omega_{12} \tilde{\sigma}_M^{(2)\dagger} \tilde{\sigma}_1^{(2)}], \end{aligned} \quad (\text{D1})$$

$$\begin{aligned} \tilde{S}_{N_2}^{(p_{21})} &\equiv (\tilde{T}_{N_2}^{(p_{21})})^M = S_{N_2}^{(p_{21})}(U_{M,1}^{(N_2, p_{21})}[\tilde{\sigma}_M^{(1)\dagger} \tilde{\sigma}_1^{(1)}])^{-1} \\ &\times U_{M,1}^{(N_2, p_{21})}[\omega_{21} \tilde{\sigma}_M^{(1)\dagger} \tilde{\sigma}_1^{(1)}] \end{aligned} \quad (\text{D2})$$

with some unitary twisted translation operator  $\tilde{T}_{N_1}^{(p_{12})}$ ,  $\tilde{T}_{N_2}^{(p_{21})}$ , where the  $\tilde{S}_{N_1}^{(p_{12})}$  incorporate a  $Z_{N_1}$  flux at the branch cut, while the  $\tilde{S}_{N_2}^{(p_{21})}$  incorporate a  $Z_{N_2}$  flux at the branch cut. Here we insert  $\omega_{12} \equiv \omega_{21} \equiv e^{i\frac{2\pi}{\text{gcd}(N_1, N_2)}}$  into the non-onsite symmetry transformation  $U_{M,1}$  at the  $M$ th and the 1st sites to capture the branch cut physics as Fig. 12. The twisted lattice translation operators solved from Eqs. (D1) and (D2) are

$$\tilde{T}_{N_1}^{(p_{12})} = T U_{M,1}^{(N_1, p_{12})}[\tilde{\sigma}_M^{(2)\dagger} \tilde{\sigma}_1^{(2)}] \tau_1^{(1)}, \quad (\text{D3})$$

$$\tilde{T}_{N_2}^{(p_{21})} = T U_{M,1}^{(N_2, p_{21})}[\tilde{\sigma}_M^{(1)\dagger} \tilde{\sigma}_1^{(1)}] \tau_1^{(2)}. \quad (\text{D4})$$

The second principle is that the twisted Hamiltonian is invariant with respect to twisted translation operators  $\tilde{T}$ , thus also invariant with respect to  $\tilde{S}$ , i.e.,

$$\begin{aligned} [\tilde{H}_N^{(p)}, \tilde{T}_{N_1}^{(p_{12})}] &= [\tilde{H}_N^{(p)}, \tilde{S}_{N_1}^{(p_{12})}] = [\tilde{H}_N^{(p)}, \tilde{T}_{N_2}^{(p_{21})}] \\ &= [\tilde{H}_N^{(p)}, \tilde{S}_{N_2}^{(p_{21})}] = 0. \end{aligned} \quad (\text{D5})$$

The twisted Hamiltonian  $\tilde{H}_{N_1, N_2}^{(p_1, p_2, p_{12})}$  for Types I and II can be readily constructed from  $H_{N_1, N_2}^{(p_1, p_2, p_{12})}$  of Eq. (37), with the condition in Eqs. (78) and (D5).

### 2. Type III

We follow the same principles to explore the Type III twisted sectors with a flux insertion (or branch cut). We will focus on Type III class with  $p_{123} \neq 0$ , and other Type I and II class indices are zeros. The first principle suggests that a string of  $M$  units of *twisted translation operator*  $\tilde{T}_{N_1, N_2, N_3}^{(p_{123})}$  modifies Eq. (20)'s  $S_{N_1, N_2, N_3}^{(p_{123})}$  to a *twisted symmetry transformation*  $\tilde{S}_{N_1, N_2, N_3}^{(p_{123})} \equiv (\tilde{T}_{N_1}^{(p_{123})})^M (\tilde{T}_{N_2}^{(p_{123})})^M (\tilde{T}_{N_3}^{(p_{123})})^M$  incorporating a  $Z_{N_1}, Z_{N_2}, Z_{N_3}$  unit flux, respectively, by

$$\begin{aligned} \tilde{S}_{N_1, N_2, N_3}^{(p_{123})} &= S_{N_1, N_2, N_3}^{(p_{123})}(W_{M,1}^{\text{III}}[\sigma_M^{(v)\dagger} \sigma_1^{(v)}])^{-1} \\ &\times W_{M,1}^{\text{III}}[\omega_{123} \sigma_M^{(v)\dagger} \sigma_1^{(v)}], \end{aligned} \quad (\text{D6})$$

where the non-onsite symmetry transformation part  $W_{j,j+1}^{\text{III}} \equiv W_{j,j+1}^{\text{III}}[\sigma_{v,j}^\dagger \sigma_{v,j+1}]$  is defined in Eq. (32) as a polynomial of  $\sigma_{v,j}^\dagger \sigma_{v,j+1}$ , and its  $\omega_{123}$  insertion

$$\begin{aligned} W_{j,j+1}^{\text{III}}[\omega_{123} \sigma_j^{(v)\dagger} \sigma_{j+1}^{(v)}] \\ \equiv \prod_{u,v,w \in \{1,2,3\}} \epsilon^{uvw} (\omega_{123} \sigma_j^{(v)\dagger} \sigma_{j+1}^{(v)})^{p_{123} \frac{\ln(\sigma_j^{(w)}) N_v N_w}{2\pi \text{gcd}(N_1, N_2, N_3)}} \end{aligned} \quad (\text{D7})$$

captures the  $Z_{N_u}$  unit flux effect by the branch cut. [In Appendix B 2, we show that Eq. (D6) is regularized on the lattice.] Adopting the notation in Eq. (32), the twisted lattice translation operator solved from Eq. (D6) is

$$\tilde{T}_{N_u}^{(p_{123})} = T W_{M,1}^{\text{III}}(\tilde{\sigma}_M^{(v)\dagger} \tilde{\sigma}_1^{(v)}) \tau_1^{(u)}, \quad (\text{D8})$$

where  $u, v, w \in \{1, 2, 3\}$ .

The second principle is that the twisted Hamiltonian is invariant with respect to twisted translation operators, thus also invariant with respect to twisted symmetry transformations

$$[\tilde{H}_N^{(p)}, \tilde{T}_{N_u}^{(p_{123})}] = 0, [\tilde{H}_N^{(p)}, \tilde{S}_{N_1, N_2, N_3}^{(p_{123})}] = 0. \quad (\text{D9})$$

Based on Eq. (D9), it is straightforward to construct a Type III twisted Hamiltonian incorporating the symmetry twist (equivalently a gauge flux) at the branch cut.

- [1] E. Noether, Invariante Variationsprobleme., Nachr. D. König. Gesellsch. D. Wiss. Zu Göttingen, Math-Phys. Klasse **1918**, 235 (1918).  
 [2] G. 't Hooft, NATO Sci. Ser. B **59**, 135 (1980).  
 [3] S. L. Adler, *Phys. Rev.* **177**, 2426 (1969).

- [4] J. S. Bell and R. Jackiw, *Nuovo Cimento A* **60**, 47 (1969).  
 [5] See Appendices for more discussions and other technical details.  
 [6] X. Chen, Z. -C. Gu, Z. -X. Liu, and X. -G. Wen, *Phys. Rev. B* **87**, 155114 (2013).

- [7] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Science* **338**, 1604 (2012).
- [8] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [9] Z.-X. Liu, Z.-C. Gu, and X.-G. Wen, *Phys. Rev. Lett.* **113**, 267206 (2014).
- [10] F. D. M. Haldane, *Phys. Lett. A* **93**, 464 (1983).
- [11] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, *Commun. Math. Phys.* **115**, 477 (1988).
- [12] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [13] B. A. Bernevig and S.-C. Zhang, *Phys. Rev. Lett.* **96**, 106802 (2006).
- [14] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 146802 (2005).
- [15] J. E. Moore and L. Balents, *Phys. Rev. B* **75**, 121306 (2007).
- [16] L. Fu, C. L. Kane, and E. J. Mele, *Phys. Rev. Lett.* **98**, 106803 (2007).
- [17] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, *Phys. Rev. B* **78**, 195424 (2008).
- [18] A. Mesaros and Y. Ran, *Phys. Rev. B* **87**, 155115 (2013).
- [19] M. Levin and Z.-C. Gu, *Phys. Rev. B* **86**, 115109 (2012).
- [20] X. Chen, Z.-X. Liu, and X.-G. Wen, *Phys. Rev. B* **84**, 235141 (2011).
- [21] X. Chen and X.-G. Wen, *Phys. Rev. B* **86**, 235135 (2012).
- [22] L. H. Santos and J. Wang, *Phys. Rev. B* **89**, 195122 (2014).
- [23] X. Chen, Y.-M. Lu, and A. Vishwanath, *Nat. Commun.* **5**, 3507 (2014).
- [24] Y.-M. Lu and A. Vishwanath, *Phys. Rev. B* **86**, 125119 (2012).
- [25] A. Vishwanath and T. Senthil, *Phys. Rev. X* **3**, 011016 (2013).
- [26] P. Ye and J. Wang, *Phys. Rev. B* **88**, 235109 (2013).
- [27] Z. Bi, A. Rasmussen, and C. Xu, *Phys. Rev. B* **91**, 134404 (2015).
- [28] A. M. Essin and M. Hermele, *Phys. Rev. B* **87**, 104406 (2013).
- [29] P. Ye and X.-G. Wen, *Phys. Rev. B* **89**, 045127 (2014).
- [30] T. Grover and A. Vishwanath, *Phys. Rev. B* **87**, 045129 (2013).
- [31] Y.-M. Lu and D.-H. Lee, *Phys. Rev. B* **89**, 195143 (2014).
- [32] Y.-M. Lu and D.-H. Lee, *Phys. Rev. B* **89**, 184417 (2014).
- [33] R. Dijkgraaf and E. Witten, *Commun. Math. Phys.* **129**, 393 (1990).
- [34] R. Jackiw, *Phys. Rev. Lett.* **54**, 159 (1985).
- [35] S. B. Treiman, E. Witten, R. Jackiw, and B. Zumino, *Current Algebra and Anomalies* (World Scientific, Singapore, 1985), p. 537.
- [36] A. Kapustin, [arXiv:1403.1467](https://arxiv.org/abs/1403.1467).
- [37] X.-G. Wen, *Phys. Rev. D* **88**, 045013 (2013).
- [38] F. J. Burnell, X. Chen, L. Fidkowski, and A. Vishwanath, [arXiv:1302.7072](https://arxiv.org/abs/1302.7072).
- [39] M. A. Metlitski, C. L. Kane, and M. P. A. Fisher, [arXiv:1306.3286](https://arxiv.org/abs/1306.3286).
- [40] C. Wang, A. C. Potter, and T. Senthil, *Phys. Rev. B* **88**, 115137 (2013).
- [41] P. Bonderson, C. Nayak, and X.-L. Qi, *J. Stat. Mech.* (2013) P09016.
- [42] K. Fujikawa, *Phys. Rev. Lett.* **42**, 1195 (1979).
- [43] J. Goldstone and F. Wilczek, *Phys. Rev. Lett.* **47**, 986 (1981).
- [44] X.-G. Wen, *Phys. Rev. B* **89**, 035147 (2014).
- [45] M. P. Zaletel, *Phys. Rev. B* **90**, 235113 (2014).
- [46] M. de Wild Propitius, *Nucl. Phys. B* **489**, 297 (1997).
- [47] V. Murg, J. I. Cirac, B. Pirvu, and F. Verstraete, *New J. Phys.* **12**, 025012 (2010).
- [48] R. Floreanini and R. Jackiw, *Phys. Rev. Lett.* **59**, 1873 (1987).
- [49] If we use the field variables  $\phi_1, \phi'_1, \phi_2, \phi'_2$ , following the lattice operator relations between  $\tilde{\sigma}_j^{(u)}$  and  $\sigma_j^{(u)}$  with the commutation relation in Eq. (38), basically we have  $\phi_u \simeq \frac{N_{12}}{N_u} \tilde{\phi}_u$  and  $\phi'_u \simeq \frac{N_u}{N_{12}} \tilde{\phi}'_u$ , the symmetry transformation operators become  $S_{N_1}^{(p_1, p_{12})} \simeq \exp[\frac{i}{N_1} (\int_0^L dx \partial_x \phi'_1 + p_1 \int_0^L dx \partial_x \phi_1 + 0 \int_0^L dx \partial_x \phi'_2 + p_{12} \frac{N_2}{N_{12}} \int_0^L dx \partial_x \phi_2)]$  and  $S_{N_2}^{(p_2, p_{21})} \simeq \exp[\frac{i}{N_2} (0 \int_0^L dx \partial_x \phi'_1 + p_{21} \frac{N_1}{N_{12}} \int_0^L dx \partial_x \phi_1 + 1 \int_0^L dx \partial_x \phi'_2 + p_2 \int_0^L dx \partial_x \phi_2)]$ , thus they transform the fields by
- $$S_{N_1}^{(p_1, p_{12})} \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} (S_{N_1}^{(p_1, p_{12})})^{-1} = \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} + \frac{2\pi}{N_1} \begin{pmatrix} 1 \\ p_1 \\ 0 \\ \frac{N_2}{N_{12}} p_{12} \end{pmatrix}.$$
- $$S_{N_2}^{(p_2, p_{21})} \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} (S_{N_2}^{(p_2, p_{21})})^{-1} = \begin{pmatrix} \phi_1(x) \\ \phi'_1(x) \\ \phi_2(x) \\ \phi'_2(x) \end{pmatrix} + \frac{2\pi}{N_2} \begin{pmatrix} 0 \\ \frac{N_1}{N_{12}} p_{21} \\ 1 \\ p_2 \end{pmatrix}.$$
- This result in principle can still capture the correct physics quantity. But the true punch line is that one should follow our fully lattice-regularized setup, while regarding our field theory approach *only* as an effective tool to easily compute observables.
- [50] M. Cheng and Z.-C. Gu, *Phys. Rev. Lett.* **112**, 141602 (2014).
- [51] R. Jackiw and C. Rebbi, *Phys. Rev. D* **13**, 3398 (1976).
- [52] Y.-M. Lu and D.-H. Lee, *Phys. Rev. B* **89**, 205117 (2014).
- [53] X. G. Wen, *Phys. Rev. B* **40**, 7387 (1989).
- [54] L.-Y. Hung and X.-G. Wen, [arXiv:1211.2767](https://arxiv.org/abs/1211.2767).
- [55] L.-Y. Hung and X.-G. Wen, *Phys. Rev. B* **89**, 075121 (2014).
- [56] Y. Hu, Y. Wan, and Y.-S. Wu, *Phys. Rev. B* **87**, 125114 (2013).
- [57] J. Wang, Z. C. Gu, and X. G. Wen, *Phys. Rev. Lett.* **114**, 031601 (2015).
- [58] M. Henkel, *Conformal Invariance and Critical Phenomena* (Springer, Berlin, 1999).
- [59] O. M. Sule, X. Chen, and S. Ryu, *Phys. Rev. B* **88**, 075125 (2013).
- [60] A. Kapustin and R. Thorngren, *Phys. Rev. Lett.* **112**, 231602 (2014).
- [61] G. Y. Cho, J. C. Teo, and S. Ryu, *Phys. Rev. B* **89**, 235103 (2014).
- [62] Z.-X. Liu and X.-G. Wen, *Phys. Rev. Lett.* **110**, 067205 (2013).
- [63] M. Koch-Janusz and Z. Ringel, *Phys. Rev. B* **89**, 075137 (2014).
- [64] A. Vaezi, *Phys. Rev. B* **87**, 035132 (2013).
- [65] Zhong Wang, [arXiv:1405.0842](https://arxiv.org/abs/1405.0842).
- [66] S. M. Kravec and J. McGreevy, *Phys. Rev. Lett.* **111**, 161603 (2013).
- [67] Some anomaly-related phenomena are discussed in other contexts in interacting bosonic SPT systems, for anomalous Hall conductance [62] or conflicting symmetry under gauging [60, 61] or in interacting fractional topological insulators [63], or fractional Majorana zero modes in fractional topological superconductors [64], or the fractionalized Weyl semimetals for bosonic model [65].
- [68] Juven Wang and X.-G. Wen, [arXiv:1307.7480](https://arxiv.org/abs/1307.7480).
- [69] P. H. Ginsparg and K. G. Wilson, *Phys. Rev. D* **25**, 2649 (1982).
- [70] One may ask that, under  $Z_{N_1}$ -symmetry-preserving flux insertion or monodromy defect, should the Type III bosonic



edge modes keep gapless instead of gapped with  $N_{123}$ -fold degeneracy? We clarify that  $N_{123}$ -fold degeneracy is due to finite-size effects where we do dimensional reduction so the 1D ring is a small circle (in Fig. 8), where the *gapless* feature is manifest only in the thermodynamic limit when the 1D ring is a large circle.

- [71] L. Santos, Y. Nishida, C. Chamon, and C. Mudry, [Phys. Rev. B \*\*83\*\*, 104522 \(2011\)](#).
- [72] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*, Graduate Texts in Contemporary Physics (Springer, Berlin, 1996).
- [73] J. S. Schwinger, [Phys. Rev. \*\*128\*\*, 2425 \(1962\)](#).