

Constraining higher derivative supergravity with scattering amplitudes

Yifan Wang*

*Center for Theoretical Physics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139, USA*

Xi Yin†

Jefferson Physical Laboratory, Harvard University, Cambridge, Massachusetts 02138, USA

(Received 22 March 2015; published 31 August 2015)

We study supersymmetry constraints on higher derivative deformations of type IIB supergravity by consideration of superamplitudes. Combining constraints of on-shell supervertices and basic results from string perturbation theory, we give a simple argument for the nonrenormalization theorem of Green and Sethi, and some of its generalizations.

DOI: [10.1103/PhysRevD.92.041701](https://doi.org/10.1103/PhysRevD.92.041701)

PACS numbers: 04.65.+e, 11.25.Db, 11.30.Pb, 04.50.-h

Supersymmetry constraints on higher derivative couplings in maximal supergravity theories have been investigated extensively in the past [1–7] and have led to remarkable exact results on the quantum effective action of string theory. The method of Refs. [3,8–10] in obtaining constraints on higher derivative terms in gauge and gravity theories with maximal supersymmetry was by explicitly analyzing supersymmetry variations of fields and the Lagrangian and their deformations, starting from the purely fermionic terms. In this paper we present a simple argument for such nonrenormalization theorems from scattering amplitudes, in the context of ten-dimensional type IIB supergravity and its deformations, largely inspired by the work of Refs. [11–13] on the classification of supergravity counterterms [14–16] using amplitudes.

To begin with, we recall the spinor helicity formulation of superamplitudes in type IIB supergravity [17,18]. A ten-dimensional null momentum p^m and the corresponding (constrained) spinor helicity variables λ_A^α are related by

$$p^m \delta_{AB} = \Gamma_{\alpha\beta}^m \lambda_A^\alpha \lambda_B^\beta, \quad (1)$$

where α is a chiral spinor index of $SO(1,9)$ and A is a spinor index of the $SO(8)$ little group. The $2^8 = 256$ states in the supergraviton multiplet are built from monomials in a set of Grassmann variables η_A . The supermomentum is then defined as

$$q^\alpha = \lambda_A^\alpha \eta^A. \quad (2)$$

A typical n -point superamplitude takes the form [19]

$$\mathcal{A} = \delta^{10}(P) \delta^{16}(Q) \mathcal{F}(\lambda_i, \eta_i), \quad (3)$$

where $P = \sum_{i=1}^n p_i$, and the 32 supercharges that act on the n -particle asymptotic states can be expressed as

$$Q^\alpha = \sum_{i=1}^n q_i^\alpha, \quad \tilde{Q}^\alpha = \sum_{i=1}^n \lambda_i^{\alpha A} \frac{\partial}{\partial \eta_i^A}. \quad (4)$$

They obey $\{Q^\alpha, \tilde{Q}^\beta\} = \frac{1}{2} \Gamma_m^{\alpha\beta} P^m$. The nontrivial supersymmetry Ward identities on \mathcal{A} are

$$\delta^{10}(P) \delta^{16}(Q) \tilde{Q}^\alpha [\mathcal{F}(\lambda_i, \eta_i)] = 0. \quad (5)$$

We can write the CPT conjugate of the amplitude \mathcal{A} as

$$\bar{\mathcal{A}} = \delta^{10}(P) \tilde{Q}^{16} \mathcal{F}(\lambda_i, \partial/\partial \eta_i) \prod_{i=1}^n \eta_i^8. \quad (6)$$

Evidently, if \mathcal{A} obeys supersymmetry Ward identities, so does $\bar{\mathcal{A}}$.

Now let us focus on supervertices, namely superamplitudes with no poles in momenta. There are three basic types of supervertices we can write down. First, we can take $\mathcal{F}(\lambda_i, \eta_i)$ to be independent of η_i , namely

$$\mathcal{F}(\lambda_i, \eta_i) = f(s_{ij}), \quad (7)$$

where $s_{ij} = -(p_i + p_j)^2 = -2p_i \cdot p_j$. The CPT conjugate of this construction gives another supervertex. We refer to these as F-term vertices [20]. A third type of supervertex (D-term) is given by

$$\delta^{10}(P) \delta^{16}(Q) \tilde{Q}^{16} h(\lambda_i, \eta_i). \quad (8)$$

Here h is an arbitrary function of the spinor helicity variables. All supervertices we know of are of these three types. We conjecture that these are in fact the only supervertices that obey supersymmetry Ward identities, and we proceed with this assumption [23].

Let us inspect a particularly simple set of $n = (4 + k)$ -point F-term vertices, with $\mathcal{F}(\lambda_i, \eta_i) = 1$,

$$\delta^{10}(P) \delta^{16}(Q). \quad (9)$$

*yifanw@mit.edu

†xiyin@fas.harvard.edu

YIFAN WANG AND XI YIN

In component fields, we expand the axion-dilaton field as $\tau = \tau_0 + \varphi$, where τ_0 is the background value. Such a vertex then corresponds to an independent set of couplings in the Lagrangian of the form [2,4]

$$\varphi^k R^4 + \dots \quad (10)$$

Similarly, the conjugate vertex

$$\delta^{10}(P)\tilde{Q}^{16}\prod_{i=1}^{4+k}\eta_i^8 \quad (11)$$

corresponds to the coupling $\bar{\varphi}^k R^4 + \dots$. Note that in the $k = 0$ case, $\delta^{16}(Q) = \tilde{Q}^{16}\prod_{i=1}^4\eta_i^8$ is self-conjugate, and corresponds to the R^4 vertex [24]. In particular, we see that there are *no independent* supervertices of the form $\varphi^k\bar{\varphi}^\ell R^4 + \dots$ with $k, \ell \geq 1$. In other words, the supersymmetry completion of such couplings must be a superamplitude nonlocal in momenta.

Note that in a superamplitude, two $SO(8)$ little group invariant monomials in η_i^A , namely 1 and η_i^8 , correspond to the i th external particle being φ and $\bar{\varphi}$, respectively. The nonlinearly realized $SL(2, \mathbb{R})$ of type IIB supergravity is broken by the expectation value of τ to a $U(1)$ [25], which acts on the amplitude by $\sum_i(\frac{1}{4}\eta_i\frac{\partial}{\partial\eta_i} - 1)$ and assigns opposite charges to φ and $\bar{\varphi}$. This $SL(2, \mathbb{R})$ is generally broken explicitly by the higher derivative supervertices of consideration here.

Now, we would like to constrain the coupling

$$f(\tau, \bar{\tau})R^4 + \dots \quad (12)$$

by type IIB supersymmetry. In a vacuum in which τ acquires a constant expectation value τ_0 , expanding $\tau = \tau_0 + \varphi$, we obtain a series of operators,

$$f(\tau_0, \bar{\tau}_0)R^4 + \partial_\tau f(\tau_0, \bar{\tau}_0)\varphi R^4 + \partial_{\bar{\tau}} f(\tau_0, \bar{\tau}_0)\bar{\varphi} R^4 + \partial_\tau\partial_{\bar{\tau}} f(\tau_0, \bar{\tau}_0)\varphi\bar{\varphi} R^4 + \dots \quad (13)$$

Since there are independent φR^4 and $\bar{\varphi} R^4$ supervertices, $\partial_\tau f$ and $\partial_{\bar{\tau}} f$ can take an arbitrary value at $\tau = \tau_0$. This reflects a freedom in adjusting $f(\tau, \bar{\tau})$ by a holomorphic and an antiholomorphic function of τ . On the other hand, $\partial_\tau\partial_{\bar{\tau}} f$ at $\tau = \tau_0$ is not independent, because there is no independent $\varphi\bar{\varphi} R^4$ vertex. This 6-point coupling, therefore, must be constrained in terms of the R^4 coefficient, namely $f(\tau_0, \bar{\tau}_0)$, by supersymmetry.

In principle, one can ask for the most general 6-point superamplitude that obeys supersymmetry Ward identities and factorization through lower point amplitudes by unitarity. By dimensional analysis, the 6-point $\varphi\bar{\varphi}R^4$ superamplitude could only factorize through a single R^4 supervertex and supergravity vertices (Fig. 1). The $\varphi\bar{\varphi}R^4$

PHYSICAL REVIEW D **92**, 041701(R) (2015)




FIG. 1. Factorization of the 6-point amplitude through one $D^n R^4$ vertex (shaded blob) and a pair of supergravity cubic vertices.

coupling itself can then be recovered by taking the soft limit on a pair of φ and $\bar{\varphi}$ scalar lines [26].

We do not know a systematic way of building higher point superamplitudes with the R^4 on-shell supervertex [27]. However, from unitarity, we know that such a relation must exist, and is linear in this case, namely

$$(\text{Im}\tau_0)^2\partial_\tau\partial_{\bar{\tau}}f(\tau_0, \bar{\tau}_0) \propto f(\tau_0, \bar{\tau}_0), \quad (14)$$

where the $(\text{Im}\tau_0)^2$ factor comes from the normalization of the dilaton-axion kinetic term. To determine the relative coefficient, it suffices to find *any* set of such couplings that solve the supersymmetry and unitarity constraints. String perturbation theory already gives such a solution. Since the tree-level effective action of type IIB string theory contains an R^4 coupling at α'^3 order, it suffices to examine this coupling in the Einstein frame, which takes the form

$$\tau_2^{3/2}R^4, \quad (15)$$

where τ_2 is the imaginary part of τ .

Since $\partial_\tau\partial_{\bar{\tau}}\tau_2^{3/2} = \frac{3}{16}\tau_2^{-1/2}$, we immediately obtain the relation

$$4(\text{Im}\tau_0)^2\partial_\tau\partial_{\bar{\tau}}f(\tau_0, \bar{\tau}_0) = \frac{3}{4}f(\tau_0, \bar{\tau}_0), \quad (16)$$

which must then hold for the general $f(\tau, \bar{\tau})$ at all values of τ_0 . This is the nonrenormalization theorem of Green and Sethi [3]. Below, we write $f_n(\tau, \bar{\tau})$ for the coefficient of $D^n R^4$, and so $f(\tau, \bar{\tau})$ is denoted $f_0(\tau, \bar{\tau})$.

Note that there is no independent $D^2 R^4$ supervertex, as the corresponding superamplitude must be proportional to $\delta^{16}(Q)(s+t+u) = 0$. We next apply the argument to the $f_4(\tau, \bar{\tau})D^4 R^4$ coupling. Once again, the holomorphic and antiholomorphic parts of $f_4(\tau, \bar{\tau})$ are unconstrained by supersymmetry, as there are independent $\varphi^k R^4$ and $\bar{\varphi}^k R^4$




FIG. 2. Factorization of the 6-point amplitude through a pair of R^4 vertices.

supervertices. On the other hand, $\partial_\tau \partial_{\bar{\tau}} f_4$ must obey a linear relation with $\tau_2^{-2} f_4(\tau, \bar{\tau})$, due to the factorization of the 6-point superamplitude. Note that the 6-point amplitude at this order in the momentum expansion does not factorize through two R^4 vertices (Fig. 2), as the latter can only contribute to the 6-point amplitude at $D^6 R^4$ order [30].

Now taking the IIB string tree-level effective action, and expanding to α'^5 order, we find in the Einstein frame the coupling

$$\tau_2^{5/2} (s^2 + t^2 + u^2) R^4. \quad (17)$$

By comparison, we then immediately obtain the relation

$$4\tau_2^2 \partial_\tau \partial_{\bar{\tau}} f_4(\tau, \bar{\tau}) = \frac{15}{4} f_4(\tau, \bar{\tau}). \quad (18)$$

At $f_6(\tau, \bar{\tau}) D^6 R^4$ order, we encounter a novelty: as already mentioned, the 6-point amplitude at this order in the momentum expansion admits a factorization into a pair of R^4 supervertices. Thus, we expect the coefficient $f_6(\tau, \bar{\tau})$ to obey a relation of the form

$$\tau_2^2 \partial_\tau \partial_{\bar{\tau}} f_6 = a f_6(\tau, \bar{\tau}) + b f_0(\tau, \bar{\tau})^2, \quad (19)$$

where a and b are two constants. More precisely, we define $f_6(\tau, \bar{\tau})$ to be the coefficient of $(s^3 + t^3 + u^3) R^4 = 3stu R^4$. Inspecting the well-known string tree-level massless 4-point amplitude,

$$\begin{aligned} \delta^{16}(Q) & \frac{\Gamma(-\frac{\alpha s}{4}) \Gamma(-\frac{\alpha t}{4}) \Gamma(-\frac{\alpha u}{4})}{\Gamma(1 + \frac{\alpha s}{4}) \Gamma(1 + \frac{\alpha t}{4}) \Gamma(1 + \frac{\alpha u}{4})} \\ & = \delta^{16}(Q) \left[-\frac{64}{\alpha^3 stu} - 2\zeta(3) - \frac{\zeta(5)}{16} \alpha^2 (s^2 + t^2 + u^2) \right. \\ & \quad \left. - \frac{\zeta(3)^2}{96} \alpha^3 (s^3 + t^3 + u^3) + \dots \right], \end{aligned} \quad (20)$$

we can identify the following couplings in the Einstein frame [31],

$$\begin{aligned} & -2\zeta(3) \tau_2^{3/2} \alpha^3 R^4 - \frac{\zeta(5)}{16} \alpha^5 \tau_2^{5/2} (s^2 + t^2 + u^2) R^4 \\ & - \frac{\zeta(3)^2}{96} \alpha^6 \tau_2^3 (s^3 + t^3 + u^3) R^4 + \dots \end{aligned} \quad (21)$$

Comparing to Eq. (19), with $f_0 \propto \tau_2^{3/2}$ and $f_6 \propto \tau_2^3$, we immediately obtain a linear relation between a and b . Another relation between a and b may be extracted from the string 1-loop effective action. The perturbative contribution to f_0 and f_6 can be expanded in the form [6]

$$f_n(\tau, \bar{\tau}) = f_n^{\text{tree}} + f_n^{1\text{-loop}} + f_n^{2\text{-loop}} + f_n^{3\text{-loop}} + \dots \quad (22)$$

In particular, at 1-loop order, we expect

$$\tau_2^2 \partial_\tau \partial_{\bar{\tau}} f_6^{1\text{-loop}} = a f_6^{1\text{-loop}}(\tau, \bar{\tau}) + 2b f_0^{\text{tree}}(\tau, \bar{\tau}) f_0^{1\text{-loop}}(\tau, \bar{\tau}). \quad (23)$$

The 4-point massless genus one string amplitude amplitude has analytic as well as nonanalytic terms in the momentum expansion. The R^4 term, with coefficient $f_0^{1\text{-loop}} \propto \tau_2^{-1/2}$, and the $D^6 R^4$ term, with coefficient $f_6^{1\text{-loop}} \propto \tau_2$, are analytic, and were computed in Ref. [35]. They give an independent linear relation which then fixes a and b , as in (5.39) of Ref. [6]. In the end, one finds

$$4\tau_2^2 \partial_\tau \partial_{\bar{\tau}} f_6 = 12f_6(\tau, \bar{\tau}) - 6f_0(\tau, \bar{\tau})^2. \quad (24)$$

As was pointed out in Ref. [6], the string 3-loop contribution $f_6^{3\text{-loop}}$ [6,36–39], proportional to τ_2^{-3} , is what solves the homogeneous version of the constraining equation (namely, it is annihilated by $4\tau_2^2 \partial_\tau \partial_{\bar{\tau}} - 12$).

Now let us consider $D^8 R^4$ terms. There is again one independent 4-point supervertex one can write down [40],

$$\delta^{16}(Q) (s^4 + t^4 + u^4). \quad (25)$$

This is, in fact, proportional to the D-term vertex

$$\delta^{16}(Q) \tilde{Q}^{16} \left[\sum_{i < j}^4 \eta_i^8 \eta_j^8 \right]. \quad (26)$$

To understand the constraints on $f_8(\tau, \bar{\tau})$, let us inspect ($n = 4 + k$)-point supervertices of the form

$$\delta^{16}(Q) \tilde{Q}^{16} F(\eta_i^8), \quad (27)$$

where $F(\eta_i^8)$ is a polynomial in the little group invariants η_i^8 , of total degree $8m$ in the η 's, for some integer $m \geq 2$. This then corresponds to a coupling of the form $\varphi^{k-m+2} \tilde{\varphi}^{m-2} D^8 R^4$. Since these D-term vertices by construction obey supersymmetry Ward identities, there are no constraints on the coefficients of $\varphi^{k-m+2} \tilde{\varphi}^{m-2} D^8 R^4$, thus, no constraint on $f_8(\tau, \bar{\tau})$ from supersymmetry alone.

At order $D^{10} R^4$, there is again just one independent 4-point supervertex $\delta^{16}(Q) (s^5 + t^5 + u^5)$. This is proportional to the D-term vertex $\delta^{16}(Q) \tilde{Q}^{16} [\sum_{i < j}^4 s_{ij} \eta_i^8 \eta_j^8]$ [41]. As in the $D^8 R^4$ case, there are no supersymmetry

constraints on the coefficient $f_{10}(\tau, \bar{\tau})$. In other words, the differential constraint proposed in Ref. [42] should be a consequence of additional properties of IIB string theory.

In conclusion, the formulation of higher derivative couplings in maximally supersymmetric gravity theories in terms of on-shell supervertices gives a simple classification of independent couplings allowed by supersymmetry. When combined with solutions to supersymmetry Ward identities provided by string perturbation theory, the consideration of supervertices then leads to a derivation of type IIB supersymmetry constraints on the F-term $f(\tau, \bar{\tau})D^n R^4$ coupling ($n = 0, 4, 6$). The result is nonetheless a consequence of maximal supersymmetry on higher derivative supergravity theories, and no longer depends on string theory. Clearly, this strategy generalizes straightforwardly to maximal supergravity theories in other dimensions as well [43]. We shall leave this to a future publication.

Finally, let us comment on the role of $SL(2, \mathbb{R})$ symmetry of type IIB supergravity, which, as already mentioned, is explicitly broken by these higher derivative terms. A coupling of the form $f_n(\tau, \bar{\tau})D^n R^4$ violates $SL(2, \mathbb{R})$ unless f_n is a constant, but the latter is incompatible with the supersymmetry constraints (a nontrivial second order

differential equation in $\tau, \bar{\tau}$) for F-term vertices. From this perspective, a role of the nonlinearly realized $SL(2, \mathbb{R})$ symmetry of type IIB supergravity is to rule out F-terms as potential counterterms. Indeed, the UV divergence in type IIB supergravity first arises at 2-loop order, corresponding to an $SL(2, \mathbb{R})$ -invariant D-term counterterm of the form $D^{10}R^4$. One may expect that the $E_{7(7)}$ symmetry of four-dimensional maximal supergravity plays a similar role in that it rules out F-terms as counterterms, but there appear to be plenty of D-term supervertices compatible with $E_{7(7)}$ that could serve as counterterms [44–52].

ACKNOWLEDGMENTS

We would like thank N. Arkani-Hamed, C. Córdova, T. Dumitrescu, H. Elvang, D. Freedman, S. Hellerman, Y.-t. Huang, H. Nicolai, A. Zhiboedov, and especially Z. Komargodski for extensive discussions. We are grateful to the hospitality of the Weizmann Institute, Jerusalem Winter School, and Kavli IPMU during the course of this work. Y. W. is supported in part by the U.S. Department of Energy under Contract No. DE-SC00012567. X. Y. is supported by Alfred P. Sloan Foundation and Simons Foundation.

-
- [1] M. B. Green and M. Gutperle, *Nucl. Phys.* **B498**, 195 (1997).
 [2] N. Berkovits, *Nucl. Phys.* **B514**, 191 (1998).
 [3] M. B. Green and S. Sethi, *Phys. Rev. D* **59**, 046006 (1999).
 [4] B. Pioline, *Phys. Lett. B* **431**, 73 (1998).
 [5] N. Berkovits and C. Vafa, *Nucl. Phys.* **B533**, 181 (1998).
 [6] M. B. Green and P. Vanhove, *J. High Energy Phys.* **01** (2006) 093.
 [7] A. Basu and S. Sethi, *J. High Energy Phys.* **09** (2008) 081.
 [8] S. Paban, S. Sethi, and M. Stern, *Nucl. Phys.* **B534**, 137 (1998).
 [9] S. Paban, S. Sethi, and M. Stern, *J. High Energy Phys.* **06** (1998) 012.
 [10] S. Sethi and M. Stern, *J. High Energy Phys.* **06** (1999) 004.
 [11] H. Elvang, D. Z. Freedman, and M. Kiermaier, *J. Phys. A* **44**, 454009 (2011).
 [12] H. Elvang, D. Z. Freedman, and M. Kiermaier, *J. High Energy Phys.* **11** (2010) 016.
 [13] H. Elvang and Y.-t. Huang, arXiv:1308.1697.
 [14] R. Kallosh, *Phys. Lett.* **99B**, 122 (1981).
 [15] P. S. Howe and U. Lindstrom, *Nucl. Phys.* **B181**, 487 (1981).
 [16] P. S. Howe, K. Stelle, and P. Townsend, *Nucl. Phys.* **B191**, 445 (1981).
 [17] S. Caron-Huot and D. O’Connell, *J. High Energy Phys.* **08** (2011) 014.
 [18] R. H. Boels and D. O’Connell, *J. High Energy Phys.* **06** (2012) 163.
 [19] The cubic vertex is special (see Ref. [18]), and may be constructed as follows. Define $\hat{N}^\alpha_\beta = p_1^m p_2^n (\Gamma_{mn})^\alpha_\beta$ which specifies the null plane spanned by the three external momenta p_1, p_2, p_3 . We have $\hat{N}\lambda_{iA} = 0$ for $i = 1, 2, 3$. We can label the $SO(1, 9)$ spinor components by $(s_0 s_1 s_2 s_3 s_4)$, $s_a = \pm$, such that $\hat{N} = (p^+)^2 \Gamma^{0-1-}$, where Γ^{0-1-} is the lowering operator on both s_0 and s_1 . Now decompose the spinor helicity variables according to their $s_0 s_1$ spin, $\lambda_{iA} = (\lambda_{iA}^{++}, \lambda_{iA}^{+-}, \lambda_{iA}^{-+}, \lambda_{iA}^{--})$, where each $\lambda_{iA}^{\pm\pm}$ is a spinor of the $SO(6)$ tiny group that acts transversely to the null plane, and the condition $\hat{N}\lambda_{iA} = 0$ amounts to $\lambda_{iA}^{++} = 0$. We define the tiny group spinor valued supermomentum $W^{\pm\pm} = \sum_i^3 \lambda_{iA}^{\pm\pm} \eta_i^A$. The cubic supervertex is then given by the boost invariant combination $\frac{1}{(p^+)^4} \delta^4(W^{+-}) \delta^4(W^{-+}) \delta^4(W^{--})$.
 [20] These vertices are also known as “maximal R -symmetry violating” (MRV) in [21,22].
 [21] R. H. Boels, *Phys. Rev. Lett.* **109**, 081602 (2012).
 [22] R. H. Boels, *Nucl. Phys.* **B876**, 215 (2013).
 [23] We observe evidences for this conjecture from IIB string perturbation theory. In lower dimensions, there could be exceptional F-term supervertices not of the given type due to a larger R -symmetry group, which we shall explore in a future publication.

- [24] In contrast, the supergravity 4-point tree amplitude is given by $\frac{\delta^{10}(P)\delta^{16}(Q)}{stuv}$ [17,18].
- [25] While this $U(1)$ acts on the target space of the axion-dilaton field locally as an isometry, in type IIB string theory it is incompatible with the $SL(2, \mathbb{Z})$ identification.
- [26] C. Cheung, K. Kampf, J. Novotny, and J. Trnka, *Phys. Rev. Lett.* **114**, 221602 (2015).
- [27] For instance, if one applies the BCFW [28] shift to a pair of external lines and tries to rewrite the higher point tree amplitude as a contour integral in the shift parameter z , one encounters nontrivial residue at $z = \infty$, which cannot be determined in a straightforward way. The all-line shift of Ref. [29] improves the behavior at $z = \infty$, but still does not appear to apply when general higher derivative vertices are present.
- [28] R. Britto, F. Cachazo, B. Feng, and E. Witten, *Phys. Rev. Lett.* **94**, 181602 (2005).
- [29] T. Cohen, H. Elvang, and M. Kiermaier, *J. High Energy Phys.* **04** (2011) 053.
- [30] This can be seen from the corresponding BCFW [18] residues: for the factorization in Fig. 2, it takes the form $\frac{\delta^{16}(Q)}{s_{123}} \int d^8\eta_P \delta^{16}(q_P + q_4 + q_5 + q_6)$.
- [31] Effective action couplings from higher-point superstring tree amplitudes have also been extracted, for example, in Refs. [32,33] for type I open strings and in Ref. [34] for type II closed strings.
- [32] L. A. Barreiro and R. Medina, *J. High Energy Phys.* **10** (2012) 108.
- [33] L. A. Barreiro and R. Medina, *Nucl. Phys.* **B886**, 870 (2014).
- [34] A. Basu, *Classical Quantum Gravity* **30**, 235028 (2013).
- [35] M. B. Green and P. Vanhove, *Phys. Rev. D* **61**, 104011 (2000).
- [36] H. Gomez and C.R. Mafra, *J. High Energy Phys.* **10** (2013) 217.
- [37] E. D'Hoker, M. B. Green, B. Pioline, and R. Russo, *J. High Energy Phys.* **01** (2015) 031.
- [38] A. Basu, *Classical Quantum Gravity* **31**, 245002 (2014).
- [39] B. Pioline, *J. High Energy Phys.* **04** (2015) 057.
- [40] Note that $(s^2 + t^2 + u^2)^2$ is proportional to $(s^4 + t^4 + u^4)$, with $s + t + u = 0$.
- [41] One may try to write down another D-term vertex using little group invariants of the form $\delta^{16}(Q)\tilde{Q}^{16}[(\eta_1^7\lambda_1)^\alpha(\eta_2^7\lambda_2)^\beta q_3^\gamma q_4^\delta(\Gamma^{mnp})_{\alpha\beta}(\Gamma_{mnp})_{\gamma\delta} + \text{permutations}]$. Nonetheless, this must be proportional to $\delta^{16}(Q)\tilde{Q}^{16}[\sum_{i<j} s_{ij}\eta_i^8\eta_j^8]$ and cannot be an independent vertex.
- [42] A. Basu, *Phys. Lett. B* **648**, 378 (2007).
- [43] Except for type IIA and eleven-dimensional supergravity, where a supermomentum formulation of the superamplitude as presented here is unavailable.
- [44] J. Broedel and L. J. Dixon, *J. High Energy Phys.* **05** (2010) 003.
- [45] N. Beisert, H. Elvang, D.Z. Freedman, M. Kiermaier, A. Morales *et al.*, *Phys. Lett. B* **694**, 265 (2010).
- [46] G. Bossard, P. Howe, and K. Stelle, *J. High Energy Phys.* **01** (2011) 020.
- [47] R. Kallosh, [arXiv:1412.7117](https://arxiv.org/abs/1412.7117).
- [48] Z. Bern, L.J. Dixon, D. Dunbar, M. Perelstein, and J. Rozowsky, *Nucl. Phys.* **B530**, 401 (1998).
- [49] Z. Bern, L. J. Dixon, and R. Roiban, *Phys. Lett. B* **644**, 265 (2007).
- [50] Z. Bern, J. J. Carrasco, L. J. Dixon, H. Johansson, D. A. Kosower, and R. Roiban, *Phys. Rev. Lett.* **98**, 161303 (2007).
- [51] Z. Bern, J. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, *Phys. Rev. Lett.* **103**, 081301 (2009).
- [52] Z. Bern, J. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, *Fortschr. Phys.* **59**, 561 (2011).