

When Does the Set of (a, b, c) -Core Partitions Have a Unique Maximal Element?

Amol Aggarwal *

Department of Mathematics
Massachusetts Institute of Technology
Massachusetts, U.S.A.

agg_a@mit.edu

Submitted: Oct 21, 2014; Accepted: May 9, 2015; Published: May 22, 2015

Mathematics Subject Classifications: 05A17

Abstract

In 2007, Olsson and Stanton gave an explicit form for the largest (a, b) -core partition, for any relatively prime positive integers a and b , and asked whether there exists an (a, b) -core that contains all other (a, b) -cores as subpartitions; this question was answered in the affirmative first by Vandehey and later by Fayers independently. In this paper we investigate a generalization of this question, which was originally posed by Fayers: for what triples of positive integers (a, b, c) does there exist an (a, b, c) -core that contains all other (a, b, c) -cores as subpartitions? We completely answer this question when a , b , and c are pairwise relatively prime; we then use this to generalize the result of Olsson and Stanton.

Keywords: Young diagram; hook length; core partition; numerical semigroup; UM-set; poset-UM

1 Introduction

A *partition* is a finite, nonincreasing sequence $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ of positive integers. The sum $\sum_{i=1}^r \lambda_i$ is the *size* of λ and is denoted by $|\lambda|$; the integer r is the *length* of λ . A partition $\mu = (\mu_1, \mu_2, \dots, \mu_s)$ is a *subpartition* of λ if $s \leq r$ and $\mu_i \leq \lambda_i$ for each integer $i \in [1, s]$; in this case, we say that $\mu \subseteq \lambda$.

We may represent λ by a *Young diagram*, which is a collection of r left-justified rows of cells with λ_i cells in row i . The *hook length* of any cell C in the Young diagram is defined to be the number of cells to the right of, below, or equal to C . For instance, Figure 1 shows the Young diagrams and hook lengths of the partitions $(6, 4, 2, 2, 1, 1)$ and $(5, 3, 1, 1)$.

*Supported by NSF Grant 1358659 and NSA Grant H98230-13-1-0273.

Let $\beta(\lambda)$ denote the set of hook lengths in the leftmost column of the Young tableaux associated with λ ; equivalently, $\beta(\lambda) = (\lambda_1 + r - 1, \lambda_2 + r - 2, \dots, \lambda_r)$. For instance, Figure 1 shows that $\beta(6, 4, 2, 2, 1, 1) = \{11, 8, 5, 4, 2, 1\}$ and $\beta(5, 3, 1, 1) = (8, 5, 2, 1)$.

For any set of positive integers $A = \{a_1, a_2, \dots, a_k\}$, a partition is an A -core if no cell of its Young diagram has hook length in A . Let the set of A -cores be $C(A)$; Figure 1 shows that $(6, 4, 2, 2, 1, 1) \in C(3, 7)$ and $(5, 3, 1, 1) \in C(3, 7, 11)$.

Core partitions are known to be related to representations of the symmetric group; for instance, Olsson and Stanton use simultaneous core partitions in [12] to prove the Navarro-Willems conjecture for symmetric groups. Core partitions are also known to be related to the alcove geometry for certain types of Coxeter groups (see [4, 8, 9]). Recently, there has been a growing interest in simultaneous core partitions because of their relationship with numerical semigroups (see [1, 2, 4, 15, 17]).




Figure 1: To the left is the Young diagram of $\kappa_{3,7} = (6, 4, 2, 2, 1, 1)$ and to the right is the Young diagram of $(5, 3, 1, 1)$; each cell contains its hook length.

During the past decade, combinatorialists have studied properties of $C(A)$ when $|A| = 2$ (see [1, 3, 4, 5, 6, 8, 9, 10, 11, 12, 15, 16]). For instance, Anderson showed that $|C(a, b)| = \binom{a+b}{a} / (a+b)$ if a and b are relatively prime; in particular, there are finitely many (a, b) -cores [3].

This implies that there is an (a, b) -core of maximum size. Auckerman, Kane, and Sze conjectured in [5] that this size is $(a^2 - 1)(b^2 - 1)/24$. This was verified in 2007 by Olsson and Stanton, who also found the core of this size explicitly in terms of a and b [12]. Specifically, they established the following result.

Theorem 1.1. *For any relatively prime positive integers a and b , there is a unique (a, b) -core $\kappa_{a,b}$ of maximum size; a positive integer is in $\beta(\kappa_{a,b})$ if and only if it is of the form $ab - ia - jb$ for some positive integers i and j .*

Figure 1 depicts the Young diagram of $\kappa_{3,7} = (6, 4, 2, 2, 1, 1)$. In their proof of Theorem 1.1, Stanton and Olsson showed that $\kappa_{n,n+1}$ contains every other $(n, n+1)$ -core as a subpartition, for each integer $n \geq 2$ [12]. They then asked whether $\kappa_{a,b}$ contains all other (a, b) -cores as subpartitions, for every pair of relatively prime positive integers (a, b) . Vandehey answered this question in the affirmative in 2009 through the use of abacus diagrams [16]. Recently, Fayers obtained the same result by analyzing actions of the affine symmetric group on the set of a -cores (and on the set of b -cores) [8].

To see an example of Vandehey's theorem, let $A = (3, 5)$. The nonempty partitions in $C(A)$ are $\{(1), (2), (1, 1), (3, 1), (2, 1, 1), (4, 2, 1, 1)\}$, and every element of $C(A)$ is contained in the $(3, 5)$ -core $(4, 2, 1, 1)$. However, this containment phenomenon does not necessarily hold when $|A| \geq 3$ and $\gcd A = 1$. For instance, if $A = \{3, 4, 5\}$, then the nonempty elements of $C(A)$ are $(1, 1)$ and (2) ; neither of these is contained in the other.

For any set of positive integers A , we say that $C(A)$ has a *unique maximal element* if there is an A -core κ_A that contains every other A -core as a subpartition; in this case, the set A is said to be *UM*. In [7], Fayers asked the following question.

Question 1.2. *What triples of positive integers (a, b, c) are UM?*

Vandehey's result implies a partial result in this direction. For any set of positive integers A , let $S(A)$ be the numerical semigroup generated by A ; equivalently, A consists of all linear combinations of elements in A with nonnegative integer coefficients. Due to the known fact that an (a, b) -core is an $(a + b)$ -core (see [2], for instance), Vandehey's result implies that (a, b, c) is UM if a and b are relatively prime and $c \in S(a, b)$. Recently, Yang, Zhong, and Zhou showed that $(2k + 1, 2k + 2, 2k + 3)$ is not UM for any positive integer k [17].

In this paper we give a partial answer to Question 1.2. We call a triple of positive integers (a, b, c) *aprimitive* if either $a \in S(b, c)$, $b \in S(a, c)$, or $c \in S(a, b)$. The following theorem gives a restriction on triples that can be UM.

Theorem 1.3. *Suppose that (a, b, c) is a triple of positive integers such that $\gcd(a, b, c) = 1$; let $p = \gcd(a, b)$, $q = \gcd(a, c)$, $r = \gcd(b, c)$, and d, e, f be integers such that $(a, b, c) = (dpq, epr, fqr)$ as ordered triples. If (a, b, c) is UM, then (d, e, f) is aprimitive.*

Not all triples of the form given by the above theorem are UM; for instance, we will see in Section 2 that $(4, 5, 6)$ is not UM. However, we may use Theorem 1.3 to answer Question 1.2 completely when a, b , and c are pairwise relatively prime.

Corollary 1.4. *If $a < b < c$ are pairwise relatively prime positive integers, then (a, b, c) is UM if and only if $c \in S(a, b)$.*

Proof. As noted previously, Vandehey's theorem implies that (a, b, c) is UM if $c \in S(a, b)$. Setting $p = q = r = 1$ in Theorem 1.3 yields the converse. \square

Corollary 1.4 can be viewed as a converse to Vandehey's theorem; it also generalizes the previously mentioned result of Yang, Zhong, and Zhou.

Using Theorem 1.3, we will also be able to express the unique maximal (a, b, c) -core $\kappa_{a,b,c}$ in terms of a, b , and c if the triple (a, b, c) is UM.

Theorem 1.5. *Suppose that $A = (a, b, c)$ is a triple of positive integers that is UM; let $p = \gcd(a, b)$, $q = \gcd(a, c)$, $r = \gcd(b, c)$, and d, e, f be integers such that $(a, b, c) = (dpq, epr, fqr)$ as ordered triples. If $f \in S(d, e)$, then a positive integer is in $\beta(\kappa_A)$ if and only if it is of the form $(de + f)pqr - ia - jb - kc$ for some positive integers i, j , and k .*

Observe that letting $p = q = r = 1$ in Theorem 1.5 yields Theorem 1.1 of Olsson and Stanton, due to Vandehey's theorem.

The proofs of Theorem 1.3 and Theorem 1.5 use a recently developed characterization of simultaneous cores through numerical semigroups, which we will explain further in the next section.

2 Proofs of Theorems 1.3 and 1.5

In this section, we first explain a bijection (originally due to Stanley and Zanello in [15] when $|A| = 2$ and later generalized to arbitrary sets A by Amdeberhan and Leven in [2]) between A -cores and order ideals of some poset $P(A)$. We will then use this bijection to obtain a preliminary necessary condition for a set A to be UM. Using this condition, we will establish Theorem 1.3 and Theorem 1.5.

Now let us define the poset $P(A)$. The elements of $P(A)$ are those of $\mathbb{Z}_{\geq 0} \setminus S(A)$, the set of positive integers not contained in the numerical semigroup generated by A . Notice that if $\gcd A = 1$, then $|P(A)| < \infty$; we will suppose that this is the case for the remainder of the section. The order on $P(A)$ is fixed by requiring $p \in P(A)$ to be greater than $q \in P(A)$ if $p - q \in S(A)$. Under this partial order, $P(A)$ is a poset; we will follow the poset terminology given in Chapter 3 of Stanley's text [13, 14]. Figure 2 depicts the Hasse diagrams of the posets $P(3, 7)$ and $P(3, 7, 11)$.



Figure 2: The Hasse diagrams of $P(3, 7)$ and $P(3, 7, 11)$ are shown on the left and right, respectively.

The following lemma is due to Amdeberhan and Leven in [2].

Lemma 2.1. *There is a bijection between $C(A)$ and the set of order ideals of $P(A)$. Specifically, for each partition λ , the set $\beta(\lambda)$ is an order ideal of $P(A)$ if and only if λ is an A -core.*

For instance, suppose that $A \subseteq \{3, 7, 11\}$; then, $(5, 3, 1, 1)$ is an A -core and its beta set $\{8, 5, 2, 1\}$ is an order ideal of $P(A)$. Furthermore, $\kappa_{3,7} = (6, 4, 2, 2, 1, 1)$ is a $(3, 7)$ -core and its beta set $\{1, 2, 4, 5, 8, 11\}$ is an order ideal of $P(3, 7)$; however, $\kappa_{3,7}$ is not a $(3, 7, 11)$ -core and its beta set is not an order ideal of $P(3, 7, 11)$.

From Lemma 2.1, there is an A -core κ'_A such that $\beta(\kappa'_A) = P(A)$. The following result states that κ'_A is the unique maximal element of $C(A)$ if A is UM.

Corollary 2.2. *If a set of positive integers A is UM, then $\kappa'_A = \kappa_A$.*

Proof. The bijection in Lemma 2.1 is length preserving; since the longest order ideal of $P(A)$ is $P(A)$, the longest A -core is κ'_A . Therefore, κ'_A is not contained in any other A -core, which implies that κ_A is the unique maximal element of $C(A)$ because A is UM. Thus, $\kappa'_A = \kappa_A$. \square

Now, we call a poset P *poset-UM* if P contains a unique maximal element. For instance, Figure 2 shows that $P(3, 7)$ is poset-UM with unique maximal element 11 and that $P(3, 7, 11)$ is not poset-UM since it has both 4 and 8 as maximal elements. The following known lemma gives examples of posets that are poset-UM.

Lemma 2.3. *Suppose that a and b are relatively prime positive integers; then, $P(a, b)$ is poset-UM with maximal element $ab - a - b$. Equivalently, a positive integer is in $P(a, b)$ if and only if it is of the form $ab - ja - hb$ for some integers $j \in [1, b - 1]$ and $h \in [1, a - 1]$.*

The following proposition yields a preliminary necessary condition for a set of positive integers to be UM.

Proposition 2.4. *If a set of positive integers A is UM, then $P(A)$ is poset-UM.*

Proof. Suppose that $P(A)$ is not poset-UM but that A is UM. Consider the element $m \in P(A)$ of maximum magnitude; for instance, if $A = (3, 7, 11)$, then $m = 8$. Since $P(A)$ is not poset-UM, there is an order ideal $I \subseteq P(A)$ containing m but not equal to $P(A)$. By Lemma 2.1, there are A -cores $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ and $\kappa = \kappa_A = (\kappa_1, \kappa_2, \dots, \kappa_s)$ such that $\beta(\lambda) = I$ and $\beta(\kappa) = P(A)$. By Corollary 2.2, κ is the unique maximal element of $C(A)$; therefore, $\lambda \subset \kappa$. Hence, $m - r + 1 = \lambda_1 \leq \kappa_1 = m - s + 1$, which is a contradiction since $s = |P(A)| > |I| = r$. \square

As an application, the above proposition implies that $(3, 7, 11)$ is not UM. Observe that the converse of Proposition 2.4 does not always hold. For instance, suppose that $A = \{4, 5, 6\}$; then, $P(A) = \{1, 2, 3, 7\}$ is poset-UM with unique maximal element 7. Therefore the longest A -core is $\kappa_A = (3, 1, 1, 1)$, which does not contain the A -core $(2, 2)$ as a subpartition.

We will now classify all triples of positive integers (a, b, c) whose associated posets $P(a, b, c)$ are poset-UM. The following proposition forms a bijection between the maximal elements of $P(a, b, c)$ and the maximal elements of $P(d, e, f)$ for particular triples (a, b, c) and (d, e, f) .

Proposition 2.5. *Suppose that (a, b, c) is a triple of positive integers with $\gcd(a, b, c) = 1$; let $\gcd(a, b) = p$, $\gcd(a, c) = q$, $\gcd(b, c) = r$, and d, e, f be integers such that $(a, b, c) = (dpq, epr, fqr)$ as ordered triples. For m and n positive integers, $(m - 1)d + (n - 1)e - f$ is a maximal element of $P(d, e, f)$ if and only if $(mr - 1)dpq + (nq - 1)epr - fqr$ is a maximal element of $P(a, b, c)$.*

Proof. Let $s = (m - 1)d + (n - 1)e - f$ and $t = (mr - 1)dpq + (nq - 1)epr - fqr$. Suppose that s is a maximal element of $P(d, e, f)$; we will show that t is a maximal element of $P(a, b, c)$. Let us first verify that $t \in P(a, b, c)$. Suppose to the contrary that $t \in S(a, b, c)$. Then,

there are nonnegative integers h, i, j such that $(mr - 1)dpq + (nq - 1)epr - fqr = hdpq + iepr + jfqr$. Since $\gcd(a, b, c) = 1$, we have that $\gcd(r, p) = \gcd(r, q) = 1 = \gcd(r, d)$. The previous equality implies that $(mr - 1 - h)dpq$ is a multiple of r , which yields $h = rh' - 1$ for some positive integer h' . Similarly, $i = qi' - 1$ for some positive integer i' ; therefore, $(m - h')dp + (n - i')ep = (j + 1)f$. Thus, $j = j'p - 1$ for some positive integer j' ; hence $s = (m - 1)d + (n - 1)e - f = (h' - 1)d + (i' - 1)e + (j' - 1)f \in S(d, e, f)$, which is a contradiction.

Now, to see that t is maximal, it suffices to verify that $t + a, t + b, t + c \in S(a, b, c)$. Observe that $t + a = mdpqr + (nq - 1)epr - fqr = pqr(s + d) + epr(q - 1) + fqr(p - 1) \in S(dpq, epr, fqr)$ because $s + d \in S(d, e, f)$ by the maximality of s . By similar reasoning, $t + b \in S(a, b, c)$; since $t + c = (mr - 1)dpq + (nq - 1)epr \in S(a, b, c)$, it follows that t is a maximal element of $P(a, b, c)$.

This implies that t is a maximal element of $P(a, b, c)$ if s is a maximal element of $P(d, e, f)$. Through similar reasoning, one may show that s is a maximal element of $P(d, e, f)$ if t is a maximal element of $P(a, b, c)$. \square

The following corollary reduces the classification of triples (a, b, c) whose associated posets $P(a, b, c)$ are poset-UM to the case when a, b , and c are pairwise relatively prime.

Corollary 2.6. *Suppose that (a, b, c) is a triple of positive integers such that $\gcd(a, b, c) = 1$; let $\gcd(a, b) = p$, $\gcd(a, c) = q$, $\gcd(b, c) = r$, and d, e, f be integers such that $(a, b, c) = (dpq, epr, fqr)$ as ordered triples. Then, $P(a, b, c)$ is poset-UM if and only if $P(d, e, f)$ is poset-UM.*

Proof. Suppose that $P(d, e, f)$ is not poset-UM and let s_1 and s_2 be two distinct maximal elements of $P(d, e, f)$. Since s_1 and s_2 are maximal, we have that $s_1 + f, s_2 + f \in S(d, e)$; thus, there are positive integers m_1, n_1, m_2, n_2 such that $s_1 = (m_1 - 1)d + (n_1 - 1)e - f$ and $s_2 = (m_2 - 1)d + (n_2 - 1)e - f$. Let $t_1 = (rm_1 - 1)dpq + (qn_1 - 1)epr - fqr$ and $t_2 = (rm_2 - 1)dpq + (qn_2 - 1)epr - fqr$; by Proposition 2.5, t_1 and t_2 are maximal elements of $P(a, b, c)$. Since s_1 and s_2 are distinct, $t_1 \neq t_2$; thus $P(a, b, c)$ has two distinct maximal elements and is therefore not poset-UM.

Now, suppose that $P(a, b, c)$ is not poset-UM and let t_1 and t_2 be two distinct maximal elements of $P(a, b, c)$. As above, there are positive integers m'_1, n'_1, m'_2, n'_2 such that $t_1 = (m'_1 - 1)dpq + (n'_1 - 1)epr - fqr$ and $t_2 = (m'_2 - 1)dpq + (n'_2 - 1)epr - fqr$. We claim that m'_1 is a multiple of r . Indeed, since t_1 is maximal, $m'_1 dpq + (n'_1 - 1)epr - fqr = t_1 + a \in S(a, b, c)$; therefore, there are nonnegative integers h, i, j such that $m'_1 dpq + (n'_1 - 1)epr - fqr = hdpq + iepr + jfqr$. Since $t_1 \notin S(a, b, c)$, we have that $h = 0$. Therefore, r divides $m'_1 dpq$; the fact that $\gcd(d, r) = \gcd(p, r) = \gcd(q, r) = 1$ thus yields r divides m'_1 . Hence, there is an integer m_1 such that $m'_1 = m_1 r$; by similar reasoning, there are integers n_1, m_2, n_2 such that $n'_1 = n_1 q$, $m'_2 = m_2 r$, and $n'_2 = n_2 q$. By Proposition 2.5, $s_1 = (m_1 - 1)d + (n_1 - 1)e - f$ and $s_2 = (m_2 - 1)d + (n_2 - 1)e - f$ are unique maximal elements of $P(d, e, f)$. Since t_1 and t_2 are distinct, $s_1 \neq s_2$; this implies that $P(d, e, f)$ has two maximal elements and is therefore not poset-UM. \square

The following proposition, which we will prove in the next section, classifies all triples of pairwise relatively prime positive integers (a, b, c) such that $P(a, b, c)$ is poset-UM.

Proposition 2.7. *If $a < b < c$ are pairwise relatively prime positive integers, then $P(a, b, c)$ is poset-UM if and only if $c \in S(a, b)$.*

Assuming Proposition 2.7, we may classify all triples of positive integers (a, b, c) for which $P(a, b, c)$ is poset-UM.

Corollary 2.8. *Suppose that (a, b, c) is a triple of positive integers such that $\gcd(a, b, c) = 1$; let $p = \gcd(a, b)$, $q = \gcd(a, c)$, $r = \gcd(b, c)$, and d, e, f be integers such that $(a, b, c) = (dpq, epr, fqr)$ as ordered triples. Then, $P(a, b, c)$ is poset-UM if and only if (d, e, f) is primitive.*

Proof. This follows from Corollary 2.6 and Proposition 2.7. □

We may now establish Theorem 1.3.

Proof of Theorem 1.3. This follows from Proposition 2.4 and Corollary 2.8. □

Using Theorem 1.3, we may now establish Theorem 1.5.

Proof of Theorem 1.5. Let $f = md + ne$ for some nonnegative integers m and n . By Lemma 2.3, the unique maximal element of $P(d, e, f)$ is $de - d - e = (d+n-1)e + (m-1)d - f$. By Proposition 2.4, $P(A)$ has a unique maximal element; by Proposition 2.5, the unique maximal element of $P(A)$ is $(de + md + ne)pqr - dpq - epr - fqr = (de + f)pqr - a - b - c$. Hence, a positive integer is in $P(A)$ if and only if it is of the form $(de + f)pqr - ia - jb - kc$ for some positive integers i, j , and k . This implies the corollary since $\beta(\kappa_A) = P(A)$. □

3 Proof of Proposition 2.7

Throughout this section, we will adopt the notation given in Proposition 2.7. If $c \in S(a, b)$, then $P(a, b, c) = P(a, b)$ because $S(a, b, c) = S(a, b)$; therefore, the proposition follows from Lemma 2.3.

So, let us suppose that $c \notin S(a, b)$. By Lemma 2.3, there are positive integers $s_1 \in [1, b - 1]$ and $t_1 \in [1, a - 1]$ such that $c = ab - s_1a - t_1b$. Let k be the largest positive integer such that $ic \notin S(a, b)$ for each integer $i \in [1, k]$; observe that $k < b$.

By Lemma 2.3, there are integers $s_1, s_2, \dots, s_k \in [1, b - 1]$ and $t_1, t_2, \dots, t_k \in [1, a - 1]$ such that $ic = ab - s_i a - t_i b$ for each integer $i \in [1, k]$; moreover, let $(s_0, t_0) = (0, a)$ and $(s_{k+1}, t_{k+1}) = (b, 0)$. Observe that the s_i are distinct, since we would otherwise have that $(j - i)c = (t_i - t_j)b$ for some integers $1 \leq i < j \leq k$, which contradicts the facts that $k < b$ and $\gcd(b, c) = 1$; similarly, we see that the t_i are distinct.

By establishing properties about the s_i and t_i , we will show that $P(a, b, c)$ has at least two distinct maximal elements, which would imply that $P(a, b, c)$ is not UM. Let us begin with the following property.

Lemma 3.1. *For all integers $i, j \in [0, k + 1]$, we have that $t_i < t_j$ if and only if $s_i > s_j$.*

Proof. Suppose to the contrary that there exist integers $i, j \in [0, k + 1]$ such that $t_i < t_j$ and $s_i < s_j$. Then, $ic = ab - s_i a - t_i b > ab - s_j a - t_j b = jc$, which implies that $i > j$. Furthermore, $i \neq k + 1$ since $s_{k+1} = b \geq s_h$ for any integer $h \in [0, k + 1]$. Therefore, $i - j \in [1, k]$, so $(i - j)c \notin S(a, b)$ by the definition of k . However, $(i - j)c = (s_j - s_i)a + (t_j - t_i)b \in S(a, b)$, which is a contradiction. Therefore, the proposition holds. \square

Now, let $m, n \in [1, k]$ be integers such that $s_m = \min_{i \in [1, k]} s_i$ and $t_n = \min_{i \in [1, k]} t_i$. By Lemma 3.1, we have that $s_n = \max_{i \in [1, k]} s_i$ and $t_m = \max_{i \in [1, k]} t_i$.

Lemma 3.2. *We have that $k = m + n - 1$.*

Proof. We will first show that $k < m + n$. Suppose otherwise, so in particular $2ab - (s_m + s_n)a - (t_m + t_n)b = (m + n)c \notin S(a, b)$. Thus $2ab - (s_m + s_n)a - (t_m + t_n)b = ab - s_{m+n}a - t_{m+n}b$, so $ab = (s_m + s_n - s_{m+n})a + (t_m + t_n - t_{m+n})b$. Since $|s_m + s_n - s_{m+n}| < 2b$, $|t_m + t_n - t_{m+n}| < 2a$, and $\gcd(a, b) = 1$, this implies that either $s_m + s_n - s_{m+n} = b$ and $t_m + t_n = t_{m+n}$ or $s_m + s_n = s_{m+n}$ and $t_m + t_n - t_{m+n} = a$. Without loss of generality, suppose that the former holds; then $s_{m+n} = s_m + s_n - b < s_m$, which contradicts the minimality of s_m . Therefore, $k \leq m + n - 1$.

To see that $k \geq m + n - 1$, observe that $(m + n - i)c = ab - (s_m + s_n - s_i)a - (t_m + t_n - t_i)b$ for each integer $i \in [1, k]$. Since $0 < s_m \leq s_i \leq s_n$ and $0 < t_n \leq t_i \leq t_m$, we obtain that $(m + n - i)c$ is of the form $ab - ah - bj$ for some positive integers h and j . Thus Lemma 2.3 implies that $ic \notin S(a, b)$ for each integer $i \in [1, m + n - 1]$, which yields $k \geq m + n - 1$. Hence, $k = m + n - 1$. \square

Now let $\{r_0, r_1, \dots, r_{k+1}\}$ be a permutation of $\{0, 1, 2, \dots, k, k + 1\}$ such that $s_{r_0} < s_{r_1} < \dots < s_{r_{k+1}}$. By Lemma 3.1, we have that $t_{r_k} < t_{r_{k-1}} < \dots < t_{r_1}$. Furthermore, observe that $r_0 = 0$; $r_{k+1} = k + 1$; $r_1 = m$; and $r_k = n$.

Let $p_i = ab - (s_{r_i} + 1)a - (t_{r_{i+1}} + 1)b$ for each integer $i \in [0, k]$. Observe that the p_i are positive since $p_i \geq ab - s_{r_{i+1}}a - t_{r_{i+1}}b - b = r_{i+1}c - b$, which is positive since $b < c$. We will show that p_j and p_{j+1} are both maximal elements of $P(a, b, c)$ for some integer $j \in [0, k]$. However, let us first show the following properties about p_i for any integer $i \in [0, k]$.

Lemma 3.3. *For any integer $i \in [0, k]$, we have that $p_i \in P(a, b, c)$ but $p_i + a, p_i + b \notin P(a, b, c)$.*

Proof. Let us first show that the former claim holds; suppose to the contrary that $p_i \in S(a, b, c)$ for some integer $i \in [0, k]$. Then, there exist integers $f \in [0, b]$, $h \in [0, a]$, and $j \in [0, k]$ such that $ab - (s_{r_i} + 1)a - (t_{r_{i+1}} + 1)b = p_i = fa + hb + jc = ab - (s_j - f)a - (t_j - h)b$. Thus $s_j > s_{r_i}$ and $t_j > t_{r_{i+1}}$. Letting $j = r_h$ for some integer $h \in [0, k + 1]$, the former inequality implies that $h > i$ and the latter inequality implies that $h < i + 1$; this is a contradiction, which yields $p_i \in P(a, b, c)$.

Now, for the second claim, observe that $p_i + a = ab - s_{r_i}a - (t_{r_{i+1}} + 1)b = r_i c + (t_{r_i} - t_{r_{i+1}} - 1)b$, which is in $S(a, b, c)$ since $t_{r_i} > t_{r_{i+1}}$. Therefore, $p_i + a \notin P(a, b, c)$. By similar reasoning, $p_i + b \notin P(a, b, c)$. \square

Let k' be such that $r_{k'} = k$. We will show that p_j and p_{j+1} are distinct maximal elements of $P(a, b, c)$ with $j = k' - 1$. First, let us observe the following property about k' .

Lemma 3.4. *We have that $r_{k'-1} = k - m$ and $r_{k'+1} = k - n$.*

Proof. We will only show the former statement since the proof of the latter is similar. Observe that, for any integer $i \in [0, k']$, we have that $ab - s_{k-r_i}a - t_{k-r_i}b = (k - r_i)c = ab - (s_{r'_k} - s_{r_i})a - (a + t_{r_{k'}} - t_{r_i})b$. Since $s_{r_i} < s_{r_{k'}}$, we have that $s_{k-r_i} = s_{r_{k'}} - s_{r_i}$; this is minimal when $i = k' - 1$. Therefore, the minimality of s_m implies that $r_{k'-1} = k - m$. \square

Now we can establish the following lemma.

Lemma 3.5. *We have that $p_{k'-1}$ and $p_{k'}$ are maximal elements of $P(a, b, c)$.*

Proof. By Lemma 3.3, it suffices to show that $p_{k'-1} + c \notin P(a, b, c)$ and that $p_{k'} + c \notin P(a, b, c)$. We will only show the first statement, since the proof of the second is similar.

Suppose to the contrary that $p_{k'-1} + c \in P(a, b, c) \subset P(a, b)$. Then, Lemma 2.3 implies that there exist positive integers j and h such that $2ab - (s_{r_{k'-1}} + s_1 + 1)a - (t_{r_{k'}} + t_1 + 1)b = p_{k'-1} + c = ab - ja - hb$. Using Lemma 3.4 and the fact that $r_{k'} = k$, we find that $(s_{k-m} + s_1 + 1 - j)a + (t_k + t_1 + 1 - h)b = ab$. Therefore, we have that either $s_{k-m} + s_1 \geq b$ or $t_k + t_1 \geq a$.

By Lemma 3.2, we have that $k = m + n - 1$. Thus, $ab - s_{k-m}a - t_{k-m}b = (k - m)c = (n - 1)c = ab - (s_n - s_1)a - (a + t_n - t_1)b$, which implies that $s_{k-m} + s_1 = s_n < b$. Hence, $t_k + t_1 \geq a$.

We claim that $t_k + t_1 \leq a$ holds as well. Indeed, applying Lemma 3.2 again, we find that $ab - s_k a - t_k b = kc = (m + n - 1)c = ab - (s_m + s_n - s_1)a - (t_m + t_n - t_1)b$. Thus, $s_k = s_m + s_n - s_1$ and $t_k = t_m + t_n - t_1$. Moreover, since $ab - (s_m + s_n)a - (t_m + t_n - a)b = (m + n)c \in S(a, b)$, we have that $t_k + t_1 = t_m + t_n \leq a$ by Lemma 2.3.

Therefore, $t_m + t_n = t_k + t_1 = a$ and hence $a(b - s_m - s_n) = (m + n)c$. Since a and c are relatively prime, $b - s_m - s_n$ is a positive multiple of c ; this contradicts the fact that $b < c$. Thus, $p_{k'-1} + c \notin P(a, b, c)$. \square

Using Lemma 3.5, we may establish Proposition 2.7.

Proof of Proposition 2.7. If $c \in S(a, b)$, then $P(a, b, c) = P(a, b)$ because $S(a, b, c) = S(a, b)$; therefore, the proposition follows from Lemma 2.3. If $c \notin S(a, b)$, then $P(a, b, c)$ has two distinct maximal elements by Lemma 3.5, which implies that $P(a, b, c)$ cannot be UM. Therefore, $P(a, b, c)$ is UM if and only if $c \in S(a, b)$. \square

Acknowledgements

This research was conducted under the supervision of Joe Gallian at the University of Minnesota Duluth REU, funded by NSF Grant 1358659 and NSA Grant H98230-13-1-0273. The author heartily thanks Matt Fayers, Rishi Nath, and Joe Gallian for suggesting the topic of this project and for their valuable advice; the author also thanks David Moulton for his insightful discussions and Aaron Abrams and the referees for their suggestions.

References

- [1] A. Aggarwal. Armstrong’s conjecture for $(k, mk + 1)$ -core partitions. *European J. Combin.*, 47:54–67, 2015.
- [2] T. Amdeberhan and E. Leven. Multi-cores, posets, and lattice paths. Preprint, 2014. [arXiv:1406.2250v2](#)
- [3] J. Anderson. Partitions which are simultaneously t_1 - and t_2 -core. *Discrete Math.*, 248:237–243, 2002.
- [4] D. Armstrong, C. Hanusa, and B. Jones. Results and conjectures on simultaneous core partitions. *European J. Combin.*, 41:205–220, 2014.
- [5] D. Auckerman, B. Kane, and L. Sze. On simultaneous s -cores/ t -cores. *Discrete Math.*, 309:2712–2720, 2009.
- [6] W. Chen, H. Huang, and L. Wang. Average size of a self-conjugate (s, t) -core partition. Preprint, 2014. [arXiv:1405.2175v1](#)
- [7] M. Fayers, Personal communication.
- [8] M. Fayers. The t -core of an s -core. *J. Combin. Theory Ser. A*, 118:1525–1539, 2011.
- [9] S. Fishel and M. Vazirani. A bijection between dominant Shi regions and core partitions. *European J. Combin.*, 31:2087–2101, 2010.
- [10] P. Johnson. Lattice points and simultaneous core partitions. Preprint, 2015. [arXiv:1502.07934v1](#)
- [11] J. Olsson. A theorem on the cores of partitions. *J. Combin. Theory Ser. A*, 116:733–740, 2009.
- [12] J. Olsson and D. Stanton. Block inclusions and cores of partitions. *Aequationes Math.*, 74:90–110, 2007.
- [13] R. Stanley. *Enumerative Combinatorics*. Volume 1. Second edition. Cambridge Studies in Advanced Mathematics, 49. Cambridge University Press, Cambridge, 2012.
- [14] R. Stanley. *Enumerative Combinatorics*. Volume 2. Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999.
- [15] R. Stanley and F. Zanello. The Catalan case of Armstrong’s conjecture on simultaneous core partitions. *SIAM J. Discrete Math.*, 29:658–666, 2015.
- [16] J. Vandehey. Containment in (s, t) -core partitions. Preprint, 2008. [arXiv:0809.2134](#)
- [17] J. Yang, M. Zhong, and R. Zhou. On the enumeration of $(s, s+1, s+2)$ -core partitions. Preprint, 2014. [arXiv:1406.2583v1](#)