



Computer Science and Artificial Intelligence Laboratory

Technical Report

MIT-CSAIL-TR-2016-005

May 23, 2016

Deep Learning without Poor Local Minima

Kenji Kawaguchi

Deep Learning without Poor Local Minima

Kenji Kawaguchi
Massachusetts Institute of Technology
kawaguch@mit.edu

Abstract

In this paper, we prove a conjecture published in 1989 and also partially address an open problem announced at the Conference on Learning Theory (COLT) 2015. For an expected loss function of a deep nonlinear neural network, we prove the following statements under the independence assumption adopted from recent work: 1) the function is non-convex and non-concave, 2) every local minimum is a global minimum, 3) every critical point that is not a global minimum is a saddle point, and 4) the property of saddle points differs for shallow networks (with three layers) and deeper networks (with more than three layers). Moreover, we prove that the same four statements hold for deep linear neural networks with any depth, any widths and no unrealistic assumptions. As a result, we present an instance, for which we can answer to the following question: how difficult to directly train a deep model in theory? It is more difficult than the classical machine learning models (because of the non-convexity), but not too difficult (because of the nonexistence of poor local minima and the property of the saddle points). We note that even though we have advanced the theoretical foundations of deep learning, there is still a gap between theory and practice.

1 Introduction

Deep learning has been a great practical success in many fields, including the fields of computer vision, machine learning, and artificial intelligence. In addition to its practical success, theoretical results have shown that deep learning is attractive in terms of its generalization properties (Livni *et al.*, 2014; Mhaskar *et al.*, 2016). That is, deep learning introduces good function classes that may have a low capacity in the VC sense while being able to represent target functions of interest well. However, deep learning requires us to deal with seemingly intractable optimization problems. Typically, training of a deep model is conducted via non-convex optimization. Because finding a global minimum of a *general* non-convex function is an NP-complete problem (Murty & Kabadi, 1987), a hope is that a function induced by a deep model has some structure that makes the non-convex optimization tractable. Unfortunately, it was shown in 1992 that training a very simple neural network is indeed NP-hard (Blum & Rivest, 1992). In the past, such theoretical concerns in optimization played a major role in shrinking the field of deep learning. That is, many researchers instead favored classical machine learning models (with or without a kernel approach) that require only convex optimization. While the recent great practical successes have revived the field, we do not yet know what makes optimization in deep learning tractable in theory.

In this paper, as a step toward establishing the optimization theory for deep learning, we prove a conjecture noted in (Goodfellow *et al.*, 2016) for deep *linear* networks, and also address an open problem announced in (Choromanska *et al.*, 2015b) for deep *nonlinear* networks. Moreover, for both the conjecture and the open problem, we prove more general and tighter statements than those previously given.

2 Deep linear neural networks

Given the absence of a theoretical understanding of deep nonlinear neural networks, Goodfellow *et al.* (2016) noted that it is beneficial to theoretically analyze the loss functions of simpler models, i.e., *linear* neural networks. The function class of a linear neural network only contains functions that are linear with respect to inputs. However, their loss functions are non-convex in the weight parameters and thus nontrivial. Saxe *et al.* (2014) empirically showed that the optimization of deep *linear* models exhibits similar properties to those of the optimization of deep *nonlinear* models. Ultimately, for theoretical development, it is natural to start with linear models before working with nonlinear models (Baldi & Lu, 2012), and yet even for linear models, the understanding is scarce when the models become deep.

2.1 Model and notation

We begin by defining the notation. Let H be the number of hidden layers, and let (X, Y) be the training data set, with $Y \in \mathbb{R}^{d_y \times m}$ and $X \in \mathbb{R}^{d_x \times m}$, where m is the number of data points. Here, $d_y \geq 1$ and $d_x \geq 1$ are the number of components (or dimensions) of the outputs and inputs, respectively. We denote the model (weight) parameters by W , which consists of parameter matrices corresponding to each layer: $W_{H+1} \in \mathbb{R}^{d_y \times d_H}, \dots, W_k \in \mathbb{R}^{d_k \times d_{k-1}}, \dots, W_1 \in \mathbb{R}^{d_1 \times d_x}$. Here, d_k represents the width of the k -th layer, where the 0-th layer is the input layer and the $(H+1)$ -th layer is the output layer (i.e., $d_0 = d_x$ and $d_{H+1} = d_y$). Let I_{d_k} be the $d_k \times d_k$ identity matrix. Let $p = \min(d_H, \dots, d_1)$ be the smallest width of a hidden layer. We denote the (j, i) -th entry of a matrix M by $M_{j,i}$. We also denote the j -th row vector of M by $M_{j,\cdot}$ and the i -th column vector of M by $M_{\cdot,i}$.

We can then write the output of a feedforward deep linear model, $\bar{Y}(W, X) \in \mathbb{R}^{d_y \times m}$, as

$$\bar{Y}(W, X) = W_{H+1}W_HW_{H-1} \cdots W_2W_1X.$$

We consider one of the most widely used loss functions, squared error loss:

$$\bar{\mathcal{L}}(W) = \frac{1}{2} \sum_{i=1}^m \|\bar{Y}(W, X)_{\cdot,i} - Y_{\cdot,i}\|_2^2 = \frac{1}{2} \|\bar{Y}(W, X) - Y\|_F^2,$$

where $\|\cdot\|_F$ is the Frobenius norm. Note that $\frac{2}{m}\bar{\mathcal{L}}(W)$ is the usual *mean* squared error, for which all of our theorems hold as well, since multiplying $\bar{\mathcal{L}}(W)$ by a constant in W results in an equivalent optimization problem.

2.2 Background

Recently, Goodfellow *et al.* (2016) remarked that when Baldi & Hornik (1989) stated and proved Proposition 2.1 for shallow linear networks, they also stated Conjecture 2.2 for deep linear networks.

Proposition 2.1 (Baldi & Hornik, 1989: *shallow* linear network) *Assume that $H = 1$ (i.e., $\bar{Y}(W, X) = W_2W_1X$), assume that XX^T and XY^T are invertible, and assume that $p < d_x$, $p < d_y$ and $d_y = d_x$ (e.g., an autoencoder). Then, the loss function $\bar{\mathcal{L}}(W)$ has the following properties:*

- (i) *It is convex in each matrix W_1 (or W_2) when the other W_2 (or W_1) is fixed.*
- (ii) *Every local minimum is a global minimum.*

Conjecture 2.2 (Baldi & Hornik, 1989: *deep* linear network) *Assume the same set of conditions as in Proposition 2.1 except for $H = 1$. Then, the loss function $\bar{\mathcal{L}}(W)$ has the following properties:*

- (i) *For any $k \in \{1, \dots, H+1\}$, it is convex in each matrix W_k when for all $k' \neq k$, $W_{k'}$ is fixed.*
- (ii) *Every local minimum is a global minimum.*

Baldi & Lu (2012) recently provided a proof for Conjecture 2.2 (i), leaving the proof of Conjecture 2.2 (ii) for future work. They also noted that the case of $p \geq d_x = d_x$ is of interest, but requires further analysis, even for a shallow network with $H = 1$. An informal discussion of Conjecture 2.2 can be found in (Baldi, 1989). In Appendix D in the supplementary material, we provide a more detailed discussion of this subject.

2.3 Results

We now state our main theoretical results for deep linear networks, which imply Conjecture 2.2 (ii) as well as obtain further information regarding the critical points with more generality.

Theorem 2.3 (Loss surface of deep linear networks with more generality) *Assume that XX^T and XY^T are full rank. Further, assume that $d_y \leq d_x$. Then, for any depth $H \geq 1$ and for any layer widths and any input-output dimensions $d_y, d_H, d_{H-1}, \dots, d_1, d_x$ (the widths can arbitrarily differ from each other and from d_y and d_x), the loss function $\tilde{\mathcal{L}}(W)$ has the following properties:*

- (i) *It is non-convex and non-concave.*
- (ii) *Every local minimum is a global minimum.*
- (iii) *Every critical point that is not a global minimum is a saddle point.*
- (iv) *If $\text{rank}(W_H \cdots W_2) = p$, then the Hessian at any saddle point has at least one (strictly) negative eigenvalue.¹*

Corollary 2.4 (Effect of deepness on the loss surface) *Assume the same set of conditions as in Theorem 2.3 and consider the loss function $\tilde{\mathcal{L}}(W)$. For three-layer networks (i.e., $H = 1$), the Hessian at any saddle point has at least one (strictly) negative eigenvalue. In contrast, for networks deeper than three layers (i.e., $H \geq 2$), there exist saddle points at which the Hessian does not have any negative eigenvalue.*

The full rank assumptions on XX^T and XY^T in Theorem 2.3 are realistic and practically easy to satisfy, as discussed in previous work (e.g., Baldi & Hornik, 1989). In contrast to related previous work (Baldi & Hornik, 1989; Baldi & Lu, 2012), we do not assume the invertibility of XY^T , $p < d_x$, $p < d_y$ nor $d_y = d_x$. In Theorem 2.3, $p \geq d_x$ is allowed, as well as many other relationships among the widths of the layers. Therefore, Theorem 2.3 (ii) implies Conjecture 2.2 (ii) and is more general than Conjecture 2.2 (ii). Moreover, Theorem 2.3 (iv) and Corollary 2.4 provide additional information regarding the important properties of saddle points.

Theorem 2.3 presents an instance of a deep model that is not too difficult to train with direct greedy optimization, such as gradient-based methods. If there are “bad” local minima with large loss values everywhere, we would have to search the entire space,² the volume of which increases exponentially with the number of variables. This is a major cause of NP-hardness for non-convex optimization. In contrast, if there are no poor local minima as Theorem 2.3 (ii) states, then saddle points are the remaining concern in terms of tractability.³ Because the Hessian of $\tilde{\mathcal{L}}(W)$ is Lipschitz continuous, if the Hessian at a saddle point has a negative eigenvalue, it starts appearing as we approach the saddle point. Thus, Theorem 2.3 and Corollary 2.4 suggest that for 1-hidden layer networks, training can be done in polynomial time with a second order method or even with a modified stochastic gradient decent method, as discussed in (Ge et al., 2015). For deeper networks, Corollary 2.4 states that there exist “bad” saddle points in the sense that the Hessian at the point has no negative eigenvalue. However, from Theorem 2.3 (iv), we know exactly when this can happen, and from the proof of Theorem 2.3, we see that some perturbation is sufficient to escape such bad saddle points.

3 Deep nonlinear neural networks

Given this understanding of the loss surface of deep linear models, we discuss deep nonlinear models.

3.1 Model

We use the same notation as for the deep linear models, defined in the beginning of Section 2.1. The output of deep nonlinear neural network, $\hat{Y}(W, X) \in \mathbb{R}^{d_y \times m}$, is defined as

$$\hat{Y}(W, X) = q\sigma_{H+1}(W_{H+1}\sigma_H(W_H\sigma_{H-1}(W_{H-1}\cdots\sigma_2(W_2\sigma_1(W_1X))\cdots))),$$

¹If $H = 1$, to be succinct, we define $W_H \cdots W_2 = W_1 \cdots W_2 \triangleq I_{d_1}$, with a slight abuse of notation.

²Typically, we do this by assuming smoothness in the values of the loss function.

³Other problems such as the ill-conditioning can make it difficult to obtain a fast convergence rate.

where $q \in \mathbb{R}$ is simply a normalization factor, the value of which is specified later. Here, $\sigma_k : \mathbb{R}^{d_k \times m} \rightarrow \mathbb{R}^{d_k \times m}$ is the element-wise rectified linear function:

$$\sigma_k \left(\begin{bmatrix} b_{11} & \cdots & b_{1m} \\ \vdots & \ddots & \vdots \\ b_{d_k 1} & \cdots & b_{d_k m} \end{bmatrix} \right) = \begin{bmatrix} \bar{\sigma}(b_{11}) & \cdots & \bar{\sigma}(b_{1m}) \\ \vdots & \ddots & \vdots \\ \bar{\sigma}(b_{d_k 1}) & \cdots & \bar{\sigma}(b_{d_k m}) \end{bmatrix},$$

where $\bar{\sigma}(b_{ij}) = \max(0, b_{ij})$. In practice, we usually set σ_{H+1} to be an identity map in the last layer, in which case all our theoretical results still hold true.

3.2 Background

Following the work by Dauphin *et al.* (2014), Choromanska *et al.* (2015a) investigated the connection between the loss functions of deep nonlinear networks and a function well-studied via random matrix theory (i.e., the Hamiltonian of the spherical spin-glass model). They explained that their theoretical results relied on several *unrealistic* assumptions. Later, Choromanska *et al.* (2015b) suggested at the Conference on Learning Theory (COLT) 2015 that discarding these assumptions is an important open problem. The assumptions were labeled A1p, A2p, A3p, A4p, A5u, A6u, and A7p.

Here, we discuss the most relevant assumptions: A1p, A5u, and A6u. We refer to the part of assumption A1p (resp. A5u) that corresponds only to the *model* assumption as A1p-m (resp. A5u-m). Note that assumptions A1p-m and A5u-m are explicitly used in the previous work (Choromanska *et al.*, 2015a) and included in A1p and A5u (i.e., we are *not* making new assumptions here). As the model $\hat{Y}(W, X) \in \mathbb{R}^{d_y \times m}$ represents a directed acyclic graph, we can express an output from one of the units in the output layer as

$$\hat{Y}(W, X)_{j,i} = q \sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} [Z_i]_{(j,p)} \prod_{k=1}^{H+1} w_{(j,p)}^{(k)},$$

where Ψ_j is the total number of paths from the inputs to the j -th output in the directed acyclic graph. In addition, $[X_i]_{(j,p)} \in \mathbb{R}$ represents the entry of the i -th sample input datum that is used in the p -th path of the j -th output. For each layer k , $w_{(j,p)}^{(k)} \in \mathbb{R}$ is the entry of W_k that is used in the p -th path of the j -th output. Finally, $[Z_i]_{(j,p)} \in \{0, 1\}$ represents whether the p -th path of the j -th output is active ($[Z_i]_{(j,p)} = 1$) or not ($[Z_i]_{(j,p)} = 0$) for each sample i because of the rectified linear activation.

Assumption A1p-m assumes that the Z 's are Bernoulli random variables with the same probability of success, $\Pr([Z_i]_{(j,p)} = 1) = \rho$ for all i and (j, p) . Assumption A5u-m assumes that the Z 's are independent from the input X 's, parameters w 's, and each other (the independence was required, for example, in the first equation of the proof of Theorem 3.3 in (Choromanska *et al.*, 2015a)). With assumptions A1p-m and A5u-m, we can write $\mathbb{E}_Z[\hat{Y}(W, X)_{j,i}] = q \sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \rho \prod_{k=1}^H w_{(j,p)}^{(k)}$.

The previous work also assumes the use of ‘‘independent random’’ loss functions. Consider the hinge loss, $\mathcal{L}_{\text{hinge}}(W)_{j,i} = \max(0, 1 - Y_{j,i} \hat{Y}(W, X)_{j,i})$. By modeling the max operator as a Bernoulli random variable ξ , we can then write $\mathcal{L}_{\text{hinge}}(W)_{j,i} = \xi - q \sum_{p=1}^{\Psi_j} Y_{j,i} [X_i]_{(j,p)} \xi [Z_i]_{(j,p)} \prod_{k=1}^{H+1} w_{(j,p)}^{(k)}$. A1p then assumes that for all i and (j, p) , the $\xi [Z_i]_{(j,p)}$ are Bernoulli random variables with equal probabilities of success. Furthermore, A5u assumes that the independence of $\xi [Z_i]_{(j,p)}$, $Y_{j,i} [X_i]_{(j,p)}$, and $w_{(j,p)}$. Finally, A6u assumes that $Y_{j,i} [X_i]_{(j,p)}$ for all (j, p) and i are independent.

Proposition 3.1 (High-level description of a main result in Choromanska *et al.*, 2015a) *Assume A1p (including A1p-m), A2p, A3p, A4p, A5u (including A5u-m), A6u, and A7p (Choromanska et al., 2015b). Furthermore, assume that $d_y = 1$. Then, the expected loss of each sample datum, $\mathbb{E}_{\xi, Z}[\mathcal{L}_{\text{hinge}}(W)_{i,1}]$, has the following property: above a certain loss value, the number of local minima diminishes exponentially as the loss value increases.*

Choromanska *et al.* (2015b) noted that A6u is unrealistic because it implies that the inputs are not shared among the paths. In addition, A5u is unrealistic because it implies that the activation of any path is independent of the input data.

3.3 Results

We now state our main theoretical results for deep nonlinear networks, which partially address the aforementioned open problem and lead to more general and tighter results. Unlike the pre-

vious work, we do not assume that we can take the expectation over random variable ξ . Moreover, we consider loss functions for all the data points and all possible output dimensionalities (i.e., vectored-valued output). More concretely, we consider the expected squared error loss, $\mathbb{E}_Z[\mathcal{L}(W)] = \mathbb{E}_Z[\frac{1}{2}\|\hat{Y}(W, X) - Y\|_F^2]$. We also consider the squared error loss of the expected model, $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W) = \frac{1}{2}\|E[\hat{Y}(W, X)] - Y\|_F^2$.

Theorem 3.2 (Loss surface of deep nonlinear networks) *Assume A1p-m and A5u-m. Further assume that $d_y \leq d_x$ and that XX^T and XY^T are full rank. Let $q = \rho^{-1}$. Then, for any depth $H \geq 1$ and for any layer widths and any input-output dimensions $d_y, d_H, d_{H-1}, \dots, d_1, d_x$ (the widths can arbitrarily differ from each other and from d_y and d_x), both the expected loss function $\mathbb{E}_Z[\mathcal{L}(W)]$ and the loss function of the expected model $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$ have the following properties:*

- (i) *They are non-convex and non-concave.*
- (ii) *Every local minimum is a global minimum.*
- (iii) *Every critical point that is not a global minimum is a saddle point.*
- (iv) *If $\text{rank}(W_H \cdots W_2) = p$, then the Hessian at any saddle point has at least one (strictly) negative eigenvalue.⁴*

Corollary 3.3 (Effect of deepness on the loss surface) *Assume the same set of conditions as in Theorem 3.2. Consider the loss function $\mathbb{E}_Z[\mathcal{L}(W)]$ or $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$. Then, for three-layer networks (i.e., $H = 1$), the Hessian at any saddle point has some (strictly) negative eigenvalue. In contrast, for networks deeper than three layers (i.e., $H \geq 2$), there exist saddle points at which the Hessian does not have a negative eigenvalue.*

Comparing Theorem 3.2 and Proposition 3.1, we can see that we successfully discarded assumptions A2p, A3p, A4p, A6u, and A7p while obtaining a tighter statement in general. Again, note that the full rank assumptions on XX^T and XY^T in Theorem 3.2 are realistic and practically easy to satisfy, as discussed in previous work (e.g., Baldi & Hornik, 1989). Furthermore, our model \hat{Y} is strictly more general than the model analyzed in (Choromanska *et al.*, 2015a,b) (i.e., this paper’s model class contains the previous work’s model class but not vice versa).

4 Important lemmas

In this section, we provide additional theoretical results as lemmas that lead to further insights. The proofs of the lemmas are in the appendix in the supplementary material.

Let $M \otimes M'$ be the Kronecker product of M and M' . Let $\mathcal{D}_{\text{vec}(W_k^T)} f(\cdot) = \frac{\partial f(\cdot)}{\partial \text{vec}(W_k^T)}$ be the partial derivative of f with respect to $\text{vec}(W_k^T)$ in the numerator layout. That is, if $f : \mathbb{R}^{d_{in}} \rightarrow \mathbb{R}^{d_{out}}$, we have $\mathcal{D}_{\text{vec}(W_k^T)} f(\cdot) \in \mathbb{R}^{d_{out} \times (d_k d_{k-1})}$. Let $\mathcal{R}(M)$ be the range (or the column space) of a matrix M . Let M^- be any generalized inverse of M . When we write a generalized inverse in a condition or statement, we mean it for any generalized inverse (i.e., we omit the universal quantifier over generalized inverses, as this is clear). Let $r = (\bar{Y}(W, X) - Y)^T \in \mathbb{R}^{m \times d_y}$ be an error matrix. Let $C = W_{H+1} \cdots W_2 \in \mathbb{R}^{d_y \times d_1}$. When we write $W_k \cdots W'_k$, we generally intend that $k > k'$ and the expression denotes a product over W_j for integer $k \geq j \geq k'$. For notational compactness, two additional cases can arise: when $k = k'$, the expression denotes simply W_k , and when $k < k'$, it denotes I_{d_k} . For example, in the statement of Lemma 4.1, if we set $k := H + 1$, we have that $W_{H+1} W_H \cdots W_{H+2} \triangleq I_{d_y}$.

In Lemma 4.6 and the proofs of Theorems 2.3 and 3.2, we use the following additional notation. Let $\Sigma = YX^T(XX^T)^{-1}XY^T$ and its eigendecomposition be $U\Lambda U^T = \Sigma$, where the entries of the eigenvalues are ordered as $\Lambda_{1,1} \geq \dots \geq \Lambda_{d_y, d_y}$ with corresponding orthogonal eigenvector matrix $U = [u_1, \dots, u_{d_y}]$. For each $k \in \{1, \dots, d_y\}$, $u_k \in \mathbb{R}^{d_y \times 1}$ is a column eigenvector. As Σ is real symmetric, we can always make U orthogonal. Let $\bar{p} = \text{rank}(C) \in \{1, \dots, \min(d_y, p)\}$. We define a matrix containing the subset of the \bar{p} largest eigenvectors as $U_{\bar{p}} = [u_1, \dots, u_{\bar{p}}]$. Given any ordered

⁴If $H = 1$, to be succinct, we define $W_H \cdots W_2 = W_1 \cdots W_2 \triangleq I_{d_1}$, with a slight abuse of notation.

set $\mathcal{I}_{\bar{p}} = \{i_1, \dots, i_{\bar{p}} \mid 1 \leq i_1 < \dots < i_{\bar{p}} \leq \min(d_y, p)\}$, we define a matrix containing the subset of the corresponding eigenvectors as $U_{\mathcal{I}_{\bar{p}}} = [u_{i_1}, \dots, u_{i_{\bar{p}}}]$. Note the difference between $U_{\bar{p}}$ and $U_{\mathcal{I}_{\bar{p}}}$.

Lemma 4.1 (Critical point necessary and sufficient condition) *W is a critical point of $\bar{\mathcal{L}}(W)$ if and only if for all $k \in \{1, \dots, H+1\}$,*

$$\left(\mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W)\right)^T = (W_{H+1}W_H \cdots W_{k+1} \otimes (W_{k-1} \cdots W_2W_1X)^T)^T \text{vec}(r) = 0.$$

Lemma 4.2 (Representation at critical point) *If W is a critical point of $\bar{\mathcal{L}}(W)$, then*

$$W_{H+1}W_H \cdots W_2W_1 = C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1}.$$

Lemma 4.3 (Block Hessian with Kronecker product) *Write the entries of $\nabla^2 \bar{\mathcal{L}}(W)$ in a block form as*

$$\nabla^2 \bar{\mathcal{L}}(W) = \begin{bmatrix} \mathcal{D}_{\text{vec}(W_{H+1}^T)} \left(\mathcal{D}_{\text{vec}(W_{H+1}^T)} \bar{\mathcal{L}}(W)\right)^T & \cdots & \mathcal{D}_{\text{vec}(W_1^T)} \left(\mathcal{D}_{\text{vec}(W_{H+1}^T)} \bar{\mathcal{L}}(W)\right)^T \\ \vdots & \ddots & \vdots \\ \mathcal{D}_{\text{vec}(W_{H+1}^T)} \left(\mathcal{D}_{\text{vec}(W_1^T)} \bar{\mathcal{L}}(W)\right)^T & \cdots & \mathcal{D}_{\text{vec}(W_1^T)} \left(\mathcal{D}_{\text{vec}(W_1^T)} \bar{\mathcal{L}}(W)\right)^T \end{bmatrix}.$$

Then, for any $k \in \{1, \dots, H+1\}$,

$$\begin{aligned} & \mathcal{D}_{\text{vec}(W_k^T)} \left(\mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W)\right)^T \\ &= ((W_{H+1} \cdots W_{k+1})^T (W_{H+1} \cdots W_{k+1}) \otimes (W_{k-1} \cdots W_1 X) (W_{k-1} \cdots W_1 X)^T), \end{aligned}$$

and, for any $k \in \{2, \dots, H+1\}$,

$$\begin{aligned} & \mathcal{D}_{\text{vec}(W_k^T)} \left(\mathcal{D}_{\text{vec}(W_1^T)} \bar{\mathcal{L}}(W)\right)^T \\ &= (C^T (W_{H+1} \cdots W_{k+1}) \otimes X (W_{k-1} \cdots W_1 X)^T) + \\ & \quad [(W_{k-1} \cdots W_2)^T \otimes X] [I_{d_{k-1}} \otimes (r W_{H+1} \cdots W_{k+1})_{\cdot, 1} \cdots I_{d_{k-1}} \otimes (r W_{H+1} \cdots W_{k+1})_{\cdot, d_k}]. \end{aligned}$$

Lemma 4.4 (Hessian semidefinite necessary condition) *If $\nabla^2 \bar{\mathcal{L}}(W)$ is positive semidefinite or negative semidefinite at a critical point, then for any $k \in \{2, \dots, H+1\}$,*

$$\mathcal{R}((W_{k-1} \cdots W_3 W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad X r W_{H+1} W_H \cdots W_{k+1} = 0.$$

Corollary 4.5 *If $\nabla^2 \bar{\mathcal{L}}(W)$ is positive semidefinite or negative semidefinite at a critical point, then for any $k \in \{2, \dots, H+1\}$,*

$$\text{rank}(W_{H+1} W_H \cdots W_k) \geq \text{rank}(W_{k-1} \cdots W_3 W_2) \quad \text{or} \quad X r W_{H+1} W_H \cdots W_{k+1} = 0.$$

Lemma 4.6 (Hessian positive semidefinite necessary condition) *If $\nabla^2 \bar{\mathcal{L}}(W)$ is positive semidefinite at a critical point, then*

$$C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}^T \quad \text{or} \quad X r = 0.$$

5 Proof sketches of theorems

We now provide overviews of the proofs of Theorems 2.3 and 3.2. We complete the proofs of the theorems in the appendix in the supplementary material.

Our proof approach largely differs from those in previous work (Baldi & Hornik, 1989; Baldi & Lu, 2012; Choromanska *et al.*, 2015a,b). In contrast to (Baldi & Hornik, 1989; Baldi & Lu, 2012), we need a different approach to deal with the ‘‘bad’’ saddle points that start appearing when the model becomes deeper (see Section 2.3), as well as to obtain more comprehensive properties of the critical points with more generality. While the previous proofs heavily rely on the first-order information, the main parts of our proofs take advantage of the second order information. In contrast, Choromanska *et al.* (2015a,b) used the seven assumptions to relate the loss functions of deep models to

a function previously analyzed with a tool of random matrix theory (i.e., Gaussian orthogonal ensemble). With no reshaping assumptions (A3p, A4p, and A6u), we cannot relate our loss function to such a function. Moreover, with no distributional assumptions (A2p and A6u) (except the activation), our Hessian is deterministic, and therefore, even random matrix theory itself is insufficient for our purpose. Furthermore, with no spherical constraint assumption (A7p), the number of local minima in our loss function can be uncountable.

One natural strategy to proceed toward Theorems 2.3 and 3.2 would be to use the first order and the second order necessary conditions of local minima (e.g., the gradient is zero and the Hessian is positive semidefinite).⁵ However, are the first-order and second-order conditions sufficient to prove Theorems 2.3 and 3.2? Corollaries 2.4 and 3.3 show that the answer is negative for *deep* models with $H \geq 2$, while it is affirmative for shallow models with $H = 1$. Thus, for deep models, a simple use of the first-order and second-order information is insufficient to characterize the properties of each critical point. In addition to the complexity of the Hessian of the *deep* models, this suggests that we must strategically extract the second order information. Accordingly, we obtained an organized representation of the Hessian in Lemma 4.3 and strategically extracted the information in Lemmas 4.4 and 4.6, with which we are ready to prove Theorems 2.3 and 3.2.

5.1 Proof sketch of Theorem 2.3 (ii)

By case analysis, we show that any point that satisfies the necessary conditions and the definition of a local minimum is a global minimum.

Case I: $\text{rank}(W_H \cdots W_2) = p$ and $d_y \leq p$: Assume that $\text{rank}(W_H \cdots W_2) = p$. If $d_y < p$, Corollary 4.5 with $k = H + 1$ implies the necessary condition that $Xr = 0$. If $d_y = p$, Lemma 4.6 with $k = H + 1$ and $k = 2$, combined with the fact that $\mathcal{R}(C) \subseteq \mathcal{R}(YX^T)$, implies the necessary condition that $Xr = 0$. Therefore, we have the necessary condition, $Xr = 0$. Interpreting condition $Xr = 0$, we conclude that W achieving $Xr = 0$ is indeed a global minimum.

Case II: $\text{rank}(W_H \cdots W_2) = p$ and $d_y > p$: From Lemma 4.6, we have the necessary condition that $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}^T$ or $Xr = 0$. If $Xr = 0$, using the exact same proof as in Case I, it is a global minimum. Suppose then that $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}^T$. From Lemma 4.4 with $k = H + 1$, we conclude that $\bar{p} \triangleq \text{rank}(C) = p$. Then, from Lemma 4.2, we write $W_{H+1} \cdots W_1 = U_p U_p^T Y X^T (X X^T)^{-1}$, which is the orthogonal projection onto the subspace spanned by the p eigenvectors corresponding to the p largest eigenvalues following the ordinary least square regression matrix. This is indeed the expression of a global minimum.

Case III: $\text{rank}(W_H \cdots W_2) < p$: Suppose that $\text{rank}(W_H \cdots W_2) < p$. From Lemma 4.4, we have the following necessary condition for the Hessian to be (positive or negative) semidefinite at a critical point: for any $k \in \{2, \dots, H + 1\}$,

$$\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad Xr W_{H+1} \cdots W_{k+1} = 0,$$

where the first condition is shown to imply $\text{rank}(W_{H+1} \cdots W_k) \geq \text{rank}(W_{k-1} \cdots W_2)$ in Corollary 4.5. We repeatedly apply these conditions for $k = 2, \dots, H + 1$ to claim that with arbitrarily small $\epsilon > 0$, we can perturb each parameter (i.e., each entry of W_H, \dots, W_2) such that $\text{rank}(W_{H+1} \cdots W_2) \geq \min(p, d_x)$ without changing the value of $\tilde{\mathcal{L}}(W)$. We prove this by induction on k , using Lemmas 4.2, 4.4, and 4.6.

We consider the base case, $k = 2$. From the condition with $k = 2$ of Lemma 4.4, we have that $\text{rank}(W_{H+1} \cdots W_2) \geq d_1 \geq p$ or $Xr W_{H+1} \cdots W_3 = 0$ (note that $d_1 \geq p \geq \bar{p}$ by their definitions). The former condition is false since $\text{rank}(W_{H+1} \cdots W_2) \leq \text{rank}(W_H \cdots W_2) < p$. From the latter condition, for an arbitrary L_2 , with $A_2 = W_{H+1} \cdots W_3$,

$$\begin{aligned} 0 &= Xr W_{H+1} \cdots W_3 \\ &\Leftrightarrow W_2 W_1 = (A_2^T A_2)^{-1} A_2^T Y X^T (X X^T)^{-1} + (I - (A_2^T A_2)^{-1} A_2^T A_2) L_2 \\ &\Leftrightarrow W_{H+1} \cdots W_1 = A_2 (A_2^T A_2)^{-1} A_2^T Y X^T (X X^T)^{-1} \\ &= C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}, \end{aligned}$$

⁵For a non-convex and *non-differentiable* function, we can still have a first-order and second-order necessary condition (e.g., Rockafellar & Wets, 2009, theorem 13.24, p. 606).

where the last two equalities follow Lemmas 4.2 and 4.6 (since if $Xr = 0$, we immediately obtain the desired result). Since XY^T is full rank with $d_y \leq d_x$ (i.e., $\text{rank}(XY^T) = d_y$),

$$A_2 (A_2^T A_2)^{-1} A_2 = U_{\bar{p}} U_{\bar{p}}^T = U_{\bar{p}} (U_{\bar{p}}^T U_{\bar{p}})^{-1} U_{\bar{p}}^T.$$

From this, with extra steps, we can deduce that we can have $\text{rank}(W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each entry of W_2 while retaining the loss value.

Thus, we conclude the proof for the base case with $k = 2$. For the inductive step with $k \in \{3, \dots, H + 1\}$, we essentially use the same proof procedure but with the inductive hypothesis that we can have $\text{rank}(W_{k-1} \cdots W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each entry of W_{k-1}, \dots, W_2 without changing the loss value. We need the inductive hypothesis to conclude that the first condition in $(\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C)$ or $XrW_{H+1} \cdots W_{k+1} = 0$) is false, and thus the second condition must be satisfied at a candidate point of a local minima.

We then conclude the induction, proving that we can have $\text{rank}(W_H \cdots W_2) \geq \text{rank}(W_{H+1} \cdots W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each parameter without changing the value of $\mathcal{L}(W)$. If $p \leq d_x$, this means that upon such a perturbation, we have the case of $\text{rank}(W_H \cdots W_2) = p$. Thus, such a critical point is not a local minimum unless it is a global minimum. If $p > d_x$, upon such a perturbation, we have $\text{rank}(W_{H+1} \cdots W_2) \geq d_x$. Thus, $W_{H+1} \cdots W_1 = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1} = U U^T Y X^T (X X^T)^{-1}$, which is a global minimum.

Summarizing the above, any point that satisfies the definition (and necessary conditions) of a local minimum is indeed a global minimum. Therefore, we conclude the proof sketch of Theorem 2.3 (ii).

5.2 Proof sketch of Theorem 2.3 (i), (iii) and (iv)

We can prove the non-convexity and non-concavity of this function simply from its Hessian (Theorem 2.3 (i)). That is, we can show that in the domain of the function, there exist points at which the Hessian becomes indefinite. Indeed, The domain contains uncountably many points at which the Hessian is indefinite.

We now consider Theorem 2.3 (iii): every critical point that is not a global minimum is a saddle point. Combined with Theorem 2.3 (ii), which is proven independently, this is equivalent to the statement that there are no local maxima. We first show that if $W_H \cdots W_1 \neq 0$, the loss function is strictly convex in one of the coordinates. This means that there is always an increasing direction and hence no local maximum. If $W_H \cdots W_1 = 0$, we show that at a critical point, if the Hessian is negative semidefinite, we can have $W_H \cdots W_1 \neq 0$ with arbitrarily small perturbation without changing the loss value. We can prove this by induction on $k = 1, \dots, H$, similar to the induction in the proof of Theorem 2.3 (ii).

Theorem 2.3 (iv) follows Theorem 2.3 (ii)-(iii) and the fact that when $\text{rank}(W_H \cdots W_2) = p$, if $\nabla^2 \mathcal{L}(W) \succeq 0$ at a critical point, W is a global minimum (this is the statement obtained in the proof of Theorem 2.3 (ii) for the case, $\text{rank}(W_H \cdots W_2) = p$).

5.3 Proof sketch of Theorem 3.2

Similarly to the previous work (Choromanska *et al.*, 2015a,b), we relate our loss function to another function under the adopted assumptions. More concretely, we show that all the theoretical results developed so far for the loss function of the deep linear models, $\tilde{\mathcal{L}}(W)$, hold true for the loss functions of the deep nonlinear models, $\mathbb{E}_Z[\mathcal{L}(W)]$ and $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$.

6 Conclusion

In this paper, we addressed some open problems, pushing forward the theoretical foundations of deep learning and non-convex optimization. For deep *linear* neural networks, we proved the aforementioned conjecture and more detailed statements with more generality. For deep *nonlinear* neural networks with rectified linear activation, when compared with the previous work, we proved a tighter statement with more generality (d_y can vary) and with strictly weaker model assumptions (only two assumptions out of seven). However, our theory does not yet directly apply to the practical situation. To fill the gap between theory and practice, future work would further discard the remaining two out of the seven assumptions made in previous work.

Acknowledgments

The author would like to thank Prof. Leslie Kaelbling for her thoughtful comments on the paper. We gratefully acknowledge support from NSF grant 1420927, from ONR grant N00014-14-1-0486, and from ARO grant W911NF1410433. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of our sponsors.

References

- Baldi, Pierre. 1989. Linear learning: Landscapes and algorithms. In *Advances in neural information processing systems*. pp. 65–72.
- Baldi, Pierre, & Hornik, Kurt. 1989. Neural networks and principal component analysis: Learning from examples without local minima. *Neural networks*, **2**(1), 53–58.
- Baldi, Pierre, & Lu, Zhiqin. 2012. Complex-valued autoencoders. *Neural Networks*, **33**, 136–147.
- Blum, Avrim L, & Rivest, Ronald L. 1992. Training a 3-node neural network is NP-complete. *Neural Networks*, **5**(1), 117–127.
- Choromanska, Anna, Henaff, Mikael, Mathieu, Michael, Ben Arous, Gerard, & LeCun, Yann. 2015a. The Loss Surfaces of Multilayer Networks. In *Proceedings of the Eighteenth International Conference on Artificial Intelligence and Statistics*. pp. 192–204.
- Choromanska, Anna, LeCun, Yann, & Arous, Gérard Ben. 2015b. Open Problem: The landscape of the loss surfaces of multilayer networks. In *Proceedings of The 28th Conference on Learning Theory*. pp. 1756–1760.
- Dauphin, Yann N, Pascanu, Razvan, Gulcehre, Caglar, Cho, Kyunghyun, Ganguli, Surya, & Bengio, Yoshua. 2014. Identifying and attacking the saddle point problem in high-dimensional non-convex optimization. In *Advances in Neural Information Processing Systems*. pp. 2933–2941.
- Ge, Rong, Huang, Furong, Jin, Chi, & Yuan, Yang. 2015. Escaping From Saddle Points—Online Stochastic Gradient for Tensor Decomposition. In *Proceedings of The 28th Conference on Learning Theory*. pp. 797–842.
- Goodfellow, Ian, Bengio, Yoshua, & Courville, Aaron. 2016. *Deep Learning*. Book in preparation for MIT Press. <http://www.deeplearningbook.org>.
- Livni, Roi, Shalev-Shwartz, Shai, & Shamir, Ohad. 2014. On the computational efficiency of training neural networks. In *Advances in Neural Information Processing Systems*. pp. 855–863.
- Mhaskar, Hrushikesh, Liao, Qianli, & Poggio, Tomaso. 2016. Learning Real and Boolean Functions: When Is Deep Better Than Shallow. *Massachusetts Institute of Technology CBMM Memo No. 45*.
- Murty, Katta G, & Kabadi, Santosh N. 1987. Some NP-complete problems in quadratic and nonlinear programming. *Mathematical programming*, **39**(2), 117–129.
- Rockafellar, R Tyrrell, & Wets, Roger J-B. 2009. *Variational analysis*. Vol. 317. Springer Science & Business Media.
- Saxe, Andrew M, McClelland, James L, & Ganguli, Surya. 2014. Exact solutions to the nonlinear dynamics of learning in deep linear neural networks. In *International Conference on Learning Representations*.
- Zhang, Fuzhen. 2006. *The Schur complement and its applications*. Vol. 4. Springer Science & Business Media.

Deep Learning without Poor Local Minima

Supplementary Material

Appendix

A Proofs of lemmas and corollary in Section 4

We complete the proofs of the lemmas and corollary in Section 4.

A.1 Proof of Lemma 4.1

Proof Since $\bar{\mathcal{L}}(W) = \frac{1}{2} \|\bar{Y}(W, X) - Y\|_F^2 = \frac{1}{2} \text{vec}(r)^T \text{vec}(r)$,

$$\begin{aligned} \mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W) &= (\mathcal{D}_{\text{vec}(r)} \bar{\mathcal{L}}(W)) \left(\mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(r) \right) \\ &= \text{vec}(r)^T \left(\mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(X^T I_{d_x} W_1^T \cdots W_{H+1}^T I_{d_y}) - \mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(Y^T) \right) \\ &= \text{vec}(r)^T \left(\mathcal{D}_{\text{vec}(W_k^T)} (W_{H+1} \cdots W_{k+1} \otimes (W_{k-1} \cdots W_1 X)^T) \text{vec}(W_k^T) \right) \\ &= \text{vec}(r)^T (W_{H+1} \cdots W_{k+1} \otimes (W_{k-1} \cdots W_1 X)^T). \end{aligned}$$

By setting $\left(\mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W) \right)^T = 0$ for all $k \in \{1, \dots, H+1\}$, we obtain the statement of Lemma 4.1. For the boundary conditions (i.e., $k = H+1$ or $k = 1$), it can be seen from the second to the third lines that we obtain the desired results with the definition, $W_k \cdots W_{k+1} \triangleq I_{d_k}$ (i.e., $W_{H+1} \cdots W_{H+2} \triangleq I_{d_y}$ and $W_0 \cdots W_1 \triangleq I_{d_x}$). \square

A.2 Proof of Lemma 4.2

Proof From the critical point condition with respect to W_1 (Lemma 4.1),

$$0 = \left(\mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W) \right)^T = (W_{H+1} \cdots W_2 \otimes X^T)^T \text{vec}(r) = \text{vec}(XrW_{H+1} \cdots W_2),$$

which is true if and only if $XrW_{H+1} \cdots W_2 = 0$. By expanding r , $0 = XX^T W_1^T C^T C - XY^T C$. By solving for W_1 ,

$$W_1 = (C^T C)^{-1} C^T Y X^T (X X^T)^{-1} + (I - (C^T C)^{-1} C^T C) L,$$

for an arbitrary matrix L . Due to the property of any generalized inverse (Zhang, 2006, p. 41), we have that $C(C^T C)^{-1} C^T C = C$. Thus,

$$C W_1 = C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1} + (C - C(C^T C)^{-1} C^T C) L = C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1}. \quad \square$$

A.3 Proof of Lemma 4.3

Proof For the diagonal blocks: the entries of diagonal blocks are obtained simply using the result of Lemma 4.1 as

$$\mathcal{D}_{\text{vec}(W_k^T)} \left(\mathcal{D}_{\text{vec}(W_k^T)} \bar{\mathcal{L}}(W) \right)^T = (W_{H+1} \cdots W_{k+1} \otimes (W_{k-1} \cdots W_1 X)^T)^T \mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(r).$$

Using the formula of $\mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(r)$ computed in the proof of Lemma 4.1 yields the desired result.

For the off-diagonal blocks with $k = 2, \dots, H$:

$$\begin{aligned} &\mathcal{D}_{\text{vec}(W_k^T)} [\mathcal{D}_{\text{vec}(W_1^T)} \bar{\mathcal{L}}(W)]^T \\ &= (W_{H+1} \cdots W_2 \otimes X)^T \mathcal{D}_{\text{vec}(W_k^T)} \text{vec}(r) + \left(\mathcal{D}_{\text{vec}(W_k^T)} W_{H+1} \cdots W_{k+1} \otimes X^T \right)^T \text{vec}(r) \end{aligned}$$

The first term above is reduced to the first term of the statement in the same way as the diagonal blocks. For the second term,

$$\begin{aligned}
& \left(\mathcal{D}_{\text{vec}(W_k^T)} W_{H+1} \cdots W_2 \otimes X^T \right)^T \text{vec}(r) \\
&= \sum_{i=1}^m \sum_{j=1}^{d_y} \left(\left(\mathcal{D}_{\text{vec}(W_k^T)} W_{H+1,j} W_H \cdots W_2 \right) \otimes X_i^T \right)^T r_{i,j} \\
&= \sum_{i=1}^m \sum_{j=1}^{d_y} \left((A_k)_{j,\cdot} \otimes B_k^T \otimes X_i^T \right)^T r_{i,j} \\
&= \sum_{i=1}^m \sum_{j=1}^{d_y} \left[(A_k)_{j,1} (B_k^T \otimes X_i) \cdots (A_k)_{j,d_k} (B_k^T \otimes X_i) \right] r_{i,j} \\
&= \left[\left(B_k^T \otimes \sum_{i=1}^m \sum_{j=1}^{d_y} r_{i,j} (A_k)_{j,1} X_i \right) \cdots \left(B_k^T \otimes \sum_{i=1}^m \sum_{j=1}^{d_y} r_{i,j} (A_k)_{j,d_k} X_i \right) \right].
\end{aligned}$$

where $A_k = W_{H+1} \cdots W_{k+1}$ and $B_k = W_{k-1} \cdots W_2$. The third line follows the fact that $(W_{H+1,j} W_H \cdots W_2)^T = \text{vec}(W_2^T \cdots W_H^T W_{H+1,j}^T) = (W_{H+1,j} \cdots W_{k+1} \otimes W_2^T \cdots W_{k-1}^T) \text{vec}(W_k^T)$. In the last line, we have the desired result by rewriting $\sum_{i=1}^m \sum_{j=1}^{d_y} r_{i,j} (A_k)_{j,t} X_i = X(rW_{H+1} \cdots W_{k+1})_{\cdot,t}$.

For the off-diagonal blocks with $k = H + 1$: The first term in the statement is obtained in the same way as above (for the off-diagonal blocks with $k = 2, \dots, H$). For the second term, notice that $\text{vec}(W_{H+1}^T) = [(W_{H+1})_{1,\cdot}^T \cdots (W_{H+1})_{d_y,\cdot}^T]^T$ where $(W_{H+1})_{j,\cdot}$ is the j -th row vector of W_{H+1} or the vector corresponding to the j -th output component. That is, it is conveniently organized as the blocks, each of which corresponds to each output component (or rather we chose $\text{vec}(W_k^T)$ instead of $\text{vec}(W_k)$ for this reason, among others). Also,

$$\begin{aligned}
& \left(\mathcal{D}_{\text{vec}(W_{H+1}^T)} W_{H+1} \cdots W_2 \otimes X^T \right)^T \text{vec}(r) = \\
&= \left[\sum_{i=1}^m \left(\left(\mathcal{D}_{(W_{H+1})_{1,\cdot}^T} C_{1,\cdot} \right) \otimes X_i^T \right)^T r_{i,1} \cdots \sum_{i=1}^m \left(\left(\mathcal{D}_{(W_{H+1})_{d_y,\cdot}^T} C_{d_y,\cdot} \right) \otimes X_i^T \right)^T r_{i,d_y} \right],
\end{aligned}$$

where we also used the fact that

$$\sum_{i=1}^m \sum_{j=1}^{d_y} \left(\left(\mathcal{D}_{\text{vec}((W_{H+1})_{t,\cdot}^T)} C_{j,\cdot} \right) \otimes X_i^T \right)^T r_{i,j} = \sum_{i=1}^m \left(\left(\mathcal{D}_{\text{vec}((W_{H+1})_{t,\cdot}^T)} C_{t,\cdot} \right) \otimes X_i^T \right)^T r_{i,t}.$$

For each block entry $t = 1, \dots, d_y$ in the above, similarly to the case of $k = 2, \dots, H$,

$$\sum_{i=1}^m \left(\left(\mathcal{D}_{\text{vec}((W_{H+1})_{t,\cdot}^T)} C_{j,\cdot} \right) \otimes X_i^T \right)^T r_{i,t} = \left(B_{H+1}^T \otimes \sum_{i=1}^m r_{i,t} (A_{H+1})_{j,t} X_i \right).$$

Here, we have the desired result by rewriting $\sum_{i=1}^m r_{i,t} (A_{H+1})_{j,1} X_i = X(rI_{d_y})_{\cdot,t} = Xr_{\cdot,t}$. \square

A.4 Proof of Lemma 4.4

Proof Note that a similarity transformation preserves the eigenvalues of a matrix. For each $k \in \{2, \dots, H + 1\}$, we take a similarity transform of $\nabla^2 \tilde{\mathcal{L}}(W)$ (whose entries are organized as in Lemma 4.3) as

$$P_k^{-1} \nabla^2 \tilde{\mathcal{L}}(W) P_k = \begin{bmatrix} \mathcal{D}_{\text{vec}(W_1^T)} \left(\mathcal{D}_{\text{vec}(W_1^T)} \tilde{\mathcal{L}}(W) \right)^T & \mathcal{D}_{\text{vec}(W_k^T)} \left(\mathcal{D}_{\text{vec}(W_1^T)} \tilde{\mathcal{L}}(W) \right)^T & \cdots \\ \mathcal{D}_{\text{vec}(W_1^T)} \left(\mathcal{D}_{\text{vec}(W_k^T)} \tilde{\mathcal{L}}(W) \right)^T & \mathcal{D}_{\text{vec}(W_k^T)} \left(\mathcal{D}_{\text{vec}(W_k^T)} \tilde{\mathcal{L}}(W) \right)^T & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

Here, $P_k = [\mathbf{e}_{H+1} \quad \mathbf{e}_k \quad \tilde{P}_k]$ is the permutation matrix where \mathbf{e}_i is the i -th element of the standard basis (i.e., a column vector with 1 in the i -th entry and 0 in every other entries), and \tilde{P}_k is any

arbitrarily matrix that makes P_k to be a permutation matrix. Let M_k be the principal submatrix of $P_k^{-1}\nabla^2\bar{\mathcal{L}}(W)P_k$ that consists of the first four blocks appearing in the above equation. Then,

$$\begin{aligned} \nabla^2\bar{\mathcal{L}}(W) &\succeq 0 \\ \Rightarrow \forall k \in \{2, \dots, H+1\}, M_k &\succeq 0 \\ \Rightarrow \forall k \in \{2, \dots, H+1\}, \mathcal{R}(\mathcal{D}_{\text{vec}(W_k^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T) &\subseteq \mathcal{R}(\mathcal{D}_{\text{vec}(W_1^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T), \end{aligned}$$

Here, the first implication follows the necessary condition with any principal submatrix and the second implication follows the necessary condition with the Schur complement (Zhang, 2006, theorem 1.20, p. 44).

Note that $\mathcal{R}(M') \subseteq \mathcal{R}(M) \Leftrightarrow (I - MM^{-})M' = 0$ (Zhang, 2006, p. 41). Thus, by plugging in the formulas of $\mathcal{D}_{\text{vec}(W_k^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T$ and $\mathcal{D}_{\text{vec}(W_1^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T$ that are derived in Lemma 4.3, $\nabla^2\bar{\mathcal{L}}(W) \succeq 0 \Rightarrow \forall k \in \{2, \dots, H+1\}$,

$$\begin{aligned} 0 = & \left(I - (C^T C \otimes (XX^T))(C^T C \otimes (XX^T))^{-} \right) (C^T A_k \otimes B_k W_1 X) \\ & + \left(I - (C^T C \otimes (XX^T))(C^T C \otimes (XX^T))^{-} \right) [B_k^T \otimes X] \begin{bmatrix} I_{d_{k-1}} \otimes (rA_k)_{\cdot,1} & \dots & I_{d_{k-1}} \otimes (rA_k)_{\cdot,d_k} \end{bmatrix} \end{aligned}$$

where $A_k = W_{H+1} \cdots W_{k+1}$ and $B_k = W_{k-1} \cdots W_2$. Here, we can replace $(C^T C \otimes (XX^T))^{-}$ by $((C^T C)^{-} \otimes (XX^T)^{-1})$ (see Appendix A.7). Thus, $I - (C^T C \otimes (XX^T))(C^T C \otimes (XX^T))^{-}$ can be replaced by $(I_{d_1} \otimes I_{d_y}) - (C^T C (C^T C)^{-} \otimes I_{d_y}) = (I_{d_1} - C^T C (C^T C)^{-}) \otimes I_{d_y}$. Accordingly, the first term is reduced to zero as

$$\left((I_{d_1} - C^T C (C^T C)^{-}) \otimes I_{d_y} \right) (C^T A_k \otimes B_k W_1 X) = ((I_{d_1} - C^T C (C^T C)^{-}) C^T A_k) \otimes B_k W_1 X = 0,$$

since $C^T C (C^T C)^{-} C^T = C^T$ (Zhang, 2006, p. 41). Thus, with the second term remained, the condition is reduced to

$$\forall k \in \{2, \dots, H+1\}, \forall t \in \{1, \dots, d_y\}, (B_k^T - C^T C (C^T C)^{-} B_k^T) \otimes X (rA_k)_{\cdot,t} = 0.$$

This implies

$$\forall k \in \{2, \dots, H+1\}, (R(B_k^T) \subseteq \mathcal{R}(C^T C) \text{ or } X rA_k = 0),$$

which concludes the proof for the positive semidefinite case. For the necessary condition of the negative semidefinite, we obtain the same condition since

$$\begin{aligned} \nabla^2\bar{\mathcal{L}}(W) &\preceq 0 \\ \Rightarrow \forall k \in \{2, \dots, H+1\}, M_k &\preceq 0 \\ \Rightarrow \forall k \in \{2, \dots, H+1\}, \mathcal{R}(-\mathcal{D}_{\text{vec}(W_k^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T) &\subseteq \mathcal{R}(-\mathcal{D}_{\text{vec}(W_1^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T) \\ \Rightarrow \forall k \in \{2, \dots, H+1\}, \mathcal{R}(\mathcal{D}_{\text{vec}(W_k^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T) &\subseteq \mathcal{R}(\mathcal{D}_{\text{vec}(W_1^T)}(\mathcal{D}_{\text{vec}(W_1^T)}\bar{\mathcal{L}}(W))^T). \end{aligned}$$

□

A.5 Proof of Corollary 4.5

Proof From the first condition in the statement of Lemma 4.4,

$$\begin{aligned} \mathcal{R}(W_2^T \cdots W_{k-1}^T) &\subseteq \mathcal{R}(W_2^T \cdots W_{H+1}^T W_{H+1} \cdots W_2) \\ \Rightarrow \text{rank}(W_k^T \cdots W_{H+1}^T) &\geq \text{rank}(W_2^T \cdots W_{k-1}^T) \Rightarrow \text{rank}(W_{H+1} \cdots W_k) \geq \text{rank}(W_{k-1} \cdots W_2). \end{aligned}$$

The first implication follows the fact that the rank of a product of matrices is at most the minimum of the ranks of the matrices, and the fact that the column space of $W_2^T \cdots W_{H+1}^T$ is subspace of the column space of $W_2^T \cdots W_{k-1}^T$. □

A.6 Proof of Lemma 4.6

Proof For the $(Xr = 0)$ condition: Let M_{H+1} be the principal submatrix as defined in the proof of Lemma 4.4 (the principal submatrix of $P_{H+1}^{-1}\nabla^2\bar{\mathcal{L}}(W)P_{H+1}$ that consists of the first four blocks of

it). Let $B_k = W_{k-1} \cdots W_2$. Let $F = B_{H+1} W_1 X X^T W_1^T B_{H+1}^T$. Using Lemma 4.3 for the blocks corresponding to W_1 and W_{H+1} ,

$$M_{H+1} = \begin{bmatrix} C^T C \otimes X X^T & (C^T \otimes X X^T (B_{H+1} W_1)^T) + E \\ (C \otimes B_{H+1} W_1 X X^T) + E^T & I_{d_y} \otimes F \end{bmatrix}$$

where $E = [B_{H+1}^T \otimes X r_{\cdot,1} \ \cdots \ B_{H+1}^T \otimes X r_{\cdot,d_y}]$. Then, by the necessary condition with the Schur complement (Zhang, 2006, theorem 1.20, p. 44), $M_{H+1} \succeq 0$ implies

$$\begin{aligned} 0 &= ((I_{d_y} \otimes I_{d_H}) - (I_{d_y} \otimes F)(I_{d_y} \otimes F)^{-})((C \otimes B_{H+1} W_1 X X^T) + E^T) \\ \Rightarrow 0 &= (I_{d_y} \otimes I_{d_H} - F F^{-})(C \otimes B_{H+1} W_1 X X^T) + (I_{d_y} \otimes I_{d_H} - F F^{-})E^T \\ &= (I_{d_y} \otimes I_{d_H} - F F^{-})E^T \\ &= \begin{bmatrix} I_{d_H} - F F^{-} \otimes I_1 & & \mathbf{0} \\ & \ddots & \\ \mathbf{0} & & I_{d_H} - F F^{-} \otimes I_1 \end{bmatrix} \begin{bmatrix} B_{H+1} \otimes (X r_{\cdot,1})^T \\ \vdots \\ B_{H+1} \otimes (X r_{\cdot,d_y})^T \end{bmatrix} \\ &= \begin{bmatrix} (I_{d_H} - F F^{-})B_{H+1} \otimes (X r_{\cdot,1})^T \\ \vdots \\ (I_{d_H} - F F^{-})B_{H+1} \otimes (X r_{\cdot,d_y})^T \end{bmatrix} \end{aligned}$$

where the second line follows the fact that $(I_{d_y} \otimes F)^{-}$ can be replaced by $(I_{d_y} \otimes F^{-})$ (see Appendix A.7). The third line follows the fact that $(I - F F^{-})B_{H+1} W_1 X = 0$ because $\mathcal{R}(B_{H+1} W_1 X) = \mathcal{R}(B_{H+1} W_1 X X^T W_1^T B_{H+1}^T) = \mathcal{R}(F)$. In the fourth line, we expanded E and used the definition of the Kronecker product. It implies

$$F F^{-} B_{H+1} = B_{H+1} \quad \text{or} \quad X r = 0.$$

Here, if $X r = 0$, we obtained the statement of the lemma. Thus, from now on, we focus on the case where $F F^{-} B_{H+1} = B_{H+1}$ and $X r \neq 0$ to obtain the other condition, $C(C^T C)^{-} C^T = U_{\bar{p}} U_{\bar{p}}^T$.

For the $(C(C^T C)^{-} C^T = U_{\bar{p}} U_{\bar{p}}^T)$ condition: By using another necessary condition of a matrix being positive semidefinite with the Schur complement (Zhang, 2006, theorem 1.20, p. 44), $M_{H+1} \succeq 0$ implies that

$$(I_{d_y} \otimes F) - \left(C \otimes B_{H+1} W_1 X X^T + E^T \right) (C^T C \otimes X X^T)^{-} \left(C^T \otimes X X^T (B_{H+1} W_1)^T + E \right) \succeq 0 \quad (1)$$

Since we can replace $(C^T C \otimes X X^T)^{-}$ by $(C^T C)^{-} \otimes (X X^T)^{-1}$ (see Appendix A.7), the second term in the left hand side is simplified as

$$\begin{aligned} & \left(C \otimes B_{H+1} W_1 X X^T + E^T \right) (C^T C \otimes X X^T)^{-} \left(C^T \otimes X X^T (B_{H+1} W_1)^T + E \right) \\ &= \left((C(C^T C)^{-} \otimes B_{H+1} W_1) + E^T \left((C^T C)^{-} \otimes (X X^T)^{-1} \right) \right) \left((C^T \otimes X X^T (B_{H+1} W_1)^T) + E \right) \\ &= \left(C(C^T C)^{-} C^T \otimes F \right) + E^T \left((C^T C)^{-} \otimes (X X^T)^{-1} \right) E \\ &= \left(C(C^T C)^{-} C^T \otimes F \right) + \left(r^T X^T (X X^T)^{-1} X r \otimes B_{H+1} (C^T C)^{-} B_{H+1} \right) \end{aligned} \quad (2)$$

In the third line, the crossed terms $-(C(C^T C)^{-} \otimes B_{H+1} W_1) E$ and its transpose $-$ are vanished to 0 because of the following. From Lemma 4.1, $(I_{d_y} \otimes (W_H \cdots W_1 X)^T)^T \text{vec}(r) = 0 \Leftrightarrow W_H \cdots W_1 X r = B_{H+1} W_1 X r = 0$ at any critical point. Thus, $(C(C^T C)^{-} \otimes B_{H+1} W_1) E = [C(C^T C)^{-} B_{H+1}^T \otimes B_{H+1} W_1 X r_{\cdot,1} \ \cdots \ C(C^T C)^{-} B_{H+1}^T \otimes B_{H+1} W_1 X r_{\cdot,d_y}] = 0$. The fourth line follows

$$\begin{aligned} & E^T \left((C^T C)^{-} \otimes (X X^T)^{-1} \right) E = \\ & \begin{bmatrix} B_{H+1} (C^T C)^{-} B_{H+1}^T \otimes (r_{\cdot,1})^T X^T (X X^T)^{-1} X r_{\cdot,1} \ \cdots \ B_{H+1} (C^T C)^{-} B_{H+1}^T \otimes (r_{\cdot,1})^T X^T (X X^T)^{-1} X r_{\cdot,d_y} \\ \vdots \\ B_{H+1} (C^T C)^{-} B_{H+1}^T \otimes (r_{\cdot,d_y})^T X^T (X X^T)^{-1} X r_{\cdot,1} \ \cdots \ B_{H+1} (C^T C)^{-} B_{H+1}^T \otimes (r_{\cdot,d_y})^T X^T (X X^T)^{-1} X r_{\cdot,d_y} \end{bmatrix} \\ &= r^T X^T (X X^T)^{-1} X r \otimes B_{H+1} (C^T C)^{-} B_{H+1}, \end{aligned}$$

where the last line is due to the fact that $\forall t, (r_{\cdot,t})^T X^T (XX^T)^{-1} X r_{\cdot,t}$ is a scalar and the fact that

$$\text{for any matrix } L, r^T L r = \begin{bmatrix} (r_{\cdot,1})^T L r_{\cdot,1} & \cdots & (r_{\cdot,1})^T L r_{\cdot,d_y} \\ \vdots & & \vdots \\ (r_{\cdot,d_y})^T L r_{\cdot,1} & \cdots & (r_{\cdot,d_y})^T L r_{\cdot,d_y} \end{bmatrix}.$$

From equations 1 and 2, $M_{H+1} \succeq 0 \Rightarrow$

$$((I_{d_y} - C(C^T C)^{-1} C^T) \otimes F) - (r^T X^T (XX^T)^{-1} X r \otimes B_{H+1} (C^T C)^{-1} B_{H+1}) \succeq 0. \quad (3)$$

In the following, we simplify equation 3 by first showing that $\mathcal{R}(C) \subseteq \mathcal{R}(\Sigma)$ and then simplifying $C(C^T C)^{-1} C^T, r^T X^T (XX^T)^{-1} X r, F$ and $B_{H+1} (C^T C)^{-1} B_{H+1}$.

Showing that $\mathcal{R}(C) \subseteq \mathcal{R}(\Sigma)$: Again, using Lemma 4.1 with $k = H + 1$,

$$0 = B_{H+1} W_1 X r \Leftrightarrow F W_{H+1}^T = B_{H+1} W_1 X Y^T \Leftrightarrow W_{H+1}^T = F^- B_{H+1} W_1 X Y^T + (I - F^- F) L,$$

for any arbitrary matrix L . Then,

$$\begin{aligned} C &= W_{H+1} B_{H+1} \\ &= Y X^T W_1^T B_{H+1}^T F^- B_{H+1} + L^T (I - F F^-) B_{H+1} \\ &= Y X^T W_1^T B_{H+1}^T F^- B_{H+1}, \end{aligned}$$

where the second equality follows the fact that we are conducting the case analysis with the case of $F F^- B_{H+1} = B_{H+1}$ here. Using Lemma 4.1 with $k = 1$,

$$0 = X r W_{H+1} \cdots W_2 \Leftrightarrow W_1 = (C^T C)^{-1} C^T Y X^T (XX^T)^{-1} + (I - (C^T C)^{-1} C^T C) L,$$

for any arbitrary matrix L . Pugging this formula of W_1 into the above,

$$\begin{aligned} C &= Y X^T ((C^T C)^{-1} C^T Y X^T (XX^T)^{-1} + (I - (C^T C)^{-1} C^T C) L)^T B_{H+1}^T F^- B_{H+1} \\ &= \Sigma C (C^T C)^{-1} B_{H+1}^T F^- B_{H+1} \end{aligned}$$

where the second line follows Lemma 4.4 with $k = H + 1$ (i.e., $C^T C (C^T C)^{-1} B_{H+1}^T = B_{H+1}^T$). Thus, we have the desired result, $\mathcal{R}(C) \subseteq \mathcal{R}(\Sigma)$.

Simplifying $C(C^T C)^{-1} C^T$: Remember that \bar{p} is the rank of C . To simplify the notation, we rearrange the entries of D and U such that the eigenvalues and eigenvectors selected by the index set $\mathcal{I}_{\bar{p}}$ comes first. That is, $U = [U_{\mathcal{I}_{\bar{p}}} \quad U_{-\mathcal{I}_{\bar{p}}}]$ and $\Lambda = \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & 0 \\ 0 & \Lambda_{-\mathcal{I}_{\bar{p}}} \end{bmatrix}$ where $U_{-\mathcal{I}_{\bar{p}}}$ consists of all the eigenvectors that are not contained in $U_{\mathcal{I}_{\bar{p}}}$, and accordingly $\Lambda_{\mathcal{I}_{\bar{p}}}$ (resp. $\Lambda_{-\mathcal{I}_{\bar{p}}}$) consists of all the eigenvalues that correspond (resp. do not correspond) to the index set $\mathcal{I}_{\bar{p}}$. Since $\mathcal{R}(C) \subseteq \mathcal{R}(\Sigma)$, we can write C in the following form: for some index set $\mathcal{I}_{\bar{p}}$, $C = [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_1$, where $\mathbf{0} \in \mathbb{R}^{d_y \times (d_1 - \bar{p})}$ and $G_1 \in GL_{d_1}(\mathbb{R})$ (a $d_1 \times d_1$ invertible matrix) (notice that $d_1 \geq p \geq \bar{p}$ by their definitions). Then,

$$(C^T C)^{-1} = (G_1^T [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}]^T [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_1)^{-1} = \left(G_1^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_1 \right)^{-1}.$$

Note that the set of all generalized inverse of $C^T C = G_1^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_1$ is as follows (Zhang, 2006, p. 41):

$$\left\{ G_1^{-1} \begin{bmatrix} I_{\bar{p}} & L_1 \\ L_2 & L_3 \end{bmatrix} G_1^{-T} \mid L_1, L_2, L_3 \text{ arbitrary} \right\}.$$

Thus, for any arbitrary L_1, L_2 and L_3 ,

$$C(C^T C)^{-1} C^T = C G_1^{-1} \begin{bmatrix} I_{\bar{p}} & L_1 \\ L_2 & L_3 \end{bmatrix} G_1^{-T} C^T = [U_{\mathcal{I}_{\bar{p}}} \quad \mathbf{0}] \begin{bmatrix} I_{\bar{p}} & L_1 \\ L_2 & L_3 \end{bmatrix} \begin{bmatrix} U_{\mathcal{I}_{\bar{p}}}^T \\ \mathbf{0} \end{bmatrix} = U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T.$$

Simplifying $r^T X^T (XX^T)^{-1} X r$:

$$\begin{aligned} r^T X^T (XX^T)^{-1} X r &= (C W_1 X - Y) X^T (XX^T)^{-1} X (X^T (C W_1)^T - Y^T) \\ &= C W_1 X X^T (C W_1)^T - C W_1 X Y^T - Y X^T (C W_1)^T + \Sigma \\ &= P_C \Sigma P_C - P_C \Sigma - \Sigma P_C + \Sigma \\ &= \Sigma - U_{\bar{p}} \Lambda_{\mathcal{I}_{\bar{p}}} U_{\bar{p}}^T \end{aligned}$$

where $P_C = C(C^T C)^{-1} C^T = U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T$ and the last line follows the facts:

$$P_C \Sigma P_C = U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T U \Lambda U^T U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T = U_{\mathcal{I}_{\bar{p}}} [I_{\bar{p}} \ 0] \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & 0 \\ 0 & \Lambda_{-\mathcal{I}_{\bar{p}}} \end{bmatrix} \begin{bmatrix} I_{\bar{p}} \\ 0 \end{bmatrix} U_{\mathcal{I}_{\bar{p}}}^T = U_{\mathcal{I}_{\bar{p}}} \Lambda_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T,$$

$$P_C \Sigma = U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T U \Lambda U^T = U_{\mathcal{I}_{\bar{p}}} [I_{\bar{p}} \ 0] \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & 0 \\ 0 & \Lambda_{-\mathcal{I}_{\bar{p}}} \end{bmatrix} \begin{bmatrix} U_{\mathcal{I}_{\bar{p}}}^T \\ U_{-\mathcal{I}_{\bar{p}}} \end{bmatrix} = U_{\mathcal{I}_{\bar{p}}}^T \Lambda_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}},$$

and similarly, $\Sigma P_C = U_{\mathcal{I}_{\bar{p}}}^T \Lambda_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}$.

Simplifying F : In the proof of Lemma 4.2, by using Lemma 4.1 with $k = 1$, we obtained that $W_1 = (C^T C)^{-1} C^T Y X^T (X X^T)^{-1} + (I - (C^T C)^{-1} C^T C) L$. Also, from Lemma 4.4, we have that $X r = 0$ or $B_{H+1} (C^T C)^{-1} C^T C = (C^T C (C^T C)^{-1} B_{H+1}^T)^T = B_{H+1}$. If $X r = 0$, we got the statement of the lemma, and so we consider the case of $B_{H+1} (C^T C)^{-1} C^T C = B_{H+1}$. Therefore,

$$B_{H+1} W_1 = B_{H+1} (C^T C)^{-1} C^T Y X^T (X X^T)^{-1}.$$

Since $F = B_{H+1} W_1 X X^T W_1^T B_{H+1}^T$,

$$F = B_{H+1} (C^T C)^{-1} C^T \Sigma C (C^T C)^{-1} B_{H+1}^T.$$

From Lemma 4.4 with $k = H + 1$, $\mathcal{R}(B_{H+1}^T) \subseteq \mathcal{R}(C^T C) = \mathcal{R}(B_{H+1}^T W_{H+1}^T W_{H+1} B_{H+1}) \subseteq \mathcal{R}(B_{H+1}^T)$, which implies that $\mathcal{R}(B_{H+1}^T) = \mathcal{R}(C^T C)$. Therefore, $\mathcal{R}(C (C^T C)^{-1} B_{H+1}^T) = \mathcal{R}(C (C^T C)^{-1}) = \mathcal{R}(C) \subseteq \mathcal{R}(\Sigma)$. Accordingly, we can write it in the form, $C (C^T C)^{-1} B_{H+1}^T = [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_2$, where $\mathbf{0} \in \mathbb{R}^{d_y \times (d_1 - \bar{p})}$ and $G_2 \in GL_{d_1}(\mathbb{R})$ (we can write it in the form of $[U_{\mathcal{I}_{\bar{p}'}}], \mathbf{0}] G_2$ for some $\mathcal{I}_{\bar{p}'}$ because of the inclusion $\subseteq \mathcal{R}(\Sigma)$ and $\mathcal{I}_{\bar{p}'} = \mathcal{I}_{\bar{p}}$ because of the equality $= \mathcal{R}(C)$). Thus,

$$F = G_2^T \begin{bmatrix} U_{\mathcal{I}_{\bar{p}}}^T \\ \mathbf{0} \end{bmatrix} U \Lambda U^T [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_2 = G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \Lambda \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2 = G_2^T \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2.$$

Simplifying $B_{H+1} (C^T C)^{-1} B_{H+1}$: From Lemma 4.4, $C^T C (C^T C)^{-1} B_{H+1} = B_{H+1}$ (again since we are done if $X r = 0$). Thus, $B_{H+1} (C^T C)^{-1} B_{H+1} = B_{H+1} (C^T C)^{-1} C^T C (C^T C)^{-1} B_{H+1}^T$. As discussed above, we write $C (C^T C)^{-1} B_{H+1}^T = [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_2$. Thus,

$$B_{H+1} (C^T C)^{-1} B_{H+1} = G_2^T \begin{bmatrix} U_{\mathcal{I}_{\bar{p}}}^T \\ \mathbf{0} \end{bmatrix} [U_{\mathcal{I}_{\bar{p}}}, \mathbf{0}] G_2 = G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2.$$

Putting results together: We use the simplified formulas of $C (C^T C)^{-1} C^T$, $r^T X^T (X X^T)^{-1} X r$, F and $B_{H+1} (C^T C)^{-1} B_{H+1}$ in equation 3, obtaining

$$\left((I_{d_y} - U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T) \otimes G_2^T \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2 \right) - \left((\Sigma - U_{\bar{p}} \Lambda_{\mathcal{I}_{\bar{p}}} U_{\bar{p}}^T) \otimes G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2 \right) \succeq 0.$$

Due to the Sylvester's law of inertia (Zhang, 2006, theorem 1.5, p. 27), with a nonsingular matrix $U \otimes G_2^{-1}$ (it is nonsingular because each of U and G_2^{-1} is nonsingular), the necessary condition is reduced to

$$\begin{aligned} & (U \otimes G_2^{-1})^T \left(\left((I_{d_y} - U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T) \otimes G_2^T \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2 \right) - \left((\Sigma - U_{\bar{p}} \Lambda_{\mathcal{I}_{\bar{p}}} U_{\bar{p}}^T) \otimes G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2 \right) \right) (U \otimes G_2^{-1}) \\ &= \left(\left(I_{d_y} - \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) \otimes \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) - \left(\left(\Lambda - \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) \otimes \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) \\ &= \left(\begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & I_{(d_y - \bar{p})} \end{bmatrix} \otimes \begin{bmatrix} \Lambda_{\mathcal{I}_{\bar{p}}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) - \left(\begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \Lambda_{-\mathcal{I}_{\bar{p}}} \end{bmatrix} \otimes \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) \\ &= \left[\begin{array}{c|c} \mathbf{0} & \mathbf{0} \\ \hline \Lambda_{\mathcal{I}_{\bar{p}}} - (\Lambda_{-\mathcal{I}_{\bar{p}}})_{1,1} I_{\bar{p}} & \mathbf{0} \\ \hline \mathbf{0} & \ddots \\ \mathbf{0} & \Lambda_{\mathcal{I}_{\bar{p}}} - (\Lambda_{-\mathcal{I}_{\bar{p}}})_{(d_y - \bar{p}), (d_y - \bar{p})} I_{\bar{p}} \end{array} \right] \succeq 0, \end{aligned}$$

which implies that for all $(i, j) \in \{(i, j) \mid i \in \{1, \dots, \bar{p}\}, j \in \{1, \dots, (d_y - \bar{p})\}\}$, $(\Lambda_{\mathcal{I}_{\bar{p}}})_{i,i} \geq (\Lambda_{-\mathcal{I}_{\bar{p}}})_{j,j}$. In other words, the index set $\mathcal{I}_{\bar{p}}$ must select the largest \bar{p} eigenvalues whatever \bar{p} is. Since $C(C^T C)^{-1} C^T = U_{\mathcal{I}_{\bar{p}}} U_{\mathcal{I}_{\bar{p}}}^T$ (which is obtained above), we have that $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}$ in this case.

Summarizing the above case analysis, if $\nabla^2 \bar{\mathcal{L}}(W) \succeq 0$ at a critical point, $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}$ or $Xr = 0$. \square

A.7 Generalized inverse of Kronecker product

$(A^- \otimes B^-)$ is a generalized inverse of $A \otimes B$.

Proof For a matrix M , the definition of a generalized inverse, M^- , is $MM^-M = M$. Setting $M := A \otimes B$, we check if $(A^- \otimes B^-)$ satisfies the definition: $(A \otimes B)(A^- \otimes B^-)(A \otimes B) = (AA^-A \otimes BB^-B) = (A \otimes B)$ as desired. \square

We avoid discussing the other direction as it is unnecessary in this paper (i.e., we avoid discussing if $(A^- \otimes B^-)$ is the only generalized inverse of $A \otimes B$). Notice that the necessary condition that we have in our proof (where we need a generalized inverse of $A \otimes B$) is for any generalized inverse of $A \otimes B$. Thus, replacing it by one of any generalized inverse suffices to obtain a necessary condition. Indeed, choosing Moore–Penrose pseudoinverse suffices here, with which we know $(A \otimes B)^\dagger = (A^\dagger \otimes B^\dagger)$. But, to give a simpler argument later, we keep more generality by choosing $(A^- \otimes B^-)$ as a generalized inverse of $A \otimes B$.

B Proof of Theorems 2.3 and 3.2

We complete the proofs of Theorems 2.3 and 3.2.

B.1 Proof of Theorem 2.3 (ii)

Proof By case analysis, we show that any point that satisfies the necessary conditions and the definition of a local minimum is a global minimum. When we write a statement in the proof, we often mean that a necessary condition of local minima implies the statement as it should be clear (i.e., we are not claiming that the statement must hold true unless the point is the candidate of local minima).

The case where $\text{rank}(W_H \cdots W_2) = p$ and $d_y \leq p$: Assume that $\text{rank}(W_H \cdots W_2) = p$. We first obtain a necessary condition of the Hessian being positive semidefinite at a critical point, $Xr = 0$, and then interpret the condition. If $d_y < p$, Corollary 4.5 with $k = H + 1$ implies the necessary condition that $Xr = 0$. This is because the other condition $p > \text{rank}(W_{H+1}) \geq \text{rank}(W_H \cdots W_2) = p$ is false.

If $d_y = p$, Lemma 4.6 with $k = H + 1$ implies the necessary condition that $Xr = 0$ or $\mathcal{R}(W_H \cdots W_2) \subseteq \mathcal{R}(C^T C)$. Suppose that $\mathcal{R}(W_H \cdots W_2) \subseteq \mathcal{R}(C^T C)$. Then, we have that $p = \text{rank}(W_H \cdots W_2) \leq \text{rank}(C^T C) = \text{rank}(C)$. That is, $\text{rank}(C) \geq p$.

From Corollary 4.5 with $k = 2$ implies the necessary condition that

$$\text{rank}(C) \geq \text{rank}(I_{d_1}) \quad \text{or} \quad XrW_{H+1} \cdots W_3 = 0.$$

Suppose the latter: $XrW_{H+1} \cdots W_3 = 0$. Since $\text{rank}(W_{H+1} \cdots W_3) \geq \text{rank}(C) \geq p$ and $d_{H+1} = d_y = p$, the left null space of $W_{H+1} \cdots W_3$ contains only zero. Thus,

$$XrW_{H+1} \cdots W_3 = 0 \Rightarrow Xr = 0.$$

Suppose the former: $\text{rank}(C) \geq \text{rank}(I_{d_1})$. Because $d_y = p \leq d_1$, $\text{rank}(C) \geq p$, and $\mathcal{R}(C) \subseteq \mathcal{R}(YX^T)$ as shown in the proof of Lemma 4.6, we have that $\mathcal{R}(C) = \mathcal{R}(YX^T)$.

$$\text{rank}(C) \geq \text{rank}(I_{d_1}) \Rightarrow C^T C \text{ is full rank} \Rightarrow Xr = XY^T C(C^T C)^{-1} C^T - XY^T = 0,$$

where the last equality follows the fact that $(Xr)^T = C(C^T C)^{-1} C^T YX^T - YX^T = 0$ since $\mathcal{R}(C) = \mathcal{R}(YX^T)$ and thereby the projection of YX^T onto the range of C is YX^T . Therefore, we have the condition, $Xr = 0$ when $d_y \leq p$.

To interpret the condition $Xr = 0$, consider a loss function with a linear model without any hidden layer, $f(W') = \|W'X - Y\|_F^2$ where $W' \in \mathbb{R}^{d_y \times d_x}$. Then, any point satisfying $Xr' = 0$ is a global minimum of f , where $r' = (W'X - Y)^T$ is an error matrix.⁶ For any values of $W_{H+1} \cdots W_1$, there exists W' such that $W' = W_{H+1} \cdots W_1$ (the opposite is also true when $d_y \leq p$ although we don't need it in our proof). That is, $\mathcal{R}(\tilde{\mathcal{L}}) \subseteq \mathcal{R}(f)$ and $\mathcal{R}(r) \subseteq \mathcal{R}(r')$ (as functions of W and W' respectively) (the equality is also true when $d_y \leq p$ although we don't need it in our proof). Summarizing the above, whenever $Xr = 0$, there exists $W' = W_{H+1} \cdots W_1$ such that $Xr = Xr' = 0$, which achieves the global minimum value of f , f^* and $f^* \leq \tilde{\mathcal{L}}^*$ (i.e., the global minimum value of f is at most the global minimum value of $\tilde{\mathcal{L}}$ since $\mathcal{R}(\tilde{\mathcal{L}}) \subseteq \mathcal{R}(f)$). In other words, $W_{H+1} \cdots W_1$ achieving $Xr = 0$ attains a global minimum value of f that is at most the global minimum value of $\tilde{\mathcal{L}}$. This means that $W_{H+1} \cdots W_1$ achieving $Xr = 0$ is a global minimum.

Thus, we have proved that when $\text{rank}(W_H \cdots W_2) = p$ and $d_y \leq p$, if $\nabla^2 \tilde{\mathcal{L}}(W) \succeq 0$ at a critical point, it is a global minimum.

The case where $\text{rank}(W_H \cdots W_2) = p$ and $d_y > p$: We first obtain a necessary condition of the Hessian being positive semidefinite at a critical point and then interpret the condition. From Lemma 4.6, we have that $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}^T$ or $Xr = 0$. If $Xr = 0$, with the exact same proof as in the case of $d_y \leq p$, it is a global minimum. Suppose that $C(C^T C)^{-1} C^T = U_{\bar{p}} U_{\bar{p}}^T$. Combined with Lemma 4.2, we have a necessary condition:

$$W_{H+1} \cdots W_1 = C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}.$$

From Lemma 4.4 with $k = H + 1$, $\mathcal{R}(W_2^T \cdots W_H^T) \subseteq \mathcal{R}(C^T C) = \mathcal{R}(C^T)$, which implies that $\bar{p} \triangleq \text{rank}(C) = p$ (since $\text{rank}(W_H \cdots W_2) = p$). Thus, we can rewrite the above equation as $W_{H+1} \cdots W_1 = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}$, which is the orthogonal projection on to subspace spanned by the p eigenvectors corresponding to the p largest eigenvalues following the ordinary least square regression matrix. This is indeed the expression of a global minimum (Baldi & Hornik, 1989; Baldi & Lu, 2012).

Thus, we have proved that when $\text{rank}(W_H \cdots W_2) = p$, if $\nabla^2 \tilde{\mathcal{L}}(W) \succeq 0$ at a critical point, it is a global minimum.

The case where $\text{rank}(W_H \cdots W_2) < p$: Suppose that $\text{rank}(W_H \cdots W_2) < p$. From Lemma 4.4, we have a following necessary condition for the Hessian to be (positive or negative) semidefinite at a critical point: for any $k \in \{2, \dots, H + 1\}$,

$$\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad Xr W_{H+1} \cdots W_{k+1} = 0,$$

where the first condition is shown to imply $\text{rank}(W_{H+1} \cdots W_k) \geq \text{rank}(W_{k-1} \cdots W_2)$ in Corollary 4.5. We repeatedly apply these conditions for $k = 2, \dots, H + 1$ to claim that with arbitrarily small $\epsilon > 0$, we can perturb each parameter (i.e., each entry of W_H, \dots, W_2) such that $\text{rank}(W_{H+1} \cdots W_2) \geq \min(p, d_x)$ without changing the value of $\tilde{\mathcal{L}}(W)$.

Let $A_k = W_{H+1} \cdots W_{k+1}$. From Corollary 4.5 with $k = 2$, we have that $\text{rank}(W_{H+1} \cdots W_2) \geq d_1 \geq p$ or $Xr W_{H+1} \cdots W_3 = 0$ (note that $d_1 \geq p \geq \bar{p}$ by their definitions). The former condition is false since $\text{rank}(W_{H+1} \cdots W_2) \leq \text{rank}(W_H \cdots W_2) < p$. From the latter condition, for an arbitrary L_2 ,

$$\begin{aligned} 0 &= Xr W_{H+1} \cdots W_3 \\ \Leftrightarrow W_2 W_1 &= (A_2^T A_2)^{-} A_2^T Y X^T (X X^T)^{-1} + (I - (A_2^T A_2)^{-} A_2^T A_2) L_2 \\ \Leftrightarrow W_{H+1} \cdots W_1 &= A_2 (A_2^T A_2)^{-} A_2^T Y X^T (X X^T)^{-1} \\ &= C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}, \end{aligned} \quad (4)$$

where the last two equalities follow Lemmas 4.2 and 4.6 (since if $Xr = 0$, we immediately obtain the desired result as discussed above). Taking transpose,

$$(X X^T)^{-1} X Y^T A_2 (A_2^T A_2)^{-} A_2^T = (X X^T)^{-1} X Y^T U_{\bar{p}} U_{\bar{p}}^T,$$

⁶Proof: Any point satisfying $Xr' = 0$ is a critical point of f , which directly follows the proof of Lemma 4.1. Also, f is convex since its Hessian is positive semidefinite for all input W_{H+1} , and thus any critical point of f is a global minimum. Combining the previous two statements results in the desired claim.

which implies that

$$XY^T A_2 (A_2^T A_2)^{-1} A_2 = XY^T U_{\bar{p}} U_{\bar{p}}^T.$$

Since XY^T is full rank with $d_y \leq d_x$ (i.e., $\text{rank}(XY^T) = d_y$), there exists a left inverse and the solution of the above linear system is unique as $((XY^T)^T XY^T)^{-1} (XY^T)^T XY^T = I$, yielding,

$$A_2 (A_2^T A_2)^{-1} A_2 = U_{\bar{p}} U_{\bar{p}}^T (= U_{\bar{p}} (U_{\bar{p}}^T U_{\bar{p}})^{-1} U_{\bar{p}}^T).$$

In other words, $\mathcal{R}(A_2) = \mathcal{R}(C) = \mathcal{R}(U_{\bar{p}})$.

Suppose that $(A_2^T A_2) \in \mathbb{R}^{d_2 \times d_2}$ is nonsingular. Then, since $\mathcal{R}(A_2) = \mathcal{R}(C)$, we have that $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) = \text{rank}(A_2) = d_2 \geq p$, which is false in the case being analyzed (the case of $\text{rank}(W_H \cdots W_2) < p$). Thus, $A_2^T A_2$ is singular.

If $A_2^T A_2$ is singular, from equation 4, it is inferred that we can perturb W_2 to have $\text{rank}(W_2 W_1) \geq \min(p, d_x)$. To see this in a concrete algebraic way, first note that since $\mathcal{R}(A_2) = \mathcal{R}(U_{\bar{p}})$, we can write $A_2 = [U_{\bar{p}} \ \mathbf{0}] G_2$ for some $G_2 \in GL_{d_2}(\mathbb{R})$ where $\mathbf{0} \in \mathbb{R}^{d_y \times (d_2 - \bar{p})}$. Thus,

$$A_2^T A_2 = G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2.$$

Again, note that the set of all generalized inverse of $G_2^T \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2$ is as follows (Zhang, 2006, p. 41):

$$\left\{ G_2^{-1} \begin{bmatrix} I_{\bar{p}} & L'_1 \\ L'_2 & L'_3 \end{bmatrix} G_2^{-T} \mid L'_1, L'_2, L'_3 \text{ arbitrary} \right\}.$$

Since equation 4 must hold for any generalized inverse, we choose a generalized inverse with $L'_1 = L'_2 = L'_3 = 0$ for simplicity. That is,

$$(A_2^T A_2)^{-} := G_2^{-1} \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2^{-T}.$$

Then, plugging this into equation 4, for an arbitrary L_2 ,

$$\begin{aligned} W_2 W_1 &= G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T \\ \mathbf{0} \end{bmatrix} Y X^T (X X^T)^{-1} + (I_{d_2} - G_2^{-1} \begin{bmatrix} I_{\bar{p}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} G_2) L_2 \\ &= G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ \mathbf{0} \end{bmatrix} + G_2^{-1} \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & I_{(d_2 - \bar{p})} \end{bmatrix} G_2 L_2 \\ &= G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ \mathbf{0} \ I_{(d_2 - \bar{p})} \end{bmatrix} G_2 L_2. \end{aligned}$$

Here, $[\mathbf{0} \ I_{(d_2 - \bar{p})}] G_2 L_2 \in \mathbb{R}^{(d_2 - \bar{p}) \times d_x}$ is the last $(d_2 - \bar{p})$ rows of $G_2 L_2$. Since $\text{rank}(Y X^T (X X^T)^{-1}) = d_y$ (because the multiplication with the invertible matrix preserves the rank), the first \bar{p} rows in the above have rank \bar{p} . Thus, $W_2 W_1$ has rank at least \bar{p} , and the possible rank deficiency comes from the last $(d_2 - \bar{p})$ rows, $[\mathbf{0} \ I_{(d_2 - \bar{p})}] G_2 L_2$. Since $W_{H+1} \cdots W_1 = A_2 W_2 W_1 = [U_{\bar{p}} \ \mathbf{0}] G_2 W_2 W_1$,

$$W_{H+1} \cdots W_1 = [U_{\bar{p}} \ \mathbf{0}] \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ \mathbf{0} \ I_{(d_2 - \bar{p})} \end{bmatrix} G_2 L_2 = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}.$$

This means that changing the values of the last $(d_2 - \bar{p})$ rows of $G_2 L_2$ (i.e., $[\mathbf{0} \ I_{(d_2 - \bar{p})}] G_2 L_2$) does not change the value of $\tilde{\mathcal{L}}(W)$. Therefore, the original necessary condition implies a necessary condition that without changing the loss value, we can make $W_2 W_1$ to have full rank with arbitrarily small perturbation of the last $(d_2 - \bar{p})$ rows as $[\mathbf{0} \ I_{(d_2 - \bar{p})}] G_2 L_2 + \epsilon M_{\text{ptb}}$ where ϵM_{ptb} is a perturbation matrix with arbitrarily small $\epsilon > 0$.⁷

⁷We have only proved that the submatrix of the first \bar{p} rows has rank \bar{p} and that changing the value of the last $d_2 - \bar{p}$ rows does not change the loss value. That is, we have not proven the existence of ϵM_{ptb} that makes $W_2 W_1$ full rank. Although this is trivial since the set of full matrices is dense, we show a proof in the following

Now, we show that such a perturbation can be done via a perturbation of the entries of W_2 . From the above equation for W_2W_1 , all the possible solutions of W_2 can be written as: for an arbitrary L_0 and L_2 ,

$$W_2 = G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 \end{bmatrix} W_1^\dagger + L_0^T (I - W_1 W_1^\dagger),$$

where M^\dagger is the the Moore—Penrose pseudoinverse of M . Thus, we perturb W_2 as

$$W_2 := W_2 + \epsilon G_2^{-1} \begin{bmatrix} 0 \\ M_{\text{ptb}} \end{bmatrix} W_1^\dagger = G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 + \epsilon M_{\text{ptb}} \end{bmatrix} W_1^\dagger + L_0^T (I - W_1 W_1^\dagger).$$

Note that upon such a perturbation, equation 4 may not hold anymore; i.e.,

$$G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 + \epsilon M_{\text{ptb}} \end{bmatrix} W_1^\dagger W_1 \neq G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 + \epsilon M_{\text{ptb}} \end{bmatrix}.$$

This means that the original necessary condition that implies equation 4 no longer holds. In this case, we immediately conclude that the Hessian is no longer positive semidefinite and thus the point is a saddle point. We thereby consider the remaining case: equation 4 still holds. Then, with the perturbation on the entries of W_1 ,

$$W_2 W_1 = G_2^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 + \epsilon M_{\text{ptb}} \end{bmatrix},$$

as desired.

Thus, we showed that we can have $\text{rank}(W_2) \geq \text{rank}(W_2 W_1) \geq \min(p, d_x)$, with arbitrarily small perturbation of each entry of W_2 with the loss value being remained. To prove the corresponding results for $W_k \cdots W_2$ for any $k = 2, \dots, H + 1$, we conduct induction on $k = 2, \dots, H + 1$ with the same proof procedure. The proposition $P(k)$ to be proven is as follows: the necessary conditions with $j \leq k$ imply that we can have $\text{rank}(W_k \cdots W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each entry of W_k, \dots, W_2 without changing the loss value. For the base case $k = 2$, we have already proved the proposition in the above.

For the inductive step with $k \in \{3, \dots, H + 1\}$, we have the inductive hypothesis that we can have $\text{rank}(W_{k-1} \cdots W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each entry of W_{k-1}, \dots, W_2 without changing the loss value. Accordingly, suppose that $\text{rank}(W_{k-1} \cdots W_1) \geq \min(p, d_x)$. Again, from Lemma 4.4, for any $k \in \{3, \dots, H + 1\}$,

$$\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad X r W_{H+1} \cdots W_{k+1} = 0.$$

If the former is true, $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) \geq \text{rank}(W_{k-1} \cdots W_2) \geq \min(p, d_x)$, which is the desired statement (it immediately implies the proposition $P(k)$ for any k). If the latter is true, for an arbitrary L_k ,

$$\begin{aligned} 0 &= X r W_{H+1} \cdots W_{k+1} \\ \Leftrightarrow W_k \cdots W_1 &= (A_k^T A_k)^{-1} A_k^T Y X^T (X X^T)^{-1} + (I - (A_k^T A_k)^{-1} A_k^T A_k) L_k \\ \Leftrightarrow W_{H+1} \cdots W_1 &= A_k (A_k^T A_k)^{-1} A_k^T Y X^T (X X^T)^{-1} \\ &= C (C^T C)^{-1} C^T Y X^T (X X^T)^{-1} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}, \end{aligned} \tag{5}$$

where the last two equalities follow Lemmas 4.2 and 4.6. Taking transpose,

$$(X X^T)^{-1} X Y^T A_k (A_k^T A_k)^{-1} A_k^T = (X X^T)^{-1} X Y^T U_{\bar{p}} U_{\bar{p}}^T,$$

to be complete. Let $\bar{p}' \geq \bar{p}$ be the rank of $W_2 W_1$. That is, in $\begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_2 - \bar{p})}] G_2 L_2 \end{bmatrix}$, there exist \bar{p}' linearly independent row vectors including the first \bar{p} row vectors, denoted by $b_1, \dots, b_{\bar{p}'} \in \mathbb{R}^{1 \times d_x}$. Then, we denote the rest of row vectors by $v_1, v_2, \dots, v_{d_2 - \bar{p}'} \in \mathbb{R}^{1 \times d_x}$. Let $c = \min(d_2 - \bar{p}', d_x - \bar{p}')$. There exist linearly independent vectors $\bar{v}_1, \bar{v}_2, \dots, \bar{v}_c$ such that the set, $\{b_1, \dots, b_{\bar{p}'}, \bar{v}_1, \bar{v}_2, \dots, \bar{v}_c\}$, is linearly independent. Setting $v_i := v_i + \epsilon \bar{v}_i$ for all $i \in \{1, \dots, c\}$ makes $W_2 W_1$ full rank since $\epsilon \bar{v}_i$ cannot be expressed as a linear combination of other vectors. Thus, a desired perturbation matrix ϵM_{ptb} can be obtained by setting ϵM_{ptb} to consists of $\epsilon \bar{v}_1, \epsilon \bar{v}_2, \dots, \epsilon \bar{v}_c$ row vectors for the corresponding rows and 0 row vectors for other rows.

which implies that $XY^T A_k (A_k^T A_k)^{-1} A_k = XY^T U_{\bar{p}} U_{\bar{p}}^T$. Since XY^T is full rank with $d_y \leq d_x$ (i.e., $\text{rank}(XY^T) = d_y$), there exists a left inverse and the solution of the above linear system is unique as $((XY^T)^T XY^T)^{-1} (XY^T)^T XY^T = I$, yielding,

$$A_k (A_k^T A_k)^{-1} A_k = U_{\bar{p}} U_{\bar{p}}^T (= U_{\bar{p}} (U_{\bar{p}}^T U_{\bar{p}})^{-1} U_{\bar{p}}^T).$$

In other words, $\mathcal{R}(A_k) = \mathcal{R}(C) = \mathcal{R}(U_{\bar{p}})$.

Suppose that $(A_k^T A_k) \in \mathbb{R}^{d_k \times d_k}$ is nonsingular. Then, since $\mathcal{R}(A_k) = \mathcal{R}(C)$, $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) = \text{rank}(A_k) = d_k \geq p$, which is false in the case being analyzed (the case of $\text{rank}(W_H \cdots W_2) < p$). Thus, $A_k^T A_k$ is singular. Notice that for the boundary case with $k = H + 1$, $A_k^T A_k = I_{d_y}$, which is always nonsingular and thus the proof ends here (i.e., For the case with $k = H + 1$, since the latter condition, $X^T W_{H+1} \cdots W_{k+1} = 0$, implies a false statement, the former condition, $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) \geq \min(p, d_x)$, which is the desired statement, must be true).

If $A_k^T A_k$ is singular, from equation 5, it is inferred that we can perturb W_k to have $\text{rank}(W_k \cdots W_1) \geq \min(p, d_x)$. To see this in a concrete algebraic way, first note that since $\mathcal{R}(A_k) = \mathcal{R}(U_{\bar{p}})$, we can write $A_k = [U_{\bar{p}} \ \mathbf{0}] G_k$ for some $G_k \in GL_{d_k}(\mathbb{R})$ where $\mathbf{0} \in \mathbb{R}^{d_y \times (d_k - \bar{p})}$. Then, similarly to the base case with $k = 2$, plugging this into the condition in equation 5: for an arbitrary L_k ,

$$W_k \cdots W_1 = G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k \end{bmatrix}.$$

Since $\text{rank}(Y X^T (X X^T)^{-1}) = d_y$, the first \bar{p} rows in the above have rank \bar{p} . Thus, $W_k \cdots W_1$ has rank at least \bar{p} . On the other hand, since $W_{H+1} \cdots W_1 = A_k W_k \cdots W_1 = [U_{\bar{p}} \ \mathbf{0}] G_k W_k \cdots W_1$,

$$W_{H+1} \cdots W_1 = [U_{\bar{p}} \ \mathbf{0}] \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k \end{bmatrix} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1},$$

which means that changing the values of the last $(d_k - \bar{p})$ rows of $W_k \cdots W_1$ does not change the value of $\bar{\mathcal{L}}(W)$. Therefore, the original necessary condition implies a necessary condition that without changing the loss value, we can make $W_k \cdots W_1$ to have full rank with arbitrarily small perturbation on the last $(d_k - \bar{p})$ rows as $[0 \ I_{(d_k - \bar{p})}] G_k L_k + \epsilon M_{\text{ptb}}$ where ϵM_{ptb} is a perturbation matrix with arbitrarily small $\epsilon > 0$ (a proof of the existence of a corresponding perturbation matrix is exactly the same as the proof in the base case with $k = 2$, which is in footnote 7).

Similarly to the base case with $k = 2$, we can conclude that this perturbation can be done via a perturbation on each entry of W_k . From the above equation for $W_k \cdots W_1$, all the possible solutions of W_k can be written as: for an arbitrary L_0 and L_k ,

$$W_k = G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k \end{bmatrix} (W_{k-1} \cdots W_1)^\dagger + L_0^T (I - (W_{k-1} \cdots W_1)(W_{k-1} \cdots W_1)^\dagger).$$

Thus, we perturb W_k as

$$\begin{aligned} W_k &:= W_k + \epsilon G_k^{-1} \begin{bmatrix} 0 \\ M_{\text{ptb}} \end{bmatrix} (W_{k-1} \cdots W_1)^\dagger \\ &= G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k + \epsilon M_{\text{ptb}} \end{bmatrix} (W_{k-1} \cdots W_1)^\dagger + L_0^T (I - (W_{k-1} \cdots W_1)(W_{k-1} \cdots W_1)^\dagger). \end{aligned}$$

Note that upon such a perturbation, equation 5 may not hold anymore; i.e.,

$$G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k + \epsilon M_{\text{ptb}} \end{bmatrix} (W_{k-1} \cdots W_1)^\dagger (W_{k-1} \cdots W_1) \neq G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_2 + \epsilon M_{\text{ptb}} \end{bmatrix}.$$

This means that the original necessary condition that implies equation 5 no longer holds. In this case, we immediately conclude that the Hessian is no longer positive semidefinite and thus the point is a saddle point. We thereby consider the remaining case: equation 5 still holds. Then, with the perturbation on the entries of W_k ,

$$W_{H+1} \cdots W_1 = G_k^{-1} \begin{bmatrix} U_{\bar{p}}^T Y X^T (X X^T)^{-1} \\ [0 \ I_{(d_k - \bar{p})}] G_k L_k + \epsilon M_{\text{ptb}} \end{bmatrix},$$

as desired. Therefore, we have that $\text{rank}(W_k \cdots W_2) \geq \text{rank}(W_k \cdots W_1) \geq \min(p, d_x)$ upon such a perturbation.

Thus, we conclude the induction, proving that we can have $\text{rank}(W_H \cdots W_2) \geq \text{rank}(W_{H+1} \cdots W_2) \geq \min(p, d_x)$ with arbitrarily small perturbation of each parameter without changing the value of $\bar{\mathcal{L}}(W)$. If $p \leq d_x$, this means that upon such a perturbation, we have the case of $\text{rank}(W_H \cdots W_2) = p$ (since we have that $p \geq \text{rank}(W_H \cdots W_2) \geq p$ where the first inequality follows the definition of p), with which we have already proved the existence of some negative eigenvalue of the Hessian unless it is a global minimum. Thus, such a critical point is not a local minimum unless it is a global minimum. On the other hand, if $p > d_x$, upon such a perturbation, we have $\bar{p} \triangleq \text{rank}(W_{H+1} \cdots W_2) \geq d_x \geq d_y$. Thus, $W_{H+1} \cdots W_1 = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1} = U U^T Y X^T (X X^T)^{-1}$, which is a global minimum. We can see this in various ways. For example, $Xr = XY^T U U^T - XY^T = 0$, which means that it is a global minimum as discussed above.

Summarizing the above, any point that satisfies the definition (and necessary conditions) of a local minimum is a global minimum, concluding the proof of **Theorem 2.3 (ii)**. \square

B.2 Proof of Theorem 2.3 (i)

Proof We can prove the non-convexity and non-concavity from its Hessian (Theorem 2.3 (i)). First, consider $\bar{\mathcal{L}}(W)$. For example, from Corollary 4.5 with $k = H + 1$, it is necessary for the Hessian to be positive or negative semidefinite at a critical point that $\text{rank}(W_{H+1}) \geq \text{rank}(W_H \cdots W_2)$ or $Xr = 0$. The instances of W unsatisfying this condition at critical points form some uncountable set. For example, consider a uncountable set that consists of the points with $W_{H+1} = W_1 = 0$ and with any W_H, \dots, W_2 . Then, every point in the set defines a critical point from Lemma 4.1. Also, $Xr = XY^T \neq 0$ as $\text{rank}(XY^T) \geq 1$. So, it does not satisfy the first semidefinite condition. On the other hand, with any instance of $W_H \cdots W_2$ such that $\text{rank}(W_H \cdots W_2) \geq 1$, we have that $0 = \text{rank}(W_{H+1}) \not\geq \text{rank}(W_H \cdots W_2)$. So, it does not satisfy the second semidefinite condition as well. Thus, we have proved that in the domain of the loss function, there exist points, at which the Hessian becomes indefinite. **This implies Theorem 2.3 (i): the functions are non-convex and non-concave.** \square

B.3 Proof of Theorem 2.3 (iii)

Proof We now prove Theorem 2.3 (iii): every critical point that is not a global minimum is a saddle point. Here, we want to show that if the Hessian is negative semidefinite at a critical point, then there is an increasing direction so that there is no local maximum. Since $\bar{\mathcal{L}}(W) = \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^{d_y} ((W_{H+1})_{j,\cdot} \cdots W_1 X_{\cdot,i} - Y_{j,i})^2$,

$$\begin{aligned} \mathcal{D}_{(W_{H+1})_{1,t}} \bar{\mathcal{L}}(W) &= \frac{1}{2} \sum_{i=1}^m \mathcal{D}_{(W_{H+1})_{1,t}} ((W_{H+1})_{1,\cdot} \cdots W_1 X_{\cdot,i} - Y_{1,i})^2 \\ &= \sum_{i=1}^m ((W_{H+1})_{1,\cdot} \cdots W_1 X_{\cdot,i} - Y_{1,i}) \left(\mathcal{D}_{(W_{H+1})_{1,t}} \sum_{l=1}^{d_H} (W_{H+1})_{1,l} (W_H)_{l,\cdot} \cdots W_1 X_{\cdot,i} \right) \\ &= \sum_{i=1}^m ((W_{H+1})_{1,\cdot} \cdots W_1 X_{\cdot,i} - Y_{1,i}) ((W_H)_{t,\cdot} \cdots W_1 X_{\cdot,i}). \end{aligned}$$

Similarly,

$$\mathcal{D}_{(W_{H+1})_{1,t}} \mathcal{D}_{(W_{H+1})_{1,t}} \bar{\mathcal{L}}(W) = \sum_{i=1}^m ((W_H)_{t,\cdot} \cdots W_1 X_{\cdot,i})^2 \in \mathbb{R}.$$

Therefore, with other variables being fixed, $\bar{\mathcal{L}}$ is strictly convex in $(W_{H+1})_{t,1} \in \mathbb{R}$ coordinate for some t unless $(W_H)_{t,\cdot} \cdots W_1 X_{\cdot,i} = 0$ for all $i = 1, \dots, m$ and for all $t = 1, \dots, d_H$. Since $\text{rank}(X) = d_x$, in order to have $(W_H)_{t,\cdot} \cdots W_1 X_{\cdot,i} = 0$ for all $i = 1, \dots, m$, the dimension of the null space of $(W_H)_{t,\cdot} \cdots W_1$ must be at least d_x for each t . Since $(W_H)_{t,\cdot} \cdots W_1 \in \mathbb{R}^{1 \times d_x}$ for each t , this means that $(W_H)_{t,\cdot} \cdots W_1 = 0$ for all t . Therefore, with other variables being fixed, $\bar{\mathcal{L}}$ is strictly convex in $(W_{H+1})_{1,t} \in \mathbb{R}$ coordinate for some t if $W_H \cdots W_1 \neq 0$.

If $W_H \cdots W_1 = 0$, we claim that at a critical point, if the Hessian is negative semidefinite, we can make $W_H \cdots W_1 \neq 0$ with arbitrarily small perturbation of each parameter without changing the loss value. We can prove this by using the similar proof procedure to that used for Theorem 2.3 (ii) in the case of $\text{rank}(W_H \cdots W_2) < p$. Suppose that $W_H \cdots W_1 = 0$ and thus $\text{rank}(W_H \cdots W_1) = 0$. From Lemma 4.4, we have a following necessary condition for the Hessian to be (positive or negative) semidefinite at a critical point: for any $k \in \{2, \dots, H+1\}$,

$$\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad XrW_{H+1} \cdots W_{k+1} = 0,$$

where the first condition is shown to imply $\text{rank}(W_{H+1} \cdots W_k) \geq \text{rank}(W_{k-1} \cdots W_2)$ in Corollary 4.5.

Let $A_k = W_{H+1} \cdots W_{k+1}$. From the condition with $k = 2$, we have that $\text{rank}(W_{H+1} \cdots W_2) \geq d_1 \geq 1$ or $XrW_{H+1} \cdots W_3 = 0$. The former condition is false since $\text{rank}(W_H \cdots W_2) < 1$. From the latter condition, for an arbitrary L_2 ,

$$\begin{aligned} 0 &= XrW_{H+1} \cdots W_3 \\ \Leftrightarrow W_2 W_1 &= (A_2^T A_2)^- A_2^T Y X^T (X X^T)^{-1} + (I - (A_2^T A_2)^- A_2^T A_2) L_2 \\ \Leftrightarrow W_{H+1} \cdots W_1 &= A_2 (A_2^T A_2)^- A_2^T Y X^T (X X^T)^{-1} \\ &= C(C^T C)^- C^T Y X^T (X X^T)^{-1} \end{aligned} \quad (6)$$

where the last follow the critical point condition (Lemma 4.2). Then, similarly to the proof of Theorem 2.3 (ii),

$$A_2 (A_2^T A_2)^- A_2 = C(C^T C)^- C^T.$$

In other words, $\mathcal{R}(A_2) = \mathcal{R}(C)$.

Suppose that $\text{rank}(A_2^T A_2) \geq 1$. Then, since $\mathcal{R}(A_2) = \mathcal{R}(C)$, we have that $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) \geq 1$, which is false (or else the desired statement). Thus, $\text{rank}(A_2^T A_2) = 0$, which implies that $A_2 = 0$. Then, since $W_{H+1} \cdots W_1 = A_2 W_2 W_1$ with $A_2 = 0$, we can have $W_2 W_1 \neq 0$ without changing the loss value with arbitrarily small perturbation of W_2 and W_1 .

Thus, we showed that we can have $W_2 W_1 \neq 0$, with arbitrarily small perturbation of each parameter with the loss value being unchanged. To prove the corresponding results for $W_k \cdots W_2$ for any $k = 2, \dots, H$, we conduct induction on $k = 2, \dots, H$ with the same proof procedure. The proposition $P(k)$ to be proven is as follows: the necessary conditions with $j \leq k$ implies that we can have $W_k \cdots W_2 \neq 0$ with arbitrarily small perturbation of each parameter without changing the loss value. For the base case $k = 2$, we have already proved the proposition in the above.

For the inductive step with $k \geq 3$, we have the inductive hypothesis that we can have $W_{k-1} \cdots W_2 \neq 0$ with arbitrarily small perturbation of each parameter without changing the loss value. Accordingly, suppose that $W_{k-1} \cdots W_1 \neq 0$. Again, from Lemma 4.4, for any $k \in \{2, \dots, H+1\}$,

$$\mathcal{R}((W_{k-1} \cdots W_2)^T) \subseteq \mathcal{R}(C^T C) \quad \text{or} \quad XrW_{H+1} \cdots W_{k+1} = 0.$$

If the former is true, $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) \geq \text{rank}(W_{k-1} \cdots W_2) \geq \text{rank}(W_{k-1} \cdots W_2 W_1) \geq 1$, which is false (or the desired statement). If the latter is true, for an arbitrary L_1 ,

$$\begin{aligned} 0 &= XrW_{H+1} \cdots W_{k+1} \\ \Leftrightarrow W_k \cdots W_1 &= (A_k^T A_k)^- A_k^T Y X^T (X X^T)^{-1} + (I - (A_k^T A_k)^- A_k^T A_k) L_1 \\ \Leftrightarrow W_{H+1} \cdots W_1 &= A_k (A_k^T A_k)^- A_k^T Y X^T (X X^T)^{-1} \\ &= C(C^T C)^- C^T Y X^T (X X^T)^{-1} = U_{\bar{p}} U_{\bar{p}}^T Y X^T (X X^T)^{-1}, \end{aligned}$$

where the last follow the critical point condition (Lemma 4.2). Then, similarly to the above,

$$A_k (A_k^T A_k)^- A_k = C(C^T C)^- C^T.$$

In other words, $\mathcal{R}(A_k) = \mathcal{R}(C)$.

Suppose that $\text{rank}(A_k^T A_k) \geq 1$. Then, since $\mathcal{R}(A_k) = \mathcal{R}(C)$, we have that $\text{rank}(W_H \cdots W_2) \geq \text{rank}(C) = \text{rank}(A_k) \geq 1$, which is false (or the desired statement). Thus, $\text{rank}(A_k^T A_k) = 0$,

which implies that $A_k = 0$. Then, since $W_{H+1} \cdots W_1 = A_k W_k \cdots W_1$ with $A_k = 0$, we can have $W_k \cdots W_1 \neq 0$ without changing the loss value with arbitrarily small perturbation of each parameter.

Thus, we conclude the induction, proving that if $W_H \cdots W_1 = 0$, with arbitrarily small perturbation of each parameter without changing the value of $\bar{\mathcal{L}}(W)$, we can have $W_H \cdots W_2 \neq 0$. Thus, upon such a perturbation at any critical point with the negative semidefinite Hessian, the loss function is strictly convex in $(W_{H+1})_{1,t} \in \mathbb{R}$ coordinate for some t . That is, at any candidate point for a local maximum, there exists a strictly increasing direction in an arbitrarily small neighbourhood. This means that there is no local maximum. **Thus, we obtained the statement of Theorem 2.3 (i).**

□

B.4 Proof of Theorem 2.3 (iv)

Proof In the proof of Theorem 2.3 (ii), the case analysis with the case, $\text{rank}(W_H \cdots W_2) = p$, revealed that when $\text{rank}(W_H \cdots W_2) = p$, if $\nabla^2 \bar{\mathcal{L}}(W) \succeq 0$ at a critical point, W is a global minimum. Thus, when $\text{rank}(W_H \cdots W_2) = p$, if W is not a global minimum at a critical point, its Hessian is not positive semidefinite, containing some negative eigenvalue. From Theorem 2.3 (ii), if it is not a global minimum, it is not a local minimum. From Theorem 2.3 (iii), it is a saddle point. Thus, if $\text{rank}(W_H \cdots W_2) = p$, the Hessian at any saddle point has some negative eigenvalue, **which is the statement of Theorem 2.3 (iv).**

□

B.5 Proof of Theorem 3.2 and discussion of the assumptions

Proof

$$\begin{aligned} \mathbb{E}_Z[\mathcal{L}(W)] &= \mathbb{E}_Z \left[\frac{1}{2} \sum_{i=1}^m \sum_{j=1}^{d_y} (\hat{Y}(W, X)_{j,i} - Y_{j,i})^2 \right] \\ &= \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^{d_y} \mathbb{E}_Z[\hat{Y}(W, X)_{j,i}^2] - 2Y_{j,i} \mathbb{E}_Z[\hat{Y}(W, X)_{j,i}] + Y_{j,i}^2 \\ &= \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^{d_y} \rho^2 q^2 \left(\sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)} \right)^2 - 2\rho q Y_{j,i} \left(\sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)} \right) + Y_{j,i}^2 \end{aligned}$$

The first line follows the definition of the Frobenius norm. In the second line, we used the linearity of the expectation. The third line follows the independence assumption (A1p-m and A5u-m in (Choromanska *et al.*, 2015b,a)). That is, we have that $\mathbb{E}_Z[\hat{Y}(W, X)_{j,i}] = \rho q \sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)}$. Also, since $(\sum_{p=1}^k a_p)^2 = \sum_{p=1}^k a_p^2 + 2 \sum_{p < p'} a_p a_{p'}$ for any a and k , by denoting $a_{i,j,p} = [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)}$,

$$\begin{aligned} \mathbb{E}_Z[\hat{Y}(W, X)_{j,i}^2] &= \mathbb{E}_Z \left[\left(\sum_{p=1}^{\Psi_j} a_{i,j,p} [Z_i]_{(j,p)} \right)^2 \right] \\ &= \sum_{p=1}^{\Psi_j} a_{i,j,p}^2 \mathbb{E}_Z[[Z_i]_{(j,p)}^2] + 2 \sum_{p < p'} a_{i,j,p} a_{i,j,p'} \mathbb{E}_Z[[Z_i]_{(j,p)} [Z_i]_{(j,p')}] \\ &= \rho^2 \sum_{p=1}^{\Psi_j} a_{i,j,p}^2 + 2\rho^2 \sum_{p < p'} a_{i,j,p} a_{i,j,p'} \\ &= \rho^2 \left(\sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)} \right)^2 \end{aligned}$$

All the assumptions used above are subset of assumptions that were used, for example, in the first equation of the proof of theorem 3.3 in (Choromanska *et al.*, 2015a). Finally, since $q = \rho^{-1}$ and $\sum_{p=1}^{\Psi_j} [X_i]_{(j,p)} \prod_{k=1}^H w_{(j,p)} = (W_{H+1} W_H W_{H-1} \cdots W_2 W_1 X)_{j,i} = \bar{Y}_{j,i}$, the last line of the above equation for $\mathbb{E}_Z[\mathcal{L}(W)]$ is equal to $\frac{1}{2} \|\bar{Y} - Y\|_F^2 = \bar{\mathcal{L}}(W)$. Also, $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W) = \frac{1}{2} \|E[\hat{Y}(W, X)] - Y\|_F^2 = \frac{1}{2} \|E[\hat{Y}(W, X)] - Y\|_F^2 = \frac{1}{2} \|\bar{Y} - Y\|_F^2 = \bar{\mathcal{L}}(W)$.

Therefore, what we have proved to be true for $\bar{\mathcal{L}}(W)$ is also true for $\mathbb{E}_Z[\mathcal{L}(W)]$ and $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$. We conclude the proof of Theorem 3.2. \square

Note that we could reduce the loss functions $\mathbb{E}_Z[\mathcal{L}(W)]$ and $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$ to $\bar{\mathcal{L}}(W)$ only with a strict subset of the assumptions used in the previous work. Accordingly, a question might arise as to how much we can reshape the loss function with all the assumptions used in the previous work. To answer this question, we note that Choromanska *et al.* (2015b,a) reduced their loss functions of nonlinear neural networks to:

$$\mathbb{E}_{\xi, Z}[\mathcal{L}_{\text{hinge}}(W)_{1,1}] = \frac{1}{\Lambda^{(H-1)/2}} \sum_{i_1, i_2, \dots, i_H=1}^{\Lambda} X_{i_1, i_2, \dots, i_H} \tilde{w}_{i_1} \tilde{w}_{i_2} \cdots \tilde{w}_{i_H} \quad \text{s.t.} \quad \frac{1}{\Lambda} \sum_{i=1}^{\Lambda} \tilde{w}_i^2 = 1,$$

where $\Lambda \in \mathbb{R}$ is some constant related to the size of the network (i.e., *not* the matrix containing the eigenvalues). While we refer to (Choromanska *et al.*, 2015b,a) for the detailed definitions of the symbols, X and w are defined in the same way as ours are, and \tilde{w} is a modified version due to other assumptions that we did not adopt. Here, we observe that not only the model but also the loss function is linear in the inputs (the nonlinear activation function has disappeared—The inputs are simply multiplied by some coefficients and then summed). Moreover, the target function Y has disappeared (i.e., the loss value does not depend on the target function). That is, whatever the data points of Y are, their loss values are the same. Thus, we see that the loss functions can be reduced to much different functions with all the assumptions used in the previous work (i.e, A1p, A2p, A3p, A4p, A5u, A6u, and A7p). We adopted a strict subset of the assumptions, with which we reduced our loss function to a more realistic loss function of a deep neural network.

C Proofs of Corollaries 2.4 and 3.3

We complete the proofs of Corollaries 2.4 and 3.3.

Proof If $H = 1$, the condition in Theorem 2.3 (iv) reads "if $\text{rank}(W_1 \cdots W_2) = \text{rank}(I_{d_1}) = d_1 = p$ ", which is always true. This is because p is the smallest width of hidden layers and there is only one hidden layer, the width of which is d_1 . Thus, Theorem 2.3 (iv) immediately implies the statement of Corollary 2.4. For the statement of Corollary 2.4 with $H \geq 2$, it is suffice to show the existence of a simple set containing saddle points with the Hessian having no negative eigenvalue. Suppose that $W_H = W_{H-1} = \cdots = W_2 = W_1 = 0$. Then, from Lemma 4.1, it defines a uncountable set of critical points, in which W_{H+1} can vary in $\mathbb{R}^{d_y \times d_H}$. Since $r = Y^T \neq 0$ due to $\text{rank}(Y) \geq 1$, it is not a global minimum. To see this, we write

$$\begin{aligned} \bar{\mathcal{L}}(W) &= \frac{1}{2} \|\bar{Y}(W, X) - Y\|_F^2 = \frac{1}{2} \text{tr}(r^T r) \\ &= \frac{1}{2} \text{tr}(Y Y^T) - \frac{1}{2} \text{tr}(W_{H+1} \cdots W_1 X Y^T) - \frac{1}{2} \text{tr}((W_{H+1} \cdots W_1 X Y^T)^T) \\ &\quad + \frac{1}{2} \text{tr}(W_{H+1} \cdots W_1 X X^T (W_{H+1} \cdots W_1)^T). \end{aligned}$$

For example, with $W_{H+1} \cdots W_1 = \pm U_p U_p^T Y X^T (X X)^{-1}$,

$$\begin{aligned} \bar{\mathcal{L}}(W) &= \frac{1}{2} (\text{tr}(Y Y^T) - \text{tr}(U_p U_p^T \Sigma) - \text{tr}(\Sigma U_p U_p^T) + \text{tr}(U_p U_p^T \Sigma U_p U_p^T)) \\ &= \frac{1}{2} (\text{tr}(Y Y^T) - \text{tr}(U_p \Lambda_{1:p} U_p^T)) = \frac{1}{2} \left(\text{tr}(Y Y^T) \pm \sum_{k=1}^p \Lambda_{k,k} \right), \end{aligned}$$

where we can see that there exists a strictly lower value of $\bar{\mathcal{L}}(W)$ than the loss value with $r = Y^T$, which is $\frac{1}{2} \text{tr}(YY^T)$ (since $X \neq 0$ and $\text{rank}(\Sigma) \neq 0$).

Thus, these are not global minima, and thereby these are saddle points by Theorem 2.3 (ii) and (iii). On the other hand, from the proof of Lemma 4.3, every diagonal and off-diagonal element of the Hessian is zero if $W_H = W_{H-1} = \dots = W_2 = W_1 = 0$. Thus, the Hessian is simply a zero matrix, which has no negative eigenvalue. Using the argument in the proof of Theorem 3.2, we can deduce that the same results hold for $\mathbb{E}_Z[\mathcal{L}(W)]$ and $\mathcal{L}_{\mathbb{E}_Z[\hat{Y}]}(W)$.

□

D Discussion of the 1989 conjecture

The 1989 conjecture is based on the result for a 1-hidden layer network with $p < d_y = d_x$ (e.g., an autoencoder). That is, the previous work considered $\bar{Y} = W_2 W_1$ with the same loss function as ours with the additional assumption $p < d_y = d_x$. The previous work denotes $A \triangleq W_2$ and $B \triangleq W_1$.

The conjecture was expressed by Baldi & Hornik (1989) as

Our results, and in particular the main features of the landscape of E , hold true in the case of linear networks with several hidden layers.

Here, the “main features of the landscape of E ” refers to the following features, among other minor technical facts: 1) the function is convex in each matrix A (or B) when fixing other B (or A), and 2) every local minimum is a global minimum. No proof was provided in this work for this conjecture.

In 2012, the proof for the conjecture corresponding to the first feature (convexity in each matrix A (or B) when fixing other B (or A)) was provided in (Baldi & Lu, 2012) for both real-valued and complex-valued cases, while the proof for the conjecture for the second feature (every local minimum being a global minimum) was left for future work.

In (Baldi, 1989), there is an informal discussion regarding the conjecture. Let $i \in \{1, \dots, H\}$ be an index of a layer with the smallest width p . That is, $d_i = p$. We write

$$\begin{aligned} A &:= W_{H+1} \cdots W_{i+1} \\ B &:= W_i \cdots W_1. \end{aligned}$$

Then, what A and B can represent is the same as what the original $A := W_2$ and $B := W_1$, respectively, can represent in the 1-hidden layer case, assuming that $p < d_y = d_x$ (i.e., any element in $\mathbb{R}^{d_y \times p}$ and any element in $\mathbb{R}^{p \times d_x}$). Thus, we *would* conclude that all the local minima in the deeper models always correspond to the local minima of the collapsed 1-hidden layer version with $A := W_{H+1} \cdots W_{i+1}$ and $B := W_i \cdots W_1$.

However, the above reasoning turns out to be incomplete. Let us prove the incompleteness of the reasoning by contradiction in a way in which we can clearly see what goes wrong. Suppose that the reasoning is complete (i.e., the following statement is true: if we can collapse the model with the same expressiveness with the same rank restriction, then the local minima of the model correspond to the local minima of the collapsed model). Consider $f(w) = W_3 W_2 W_1 = 2w^2 + w^3$, where $W_1 = [w \ w \ w]$, $W_2 = [1 \ 1 \ w]^T$ and $W_3 = w$. Then, let us collapse the model as $a := W_3 W_2 W_1$ and $g(a) = a$. As a result, what $f(w)$ can represent is the same as what $g(a)$ can represent (i.e., any element in \mathbb{R}) with the same rank restriction (with a rank of at most one). Thus, with the same reasoning, we can conclude that every local minimum of $f(w)$ corresponds to a local minimum of $g(a)$. However, this is clearly false, as $f(w)$ is a non-convex function with a local minimum at $w = 0$ that is not a global minimum, while $g(a)$ is linear (convex and concave) without any local minima. The convexity for $g(a)$ is preserved after the composition with any norm. Thus, we have a contradiction, proving the incompleteness of the reasoning. What is missed in the reasoning is that even if what a model can represent is the same, the different parameterization creates different local structure in the loss surface, and thus different properties of the critical points (global minima, local minima, saddle points, and local maxima).

Now that we have proved the incompleteness of this reasoning, we discuss where the reasoning actually breaks down in a more concrete example. From Lemmas 4.1 and 4.2, if $H = 1$, we have

the following representation at critical points:

$$AB = A(A^T A)^{-1} A^T Y X^T (X X^T)^{-1}.$$

where $A := W_2$ and $B := W_1$. In contrast, from Lemmas 4.1 and 4.2, if H is arbitrary,

$$AB = C(C^T C)^{-1} C^T Y X^T (X X^T)^{-1}.$$

where $A := W_{H+1} \cdots W_{i+1}$ and $B := W_i \cdots W_1$ as discussed above, and $C = W_{H+1} \cdots W_2$. Note that by using other critical point conditions from Lemmas 4.1, we cannot obtain an expression such that $C = A$ in the above expression unless $i = 1$. Therefore, even though what A and B can represent is the same, the critical condition becomes different (and similarly, the conditions from the Hessian). Because the proof in the previous work with $H = 1$ heavily relies on the fact that $AB = A(A^T A)^{-1} A^T Y X^T (X X^T)^{-1}$, the same proof does not apply for deeper models (we may continue providing more evidence as to why the same proof does not work for deeper models, but one such example suffices for the purpose here).

In this respect, we have completed the proof of the conjecture and also provided a complete analytical proof for more general and detailed statements; that is, we did not assume that $p < d_y = d_x$, and we also proved saddle point properties with negative eigenvalue information.

