

On the classical r -matrix structure of the rational BC_n Ruijsenaars–Schneider–van Diejen system

B.G. Pusztai

Bolyai Institute, University of Szeged,
Aradi vértanúk tere 1, H-6720 Szeged, Hungary, *and*
MTA Lendület Holographic QFT Group, Wigner RCP,
H-1525 Budapest 114, P.O.B. 49, Hungary
e-mail: gpusztai@math.u-szeged.hu

Abstract

In this paper, we construct a quadratic r -matrix structure for the classical rational BC_n Ruijsenaars–Schneider–van Diejen system with the maximal number of three independent coupling parameters. As a byproduct, we provide a Lax representation of the dynamics as well.

Keywords: *Integrable systems; Ruijsenaars–Schneider–van Diejen models; Dynamical r -matrices*

MSC (2010): 70E40, 70G65, 70H06

PACS number: 02.30.Ik

arXiv:1508.03556v1 [math-ph] 14 Aug 2015

1 Introduction

The Ruijsenaars–Schneider–van Diejen (RSvD) models [1, 2] are among the most intensively studied integrable many particle systems, having numerous relationships with different branches of theoretical physics and pure mathematics. They had found applications first in the theory of the soliton equations [1, 3, 4, 5, 6], but soon they appeared in the Yang–Mills and the Seiberg–Witten theories as well (see e.g. [7, 8, 9, 10, 11]). Besides these well-known links, the RSvD systems and their non-relativistic limits, the Calogero–Moser–Sutherland (CMS) systems [12, 13, 14], have appeared in the context of random matrix theory, too. Making use of the action-angle duality between the different variants of the CMS and the RSvD systems, new classes of random matrix ensembles emerged in the literature [15, 16, 17], exhibiting spectacular statistical properties. Under the name of classical/quantum duality, it has also been observed that the Lax matrices of the CMS and the RSvD models encode the spectra of certain quantum spin chains, thereby the purely classical models provide an alternative way to analyze the quantum systems, without any reference to the celebrated Bethe Ansatz techniques (for details see e.g. [18, 19, 20, 21]). It is also worth mentioning that in the recent papers [22, 23] the authors have constructed new integrable tops, closely related to the CMS and the RSvD particle systems. Besides the Lax representation of the dynamics, in their studies the associated r -matrix structures also turn out to be indispensable.

The characteristic feature the above exciting new developments all share in common is the prominent role played by the Lax matrices of the CMS and the RSvD models. However, all these investigations are based on the translational invariant models associated with the A_n root system, exclusively. Apart from the technical difficulties, the probable explanation of this state of affair is the very limited knowledge about the Lax representation of the RSvD models in association with the non- A_n -type root systems. Of course, one can easily construct Lax representations for both the C_n -type and the BC_n -type RSvD models by the \mathbb{Z}_2 -folding of the A_{2n-1} and the A_{2n} root systems, respectively [24]. However, this trivial approach is only of very limited use, since the resulting models contain only a single coupling parameter. Nevertheless, working in a symplectic reduction framework, in our papers [25, 26] we succeeded in constructing Lax matrices for the rational C_n and the rational BC_n RSvD systems with the maximal number of independent coupling constants. Motivated by the plethora of potential applications outlined above, in this paper we work out the underlying classical r -matrix structures and also provide a Lax representation of the dynamics for the rational BC_n RSvD model with three independent coupling parameters.

Let us recall that the configuration space of the rational BC_n RSvD system is the open subset

$$\mathfrak{c} = \{\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n \mid \lambda_1 > \dots > \lambda_n > 0\} \subseteq \mathbb{R}^n, \quad (1.1)$$

that can be seen as an appropriate model for the standard open Weyl chamber of type BC_n . The cotangent bundle $T^*\mathfrak{c}$ is trivial, whence the phase space of the RSvD system can be identified with the product manifold

$$\mathcal{P}^R = \mathfrak{c} \times \mathbb{R}^n = \{(\lambda, \theta) \mid \lambda \in \mathfrak{c}, \theta \in \mathbb{R}^n\}, \quad (1.2)$$

that we endow with the symplectic form

$$\omega^R = 2 \sum_{c=1}^n d\theta_c \wedge d\lambda_c. \quad (1.3)$$

We mention in passing that the unusual numerical factor in ω^R is inserted purely for consistency with our earlier works [25, 26]. As for the dynamics, it is governed by the Hamiltonian

$$\begin{aligned} H^R = & \sum_{c=1}^n \cosh(2\theta_c) \left(1 + \frac{\nu^2}{\lambda_c^2}\right)^{\frac{1}{2}} \left(1 + \frac{\kappa^2}{\lambda_c^2}\right)^{\frac{1}{2}} \prod_{\substack{d=1 \\ (d \neq c)}}^n \left(1 + \frac{4\mu^2}{(\lambda_c - \lambda_d)^2}\right)^{\frac{1}{2}} \left(1 + \frac{4\mu^2}{(\lambda_c + \lambda_d)^2}\right)^{\frac{1}{2}} \\ & + \frac{\nu\kappa}{4\mu^2} \prod_{c=1}^n \left(1 + \frac{4\mu^2}{\lambda_c^2}\right) - \frac{\nu\kappa}{4\mu^2}, \end{aligned} \quad (1.4)$$

where μ , ν and κ are arbitrary real parameters satisfying $\mu < 0 < \nu$. Also, on these so-called coupling constants in this paper we impose the condition $\nu\kappa \geq 0$. As can be seen in [27], this additional requirement ensures that the particle system possesses only scattering trajectories.

Having defined the models of our interest, now we wish to outline the content of the rest of the paper. To keep our present work essentially self-contained, in Section 2 we briefly skim through the necessary Lie theoretic machinery and the symplectic reduction background, that provide the building blocks of the latter developments. Also, this section allows us to fix the notations. Starting with Section 3 we present our new results. Section 3 is the longest and the most technical part of our paper, in which we study of the r -matrix structure of the rational C_n RSvD model corresponding to the special choice $\kappa = 0$. Sticking to the Marsden–Weinstein reduction approach, in Subsection 3.1 we construct local extensions for the Lax matrix of the rational C_n RSvD model. Making use of these local sections, in Subsection 3.2 a series of short Propositions and Lemmas allows us to construct a classical r -matrix structure for the C_n -type model. In this respect our main result is Theorem 10, in which we formulate the r -matrix structure in a convenient quadratic form. The resulting quadratic r -matrices turn out to be fully dynamical, depending on all variables of the phase space \mathcal{P}^R . Subsequently, by switching to a purely algebraic approach, in Section 4 we generalize Theorem 10 to the rational BC_n RSvD system with three independent coupling constants. The quadratic r -matrix structure of the BC_n -type system is summarized in Theorem 11. To make this important result more transparent, in Theorem 12 we describe the r -matrix structure in a more convenient choice of gauge. In this gauge we also provide a Lax representation of the dynamics, as formulated in Theorem 13. Finally, in Section 5 we offer a short discussion on our results and also point out some open problems related to the RSvD systems.

2 Preliminaries

In this section we overview those Lie theoretic notions and results that underlie the geometric construction of the classical r -matrix structure for the rational C_n RSvD system. Our approach

is based on the symplectic reduction derivation of the RSvD models, that we also briefly outline. In Subsection 2.1 we closely follow the conventions of the standard reference [28], whereas in Subsection 2.2 we employ the notations introduced in our earlier work [25] on the RSvD systems.

2.1 Lie theoretic background

Take a positive integer $n \in \mathbb{N}$ and keep it fixed. Let $N = 2n$ and introduce the sets

$$\mathbb{N}_n = \{1, \dots, n\} \quad \text{and} \quad \mathbb{N}_N = \{1, \dots, N\}. \quad (2.1)$$

With the aid of the $N \times N$ matrix

$$\mathbf{C} = \begin{bmatrix} 0_n & \mathbf{1}_n \\ \mathbf{1}_n & 0_n \end{bmatrix} \quad (2.2)$$

we define the non-compact real reductive matrix Lie group

$$G = U(n, n) = \{y \in GL(N, \mathbb{C}) \mid y^* \mathbf{C} y = \mathbf{C}\}, \quad (2.3)$$

that we equip with the Cartan involution

$$\Theta: G \rightarrow G, \quad y \mapsto (y^{-1})^*. \quad (2.4)$$

Its fixed-point set

$$K = \{y \in G \mid \Theta(y) = y\} \quad (2.5)$$

is a maximal compact subgroup of G , having the identification $K \cong U(n) \times U(n)$.

On the Lie algebra

$$\mathfrak{g} = \mathfrak{u}(n, n) = \{Y \in \mathfrak{gl}(N, \mathbb{C}) \mid Y^* \mathbf{C} + \mathbf{C} Y = 0\} \quad (2.6)$$

the corresponding involution

$$\vartheta: \mathfrak{g} \rightarrow \mathfrak{g}, \quad Y \mapsto -Y^* \quad (2.7)$$

naturally induces the Cartan decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} \quad (2.8)$$

with the Lie subalgebra and the complementary subspace

$$\mathfrak{k} = \ker(\vartheta - \text{Id}_{\mathfrak{g}}) \quad \text{and} \quad \mathfrak{p} = \ker(\vartheta + \text{Id}_{\mathfrak{g}}), \quad (2.9)$$

respectively. That is, each element $Y \in \mathfrak{g}$ can be decomposed as

$$Y = Y_+ + Y_- \quad (2.10)$$

with unique components $Y_+ \in \mathfrak{k}$ and $Y_- \in \mathfrak{p}$. Notice that the \mathbb{Z}_2 -gradation (2.8) of \mathfrak{g} is actually orthogonal with respect to the non-degenerate Ad-invariant symmetric bilinear form

$$\langle \cdot, \cdot \rangle: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}, \quad (Y_1, Y_2) \mapsto \text{tr}(Y_1 Y_2). \quad (2.11)$$

To make our presentation simpler, for all $k, l \in \mathbb{N}_N$ we introduce the standard elementary matrix $e_{k,l} \in \mathfrak{gl}(N, \mathbb{C})$ with entries

$$(e_{k,l})_{k',l'} = \delta_{k,k'}\delta_{l,l'} \quad (k', l' \in \mathbb{N}_N). \quad (2.12)$$

Also, with each $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$ we associate the $N \times N$ diagonal matrix

$$\mathbf{\Lambda}(\lambda) = \text{diag}(\lambda_1, \dots, \lambda_n, -\lambda_1, \dots, -\lambda_n) \in \mathfrak{p}. \quad (2.13)$$

The set of diagonal matrices

$$\mathfrak{a} = \{\mathbf{\Lambda}(\lambda) \mid \lambda \in \mathbb{R}^n\} \quad (2.14)$$

forms a maximal Abelian subspace in \mathfrak{p} . Note that in \mathfrak{a} the family of matrices

$$D_c^- = \frac{1}{\sqrt{2}}(e_{c,c} - e_{n+c,n+c}) \quad (c \in \mathbb{N}_n) \quad (2.15)$$

forms an orthonormal basis, i.e. $\langle D_c^-, D_d^- \rangle = \delta_{c,d}$ for all $c, d \in \mathbb{N}_n$.

The centralizer of the Lie algebra \mathfrak{a} inside K is the Abelian Lie group

$$M = Z_K(\mathfrak{a}) = \{\text{diag}(e^{i\chi_1}, \dots, e^{i\chi_n}, e^{i\chi_1}, \dots, e^{i\chi_n}) \mid \chi_1, \dots, \chi_n \in \mathbb{R}\} \quad (2.16)$$

with Lie algebra

$$\mathfrak{m} = \{\text{diag}(i\chi_1, \dots, i\chi_n, i\chi_1, \dots, i\chi_n) \mid \chi_1, \dots, \chi_n \in \mathbb{R}\}. \quad (2.17)$$

In this Abelian Lie algebra the set of matrices

$$D_c^+ = \frac{i}{\sqrt{2}}(e_{c,c} + e_{n+c,n+c}) \quad (c \in \mathbb{N}_n) \quad (2.18)$$

forms a basis obeying the orthogonality relations $\langle D_c^+, D_d^+ \rangle = -\delta_{c,d}$ ($c, d \in \mathbb{N}_n$).

Let \mathfrak{m}^\perp and \mathfrak{a}^\perp denote the sets of the off-diagonal elements of \mathfrak{k} and \mathfrak{p} , respectively. With these subspaces can write the refined orthogonal decomposition

$$\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{m}^\perp \oplus \mathfrak{a} \oplus \mathfrak{a}^\perp. \quad (2.19)$$

In other words, each element $Y \in \mathfrak{g}$ can be uniquely decomposed as

$$Y = Y_{\mathfrak{m}} + Y_{\mathfrak{m}^\perp} + Y_{\mathfrak{a}} + Y_{\mathfrak{a}^\perp}, \quad (2.20)$$

where each component belongs to the subspace indicated by the subscript. In order to provide convenient bases in the subspaces \mathfrak{m}^\perp and \mathfrak{a}^\perp , for each $c \in \mathbb{N}_n$ we introduce the linear functional

$$\varepsilon_c: \mathbb{R}^n \rightarrow \mathbb{R}, \quad \lambda = (\lambda_1, \dots, \lambda_n) \mapsto \lambda_c. \quad (2.21)$$

Let us observe that the set of functionals

$$\mathcal{R}_+ = \{\varepsilon_a \pm \varepsilon_b \mid 1 \leq a < b \leq n\} \cup \{2\varepsilon_c \mid c \in \mathbb{N}_n\} \quad (2.22)$$

can be seen as a realization of a set of *positive* roots of type C_n . Now, associated with the positive root $2\varepsilon_c$ ($c \in \mathbb{N}_n$), we define the matrices

$$X_{2\varepsilon_c}^{\pm, i} = -\frac{i}{\sqrt{2}}(e_{c, n+c} \pm e_{n+c, c}). \quad (2.23)$$

In association with the other positive roots, for all $1 \leq a < b \leq n$ we define the following matrices with purely real entries:

$$\begin{aligned} X_{\varepsilon_a - \varepsilon_b}^{\pm, r} &= \frac{1}{2}(e_{a, b} \mp e_{b, a} \pm e_{n+a, n+b} - e_{n+b, n+a}), \\ X_{\varepsilon_a + \varepsilon_b}^{\pm, r} &= -\frac{1}{2}(e_{a, n+b} - e_{b, n+a} \pm e_{n+a, b} \mp e_{n+b, a}), \end{aligned} \quad (2.24)$$

together with the following ones with purely imaginary entries:

$$\begin{aligned} X_{\varepsilon_a - \varepsilon_b}^{\pm, i} &= \frac{i}{2}(e_{a, b} \pm e_{b, a} \pm e_{n+a, n+b} + e_{n+b, n+a}), \\ X_{\varepsilon_a + \varepsilon_b}^{\pm, i} &= -\frac{i}{2}(e_{a, n+b} + e_{b, n+a} \pm e_{n+a, b} \pm e_{n+b, a}). \end{aligned} \quad (2.25)$$

The point is that the set of vectors $\{X_\alpha^{+, \epsilon}\}$ forms a basis in the subspace \mathfrak{m}^\perp , whereas the family $\{X_\alpha^{-, \epsilon}\}$ provides a basis in \mathfrak{a}^\perp . Moreover, they obey the orthogonality relations

$$\langle X_\alpha^{+, \epsilon}, X_{\alpha'}^{+, \epsilon'} \rangle = -\delta_{\alpha, \alpha'} \delta_{\epsilon, \epsilon'} \quad \text{and} \quad \langle X_\alpha^{-, \epsilon}, X_{\alpha'}^{-, \epsilon'} \rangle = \delta_{\alpha, \alpha'} \delta_{\epsilon, \epsilon'}. \quad (2.26)$$

Note that the family of vectors

$$\{v_I\} \equiv \{D_c^\pm\} \cup \{X_\alpha^{\pm, \epsilon}\} \quad (2.27)$$

forms a basis in the real Lie algebra $\mathfrak{u}(n, n)$. We mention in passing that it is a basis in the complexification $\mathfrak{gl}(N, \mathbb{C}) \cong \mathfrak{u}(n, n)^\mathbb{C}$, too.

Next we turn to the linear operator

$$\text{ad}_{\mathbf{\Lambda}(\lambda)}: \mathfrak{g} \rightarrow \mathfrak{g}, \quad Y \mapsto [\mathbf{\Lambda}(\lambda), Y], \quad (2.28)$$

defined for each $\lambda \in \mathbb{R}^n$. The real convenience of the basis (2.27) stems from the commutation relations

$$\text{ad}_{\mathbf{\Lambda}(\lambda)}(D_c^\pm) = 0 \quad \text{and} \quad \text{ad}_{\mathbf{\Lambda}(\lambda)}(X_\alpha^{\pm, \epsilon}) = \alpha(\lambda) X_\alpha^{\mp, \epsilon}, \quad (2.29)$$

where $c \in \mathbb{N}_n$, $\alpha \in \mathcal{R}_+$ and $\epsilon \in \{r, i\}$. Notice that the subspace $\mathfrak{m}^\perp \oplus \mathfrak{a}^\perp$ is invariant under the linear operator $\text{ad}_{\mathbf{\Lambda}(\lambda)}$, whence the restriction

$$\widetilde{\text{ad}}_{\mathbf{\Lambda}(\lambda)} = \text{ad}_{\mathbf{\Lambda}(\lambda)}|_{\mathfrak{m}^\perp \oplus \mathfrak{a}^\perp} \in \mathfrak{gl}(\mathfrak{m}^\perp \oplus \mathfrak{a}^\perp) \quad (2.30)$$

is well-defined for all $\lambda \in \mathbb{R}^n$, with spectrum

$$\text{Spec}(\widetilde{\text{ad}}_{\mathbf{\Lambda}(\lambda)}) = \{\pm\alpha(\lambda) \mid \alpha \in \mathcal{R}_+\}. \quad (2.31)$$

The regular part of \mathfrak{a} is defined by the subset

$$\mathfrak{a}_{\text{reg}} = \{\mathbf{\Lambda}(\lambda) \mid \lambda \in \mathbb{R}^n \text{ and } \widetilde{\text{ad}}_{\mathbf{\Lambda}(\lambda)} \text{ is invertible}\}, \quad (2.32)$$

in which the standard Weyl chamber $\{\mathbf{\Lambda}(\lambda) \mid \lambda \in \mathfrak{c}\}$ is an appropriate connected component. Note that this Weyl chamber can be naturally identified with the configuration space \mathfrak{c} (1.1) of the rational BC_n RSvD system.

Having set up the algebraic stage, now we turn to some geometric results that are specific to the symplectic reduction derivation of the rational RSvD models. First, recall that the regular part of \mathfrak{p} (2.9) defined by

$$\mathfrak{p}_{\text{reg}} = \{k\mathbf{\Lambda}(\lambda)k^{-1} \mid \lambda \in \mathfrak{c} \text{ and } k \in K\} \quad (2.33)$$

is a dense and open subset of \mathfrak{p} . It is an important fact that with the smooth free right M -action

$$M \times (\mathfrak{c} \times K) \ni (m, (\lambda, k)) \mapsto (\lambda, km) \in \mathfrak{c} \times K \quad (2.34)$$

the map

$$\pi: \mathfrak{c} \times K \rightarrow \mathfrak{p}_{\text{reg}}, \quad (\lambda, k) \mapsto k\mathbf{\Lambda}(\lambda)k^{-1} \quad (2.35)$$

is a smooth principal M -bundle, providing the identification

$$\mathfrak{p}_{\text{reg}} \cong (\mathfrak{c} \times K)/M \cong \mathfrak{c} \times (K/M). \quad (2.36)$$

In the geometric construction of the dynamical r -matrix for the rational C_n RSvD model we shall utilize certain local sections of π with the characteristic properties below.

Proposition 1. *Take an arbitrary point $\lambda^{(0)} \in \mathfrak{c}$ and let $\mathbf{\Lambda}^{(0)} = \mathbf{\Lambda}(\lambda^{(0)})$. Then there is a smooth local section*

$$\check{\mathfrak{p}}_{\text{reg}} \ni Y \mapsto (e(Y), \sigma(Y)) \in \mathfrak{c} \times K \quad (2.37)$$

of π (2.35), defined on some open subset $\check{\mathfrak{p}}_{\text{reg}} \subseteq \mathfrak{p}_{\text{reg}}$, such that

$$\mathbf{\Lambda}^{(0)} \in \check{\mathfrak{p}}_{\text{reg}}, \quad (e(\mathbf{\Lambda}^{(0)}), \sigma(\mathbf{\Lambda}^{(0)})) = (\lambda^{(0)}, \mathbf{1}), \quad \text{ran}(\sigma_{*\mathbf{\Lambda}^{(0)}}) \subseteq \mathfrak{m}^\perp. \quad (2.38)$$

Moreover, under these conditions, at the point $\mathbf{\Lambda}^{(0)}$ the action of the derivatives of e and σ on the tangent vector $\delta Y \in \mathfrak{p} \cong T_{\mathbf{\Lambda}^{(0)}}\check{\mathfrak{p}}_{\text{reg}}$ takes the form

$$e_{*\mathbf{\Lambda}^{(0)}}(\delta Y) = (\delta Y_{1,1}, \dots, \delta Y_{n,n}) \in \mathbb{R}^n \cong T_{\lambda^{(0)}}\mathfrak{c}, \quad (2.39)$$

$$\sigma_{*\mathbf{\Lambda}^{(0)}}(\delta Y) = -(\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\delta Y)_{\mathfrak{a}^\perp}) \in \mathfrak{m}^\perp \subseteq \mathfrak{k} \cong T_{\mathbf{1}}K. \quad (2.40)$$

Proof. Notice that the point $(\lambda^{(0)}, \mathbf{1}) \in \mathfrak{c} \times K$ projects onto $\mathbf{\Lambda}^{(0)}$, that is,

$$\pi(\lambda^{(0)}, \mathbf{1}) = \mathbf{\Lambda}^{(0)}. \quad (2.41)$$

Differentiating π (2.35) at $(\lambda^{(0)}, \mathbf{1})$, let us observe that for each tangent vector

$$\delta\lambda \oplus \delta k \in \mathbb{R}^n \oplus \mathfrak{k} \cong T_{\lambda^{(0)}}\mathfrak{c} \oplus T_{\mathbf{1}}K \cong T_{(\lambda^{(0)}, \mathbf{1})}(\mathfrak{c} \times K) \quad (2.42)$$

we can write

$$\pi_{*(\lambda^{(0)}, \mathbf{1})}(\delta\lambda \oplus \delta k) = \left\{ \frac{d}{dt} \pi(\lambda^{(0)} + t\delta\lambda, e^{t\delta k}) \right\}_{t=0} = \mathbf{\Lambda}(\delta\lambda) - [\mathbf{\Lambda}^{(0)}, \delta k]. \quad (2.43)$$

Utilizing the linear operator (2.30), it is clear that

$$\pi_{*(\lambda^{(0)}, \mathbf{1})}(\delta\lambda \oplus \delta k) = \mathbf{\Lambda}(\delta\lambda) - \widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}}((\delta k)_{\mathfrak{m}^\perp}), \quad (2.44)$$

from where we conclude that

$$\ker(\pi_{*(\lambda^{(0)}, \mathbf{1})}) = \{0\} \oplus \mathfrak{m}. \quad (2.45)$$

Since the subspace $\mathbb{R}^n \oplus \mathfrak{m}^\perp$ is a complementary subspace of $\ker(\pi_{*(\lambda^{(0)}, \mathbf{1})})$ in the tangent space $T_{(\lambda^{(0)}, \mathbf{1})}(\mathfrak{c} \times K)$, it is evident that there exists a local section (e, σ) (2.37) satisfying the conditions imposed in (2.38). Moreover, by differentiating the equation

$$\pi \circ (e, \sigma) = \text{Id}_{\check{\mathfrak{p}}_{\text{reg}}} \quad (2.46)$$

at the point $\mathbf{\Lambda}^{(0)}$, the relationship (2.44) entails that for all $\delta Y \in \mathfrak{p} \cong T_{\mathbf{\Lambda}^{(0)}}\check{\mathfrak{p}}_{\text{reg}}$ we can write

$$\mathbf{\Lambda}(e_{*\mathbf{\Lambda}^{(0)}}(\delta Y)) - \widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}}(\sigma_{*\mathbf{\Lambda}^{(0)}}(\delta Y)) = \delta Y. \quad (2.47)$$

Projecting this equation onto the subspaces \mathfrak{a} and \mathfrak{a}^\perp , respectively, the formulae for the derivatives displayed in (2.39) and (2.40) follow at once. \square

To proceed further, we introduce the set of complex column vectors

$$S = \{V \in \mathbb{C}^N \mid \mathbf{C}V + V = 0 \text{ and } V^*V = N\}, \quad (2.48)$$

that can be naturally identified with a sphere of real dimension $2n - 1$. At each point $V \in S$ the tangent space to S can be identified with the real subspace of the complex column vectors

$$T_V S = \{\delta V \in \mathbb{C}^N \mid \mathbf{C}\delta V + \delta V = 0 \text{ and } (\delta V)^*V + V^*\delta V = 0\}, \quad (2.49)$$

that we endow with the inner product

$$\langle \delta V, \delta v \rangle_{T_V S} = \text{Re}((\delta V)^* \delta v) \quad (\delta V, \delta v \in T_V S). \quad (2.50)$$

Next, we introduce the distinguished column vector $E \in S$ with components

$$E_a = 1 \quad \text{and} \quad E_{n+a} = -1 \quad (a \in \mathbb{N}_n). \quad (2.51)$$

Also, with each vector $V \in S$ we associate the $N \times N$ matrix

$$\xi(V) = i\mu(VV^* - \mathbf{1}) + i(\mu - \nu)\mathbf{C} \in \mathfrak{k}. \quad (2.52)$$

Since the K -action on S defined by the smooth map

$$K \times S \ni (k, V) \mapsto kV \in S \quad (2.53)$$

is transitive, and since $k\xi(V)k^{-1} = \xi(kV)$ for all $k \in K$ and $V \in S$, it is clear that the adjoint orbit of K passing through the element $\xi(E) \in \mathfrak{k}$ has the form

$$\mathcal{O} = \{k\xi(E)k^{-1} \mid k \in K\} = \{\xi(V) \mid V \in S\}. \quad (2.54)$$

As is known, the orbit \mathcal{O} can be seen as an embedded submanifold of \mathfrak{k} , and for its tangent spaces we have the identifications

$$T_\rho \mathcal{O} = \{[X, \rho] \mid X \in \mathfrak{k}\} \subseteq \mathfrak{k} \quad (\rho \in \mathcal{O}). \quad (2.55)$$

In our earlier papers [29, 30, 25, 26] we have seen many times that this non-trivial minimal adjoint orbit plays a distinguished role in the symplectic reduction derivation of both the CMS and the RSvD systems. In this paper, throughout the construction of a dynamical r -matrix for the rational C_n RSvD system, we will also exploit that with the free $U(1)$ -action

$$U(1) \times S \ni (e^{i\psi}, V) \mapsto e^{i\psi}V \in S \quad (2.56)$$

the map

$$\xi: S \rightarrow \mathcal{O}, \quad V \mapsto \xi(V) \quad (2.57)$$

is a smooth principal $U(1)$ -bundle, providing the identification $\mathcal{O} \cong S/U(1)$. Recalling (2.52), it is clear that the derivative of ξ takes the form

$$\xi_{*V}(\delta V) = i\mu((\delta V)V^* + V(\delta V)^*) \in T_{\xi(V)}\mathcal{O} \quad (V \in S, \delta V \in T_V S), \quad (2.58)$$

whence it follows that

$$\ker(\xi_{*V}) = \mathbb{R}iV \quad \text{and} \quad (\ker(\xi_{*V}))^\perp = \{\delta v \in T_V S \mid (\delta v)^*V = V^*\delta v\}. \quad (2.59)$$

Let us also note that for all $X \in \mathfrak{k}$ and $V \in S$ we have $XV \in T_V S$ and

$$\xi_{*V}(XV) = [X, \xi(V)] \in T_{\xi(V)}\mathcal{O}. \quad (2.60)$$

The last two equations entail that for each $\delta V \in T_V S$ one can find a Lie algebra element $X \in \mathfrak{k}$ and a real number $t \in \mathbb{R}$ such that

$$\delta V = XV + tiV. \quad (2.61)$$

Having determined the derivative of ξ , now we shall work out the derivatives of certain local sections, that find applications in the latter developments.

Proposition 2. *Let $\mathcal{V}^{(0)} \in S$ be an arbitrary point and define $\rho^{(0)} = \xi(\mathcal{V}^{(0)}) \in \mathcal{O}$. Take a smooth local section*

$$W: \check{\mathcal{O}} \rightarrow S, \quad \rho \mapsto W(\rho) \quad (2.62)$$

of ξ (2.57), defined on some open subset $\check{\mathcal{O}} \subseteq \mathcal{O}$, satisfying the conditions

$$\rho^{(0)} \in \check{\mathcal{O}}, \quad W(\rho^{(0)}) = \mathcal{V}^{(0)}, \quad \text{ran}(W_{*\rho^{(0)}}) \subseteq (\ker(\xi_{*\mathcal{V}^{(0)}}))^\perp. \quad (2.63)$$

Then for the derivative of W at the point $\rho^{(0)}$ we have

$$W_{*\rho^{(0)}}([X, \rho^{(0)}]) = X\mathcal{V}^{(0)} - \frac{(\mathcal{V}^{(0)})^* X \mathcal{V}^{(0)}}{N} \mathcal{V}^{(0)} \quad (X \in \mathfrak{k}). \quad (2.64)$$

Proof. It is evident that there is a smooth local section W of the principal $U(1)$ -bundle ξ that satisfies the conditions displayed in (2.63). Take an arbitrary tangent vector $[X, \rho^{(0)}] \in T_{\rho^{(0)}}\mathcal{O}$ generated by some $X \in \mathfrak{k}$, and introduce the shorthand notation

$$\delta W = W_{*\rho^{(0)}}([X, \rho^{(0)}]) \in (\ker(\xi_{*\mathcal{V}^{(0)}}))^{\perp}. \quad (2.65)$$

By taking the derivative of the relationship $\xi \circ W = \text{Id}_{\mathcal{O}}$ at the point $\rho^{(0)}$, we find that

$$\xi_{*\mathcal{V}^{(0)}}(\delta W) = \xi_{*\mathcal{V}^{(0)}} \circ W_{*\rho^{(0)}}([X, \rho^{(0)}]) = [X, \rho^{(0)}] = \xi_{*\mathcal{V}^{(0)}}(X\mathcal{V}^{(0)}), \quad (2.66)$$

therefore $\delta W - X\mathcal{V}^{(0)} \in \ker(\xi_{*\mathcal{V}^{(0)}})$. However, due to (2.59) we can write that

$$\delta W = X\mathcal{V}^{(0)} + xi\mathcal{V}^{(0)} \quad (2.67)$$

with a unique real number x . Its value can be determined by the fact that the tangent vector δW belongs to subspace $(\ker(\xi_{*\mathcal{V}^{(0)}}))^{\perp}$, leading to the formula (2.64). \square

2.2 The rational C_n RSvD model from symplectic reduction

Based on our earlier results, in this subsection we review the symplectic reduction derivation of the rational C_n RSvD system. The surrounding ideas and the proofs can be found in [25]. An important ingredient of the symplectic reduction derivation of the RSvD system of our interest is the cotangent bundle T^*G of the Lie group G (2.3). For convenience, we trivialize T^*G by the left translations. Moreover, by identifying the dual space \mathfrak{g}^* with the Lie algebra \mathfrak{g} (2.6) via the bilinear form (2.11), it is clear that the product manifold $\mathcal{P} = G \times \mathfrak{g}$ provides an appropriate model for T^*G . For the tangent spaces of the manifold \mathcal{P} we have the natural identifications

$$T_{(y,Y)}\mathcal{P} \cong T_yG \oplus T_Y\mathfrak{g} \cong T_yG \oplus \mathfrak{g} \quad ((y, Y) \in \mathcal{P}), \quad (2.68)$$

and for the canonical symplectic form $\omega \in \Omega^2(\mathcal{P})$ we can write

$$\omega_{(y,Y)}(\Delta y \oplus \Delta Y, \delta y \oplus \delta Y) = \langle y^{-1}\Delta y, \delta Y \rangle - \langle y^{-1}\delta y, \Delta Y \rangle + \langle [y^{-1}\Delta y, y^{-1}\delta y], Y \rangle, \quad (2.69)$$

where $(y, Y) \in \mathcal{P}$ is an arbitrary point and $\Delta y \oplus \Delta Y, \delta y \oplus \delta Y \in T_yG \oplus \mathfrak{g}$ are arbitrary tangent vectors. An equally important building block in the geometric picture underlying reduction derivation of the RSvD model is the adjoint orbit \mathcal{O} (2.54). Of course, it carries the Kirillov–Kostant–Souriau symplectic form $\omega^{\mathcal{O}} \in \Omega^2(\mathcal{O})$, that can be written as

$$\omega_{\rho}^{\mathcal{O}}([X, \rho], [Z, \rho]) = \langle \rho, [X, Z] \rangle \quad (\rho \in \mathcal{O}, X, Z \in \mathfrak{k}). \quad (2.70)$$

Making use of the bundle ξ (2.57) and the equations (2.60) and (2.61), one can easily see that

$$\omega_{\xi(V)}^{\mathcal{O}}(\xi_{*V}(\delta V), \xi_{*V}(\delta v)) = 2\mu \text{Im}((\delta V)^*\delta v) \quad (V \in S, \delta V, \delta v \in T_V S). \quad (2.71)$$

Now, by taking the symplectic product of the symplectic manifolds (\mathcal{P}, ω) and $(\mathcal{O}, \omega^{\mathcal{O}})$, we introduce the extended phase space

$$(\mathcal{P}^{\text{ext}}, \omega^{\text{ext}}) = (\mathcal{P} \times \mathcal{O}, \omega + \omega^{\mathcal{O}}). \quad (2.72)$$

To describe the Poisson bracket on this space, for each smooth function $F \in C^\infty(\mathcal{P}^{\text{ext}})$, at each point $u = (y, Y, \rho) \in \mathcal{P}^{\text{ext}}$, we define the gradients

$$\nabla^G F(u) \in \mathfrak{g}, \quad \nabla^{\mathfrak{g}} F(u) \in \mathfrak{g}, \quad \nabla^{\mathcal{O}} F(u) \in T_\rho \mathcal{O} \quad (2.73)$$

by the natural requirement

$$F_{*u}(\delta y \oplus \delta Y \oplus [X, \rho]) = \langle \nabla^G F(u), y^{-1} \delta y \rangle + \langle \nabla^{\mathfrak{g}} F(u), \delta Y \rangle + \langle \nabla^{\mathcal{O}} F(u), X \rangle, \quad (2.74)$$

where $\delta y \in T_y G$, $\delta Y \in \mathfrak{g}$ and $X \in \mathfrak{k}$ are arbitrary elements. Now, one can easily verify that the Poisson bracket on \mathcal{P}^{ext} induced by the symplectic form ω^{ext} can be cast into the form

$$\begin{aligned} \{F, H\}^{\text{ext}}(u) &= \langle \nabla^G F(u), \nabla^{\mathfrak{g}} H(u) \rangle - \langle \nabla^{\mathfrak{g}} F(u), \nabla^G H(u) \rangle \\ &\quad - \langle [\nabla^{\mathfrak{g}} F(u), \nabla^{\mathfrak{g}} H(u)], Y \rangle + \omega_\rho^{\mathcal{O}}(\nabla^{\mathcal{O}} F(u), \nabla^{\mathcal{O}} H(u)), \end{aligned} \quad (2.75)$$

for all $F, H \in C^\infty(\mathcal{P}^{\text{ext}})$. To proceed further, let us note that the smooth map

$$\Phi^{\text{ext}}: (K \times K) \times \mathcal{P}^{\text{ext}} \rightarrow \mathcal{P}^{\text{ext}}, \quad ((k_L, k_R), (y, Y, \rho)) \mapsto (k_L y k_R^{-1}, k_R Y k_R^{-1}, k_L \rho k_L^{-1}) \quad (2.76)$$

is a symplectic left action of the product Lie group $K \times K$ on the extended phase space \mathcal{P}^{ext} , and it admits a $K \times K$ -equivariant momentum map

$$J^{\text{ext}}: \mathcal{P}^{\text{ext}} \rightarrow \mathfrak{k} \oplus \mathfrak{k}, \quad (y, Y, \rho) \mapsto ((y Y y^{-1})_+ + \rho) \oplus (-Y_+). \quad (2.77)$$

As we proved in [25], the rational C_n RSvD model can be derived by reducing the symplectic manifold \mathcal{P}^{ext} at the zero value of the momentum map J^{ext} .

Let us recall that the standard Marsden–Weinstein reduction consists of two major steps. At the outset, we need control over the level set

$$\mathfrak{L}_0 = (J^{\text{ext}})^{-1}(\{0\}) = \{u \in \mathcal{P}^{\text{ext}} \mid J^{\text{ext}}(u) = 0\}, \quad (2.78)$$

that turns out to be an embedded submanifold of \mathcal{P}^{ext} (2.72). However, to get a finer picture, we still need some more background material. First, for each $a \in \mathbb{N}_n$ we define the rational function

$$\mathfrak{c} \ni \lambda \mapsto z_a(\lambda) = - \left(1 + \frac{i\nu}{\lambda_a}\right) \prod_{\substack{d=1 \\ (d \neq a)}}^n \left(1 + \frac{2i\mu}{\lambda_a - \lambda_d}\right) \left(1 + \frac{2i\mu}{\lambda_a + \lambda_d}\right) \in \mathbb{C}. \quad (2.79)$$

Also, we need the vector-valued function $\mathcal{F}: \mathcal{P}^R \rightarrow \mathbb{C}^N$ with components

$$\mathcal{F}_a = e^{\theta_a} |z_a|^{\frac{1}{2}} \quad \text{and} \quad \mathcal{F}_{n+a} = e^{-\theta_a} \bar{z}_a |z_a|^{-\frac{1}{2}} \quad (a \in \mathbb{N}_n), \quad (2.80)$$

that allows us to introduce the function $\mathcal{A}: \mathcal{P}^R \rightarrow \exp(\mathfrak{p})$ with the matrix entries

$$\begin{aligned} \mathcal{A}_{a,b} &= \frac{2i\mu \mathcal{F}_a \bar{\mathcal{F}}_b}{2i\mu + \lambda_a - \lambda_b}, \quad \mathcal{A}_{n+a, n+b} = \frac{2i\mu \mathcal{F}_{n+a} \bar{\mathcal{F}}_{n+b}}{2i\mu - \lambda_a + \lambda_b}, \\ \mathcal{A}_{a, n+b} &= \bar{\mathcal{A}}_{n+b, a} = \frac{2i\mu \mathcal{F}_a \bar{\mathcal{F}}_{n+b}}{2i\mu + \lambda_a + \lambda_b} + \frac{i(\mu - \nu)}{i\mu + \lambda_a} \delta_{a,b}, \end{aligned} \quad (2.81)$$

where $a, b \in \mathbb{N}_n$. As we have seen in [25], function \mathcal{A} provides a Lax matrix for the rational C_n RSvD model with the two independent parameters μ and ν . Next, let us consider the smooth function $\mathcal{V}: \mathcal{P}^R \rightarrow S$ defined by the equation

$$\mathcal{V} = \mathcal{A}^{-\frac{1}{2}} \mathcal{F}, \quad (2.82)$$

and also introduce the product manifold

$$\mathcal{M}^R = \mathcal{P}^R \times (K \times K)/U(1)_*, \quad (2.83)$$

where $U(1)_*$ stands for the diagonal embedding of $U(1)$ in the product group $K \times K$. Having equipped with the above objects, now we are in a position to provide a convenient parametrization of the level set \mathfrak{L}_0 (2.78). Indeed, in [25] we proved that the map

$$\Upsilon^R: \mathcal{M}^R \rightarrow \mathcal{P}^{\text{ext}} \quad (2.84)$$

defined by the assignment

$$(\lambda, \theta, (\eta_L, \eta_R)U(1)_*) \mapsto (\eta_L \mathcal{A}(\lambda, \theta)^{\frac{1}{2}} \eta_R^{-1}, \eta_R \mathbf{\Lambda}(\lambda) \eta_R^{-1}, \eta_L \xi(\mathcal{V}(\lambda, \theta)) \eta_L^{-1}) \quad (2.85)$$

is a smooth injective immersion with image $\Upsilon^R(\mathcal{M}^R) = \mathfrak{L}_0$. Moreover, in [25] we also proved that Υ^R gives rise to a diffeomorphism from \mathcal{M}^R onto the embedded submanifold \mathfrak{L}_0 . In other words, the pair $(\mathcal{M}^R, \Upsilon^R)$ provides a model for the level set \mathfrak{L}_0 (2.78).

To complete the Marsden–Weinstein reduction, notice that the (residual) $K \times K$ -action on the model space \mathcal{M}^R (2.83) takes the form

$$(k_L, k_R) \cdot (\lambda, \theta, (\eta_L, \eta_R)U(1)_*) = (\lambda, \theta, (k_L \eta_L, k_R \eta_R)U(1)_*), \quad (2.86)$$

thus the orbit space $\mathcal{M}^R/(K \times K)$ can be naturally identified with the base manifold of the trivial principal $(K \times K)/U(1)_*$ -bundle

$$\pi^R: \mathcal{M}^R \rightarrow \mathcal{P}^R, \quad (\lambda, \theta, (\eta_L, \eta_R)U(1)_*) \mapsto (\lambda, \theta). \quad (2.87)$$

Now, the crux of the matter is the relationship

$$(\pi^R)^* \omega^R = (\Upsilon^R)^* \omega^{\text{ext}}, \quad (2.88)$$

that we proved in [25] by applying a chain of delicate arguments. Therefore, for the symplectic quotient in question we obtain the identification

$$(\mathcal{P}^{\text{ext}} //_0 (K \times K), \omega^{\text{red}}) \cong (\mathcal{P}^R, \omega^R). \quad (2.89)$$

Finally, note that the $K \times K$ -invariant function

$$f_1: \mathcal{P}^{\text{ext}} \rightarrow \mathbb{R}, \quad (y, Y, \rho) \mapsto \frac{1}{2} \text{tr}(y y^*) \quad (2.90)$$

survives the reduction, and by applying straightforward algebraic manipulations one can verify that the corresponding reduced function coincides with the Hamiltonian of the rational C_n RSvD system with two independent coupling parameters μ and ν , that can be obtained from the BC_n -type Hamiltonian (1.4) by setting $\kappa = 0$.

3 Dynamical r -matrix for the C_n -type model

Building on the symplectic reduction picture outlined in the previous subsection, our goal is to construct a classical r -matrix structure for the C_n -type rational RSvD system with two independent coupling parameters. In the context of the CMS models, this geometric approach goes back to the work of Avan, Babelon, and Talon [31]. Eventually, in our paper [32], we succeeded to construct a dynamical r -matrix for the most general hyperbolic BC_n Sutherland model with three independent coupling constants, too. It is worth mentioning that the surrounding ideas proves to be fruitful in the broader context of integrable field theories as well. For a systematic review see e.g. [33].

As we have seen in [25], the eigenvalues of the Lax matrix \mathcal{A} (2.81) do commute, whence it follows from general principles that \mathcal{A} obeys an r -matrix Poisson bracket (for proof, see e.g. [34, 35]). However, we wish to make this r -matrix structure as explicit as possible. For this reason, Subsection 3.1 is devoted to the study of certain local extensions for the Lax matrix of the rational C_n RSvD model. As it turns out, these local extensions are at the heart of the construction of the dynamical r -matrix structure for the RSvD system, that we elaborate in Subsection 3.2.

3.1 Local extensions of the Lax matrix \mathcal{A}

The backbone of our reduction approach is the construction of the so-called local extensions of the Lax operator \mathcal{A} (2.81), that we wish to describe below. For this reason, take an arbitrary point

$$(\lambda^{(0)}, \theta^{(0)}) \in \mathcal{P}^R \quad (3.1)$$

and keep it fixed. Clearly the point

$$s^{(0)} = (\lambda^{(0)}, \theta^{(0)}, (\mathbf{1}, \mathbf{1})U(1)_*) \in \mathcal{M}^R \quad (3.2)$$

is one of the representatives of $(\lambda^{(0)}, \theta^{(0)})$ in \mathcal{M}^R (2.83), that is, $\pi^R(s^{(0)}) = (\lambda^{(0)}, \theta^{(0)})$. Moreover, let us introduce the shorthand notations

$$\mathcal{A}^{(0)} = \mathcal{A}(\lambda^{(0)}, \theta^{(0)}), \quad \mathcal{F}^{(0)} = \mathcal{F}(\lambda^{(0)}, \theta^{(0)}), \quad \mathcal{V}^{(0)} = \mathcal{V}(\lambda^{(0)}, \theta^{(0)}), \quad (3.3)$$

together with

$$y^{(0)} = (\mathcal{A}^{(0)})^{\frac{1}{2}}, \quad Y^{(0)} = \mathbf{\Lambda}^{(0)} = \mathbf{\Lambda}(\lambda^{(0)}), \quad \rho^{(0)} = \xi(\mathcal{V}^{(0)}). \quad (3.4)$$

Corresponding to $s^{(0)}$ (3.2), in the extended phase space we also introduce the reference point

$$u^{(0)} = \Upsilon^R(s^{(0)}) = (y^{(0)}, Y^{(0)}, \rho^{(0)}) \in \mathcal{P}^{\text{ext}}. \quad (3.5)$$

Now, associated with the elements given in (3.3-3.4), let us choose a local section (e, σ) of π (2.35), and also a local section W of ξ (2.57), as described in Propositions 1 and 2, respectively. Upon defining the open subset

$$\check{\mathfrak{g}} = \{Y \in \mathfrak{g} \mid Y_- \in \check{\mathfrak{p}}_{\text{reg}}\} \subseteq \mathfrak{g}, \quad (3.6)$$

it is clear that

$$\Psi: G \times \check{\mathfrak{g}} \times \check{\mathcal{O}} \rightarrow \mathbb{C}^N, \quad (y, Y, \rho) \mapsto \sigma(Y_-)^{-1} y^* W(\rho) \quad (3.7)$$

is a well-defined smooth function. Due to the conditions imposed in the equations (2.38) and (2.63), at the point $u^{(0)}$ (3.5) for the first n components of Ψ we have

$$\Psi_a(u^{(0)}) = \mathcal{F}_a^{(0)} \quad (a \in \mathbb{N}_n). \quad (3.8)$$

Since these components are strictly positive, there is an open subset $\check{\mathcal{P}}^{\text{ext}} \subseteq G \times \check{\mathfrak{g}} \times \check{\mathcal{O}}$ containing the distinguished point $u^{(0)}$, such that for all $a \in \mathbb{N}_n$ the map

$$m_a: \check{\mathcal{P}}^{\text{ext}} \rightarrow U(1), \quad u \mapsto \frac{\Psi_a(u)}{|\Psi_a(u)|} \quad (3.9)$$

is well-defined and smooth. Let us keep in mind that by construction $m_a(u^{(0)}) = 1$.

Now we are in a position to define those group-valued functions that play the most important role in the construction of a dynamical r -matrix for the rational C_n RSvD system. First, making use of the functions m_a (3.9), we build up the M -valued function

$$m: \check{\mathcal{P}}^{\text{ext}} \rightarrow M, \quad u \mapsto \text{diag}(m_1(u), \dots, m_n(u), m_1(u), \dots, m_n(u)), \quad (3.10)$$

which satisfies $m(u^{(0)}) = \mathbf{1}$. Next, we introduce the K -valued functions

$$k: \check{\mathcal{P}}^{\text{ext}} \rightarrow K, \quad (y, Y, \rho) \mapsto \sigma(Y_-), \quad (3.11)$$

$$\varphi: \check{\mathcal{P}}^{\text{ext}} \rightarrow K, \quad u \mapsto k(u)m(u). \quad (3.12)$$

Clearly we have $k(u^{(0)}) = \varphi(u^{(0)}) = \mathbf{1}$. Finally, we define the functions

$$\mathbf{A}^{(0)}: \check{\mathcal{P}}^{\text{ext}} \rightarrow G, \quad (y, Y, \rho) \mapsto y^* y, \quad (3.13)$$

$$\mathbf{A}: \check{\mathcal{P}}^{\text{ext}} \rightarrow G, \quad u \mapsto \varphi(u)^{-1} \mathbf{A}^{(0)}(u) \varphi(u). \quad (3.14)$$

Notice that at the point $u^{(0)}$ (3.5) we have $\mathbf{A}^{(0)}(u^{(0)}) = \mathbf{A}(u^{(0)}) = \mathcal{A}^{(0)}$. Having equipped with the above objects, now we can formulate the central result of this subsection.

Lemma 3. *The G -valued smooth function \mathbf{A} (3.14) is a local extension of the Lax matrix \mathcal{A} (2.81) around the point $u^{(0)}$ in the sense that $\mathbf{A}(u^{(0)}) = \mathcal{A}(\lambda^{(0)}, \theta^{(0)})$ and*

$$\mathbf{A} \circ \Upsilon^R \Big|_{(\Upsilon^R)^{-1}(\check{\mathcal{P}}^{\text{ext}})} = \mathcal{A} \circ \pi^R \Big|_{(\Upsilon^R)^{-1}(\check{\mathcal{P}}^{\text{ext}})}. \quad (3.15)$$

Proof. It is enough to verify (3.15). For, take an arbitrary point

$$s = (\lambda, \theta, (\eta_L, \eta_R)U(1)_*) \in (\Upsilon^R)^{-1}(\check{\mathcal{P}}^{\text{ext}}) \quad (3.16)$$

with some $\lambda \in \mathfrak{c}$, $\theta \in \mathbb{R}^n$ and $\eta_L, \eta_R \in K$. Also, for brevity we define

$$u = (y, Y, \rho) = \Upsilon^R(s) \in \check{\mathcal{P}}^{\text{ext}}. \quad (3.17)$$

Recalling (2.35) and (2.85), it is clear that

$$Y = \eta_R \mathbf{A}(\lambda) \eta_R^{-1} = \pi(\lambda, \eta_R) \in \mathfrak{p}_{\text{reg}}. \quad (3.18)$$

On the other hand, since $Y \in \mathfrak{g}$, we have $Y_- \in \check{\mathfrak{p}}_{\text{reg}}$. Thus, making use of the local section (e, σ) introduced in (2.37), we see that

$$Y = Y_- = \pi(e(Y_-), \sigma(Y_-)) = \pi(e(Y_-), k(u)) \quad (3.19)$$

also holds. Recalling (2.34), the comparison of (3.18) and (3.19) yields that there is a unique element

$$\tilde{m} = \text{diag}(\tilde{m}_1, \dots, \tilde{m}_n, \tilde{m}_1, \dots, \tilde{m}_n) \in M \quad (3.20)$$

such that

$$(\lambda, \eta_R) = (e(Y_-), k(u)\tilde{m}). \quad (3.21)$$

Next, remembering the parametrization (2.85), we can write

$$\xi(\eta_L \mathcal{V}(\lambda, \theta)) = \eta_L \xi(\mathcal{V}(\lambda, \theta)) \eta_L^{-1} = \rho \in \check{\mathcal{O}}. \quad (3.22)$$

However, utilizing the local section W introduced in (2.62), we also have $\xi(W(\rho)) = \rho$, whence by (2.56) we can write that

$$\eta_L \mathcal{V}(\lambda, \theta) = e^{i\psi} W(\rho) \quad (3.23)$$

with some constant $\psi \in \mathbb{R}$. From the above observations it readily follows that

$$e^{i\psi} W(\rho) = \eta_L \mathcal{A}(\lambda, \theta)^{-\frac{1}{2}} \mathcal{F}(\lambda, \theta) = (y^*)^{-1} \eta_R \mathcal{F}(\lambda, \theta) = (y^*)^{-1} k(u) \tilde{m} \mathcal{F}(\lambda, \theta), \quad (3.24)$$

from where we get $e^{i\psi} \Psi(u) = \tilde{m} \mathcal{F}(\lambda, \theta)$. Componentwise, for each $a \in \mathbb{N}_n$ we can write

$$e^{i\psi} \Psi_a(u) = \tilde{m}_a \mathcal{F}_a(\lambda, \theta), \quad (3.25)$$

thus the relationship $|\Psi_a(u)| = \mathcal{F}_a(\lambda, \theta)$ and also

$$\tilde{m}_a = e^{i\psi} \frac{\Psi_a(u)}{\mathcal{F}_a(\lambda, \theta)} = e^{i\psi} \frac{\Psi_a(u)}{|\Psi_a(u)|} = e^{i\psi} m_a(u) \quad (3.26)$$

are evident. In other words, $\tilde{m} = e^{i\psi} m(u)$, whence from (3.21) and (3.12) we conclude that

$$\eta_R = k(u) \tilde{m} = e^{i\psi} k(u) m(u) = e^{i\psi} \varphi(u). \quad (3.27)$$

Now, turning to the functions (3.13) and (3.14), notice that

$$\mathbf{A}^{(0)}(u) = y^* y = \eta_R \mathcal{A}(\lambda, \theta) \eta_R^{-1}, \quad (3.28)$$

so from (3.27) we infer that

$$\mathbf{A} \circ \Upsilon^R(s) = \mathbf{A}(u) = \varphi(u)^{-1} \mathbf{A}^{(0)}(u) \varphi(y) = \mathcal{A}(\lambda, \theta) = \mathcal{A} \circ \pi^R(s). \quad (3.29)$$

Since s (3.16) is an arbitrary element of $(\Upsilon^R)^{-1}(\check{\mathcal{P}}^{\text{ext}})$, the Lemma follows. \square

3.2 Computing the r -matrix

The natural idea impregnated by Lemma 3 is that the Poisson brackets of the components of the Lax matrix \mathcal{A} (2.81) can be computed by inspecting the Poisson brackets of the components of the locally defined function \mathbf{A} (3.14). Indeed, since we reduce the symplectic manifold \mathcal{P}^{ext} (2.72) at the zero value of the $K \times K$ -equivariant momentum map J^{ext} (2.77), and since the local extension \mathbf{A} is (locally) $K \times K$ -invariant on the level set \mathfrak{L}_0 (2.78), using the St. Petersburg tensorial notation we can simply write

$$\{\mathcal{A} \otimes \mathcal{A}\}^R(\lambda^{(0)}, \theta^{(0)}) = \{\mathbf{A} \otimes \mathbf{A}\}^{\text{ext}}(u^{(0)}). \quad (3.30)$$

However, for the function $\mathbf{A}^{(0)}$ (3.13) we clearly have

$$\{\mathbf{A}^{(0)} \otimes \mathbf{A}^{(0)}\}^{\text{ext}} = 0, \quad (3.31)$$

that is, $\mathbf{A}^{(0)}$ obeys an r -matrix bracket with the trivial zero r -matrix. Therefore, due to the relationship $\mathbf{A} = \varphi^{-1} \mathbf{A}^{(0)} \varphi$ (3.14), it is clear that \mathbf{A} also obeys a linear r -matrix bracket

$$\{\mathbf{A} \otimes \mathbf{A}\}^{\text{ext}} = [\tilde{r}_{12}, \mathbf{A} \otimes \mathbf{1}] - [\tilde{r}_{21}, \mathbf{1} \otimes \mathbf{A}] \quad (3.32)$$

with the transformed r -matrix

$$\tilde{r}_{12} = \varphi_1^{-1} \varphi_2^{-1} \left(-\{\varphi_1, \mathbf{A}_2^{(0)}\}^{\text{ext}} \varphi_1^{-1} + \frac{1}{2} [\{\varphi_1, \varphi_2\}^{\text{ext}} \varphi_1^{-1} \varphi_2^{-1}, \mathbf{A}_2^{(0)}] \right) \varphi_1 \varphi_2. \quad (3.33)$$

Now, recalling that $\varphi(u^{(0)}) = \mathbf{1}$, from the relationships (3.30) and (3.33) we infer that for the Lax matrix \mathcal{A} we can write

$$\{\mathcal{A}_1, \mathcal{A}_2\}^R(\lambda^{(0)}, \theta^{(0)}) = [r_{12}(\lambda^{(0)}, \theta^{(0)}), \mathcal{A}_1(\lambda^{(0)}, \theta^{(0)})] - [r_{21}(\lambda^{(0)}, \theta^{(0)}), \mathcal{A}_2(\lambda^{(0)}, \theta^{(0)})] \quad (3.34)$$

with the r -matrix

$$r_{12}(\lambda^{(0)}, \theta^{(0)}) = -\{\varphi_1, \mathbf{A}_2^{(0)}\}^{\text{ext}}(u^{(0)}) + \frac{1}{2} [\{\varphi_1, \varphi_2\}^{\text{ext}}(u^{(0)}), \mathcal{A}_2(\lambda^{(0)}, \theta^{(0)})]. \quad (3.35)$$

However, since $\varphi = km$ (3.12), Leibniz rule yields

$$\{\varphi_1, \mathbf{A}_2^{(0)}\}^{\text{ext}}(u^{(0)}) = \{k_1, \mathbf{A}_2^{(0)}\}^{\text{ext}}(u^{(0)}) + \{m_1, \mathbf{A}_2^{(0)}\}^{\text{ext}}(u^{(0)}), \quad (3.36)$$

together with

$$\begin{aligned} \{\varphi_1, \varphi_2\}^{\text{ext}}(u^{(0)}) &= \{k_1, k_2\}^{\text{ext}}(u^{(0)}) + \{k_1, m_2\}^{\text{ext}}(u^{(0)}) \\ &\quad + \{m_1, k_2\}^{\text{ext}}(u^{(0)}) + \{m_1, m_2\}^{\text{ext}}(u^{(0)}). \end{aligned} \quad (3.37)$$

Thus, in order to provide an explicit formula for the above r -matrix (3.35), we still have to work out the Poisson brackets appearing in (3.36) and (3.37). However, recalling (2.75), it essentially boils down to the computation of the gradients (2.73) of the components of the functions $\mathbf{A}^{(0)}$, k and m . We accomplish these tasks in the following series of short Propositions.

Proposition 4. *At the reference point $u^{(0)}$ (3.5), for all matrix $v \in \mathfrak{gl}(N, \mathbb{C})$ we have*

$$(\nabla^G \operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = \frac{(v + v^*) \mathcal{A}^{(0)} + (\mathcal{A}^{(0)})^{-1} (v + v^*)}{2}, \quad (3.38)$$

$$(\nabla^G \operatorname{Im}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = \frac{(v - v^*) \mathcal{A}^{(0)} - (\mathcal{A}^{(0)})^{-1} (v - v^*)}{2i}, \quad (3.39)$$

whereas the remaining gradients of $\operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)}))$ and $\operatorname{Im}(\operatorname{tr}(v \mathbf{A}^{(0)}))$ are trivial, i.e.

$$(\nabla^{\mathfrak{g}} \operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = (\nabla^{\mathfrak{g}} \operatorname{Im}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = 0, \quad (3.40)$$

$$(\nabla^{\mathcal{O}} \operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = (\nabla^{\mathcal{O}} \operatorname{Im}(\operatorname{tr}(v \mathbf{A}^{(0)})))(u^{(0)}) = 0. \quad (3.41)$$

Proof. Take an arbitrary tangent vector

$$\Delta u = \delta y \oplus \delta Y \oplus [X, \rho^{(0)}] \in T_{u^{(0)}} \check{\mathcal{P}}^{\text{ext}} \quad (3.42)$$

with some Lie algebra element $X \in \mathfrak{k}$. By neglecting the second and the higher order terms in the small real parameter t , one can easily find that

$$\begin{aligned} \mathbf{A}^{(0)}(u^{(0)} + t\Delta u + \dots) \\ = \mathcal{A}^{(0)} + t \left(\mathcal{A}^{(0)} ((y^{(0)})^{-1} \delta y) + ((y^{(0)})^{-1} \delta y)^* \mathcal{A}^{(0)} \right) + \dots, \end{aligned} \quad (3.43)$$

from where we infer that

$$\begin{aligned} \operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)}))_{*u^{(0)}}(\Delta u) &= \left\{ \frac{d}{dt} \operatorname{Re}(\operatorname{tr}(v \mathbf{A}^{(0)}(u^{(0)} + t\Delta u + \dots))) \right\}_{t=0} \\ &= \operatorname{tr} \left(\frac{(v + v^*) \mathcal{A}^{(0)} + (\mathcal{A}^{(0)})^{-1} (v + v^*)}{2} (y^{(0)})^{-1} \delta y \right). \end{aligned} \quad (3.44)$$

Similarly, one obtains immediately that

$$\operatorname{Im}(\operatorname{tr}(v \mathbf{A}^{(0)}))_{*u^{(0)}}(\Delta u) = \operatorname{tr} \left(\frac{(v - v^*) \mathcal{A}^{(0)} - (\mathcal{A}^{(0)})^{-1} (v - v^*)}{2i} (y^{(0)})^{-1} \delta y \right). \quad (3.45)$$

Since the $N \times N$ matrices appearing on the right hand side of both (3.38) and (3.39) do belong to the Lie algebra \mathfrak{g} (2.6), by the definition of the gradients (2.74) the Proposition follows. \square

Proposition 5. *At the distinguished point $u^{(0)}$ (3.5), for all Lie algebra element $v \in \mathfrak{g}$ we have*

$$(\nabla^{\mathfrak{g}} \operatorname{Re}(\operatorname{tr}(vk)))(u^{(0)}) = \frac{1}{2} (\widetilde{\operatorname{ad}}_{\mathbf{A}^{(0)}})^{-1} ((v - v^*)_{\mathfrak{m}^\perp}) \quad \text{and} \quad (\nabla^{\mathfrak{g}} \operatorname{Im}(\operatorname{tr}(vk)))(u^{(0)}) = 0, \quad (3.46)$$

while the remaining gradients of $\operatorname{Re}(\operatorname{tr}(vk))$ and $\operatorname{Im}(\operatorname{tr}(vk))$ are trivial, i.e.

$$(\nabla^G \operatorname{Re}(\operatorname{tr}(vk)))(u^{(0)}) = (\nabla^G \operatorname{Im}(\operatorname{tr}(vk)))(u^{(0)}) = 0, \quad (3.47)$$

$$(\nabla^{\mathcal{O}} \operatorname{Re}(\operatorname{tr}(vk)))(u^{(0)}) = (\nabla^{\mathcal{O}} \operatorname{Im}(\operatorname{tr}(vk)))(u^{(0)}) = 0. \quad (3.48)$$

Proof. As in the proof of Proposition 4, take an arbitrary tangent vector Δu as given in (3.42). By applying a first order expansion on k (3.11) in the small real parameter t , Proposition 1 tells us that

$$k(u^{(0)} + t\Delta u + \dots) = \sigma((\mathbf{\Lambda}^{(0)} + t\delta Y)_-) = \sigma(\mathbf{\Lambda}^{(0)} + t(\delta Y)_-) = \mathbf{1} + t\delta\sigma + \dots, \quad (3.49)$$

where

$$\delta\sigma = -(\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\delta Y)_{\mathfrak{a}^\perp}) \in \mathfrak{m}^\perp. \quad (3.50)$$

Since for all $v \in \mathfrak{g}$ we have

$$\text{Re}(\text{tr}(vk)) = \frac{\text{tr}(vk) + \text{tr}(v^*k^{-1})}{2} \quad \text{and} \quad \text{Im}(\text{tr}(vk)) = \frac{\text{tr}(vk) - \text{tr}(v^*k^{-1})}{2i}, \quad (3.51)$$

it is now evident that

$$\text{Re}(\text{tr}(vk(u^{(0)} + t\Delta u + \dots))) = \frac{\text{tr}(v) + \text{tr}(v^*)}{2} + t\text{tr}\left(\frac{v - v^*}{2}\delta\sigma\right) + \dots. \quad (3.52)$$

Recalling the Cartan involution (2.7) we can write $v - v^* = v + \vartheta(v) \in \mathfrak{k}$, therefore

$$\begin{aligned} (\text{Re}(\text{tr}(vk)))_{*u^{(0)}}(\Delta u) &= -\frac{1}{2} \left\langle v + \vartheta(v), (\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\delta Y)_{\mathfrak{a}^\perp}) \right\rangle \\ &= \frac{1}{2} \left\langle (\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((v - v^*)_{\mathfrak{m}^\perp}), \delta Y \right\rangle. \end{aligned} \quad (3.53)$$

In complete analogy with (3.52), for the function $\text{Im}(\text{tr}(vk))$ we can write the expansion

$$\text{Im}(\text{tr}(vk(u^{(0)} + t\Delta u + \dots))) = \frac{\text{tr}(v) - \text{tr}(v^*)}{2i} + t\text{tr}\left(\frac{v + v^*}{2i}\delta\sigma\right) + \dots. \quad (3.54)$$

However, since $v + v^* = v - \vartheta(v) \in \mathfrak{p}$, and since the subspaces \mathfrak{k} and \mathfrak{p} (2.9) are orthogonal with respect to the bilinear form (2.11), we conclude that

$$(\text{Im}(\text{tr}(vk)))_{*u^{(0)}}(\Delta u) = \frac{1}{2i} \langle v - \vartheta(v), \delta\sigma \rangle = 0, \quad (3.55)$$

thus by recalling (2.74) the Proposition follows. \square

To find the gradients of the components of m (3.10) we find it convenient to introduce the auxiliary function

$$\tau: \mathbb{C} \setminus \{0\} \rightarrow U(1), \quad z \mapsto \frac{z}{|z|}. \quad (3.56)$$

It is clearly smooth, and at each point $x > 0$ for its derivative we have

$$\tau_{*x}(w) = i \frac{\text{Im}(w)}{x} \quad (w \in \mathbb{C} \cong T_x(\mathbb{C} \setminus \{0\})). \quad (3.57)$$

Note that with the aid of τ the function m_a (3.9) can be simply written as

$$m_a = \tau \circ \Psi_a. \quad (3.58)$$

Also, for each $a \in \mathbb{N}_n$ we introduce the $N \times N$ matrix

$$\zeta_a = i \frac{e_a(\mathcal{F}^{(0)})^* + \mathbf{C}(\mathcal{F}^{(0)})e_a^* \mathbf{C}}{2}, \quad (3.59)$$

where $e_k \in \mathbb{C}^N$ denotes the column vector with components

$$(e_k)_l = \delta_{k,l} \quad (k, l \in \mathbb{N}_N). \quad (3.60)$$

As a matter of fact, the above matrix ζ_a belongs to the Lie algebra \mathfrak{g} . Moreover, utilizing the basis (2.27), we can write

$$\begin{aligned} 2\zeta_a &= \sqrt{2}\mathcal{F}_a^{(0)}D_a^+ - \sqrt{2}\text{Re}(\mathcal{F}_{n+a}^{(0)})(X_{2\varepsilon_a}^{+,i} + X_{2\varepsilon_a}^{-,i}) \\ &+ \sum_{c=1}^{a-1} \left(\mathcal{F}_c^{(0)}(X_{\varepsilon_c - \varepsilon_a}^{+,i} - X_{\varepsilon_c - \varepsilon_a}^{-,i}) - \text{Re}(\mathcal{F}_{n+c}^{(0)})(X_{\varepsilon_c + \varepsilon_a}^{+,i} + X_{\varepsilon_c + \varepsilon_a}^{-,i}) \right. \\ &\quad \left. + \text{Im}(\mathcal{F}_{n+c}^{(0)})(X_{\varepsilon_c + \varepsilon_a}^{+,r} + X_{\varepsilon_c + \varepsilon_a}^{-,r}) \right) \\ &+ \sum_{c=a+1}^n \left(\mathcal{F}_c^{(0)}(X_{\varepsilon_a - \varepsilon_c}^{+,i} + X_{\varepsilon_a - \varepsilon_c}^{-,i}) - \text{Re}(\mathcal{F}_{n+c}^{(0)})(X_{\varepsilon_a + \varepsilon_c}^{+,i} + X_{\varepsilon_a + \varepsilon_c}^{-,i}) \right. \\ &\quad \left. - \text{Im}(\mathcal{F}_{n+c}^{(0)})(X_{\varepsilon_a + \varepsilon_c}^{+,r} + X_{\varepsilon_a + \varepsilon_c}^{-,r}) \right). \end{aligned} \quad (3.61)$$

Proposition 6. *Take an arbitrary $a \in \mathbb{N}_n$, then at the reference point $u^{(0)}$ (3.5) the gradients (2.73) of the function $\text{Re}(m_a)$ are all zeros. However, for the imaginary part of the function m_a (3.9) we have the non-trivial formulae*

$$(\nabla^G(\text{Im}(m_a)))(u^{(0)}) = \frac{1}{\mathcal{F}_a^{(0)}} \zeta_a \in \mathfrak{g}, \quad (3.62)$$

$$(\nabla^{\mathfrak{g}}(\text{Im}(m_a)))(u^{(0)}) = \frac{1}{\mathcal{F}_a^{(0)}} (\widetilde{\text{ad}}_{\mathbf{A}^{(0)}})^{-1}((\zeta_a)_{\mathfrak{m}^\perp}) \in \mathfrak{a}^\perp, \quad (3.63)$$

$$(\nabla^{\mathcal{O}}(\text{Im}(m_a)))(u^{(0)}) = \xi_{*\mathcal{V}^{(0)}}(\delta V_a) \in T_{\rho^{(0)}}\mathcal{O}, \quad (3.64)$$

where

$$\delta V_a = \frac{\mathbf{C}(\mathcal{A}^{(0)})^{\frac{1}{2}}e_a - (\mathcal{A}^{(0)})^{\frac{1}{2}}e_a}{4\mu\mathcal{F}_a^{(0)}} + \frac{\mathcal{V}^{(0)}}{2\mu N} \in T_{\mathcal{V}^{(0)}}S. \quad (3.65)$$

Proof. First, take an arbitrary tangent vector $\delta y \in T_{y^{(0)}}G$. Recalling (3.7), and the conditions (2.38), (2.63), for small values of the real parameter t we can write the first order expansion

$$\begin{aligned} &\Psi(y^{(0)} + t\delta y + \dots, Y^{(0)}, \rho^{(0)}) \\ &= \sigma((Y^{(0)})_-)^{-1}(y^{(0)} + t\delta y + \dots)^* W(\rho^{(0)}) = \mathcal{F}^{(0)} - t\mathbf{C}(y^{(0)})^{-1}(\delta y)\mathbf{C}\mathcal{F}^{(0)} + \dots, \end{aligned} \quad (3.66)$$

thus for each $a \in \mathbb{N}_n$ we have

$$\Psi_a(y^{(0)} + t\delta y + \dots, Y^{(0)}, \rho^{(0)}) = \mathcal{F}_a^{(0)} + t w_a + \dots \quad (3.67)$$

with $w_a = -\text{tr}(\mathbf{C}(y^{(0)})^{-1}(\delta y)\mathbf{C}\mathcal{F}^{(0)}e_a^*) \in \mathbb{C}$. Recalling (3.57), we can write

$$m_a(y^{(0)} + t\delta y + \dots, Y^{(0)}, \rho^{(0)}) = \tau(\mathcal{F}_a^{(0)} + tw_a + \dots) = 1 + ti\frac{\text{Im}(w_a)}{\mathcal{F}_a^{(0)}} + \dots, \quad (3.68)$$

thus clearly $(\nabla^G(\text{Re}(m_a)))(u^{(0)}) = 0$. Moreover, by inspecting w_a and (3.59), we obtain

$$(\text{Im}(m_a))_{*u^{(0)}}(\delta y \oplus 0 \oplus 0) = \frac{\text{Im}(w_a)}{\mathcal{F}_a^{(0)}} = \frac{\text{tr}(\zeta_a(y^{(0)})^{-1}\delta y)}{\mathcal{F}_a^{(0)}}, \quad (3.69)$$

so (3.62) also follows immediately.

Second, take an arbitrary tangent vector $\delta Y \in \mathfrak{g} \cong T_{Y^{(0)}}\mathfrak{g}$. According to Proposition 1, for small values of $t \in \mathbb{R}$ we have the first order expansion

$$\sigma((Y^{(0)} + t\delta Y)_-) = \sigma(\mathbf{\Lambda}^{(0)} + t(\delta Y)_-) = \mathbf{1} + t\delta\sigma + \dots, \quad (3.70)$$

with the Lie algebra element $\delta\sigma \in \mathfrak{m}^\perp$ displayed in (3.50). Therefore,

$$\Psi(y^{(0)}, Y^{(0)} + t\delta Y, \rho^{(0)}) = \sigma((Y^{(0)} + t\delta Y)_-)^{-1}(y^{(0)})^*W(\rho^{(0)}) = \mathcal{F}^{(0)} - t(\delta\sigma)\mathcal{F}^{(0)} + \dots, \quad (3.71)$$

and so for each $a \in \mathbb{N}_n$ we can write

$$\Psi_a(y^{(0)}, Y^{(0)} + t\delta Y, \rho^{(0)}) = \mathcal{F}_a^{(0)} + tw'_a + \dots \quad (3.72)$$

with $w'_a = -\text{tr}((\delta\sigma)\mathcal{F}^{(0)}e_a^*) \in \mathbb{C}$. Utilizing τ (3.56) and its derivative (3.57), we obtain

$$m_a(y^{(0)}, Y^{(0)} + t\delta Y, \rho^{(0)}) = \tau(\mathcal{F}_a^{(0)} + tw'_a + \dots) = 1 + ti\frac{\text{Im}(w'_a)}{\mathcal{F}_a^{(0)}} + \dots, \quad (3.73)$$

therefore $(\nabla^{\mathfrak{g}}(\text{Re}(m_a)))(u^{(0)}) = 0$ is immediate. Remembering (3.59) it is also clear that

$$\text{Im}(w'_a) = \langle \zeta_a, \delta\sigma \rangle = -\left\langle (\zeta_a)_{\mathfrak{m}^\perp}, (\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\delta Y)_{\mathfrak{a}^\perp}) \right\rangle = \left\langle (\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\zeta_a)_{\mathfrak{m}^\perp}), \delta Y \right\rangle. \quad (3.74)$$

Thus, by combining (3.73) and (3.74), we end up with the formula

$$(\text{Im}(m_a))_{*u^{(0)}}(0 \oplus \delta Y \oplus 0) = \frac{\text{Im}(w'_a)}{\mathcal{F}_a^{(0)}} = \left\langle \frac{1}{\mathcal{F}_a^{(0)}}(\widetilde{\text{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\zeta_a)_{\mathfrak{m}^\perp}), \delta Y \right\rangle, \quad (3.75)$$

that readily implies (3.63).

Third, take an arbitrary $X \in \mathfrak{k}$. Remembering (2.38), (2.63), and (3.7), notice that

$$\begin{aligned} & \Psi(y^{(0)}, Y^{(0)}, \rho^{(0)} + t[X, \rho^{(0)}] + \dots) \\ &= \mathcal{F}^{(0)} + t \left((\mathcal{A}^{(0)})^{\frac{1}{2}} X \mathcal{V}^{(0)} - \frac{(\mathcal{V}^{(0)})^* X \mathcal{V}^{(0)}}{N} \mathcal{F}^{(0)} \right) + \dots, \end{aligned} \quad (3.76)$$

whence for all $a \in \mathbb{N}_n$ we can write

$$\Psi_a(y^{(0)}, Y^{(0)}, \rho^{(0)} + t[X, \rho^{(0)}] + \dots) = \mathcal{F}_a^{(0)} + tw''_a + \dots \quad (3.77)$$

with the complex number

$$w''_a = \text{tr}((\mathcal{A}^{(0)})^{\frac{1}{2}} X \mathcal{V}^{(0)} e_a^*) - \frac{\mathcal{F}_a^{(0)}}{N} \text{tr}(X \mathcal{V}^{(0)} (\mathcal{V}^{(0)})^*). \quad (3.78)$$

It readily follows that

$$m_a(y^{(0)}, Y^{(0)}, \rho^{(0)} + t[X, \rho^{(0)}] + \dots) = \tau(\mathcal{F}_a^{(0)} + t w''_a + \dots) = 1 + t i \frac{\text{Im}(w''_a)}{\mathcal{F}_a^{(0)}} + \dots, \quad (3.79)$$

from where we get at once that $(\nabla^{\mathcal{O}}(\text{Re}(m_a)))(u^{(0)}) = 0$ and

$$(\text{Im}(m_a))_{*u^{(0)}}(0 \oplus 0 \oplus [X, \rho^{(0)}]) = \frac{\text{Im}(w''_a)}{\mathcal{F}_a^{(0)}}. \quad (3.80)$$

At this point notice that $i\mathcal{V}^{(0)}(\mathcal{V}^{(0)})^* \in \mathfrak{k}$. Therefore, recalling (3.59), we can write

$$\begin{aligned} \text{Im}(w''_a) &= -\text{tr}((\mathcal{A}^{(0)})^{\frac{1}{2}} \zeta_a (\mathcal{A}^{(0)})^{-\frac{1}{2}} X) + \frac{\mathcal{F}_a^{(0)}}{N} \text{tr}(i\mathcal{V}^{(0)} (\mathcal{V}^{(0)})^* X) \\ &= \mathcal{F}_a^{(0)} \left\langle -\frac{1}{\mathcal{F}_a^{(0)}} \left((\mathcal{A}^{(0)})^{\frac{1}{2}} \zeta_a (\mathcal{A}^{(0)})^{-\frac{1}{2}} \right)_{\mathfrak{k}} + \frac{i}{N} \mathcal{V}^{(0)} (\mathcal{V}^{(0)})^*, X \right\rangle. \end{aligned} \quad (3.81)$$

Now, one can verify that the column vector δV_a displayed in (3.65) does belong to the tangent space $T_{\mathcal{V}^{(0)}} S$ (2.49). Furthermore, recalling (2.58) we find that

$$-\frac{1}{\mathcal{F}_a^{(0)}} \left((\mathcal{A}^{(0)})^{\frac{1}{2}} \zeta_a (\mathcal{A}^{(0)})^{-\frac{1}{2}} \right)_{\mathfrak{k}} + \frac{i}{N} \mathcal{V}^{(0)} (\mathcal{V}^{(0)})^* = \xi_{*\mathcal{V}^{(0)}}(\delta V_a), \quad (3.82)$$

thus the relationship (3.64) also follows. \square

Having the necessary gradients at our disposal, now we are ready to work out the tensorial Poisson brackets appearing in (3.36) and (3.37).

Lemma 7. *At the point $u^{(0)}$ (3.5) we can write*

$$\{k \otimes \mathbf{A}^{(0)}\}^{\text{ext}}(u^{(0)}) = \mathcal{A}_2^{(0)} \left(\sum_{\alpha, \epsilon} \frac{X_{\alpha}^{+, \epsilon} \otimes X_{\alpha}^{-, \epsilon}}{\alpha(\lambda^{(0)})} \right) + \left(\sum_{\alpha, \epsilon} \frac{X_{\alpha}^{+, \epsilon} \otimes X_{\alpha}^{-, \epsilon}}{\alpha(\lambda^{(0)})} \right) \mathcal{A}_2^{(0)}, \quad (3.83)$$

$$\{m \otimes \mathbf{A}^{(0)}\}^{\text{ext}}(u^{(0)}) = -\mathcal{A}_2^{(0)} \left(\sum_{a=1}^n D_a^+ \otimes S_a^{(0)} \right) - \left(\sum_{a=1}^n D_a^+ \otimes S_a^{(0)} \right) \mathcal{A}_2^{(0)}, \quad (3.84)$$

where for each $a \in \mathbb{N}_n$ we have

$$\begin{aligned} S_a^{(0)} &= \frac{1}{\sqrt{2}\mathcal{F}_a^{(0)}} \left\{ -\frac{\sqrt{2}\text{Re}(\mathcal{F}_{n+a}^{(0)}) X_{2\varepsilon_a}^{-, i}}{2\lambda_a^{(0)}} \right. \\ &\quad + \sum_{c=1}^{a-1} \left(\frac{\mathcal{F}_c^{(0)} X_{\varepsilon_c - \varepsilon_a}^{-, i}}{\lambda_c^{(0)} - \lambda_a^{(0)}} - \frac{\text{Re}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_c + \varepsilon_a}^{-, i}}{\lambda_c^{(0)} + \lambda_a^{(0)}} + \frac{\text{Im}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_c + \varepsilon_a}^{-, r}}{\lambda_c^{(0)} + \lambda_a^{(0)}} \right) \\ &\quad \left. + \sum_{c=a+1}^n \left(\frac{\mathcal{F}_c^{(0)} X_{\varepsilon_a - \varepsilon_c}^{-, i}}{\lambda_a^{(0)} - \lambda_c^{(0)}} - \frac{\text{Re}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_a + \varepsilon_c}^{-, i}}{\lambda_a^{(0)} + \lambda_c^{(0)}} - \frac{\text{Im}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_a + \varepsilon_c}^{-, r}}{\lambda_a^{(0)} + \lambda_c^{(0)}} \right) \right\}. \end{aligned} \quad (3.85)$$

Proof. To prove (3.83), we utilize the family of $N \times N$ matrices $\{v_I\}$ defined in (2.27), that forms a basis in the complex linear space $\mathfrak{gl}(N, \mathbb{C})$. Recalling Proposition 5, we see that among the members of the basis $\{v_I\}$ only the vectors $X_\alpha^{+, \epsilon}$ ($\alpha \in \mathcal{R}_+, \epsilon \in \{\pm\}$) generate non-trivial gradients of the form

$$\nabla^{\mathfrak{g}}(\operatorname{Re}(\operatorname{tr}(X_\alpha^{+, \epsilon} k)))(u^{(0)}) = \frac{1}{\alpha(\lambda^{(0)})} X_\alpha^{-, \epsilon}. \quad (3.86)$$

Let $\{v^I\} \subseteq \mathfrak{gl}(N, \mathbb{C})$ denote the dual basis of $\{v_I\}$ provided by the trace-pairing of $\mathfrak{gl}(N, \mathbb{C})$. Recalling Proposition 4 and the explicit expression of the gradients (3.86), the Poisson bracket formula (2.75) allows us to write

$$\begin{aligned} \{k \otimes \mathbf{A}^{(0)}\}^{\operatorname{ext}}(u^{(0)}) &= \sum_{I, J} \{\operatorname{tr}(v_I k), \operatorname{tr}(v_J \mathbf{A}^{(0)})\}^{\operatorname{ext}}(u^{(0)}) v^I \otimes v^J \\ &= \sum_{\alpha, \epsilon} \sum_J \frac{\operatorname{tr}(v_J \mathbf{A}^{(0)} X_\alpha^{-, \epsilon}) - \operatorname{tr}(v_J \mathbf{C} X_\alpha^{-, \epsilon} \mathbf{C} \mathbf{A}^{(0)})}{\alpha(\lambda^{(0)})} X_\alpha^{+, \epsilon} \otimes v^J \\ &= \sum_{\alpha, \epsilon} \frac{1}{\alpha(\lambda^{(0)})} X_\alpha^{+, \epsilon} \otimes (\mathbf{A}^{(0)} X_\alpha^{-, \epsilon} + X_\alpha^{-, \epsilon} \mathbf{A}^{(0)}), \end{aligned} \quad (3.87)$$

so (3.83) follows immediately.

Making use of Propositions 4 and 6, let us notice that the Poisson bracket formula (2.75) yields

$$\begin{aligned} \{m \otimes \mathbf{A}^{(0)}\}^{\operatorname{ext}}(u^{(0)}) &= - \sum_{a=1}^n \sum_{k, l=1}^N \{\mathbf{A}_{k, l}^{(0)}, m_a\}^{\operatorname{ext}}(u^{(0)}) (e_{a, a} + e_{n+a, n+a}) \otimes e_{k, l} \\ &= - \sum_{a=1}^n D_a^+ \otimes \sqrt{2} (\mathbf{A}^{(0)} (\nabla^{\mathfrak{g}}(\operatorname{Im}(m_a)))(u^{(0)}) + (\nabla^{\mathfrak{g}}(\operatorname{Im}(m_a)))(u^{(0)}) \mathbf{A}^{(0)}). \end{aligned} \quad (3.88)$$

Therefore, by projecting the Lie algebra element ζ_a (3.61) onto the subspace \mathfrak{m}^\perp , the application of (3.63) immediately leads to (3.84). \square

Lemma 8. *At $u^{(0)}$ (3.5) we have the trivial Poisson bracket $\{k \otimes k\}^{\operatorname{ext}}(u^{(0)}) = 0$, whereas*

$$\{m \otimes k\}^{\operatorname{ext}}(u^{(0)}) = \sum_{a=1}^n D_a^+ \otimes T_a^{(0)}, \quad (3.89)$$

where for each $a \in \mathbb{N}_n$ we have

$$\begin{aligned} T_a^{(0)} &= \frac{1}{\sqrt{2} \mathcal{F}_a^{(0)}} \left\{ \frac{\sqrt{2} \operatorname{Re}(\mathcal{F}_{n+a}^{(0)}) X_{2\varepsilon_a}^{+, i}}{2\lambda_a^{(0)}} \right. \\ &\quad + \sum_{c=1}^{a-1} \left(\frac{\mathcal{F}_c^{(0)} X_{\varepsilon_c - \varepsilon_a}^{+, i}}{\lambda_c^{(0)} - \lambda_a^{(0)}} + \frac{\operatorname{Re}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_c + \varepsilon_a}^{+, i}}{\lambda_c^{(0)} + \lambda_a^{(0)}} - \frac{\operatorname{Im}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_c + \varepsilon_a}^{+, r}}{\lambda_c^{(0)} + \lambda_a^{(0)}} \right) \\ &\quad \left. + \sum_{c=a+1}^n \left(-\frac{\mathcal{F}_c^{(0)} X_{\varepsilon_a - \varepsilon_c}^{+, i}}{\lambda_a^{(0)} - \lambda_c^{(0)}} + \frac{\operatorname{Re}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_a + \varepsilon_c}^{+, i}}{\lambda_a^{(0)} + \lambda_c^{(0)}} + \frac{\operatorname{Im}(\mathcal{F}_{n+c}^{(0)}) X_{\varepsilon_a + \varepsilon_c}^{+, r}}{\lambda_a^{(0)} + \lambda_c^{(0)}} \right) \right\}. \end{aligned} \quad (3.90)$$

Proof. Working with the basis $\{v_I\}$ (2.27) of $\mathfrak{gl}(N, \mathbb{C})$, from Proposition 5 we see that

$$\nabla^{\mathfrak{g}}(\operatorname{Re}(\operatorname{tr}(v_I k)))(u^{(0)}) \in \mathfrak{a}^\perp \subseteq \mathfrak{p}. \quad (3.91)$$

Keeping in mind the orthogonal \mathbb{Z}_2 -gradation (2.8), the Poisson bracket formula (2.75) gives rise to the relationship

$$\begin{aligned} & \{\operatorname{tr}(v_I k), \operatorname{tr}(v_J k)\}^{\operatorname{ext}}(u^{(0)}) \\ &= \left\langle [\nabla^{\mathfrak{g}}(\operatorname{Re}(\operatorname{tr}(v_I k)))(u^{(0)}), \nabla^{\mathfrak{g}}(\operatorname{Re}(\operatorname{tr}(v_J k)))(u^{(0)})], \mathbf{\Lambda}^{(0)} \right\rangle = 0, \end{aligned} \quad (3.92)$$

thus the equation $\{k \otimes k\}^{\operatorname{ext}}(u^{(0)}) = 0$ follows immediately.

As we have already observed at the beginning of the proof of Lemma 7, among the members of the basis $\{v_I\}$ only the vectors $v = X_\alpha^{+, \epsilon}$ generate non-trivial gradients for the component functions $\operatorname{tr}(vk)$ at the point $u^{(0)}$. Utilizing these gradients (3.86), Propositions 5 and 6 allow us to write

$$\begin{aligned} \{m \otimes k\}^{\operatorname{ext}}(u^{(0)}) &= \sum_{a=1}^n \sum_{\alpha, \epsilon} \{m_a, \operatorname{tr}(X_\alpha^{+, \epsilon} k)\}^{\operatorname{ext}}(u^{(0)})(e_{a,a} + e_{n+a, n+a}) \otimes (-X_\alpha^{+, \epsilon}) \\ &= - \sum_{a=1}^n \sum_{\alpha, \epsilon} \langle \nabla^G(\operatorname{Im}(m_a))(u^{(0)}), \nabla^{\mathfrak{g}}(\operatorname{Re}(\operatorname{tr}(X_\alpha^{+, \epsilon} k)))(u^{(0)}) \rangle \sqrt{2} D_a^+ \otimes X_\alpha^{+, \epsilon} \\ &= - \sum_{a=1}^n \frac{\sqrt{2}}{\mathcal{F}_a^{(0)}} D_a^+ \otimes \sum_{\alpha, \epsilon} \langle \zeta_a, (\widetilde{\operatorname{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}(X_\alpha^{+, \epsilon}) \rangle X_\alpha^{+, \epsilon} \\ &= - \sum_{a=1}^n D_a^+ \otimes \frac{\sqrt{2}}{\mathcal{F}_a^{(0)}} (\widetilde{\operatorname{ad}}_{\mathbf{\Lambda}^{(0)}})^{-1}((\zeta_a)_{\mathfrak{a}^\perp}). \end{aligned} \quad (3.93)$$

Remembering the explicit formula of ζ_a (3.61) and the commutation relations (2.29), the Poisson bracket (3.89) also follows. \square

Lemma 9. *At the distinguished point $u^{(0)}$ (3.5) we have the Poisson bracket*

$$\{m \otimes m\}^{\operatorname{ext}}(u^{(0)}) = \sum_{1 \leq a < b \leq n} \Psi_{a,b}^{(0)} (D_a^+ \otimes D_b^+ - D_b^+ \otimes D_a^+) \quad (3.94)$$

with the coefficients

$$\Psi_{a,b}^{(0)} = \frac{1}{\lambda_a^{(0)} - \lambda_b^{(0)}} + \frac{\lambda_a^{(0)} - \lambda_b^{(0)}}{(\lambda_a^{(0)} - \lambda_b^{(0)})^2 + 4\mu^2}. \quad (3.95)$$

Proof. Using the antisymmetry of the Poisson bracket, we find that

$$\begin{aligned} \{m \otimes m\}^{\operatorname{ext}}(u^{(0)}) &= \sum_{a,b=1}^n \{m_a, m_b\}^{\operatorname{ext}}(u^{(0)})(e_{a,a} + e_{n+a, n+a}) \otimes (e_{b,b} + e_{n+b, n+b}) \\ &= -2 \sum_{1 \leq a < b \leq n} \{m_a, m_b\}^{\operatorname{ext}}(u^{(0)})(D_a^+ \otimes D_b^+ - D_b^+ \otimes D_a^+). \end{aligned} \quad (3.96)$$

To proceed further, let us choose arbitrary $a, b \in \mathbb{N}_n$ satisfying $a < b$. Notice that the Poisson bracket formula (2.75) naturally leads to the expression

$$\begin{aligned} \{m_a, m_b\}^{\text{ext}}(u^{(0)}) &= - \langle (\nabla^G(\text{Im}(m_a)))(u^{(0)}), (\nabla^{\mathfrak{g}}(\text{Im}(m_b)))(u^{(0)}) \rangle \\ &\quad + \langle (\nabla^{\mathfrak{g}}(\text{Im}(m_a)))(u^{(0)}), (\nabla^G(\text{Im}(m_b)))(u^{(0)}) \rangle \\ &\quad - \omega_{\rho^{(0)}}^{\mathcal{O}}((\nabla^{\mathcal{O}}(\text{Im}(m_a)))(u^{(0)}), (\nabla^{\mathcal{O}}(\text{Im}(m_b)))(u^{(0)})). \end{aligned} \quad (3.97)$$

However, by utilizing Proposition 6, each term on the right hand side of the above equation can be cast into a fairly explicit form. Starting with the first term, the application of (3.61) gives rise to the relationship

$$\begin{aligned} \langle (\nabla^G(\text{Im}(m_a)))(u^{(0)}), (\nabla^{\mathfrak{g}}(\text{Im}(m_b)))(u^{(0)}) \rangle &= \frac{1}{\mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}} \langle \zeta_a, (\widetilde{\text{ad}}_{\mathbf{A}^{(0)}})^{-1}((\zeta_b)_{\mathfrak{m}^\perp}) \rangle \\ &= \frac{1}{4\mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}} \left(\frac{\mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}}{\lambda_a^{(0)} - \lambda_b^{(0)}} + \frac{\text{Re}(\mathcal{F}_{n+a}^{(0)}) \text{Re}(\mathcal{F}_{n+b}^{(0)})}{\lambda_a^{(0)} + \lambda_b^{(0)}} - \frac{\text{Im}(\mathcal{F}_{n+a}^{(0)}) \text{Im}(\mathcal{F}_{n+b}^{(0)})}{\lambda_a^{(0)} + \lambda_b^{(0)}} \right). \end{aligned} \quad (3.98)$$

Keeping in mind that $a < b$, a similar argument provides

$$\begin{aligned} \langle (\nabla^{\mathfrak{g}}(\text{Im}(m_a)))(u^{(0)}), (\nabla^G(\text{Im}(m_b)))(u^{(0)}) \rangle &= \frac{1}{4\mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}} \left(-\frac{\mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}}{\lambda_a^{(0)} - \lambda_b^{(0)}} + \frac{\text{Re}(\mathcal{F}_{n+a}^{(0)}) \text{Re}(\mathcal{F}_{n+b}^{(0)})}{\lambda_a^{(0)} + \lambda_b^{(0)}} - \frac{\text{Im}(\mathcal{F}_{n+a}^{(0)}) \text{Im}(\mathcal{F}_{n+b}^{(0)})}{\lambda_a^{(0)} + \lambda_b^{(0)}} \right). \end{aligned} \quad (3.99)$$

Now, let us turn to the third appearing in (3.97). Utilizing the concise formula (2.71) for the symplectic form $\omega^{\mathcal{O}}$ (2.70), the application of the equations (3.64), (3.65) and (2.81) yields that

$$\begin{aligned} \omega_{\rho^{(0)}}^{\mathcal{O}}((\nabla^{\mathcal{O}}(\text{Im}(m_a)))(u^{(0)}), (\nabla^{\mathcal{O}}(\text{Im}(m_b)))(u^{(0)})) &= \omega_{\xi^{(\nu^{(0)})}}^{\mathcal{O}}(\xi_{*\nu^{(0)}}(\delta V_a), \xi_{*\nu^{(0)}}(\delta V_b)) \\ &= 2\mu \text{Im}((\delta V_a)^* \delta V_b) = 2\mu \text{Im} \left(\frac{\mathcal{A}_{a,b}^{(0)}}{8\mu^2 \mathcal{F}_a^{(0)} \mathcal{F}_b^{(0)}} \right) = \frac{1}{2} \frac{\lambda_a^{(0)} - \lambda_b^{(0)}}{(\lambda_a^{(0)} - \lambda_b^{(0)})^2 + 4\mu^2}. \end{aligned} \quad (3.100)$$

Now, by simply putting together the above equations, the Lemma follows at once. \square

At this point we are in a position to provide an explicit formula for the r -matrix (3.35). Remembering (3.36), let us notice that Lemma 7 itself implies that r is in fact linear in \mathcal{A} , having the form

$$r_{12}(\lambda^{(0)}, \theta^{(0)}) = (p_{12}^+)^{(0)} \mathcal{A}_2^{(0)} + \mathcal{A}_2^{(0)} (p_{12}^-)^{(0)} \quad (3.101)$$

with the $\mathfrak{g} \otimes \mathfrak{g}$ -valued functions

$$(p_{12}^\pm)^{(0)} = - \sum_{\alpha, \epsilon} \frac{X_\alpha^{+, \epsilon} \otimes X_\alpha^{-, \epsilon}}{\alpha(\lambda^{(0)})} + \sum_{a=1}^n D_a^+ \otimes S_a^{(0)} \pm \frac{1}{2} \{\varphi \otimes \varphi\}^{\text{ext}}(u^{(0)}). \quad (3.102)$$

Recalling (3.37), the above expressions can be further expanded. Indeed, by simply plugging the formulae displayed in Lemmas 8 and 9 into (3.102), we may obtain explicit expressions

for both p_{12}^\pm and r . However, since r is linear in \mathcal{A} as dictated by (3.101), the linear r -matrix Poisson bracket (3.34) can be cast into a *quadratic* form. Also, since the point $(\lambda^{(0)}, \theta^{(0)})$ (3.1) we fixed at the beginning of Subsection 3.1 was an arbitrary element of \mathcal{P}^R , the zero superscripts become superfluous and can be safely omitted. With the usual conventions for the symmetric and the antisymmetric tensor products,

$$X \vee Y = X \otimes Y + Y \otimes X \quad \text{and} \quad X \wedge Y = X \otimes Y - Y \otimes X, \quad (3.103)$$

we end up with the following result.

Theorem 10. *The Lax matrix \mathcal{A} (2.81) of the rational C_n RSvD model with two independent coupling parameters obeys the quadratic Poisson bracket*

$$\{\mathcal{A} \otimes \mathcal{A}\}^R = \mathbf{a}_{12} \mathcal{A}_1 \mathcal{A}_2 + \mathcal{A}_1 \mathbf{b}_{12} \mathcal{A}_2 - \mathcal{A}_2 \mathbf{c}_{12} \mathcal{A}_1 - \mathcal{A}_1 \mathcal{A}_2 \mathbf{d}_{12} \quad (3.104)$$

with the $\mathfrak{g} \otimes \mathfrak{g}$ -valued dynamical structure coefficients

$$\mathbf{a}_{12} = \sum_{\alpha, \epsilon} \frac{X_\alpha^{-, \epsilon} \wedge X_\alpha^{+, \epsilon}}{\alpha(\lambda)} + \sum_{a=1}^n D_a^+ \wedge (S_a + T_a) + \sum_{1 \leq a < b \leq n} \Psi_{a,b} D_a^+ \wedge D_b^+, \quad (3.105)$$

$$\mathbf{b}_{12} = \sum_{\alpha, \epsilon} \frac{X_\alpha^{-, \epsilon} \vee X_\alpha^{+, \epsilon}}{\alpha(\lambda)} - \sum_{a=1}^n (D_a^+ \vee S_a + D_a^+ \wedge T_a) - \sum_{1 \leq a < b \leq n} \Psi_{a,b} D_a^+ \wedge D_b^+, \quad (3.106)$$

$$\mathbf{c}_{12} = \sum_{\alpha, \epsilon} \frac{X_\alpha^{-, \epsilon} \vee X_\alpha^{+, \epsilon}}{\alpha(\lambda)} - \sum_{a=1}^n (D_a^+ \vee S_a - D_a^+ \wedge T_a) + \sum_{1 \leq a < b \leq n} \Psi_{a,b} D_a^+ \wedge D_b^+, \quad (3.107)$$

$$\mathbf{d}_{12} = \sum_{\alpha, \epsilon} \frac{X_\alpha^{-, \epsilon} \wedge X_\alpha^{+, \epsilon}}{\alpha(\lambda)} + \sum_{a=1}^n D_a^+ \wedge (S_a - T_a) - \sum_{1 \leq a < b \leq n} \Psi_{a,b} D_a^+ \wedge D_b^+, \quad (3.108)$$

where the constituent objects are defined in Lemmas 7, 8 and 9.

Proof. Due to (3.101), the Poisson bracket (3.34) takes the quadratic form (3.104) with

$$\mathbf{a}_{12} = p_{12}^+ - p_{21}^+, \quad \mathbf{b}_{12} = -p_{12}^+ - p_{21}^-, \quad \mathbf{c}_{12} = -p_{21}^+ - p_{12}^-, \quad \mathbf{d}_{12} = p_{12}^- - p_{21}^-. \quad (3.109)$$

Remembering (3.102), (3.37), and the explicit formulae displayed in Lemmas 8 and 9, the Theorem follows. \square

We conclude this section with an important remark. Since the quadratic structure matrices (3.105-3.108) are derived from an r -matrix linear in \mathcal{A} as described in (3.101), from the relationships (3.109) it follows immediately that they satisfy the consistency conditions

$$\mathbf{a}_{21} = -\mathbf{a}_{12}, \quad \mathbf{d}_{21} = -\mathbf{d}_{12}, \quad \mathbf{b}_{21} = \mathbf{c}_{12}, \quad \mathbf{a}_{12} + \mathbf{b}_{12} = \mathbf{c}_{12} + \mathbf{d}_{12}. \quad (3.110)$$

The above observation can be paraphrased as follows. If a Lax matrix \mathcal{A} obeys a tensorial Poisson bracket (3.34), and if the governing r -matrix is itself linear in \mathcal{A} as in (3.101), then the tensorial Poisson bracket can be rewritten as a quadratic bracket (3.104) with quadratic

structure matrices obeying the consistency conditions (3.110) automatically. It is a nice, but essentially trivial algebraic fact that the converse of this statement is also true. Indeed, suppose that a Lax matrix \mathcal{A} obeys a quadratic Poisson bracket (3.104) with coefficients satisfying (3.110). Under these assumptions the quadratic bracket can be cast into a linear form (3.34). More precisely, the governing r -matrix can be written in the form of (3.101) with

$$p_{12}^+ = \frac{\mathbf{a}_{12} + \mathbf{u}_{12}}{2} \quad \text{and} \quad p_{12}^- = \frac{\mathbf{d}_{12} - \mathbf{b}_{12} - \mathbf{c}_{12} - \mathbf{u}_{12}}{2}, \quad (3.111)$$

where \mathbf{u}_{12} is an arbitrary $\mathfrak{g} \vee \mathfrak{g}$ -valued function on the phase space, i.e. it obeys the symmetry condition $\mathbf{u}_{21} = \mathbf{u}_{12}$. This observation plays a crucial role in the developments of the next section.

4 Classical r -matrix structure of the BC_n -type model

Utilizing a symplectic reduction framework, so far we have studied the classical r -matrix structure for the rational C_n RSvD model with two independent coupling parameters μ and ν . However, to handle the BC_n -type model as well, in this section we slightly change our point of view. Switching to a purely algebraic approach, we shall generalize Theorem 10 to cover the most general rational BC_n RSvD model with three independent coupling constants. As an added bonus, at the end of this section we will provide a Lax representation of the dynamics, too.

To describe the Lax matrix of the rational BC_n RSvD system with the additional third real parameter κ , we need the functions

$$\boldsymbol{\alpha}(x) = \frac{\sqrt{x + \sqrt{x^2 + \kappa^2}}}{\sqrt{2x}} \quad \text{and} \quad \boldsymbol{\beta}(x) = i\kappa \frac{1}{\sqrt{2x}} \frac{1}{\sqrt{x + \sqrt{x^2 + \kappa^2}}}, \quad (4.1)$$

where $x \in (0, \infty)$. Also, with each $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathfrak{c}$ we associate the group element

$$h(\lambda) = \begin{bmatrix} \text{diag}(\boldsymbol{\alpha}(\lambda_1), \dots, \boldsymbol{\alpha}(\lambda_n)) & \text{diag}(\boldsymbol{\beta}(\lambda_1), \dots, \boldsymbol{\beta}(\lambda_n)) \\ -\text{diag}(\boldsymbol{\beta}(\lambda_1), \dots, \boldsymbol{\beta}(\lambda_n)) & \text{diag}(\boldsymbol{\alpha}(\lambda_1), \dots, \boldsymbol{\alpha}(\lambda_n)) \end{bmatrix} \in G. \quad (4.2)$$

In [26] we proved that the smooth function $\tilde{\mathcal{A}}: \mathcal{P}^R \rightarrow G$ defined by the formula

$$\tilde{\mathcal{A}}(\lambda, \theta) = h(\lambda)^{-1} \mathcal{A}(\lambda, \theta) h(\lambda)^{-1} \quad ((\lambda, \theta) \in \mathcal{P}^R) \quad (4.3)$$

provides a Lax matrix for the rational BC_n RSvD model (1.4) with the independent coupling parameters μ , ν and κ . Our first goal in this section to construct a quadratic algebra relation for the Lax matrix $\tilde{\mathcal{A}}$ with structure coefficients satisfying the consistency conditions analogous to (3.110).

Recalling (2.15) and (2.81), we start with the observation

$$\frac{\partial \mathcal{A}}{\partial \theta_c} = \sqrt{2}(D_c^- \mathcal{A} + \mathcal{A} D_c^-) \quad (c \in \mathbb{N}_n). \quad (4.4)$$

Therefore, upon introducing the $\mathfrak{g} \otimes \mathfrak{g}$ -valued function

$$\Gamma_{12} = \frac{1}{\sqrt{2}} \sum_{c=1}^n D_c^- \otimes \frac{\partial(h^{-1})}{\partial \lambda_c}, \quad (4.5)$$

we can write the tensorial Poisson bracket

$$\{\mathcal{A}_1, h_2^{-1}\}^R = \Gamma_{12} \mathcal{A}_1 + \mathcal{A}_1 \Gamma_{12}. \quad (4.6)$$

Now, by simply applying the Leibniz rule, from (3.104) we get that

$$\{\tilde{\mathcal{A}} \otimes \tilde{\mathcal{A}}\}^R = \tilde{\mathbf{a}}_{12} \tilde{\mathcal{A}}_1 \tilde{\mathcal{A}}_2 + \tilde{\mathcal{A}}_1 \tilde{\mathbf{b}}_{12} \tilde{\mathcal{A}}_2 - \tilde{\mathcal{A}}_2 \tilde{\mathbf{c}}_{12} \tilde{\mathcal{A}}_1 - \tilde{\mathcal{A}}_1 \tilde{\mathcal{A}}_2 \tilde{\mathbf{d}}_{12} \quad (4.7)$$

with the dynamical coefficients

$$\tilde{\mathbf{a}}_{12} = h_1^{-1} h_2^{-1} \mathbf{a}_{12} h_1 h_2 + h_1^{-1} \Gamma_{12} h_1 h_2 - h_2^{-1} \Gamma_{21} h_1 h_2, \quad (4.8)$$

$$\tilde{\mathbf{b}}_{12} = h_1 h_2^{-1} \mathbf{b}_{12} h_1^{-1} h_2 + h_1 \Gamma_{12} h_1^{-1} h_2 - h_1 h_2^{-1} \Gamma_{21} h_2, \quad (4.9)$$

$$\tilde{\mathbf{c}}_{12} = h_1^{-1} h_2 \mathbf{c}_{12} h_1 h_2^{-1} - h_1^{-1} h_2 \Gamma_{12} h_1 + h_2 \Gamma_{21} h_1 h_2^{-1}, \quad (4.10)$$

$$\tilde{\mathbf{d}}_{12} = h_1 h_2 \mathbf{d}_{12} h_1^{-1} h_2^{-1} - h_1 h_2 \Gamma_{12} h_1^{-1} + h_1 h_2 \Gamma_{21} h_2^{-1}. \quad (4.11)$$

Since the decorations coming from h are ‘equally distributed’ among these new functions, we expect that likewise they satisfy the consistency conditions analogous to (3.110). Somewhat surprisingly, this naive idea is fully confirmed by the following result.

Theorem 11. *The functions (4.8-4.11) appearing in the tensorial Poisson bracket (4.7) obey the consistency conditions*

$$\tilde{\mathbf{a}}_{21} = -\tilde{\mathbf{a}}_{12}, \quad \tilde{\mathbf{d}}_{21} = -\tilde{\mathbf{d}}_{12}, \quad \tilde{\mathbf{b}}_{21} = \tilde{\mathbf{c}}_{12}, \quad \tilde{\mathbf{a}}_{12} + \tilde{\mathbf{b}}_{12} = \tilde{\mathbf{c}}_{12} + \tilde{\mathbf{d}}_{12}. \quad (4.12)$$

In other words, the Lax matrix $\tilde{\mathcal{A}}$ (4.3) of the rational BC_n RSvD system satisfies a quadratic Poisson bracket (4.7) characterized by the consistent dynamical structure coefficients (4.8-4.11).

Proof. A moment of reflection reveals that $\tilde{\mathbf{a}}_{21} = -\tilde{\mathbf{a}}_{12}$, $\tilde{\mathbf{d}}_{21} = -\tilde{\mathbf{d}}_{12}$, and $\tilde{\mathbf{b}}_{21} = \tilde{\mathbf{c}}_{12}$, whence it is enough to prove that $\tilde{\mathbf{a}}_{12} + \tilde{\mathbf{b}}_{12} = \tilde{\mathbf{c}}_{12} + \tilde{\mathbf{d}}_{12}$. Since the verification of this last equation is basically an involved algebraic computation, in the following we wish to highlight only the key steps. First, we introduce the functions

$$\mathcal{P}(x) = \sqrt{1 + \frac{\kappa^2}{x^2}} \quad \text{and} \quad \mathcal{Q}(x) = \frac{i\kappa}{x} \quad (x \in (0, \infty)). \quad (4.13)$$

Remembering (4.1), we see that

$$\mathcal{P}(x) = \boldsymbol{\alpha}(x)^2 - \boldsymbol{\beta}(x)^2 \quad \text{and} \quad \mathcal{Q}(x) = 2\boldsymbol{\alpha}(x)\boldsymbol{\beta}(x). \quad (4.14)$$

To make the presentation a slightly simpler, we also introduce the G -valued function

$$\mathcal{H} = h^2. \quad (4.15)$$

Now, recalling (4.1), (4.2) and (4.14), with the notations

$$\mathcal{P}_a = \mathcal{P}(\lambda_a) \quad \text{and} \quad \mathcal{Q}_a = \mathcal{Q}(\lambda_a) \quad (a \in \mathbb{N}_n) \quad (4.16)$$

we can clearly write

$$\mathcal{H} = \begin{bmatrix} \text{diag}(\mathcal{P}_1, \dots, \mathcal{P}_n) & \text{diag}(\mathcal{Q}_1, \dots, \mathcal{Q}_n) \\ -\text{diag}(\mathcal{Q}_1, \dots, \mathcal{Q}_n) & \text{diag}(\mathcal{P}_1, \dots, \mathcal{P}_n) \end{bmatrix}. \quad (4.17)$$

To proceed further, we also define the $\mathfrak{g} \otimes \mathfrak{g}$ -valued function

$$\Omega_{12} = -(h_2^{-1}\Gamma_{12} + \Gamma_{12}h_2^{-1})\mathcal{H}_2. \quad (4.18)$$

Remembering the form of Γ_{12} (4.5), Leibniz rule yields

$$\Omega_{12} = \frac{1}{\sqrt{2}} \sum_{c=1}^n D_c^- \otimes \left(\mathcal{H}^{-1} \frac{\partial \mathcal{H}}{\partial \lambda_c} \right), \quad (4.19)$$

where for the derivatives we can easily find that

$$\mathcal{H}^{-1} \frac{\partial \mathcal{H}}{\partial \lambda_c} = \frac{\sqrt{2}\kappa}{\lambda_c \sqrt{\lambda_c^2 + \kappa^2}} X_{2\varepsilon_c}^{-,i} \quad (c \in \mathbb{N}_n). \quad (4.20)$$

Bearing in mind the above objects, from (4.8-4.11) one can derive that

$$\begin{aligned} & h_1^{-1}h_2^{-1}(\tilde{\mathbf{a}}_{12} + \tilde{\mathbf{b}}_{12} - \tilde{\mathbf{c}}_{12} - \tilde{\mathbf{d}}_{12})h_1h_2 \\ &= \mathcal{H}_1^{-1}\mathcal{H}_2^{-1}\mathbf{a}_{12}\mathcal{H}_1\mathcal{H}_2 + \mathcal{H}_2^{-1}\mathbf{b}_{12}\mathcal{H}_2 - \mathcal{H}_1^{-1}\mathbf{c}_{12}\mathcal{H}_1 - \mathbf{d}_{12} \\ & \quad - \mathcal{H}_1^{-1}\Omega_{12}\mathcal{H}_1 + \mathcal{H}_2^{-1}\Omega_{21}\mathcal{H}_2 - \Omega_{12} + \Omega_{21}. \end{aligned} \quad (4.21)$$

To handle the right hand side of the above equation, we need the commutation relations listed below. First, for each $c \in \mathbb{N}_n$ we have

$$\mathcal{H}^{-1}D_c^+\mathcal{H} = D_c^+, \quad (4.22)$$

$$\mathcal{H}^{-1}D_c^-\mathcal{H} = (\mathcal{P}_c^2 - \mathcal{Q}_c^2)D_c^- + 2i\mathcal{P}_c\mathcal{Q}_cX_{2\varepsilon_c}^{+,i}, \quad (4.23)$$

$$\mathcal{H}^{-1}X_{2\varepsilon_c}^{+,i}\mathcal{H} = (\mathcal{P}_c^2 - \mathcal{Q}_c^2)X_{2\varepsilon_c}^{+,i} + 2i\mathcal{P}_c\mathcal{Q}_cD_c^-, \quad (4.24)$$

$$\mathcal{H}^{-1}X_{2\varepsilon_c}^{-,i}\mathcal{H} = X_{2\varepsilon_c}^{-,i}. \quad (4.25)$$

Also, if $a, b \in \mathbb{N}_n$ and $a < b$, then we can write

$$\mathcal{H}^{-1}X_{\varepsilon_a - \varepsilon_b}^{+,r}\mathcal{H} = (\mathcal{P}_a\mathcal{P}_b + \mathcal{Q}_a\mathcal{Q}_b)X_{\varepsilon_a - \varepsilon_b}^{+,r} + i(\mathcal{P}_a\mathcal{Q}_b - \mathcal{P}_b\mathcal{Q}_a)X_{\varepsilon_a + \varepsilon_b}^{-,i}, \quad (4.26)$$

$$\mathcal{H}^{-1}X_{\varepsilon_a - \varepsilon_b}^{-,r}\mathcal{H} = (\mathcal{P}_a\mathcal{P}_b - \mathcal{Q}_a\mathcal{Q}_b)X_{\varepsilon_a - \varepsilon_b}^{-,r} + i(\mathcal{P}_a\mathcal{Q}_b + \mathcal{P}_b\mathcal{Q}_a)X_{\varepsilon_a + \varepsilon_b}^{+,i}, \quad (4.27)$$

$$\mathcal{H}^{-1}X_{\varepsilon_a + \varepsilon_b}^{+,r}\mathcal{H} = (\mathcal{P}_a\mathcal{P}_b - \mathcal{Q}_a\mathcal{Q}_b)X_{\varepsilon_a + \varepsilon_b}^{+,r} - i(\mathcal{P}_a\mathcal{Q}_b + \mathcal{P}_b\mathcal{Q}_a)X_{\varepsilon_a - \varepsilon_b}^{-,i}, \quad (4.28)$$

$$\mathcal{H}^{-1}X_{\varepsilon_a + \varepsilon_b}^{-,r}\mathcal{H} = (\mathcal{P}_a\mathcal{P}_b + \mathcal{Q}_a\mathcal{Q}_b)X_{\varepsilon_a + \varepsilon_b}^{-,r} - i(\mathcal{P}_a\mathcal{Q}_b - \mathcal{P}_b\mathcal{Q}_a)X_{\varepsilon_a - \varepsilon_b}^{+,i}, \quad (4.29)$$

together with the relations

$$\mathcal{H}^{-1} X_{\varepsilon_a - \varepsilon_b}^{+,i} \mathcal{H} = (\mathcal{P}_a \mathcal{P}_b + \mathcal{Q}_a \mathcal{Q}_b) X_{\varepsilon_a - \varepsilon_b}^{+,i} - i(\mathcal{P}_a \mathcal{Q}_b - \mathcal{P}_b \mathcal{Q}_a) X_{\varepsilon_a + \varepsilon_b}^{-,r}, \quad (4.30)$$

$$\mathcal{H}^{-1} X_{\varepsilon_a - \varepsilon_b}^{-,i} \mathcal{H} = (\mathcal{P}_a \mathcal{P}_b - \mathcal{Q}_a \mathcal{Q}_b) X_{\varepsilon_a - \varepsilon_b}^{-,i} - i(\mathcal{P}_a \mathcal{Q}_b + \mathcal{P}_b \mathcal{Q}_a) X_{\varepsilon_a + \varepsilon_b}^{+,r}, \quad (4.31)$$

$$\mathcal{H}^{-1} X_{\varepsilon_a + \varepsilon_b}^{+,i} \mathcal{H} = (\mathcal{P}_a \mathcal{P}_b - \mathcal{Q}_a \mathcal{Q}_b) X_{\varepsilon_a + \varepsilon_b}^{+,i} + i(\mathcal{P}_a \mathcal{Q}_b + \mathcal{P}_b \mathcal{Q}_a) X_{\varepsilon_a - \varepsilon_b}^{-,r}, \quad (4.32)$$

$$\mathcal{H}^{-1} X_{\varepsilon_a + \varepsilon_b}^{-,i} \mathcal{H} = (\mathcal{P}_a \mathcal{P}_b + \mathcal{Q}_a \mathcal{Q}_b) X_{\varepsilon_a + \varepsilon_b}^{-,i} + i(\mathcal{P}_a \mathcal{Q}_b - \mathcal{P}_b \mathcal{Q}_a) X_{\varepsilon_a - \varepsilon_b}^{+,r}. \quad (4.33)$$

Now, let us examine the first four terms appearing on the right hand side of (4.21). Recalling (3.105-3.108), the application of (4.22) itself yields the formula

$$\begin{aligned} & \mathcal{H}_1^{-1} \mathcal{H}_2^{-1} \mathbf{a}_{12} \mathcal{H}_1 \mathcal{H}_2 + \mathcal{H}_2^{-1} \mathbf{b}_{12} \mathcal{H}_2 - \mathcal{H}_1^{-1} \mathbf{c}_{12} \mathcal{H}_1 - \mathbf{d}_{12} \\ &= \sum_{\alpha, \epsilon} \frac{(\mathcal{H}^{-1} X_{\alpha}^{-, \epsilon} \mathcal{H} + X_{\alpha}^{-, \epsilon}) \wedge (\mathcal{H}^{-1} X_{\alpha}^{+, \epsilon} \mathcal{H} - X_{\alpha}^{+, \epsilon})}{\alpha(\lambda)}. \end{aligned} \quad (4.34)$$

However, in order to further simplify this expression, we still have to exploit some functional equations obeyed by \mathcal{P} and \mathcal{Q} . By inspecting the definitions (4.13), we see immediately that

$$\mathcal{P}(x)^2 - \mathcal{Q}(x)^2 = 1 + \frac{2\kappa^2}{x^2} \quad (x \in (0, \infty)). \quad (4.35)$$

A slightly longer calculation also reveals that

$$\frac{\mathcal{P}(x)^2 \mathcal{P}(y)^2 - (\mathcal{Q}(x) \mathcal{Q}(y) - 1)^2}{x - y} + \frac{\mathcal{P}(x)^2 \mathcal{Q}(y)^2 - \mathcal{P}(y)^2 \mathcal{Q}(x)^2}{x + y} = 0, \quad (4.36)$$

$$\frac{\mathcal{P}(x)^2 \mathcal{Q}(y)^2 - \mathcal{P}(y)^2 \mathcal{Q}(x)^2}{x - y} + \frac{\mathcal{P}(x)^2 \mathcal{P}(y)^2 - (\mathcal{Q}(x) \mathcal{Q}(y) + 1)^2}{x + y} = 0, \quad (4.37)$$

where $x, y \in (0, \infty)$ and $x \neq y$. Having equipped with the relations (4.35-4.37), let us note that the application of the commutation relations (4.22-4.33) does give rise to an even greater simplification in (4.34). Indeed, we find that

$$\begin{aligned} & \sum_{\alpha, \epsilon} \frac{(\mathcal{H}^{-1} X_{\alpha}^{-, \epsilon} \mathcal{H} + X_{\alpha}^{-, \epsilon}) \wedge (\mathcal{H}^{-1} X_{\alpha}^{+, \epsilon} \mathcal{H} - X_{\alpha}^{+, \epsilon})}{\alpha(\lambda)} \\ &= \sum_{c=1}^n \left(\frac{2\kappa^2}{\lambda_c^3} X_{2\varepsilon_c}^{-,i} \wedge X_{2\varepsilon_c}^{+,i} + \frac{2\kappa}{\lambda_c^2} \sqrt{1 + \frac{\kappa^2}{\lambda_c^2}} D_c^- \wedge X_{2\varepsilon_c}^{-,i} \right). \end{aligned} \quad (4.38)$$

Now, let us turn to the last four terms appearing on the right hand side of (4.21). Recalling (4.19), (4.20), and (4.35), we can write that

$$\mathcal{H}_1^{-1} \Omega_{12} \mathcal{H}_1 + \Omega_{12} = \sum_{c=1}^n \left(-\frac{2\kappa^2}{\lambda_c^3} X_{2\varepsilon_c}^{+,i} \otimes X_{2\varepsilon_c}^{-,i} + \frac{2\kappa}{\lambda_c^2} \sqrt{1 + \frac{\kappa^2}{\lambda_c^2}} D_c^- \otimes X_{2\varepsilon_c}^{-,i} \right). \quad (4.39)$$

Now, by plugging (4.34), (4.38) and (4.39) into (4.21), we obtain at once that

$$h_1^{-1} h_2^{-1} (\tilde{\mathbf{a}}_{12} + \tilde{\mathbf{b}}_{12} - \tilde{\mathbf{c}}_{12} - \tilde{\mathbf{d}}_{12}) h_1 h_2 = 0, \quad (4.40)$$

whence the proof is complete. \square

Having completed the proof, now we offer a few remarks on the result. First, since the Lax matrix $\tilde{\mathcal{A}}$ obeys the quadratic bracket (4.7) with the dynamical objects (4.8-4.11) satisfying the consistency conditions (4.12), the quadratic bracket (4.7) can be rewritten as

$$\{\tilde{\mathcal{A}}_1, \tilde{\mathcal{A}}_2\}^R = [\tilde{r}_{12}, \tilde{\mathcal{A}}_1] - [\tilde{r}_{21}, \tilde{\mathcal{A}}_2]. \quad (4.41)$$

Indeed, recalling our discussion at the end of the previous section, an appropriate r -matrix is provided by the formula

$$\tilde{r}_{12} = \tilde{p}_{12}^+ \tilde{\mathcal{A}}_2 + \tilde{\mathcal{A}}_2 \tilde{p}_{12}^-, \quad (4.42)$$

where

$$\tilde{p}_{12}^+ = \frac{\tilde{\mathbf{a}}_{12} + \tilde{\mathbf{u}}_{12}}{2} \quad \text{and} \quad \tilde{p}_{12}^- = \frac{\tilde{\mathbf{d}}_{12} - \tilde{\mathbf{b}}_{12} - \tilde{\mathbf{c}}_{12} - \tilde{\mathbf{u}}_{12}}{2}, \quad (4.43)$$

with an arbitrary $\mathfrak{g} \vee \mathfrak{g}$ -valued dynamical object $\tilde{\mathbf{u}}_{12}$.

Second, one may raise the objection that the formulae (4.8-4.11) for the quadratic structure matrices in the BC_n case are ‘less explicit’ than the analogous objects (3.105-3.108) in the C_n case. The trouble is mainly caused by the derivatives of h^{-1} appearing in the definition of Γ_{12} (4.5). Though these derivatives can be worked out rather easily, we propose an alternative approach to cure the problem. Namely, let us apply the gauge transformation

$$\hat{\mathcal{A}}(\lambda, \theta) = h(\lambda) \tilde{\mathcal{A}}(\lambda, \theta) h(\lambda)^{-1} \quad ((\lambda, \theta) \in \mathcal{P}^R). \quad (4.44)$$

on the Lax matrix $\tilde{\mathcal{A}}$ (4.3). By applying the corresponding transformation on \tilde{r} (4.42), it turns out that the transformed r -matrix takes the form

$$\hat{r}_{12} = \hat{p}_{12}^+ \hat{\mathcal{A}}_2 + \hat{\mathcal{A}}_2 \hat{p}_{12}^-, \quad (4.45)$$

where

$$\hat{p}_{12}^+ = h_1 h_2 \tilde{p}_{12}^+ h_1^{-1} h_2^{-1} + h_1 \Gamma_{21} \quad \text{and} \quad \hat{p}_{12}^- = h_1 h_2 \tilde{p}_{12}^- h_1^{-1} h_2^{-1} + h_1 \mathcal{H}_2 \Gamma_{21} \mathcal{H}_2^{-1}. \quad (4.46)$$

Since \hat{r}_{12} is linear in $\hat{\mathcal{A}}$, the tensorial Poisson bracket for $\hat{\mathcal{A}}$ can be cast into a quadratic form with structure matrices obeying the consistency conditions analogous to (4.12). To save time on the algebraic details, we present only the resulting formulae.

Theorem 12. *For the transformed Lax matrix $\hat{\mathcal{A}} = \mathcal{A} \mathcal{H}^{-1}$ of the rational BC_n RSvD model we have*

$$\{\hat{\mathcal{A}}_1, \hat{\mathcal{A}}_2\}^R = \hat{\mathbf{a}}_{12} \hat{\mathcal{A}}_1 \hat{\mathcal{A}}_2 + \hat{\mathcal{A}}_1 \hat{\mathbf{b}}_{12} \hat{\mathcal{A}}_2 - \hat{\mathcal{A}}_2 \hat{\mathbf{c}}_{12} \hat{\mathcal{A}}_1 - \hat{\mathcal{A}}_1 \hat{\mathcal{A}}_2 \hat{\mathbf{d}}_{12}, \quad (4.47)$$

where the dynamical objects

$$\hat{\mathbf{a}}_{12} = \mathbf{a}_{12}, \quad (4.48)$$

$$\hat{\mathbf{b}}_{12} = \mathcal{H}_1 \left(\mathbf{b}_{12} + \frac{1}{\sqrt{2}} \sum_{c=1}^n \left(\mathcal{H}^{-1} \frac{\partial \mathcal{H}}{\partial \lambda_c} \right) \otimes D_c^- \right) \mathcal{H}_1^{-1}, \quad (4.49)$$

$$\hat{\mathbf{c}}_{12} = \mathcal{H}_2 \left(\mathbf{c}_{12} + \frac{1}{\sqrt{2}} \sum_{c=1}^n D_c^- \otimes \left(\mathcal{H}^{-1} \frac{\partial \mathcal{H}}{\partial \lambda_c} \right) \right) \mathcal{H}_2^{-1}, \quad (4.50)$$

$$\hat{\mathbf{d}}_{12} = \mathcal{H}_1 \mathcal{H}_2 \left(\mathbf{d}_{12} + \frac{1}{\sqrt{2}} \sum_{c=1}^n D_c^- \wedge \left(\mathcal{H}^{-1} \frac{\partial \mathcal{H}}{\partial \lambda_c} \right) \right) \mathcal{H}_1^{-1} \mathcal{H}_2^{-1} \quad (4.51)$$

are built up from the explicitly given functions (3.105-3.108), (4.17) and (4.20). Furthermore, by construction, they satisfy the consistency conditions

$$\hat{\mathbf{a}}_{21} = -\hat{\mathbf{a}}_{12}, \quad \hat{\mathbf{d}}_{21} = -\hat{\mathbf{d}}_{12}, \quad \hat{\mathbf{b}}_{21} = \hat{\mathbf{c}}_{12}, \quad \hat{\mathbf{a}}_{12} + \hat{\mathbf{b}}_{12} = \hat{\mathbf{c}}_{12} + \hat{\mathbf{d}}_{12}. \quad (4.52)$$

As an immediate consequence of the r -matrix formalism, now we can easily construct a Lax pair for the rational BC_n RSvD system. For this reason, let us recall the partial trace operation on the second factor of $\mathfrak{gl}(N, \mathbb{C}) \otimes \mathfrak{gl}(N, \mathbb{C})$, which is uniquely determined by the condition

$$\mathrm{tr}_2(X \otimes Y) = \mathrm{tr}(Y)X \quad (X, Y \in \mathfrak{gl}(N, \mathbb{C})). \quad (4.53)$$

As we proved in [26], for the Hamiltonian of the rational BC_n RSvD model we can write

$$H^R = \frac{1}{2} \mathrm{tr}(\tilde{\mathcal{A}}) = \frac{1}{2} \mathrm{tr}(\hat{\mathcal{A}}). \quad (4.54)$$

Therefore, by expanding the Lax matrix $\hat{\mathcal{A}}$ (4.44) in an arbitrary basis of the complex linear space $\mathfrak{gl}(N, \mathbb{C})$, say in the basis $\{v_I\}$ (2.27), we can write

$$H^R = \frac{1}{2} \mathrm{tr} \left(\sum_J \hat{\mathcal{A}}^J v_J \right) = \frac{1}{2} \sum_J \hat{\mathcal{A}}^J \mathrm{tr}(v_J). \quad (4.55)$$

Making use of this expansion, for the action of the Hamiltonian vector field $\mathbf{X}_{H^R} \in \mathfrak{X}(\mathcal{P}^R)$ on the Lax operator $\hat{\mathcal{A}}$ we obtain

$$\begin{aligned} \mathbf{X}_{H^R}[\hat{\mathcal{A}}] &= \sum_I \mathbf{X}_{H^R}[\hat{\mathcal{A}}^I] v_I = \sum_I \{\hat{\mathcal{A}}^I, H^R\}^R v_I \\ &= \frac{1}{2} \sum_{I,J} \{\hat{\mathcal{A}}^I, \hat{\mathcal{A}}^J\}^R \mathrm{tr}(v_J) v_I = \frac{1}{2} \mathrm{tr}_2 \left(\sum_{I,J} \{\hat{\mathcal{A}}^I, \hat{\mathcal{A}}^J\}^R v_I \otimes v_J \right) \\ &= \frac{1}{2} \mathrm{tr}_2 \left([\hat{r}_{12}, \hat{\mathcal{A}}_1] - [\hat{r}_{21}, \hat{\mathcal{A}}_2] \right) = \frac{1}{2} [\mathrm{tr}_2(\hat{r}_{12}), \hat{\mathcal{A}}]. \end{aligned} \quad (4.56)$$

Since for \hat{p}_{12}^\pm (4.46) we can write

$$\hat{p}_{12}^+ = \frac{\hat{\mathbf{a}}_{12} + \hat{\mathbf{u}}_{12}}{2} \quad \text{and} \quad \hat{p}_{12}^- = \frac{\hat{\mathbf{d}}_{12} - \hat{\mathbf{b}}_{12} - \hat{\mathbf{c}}_{12} - \hat{\mathbf{u}}_{12}}{2}, \quad (4.57)$$

where $\hat{\mathbf{u}}_{12}$ is a $\mathfrak{g} \vee \mathfrak{g}$ -valued function, the application of (4.45) and (4.52) yields that

$$\mathrm{tr}_2(\hat{r}_{12}) = \mathrm{tr}_2((\hat{\mathbf{a}}_{12} - \hat{\mathbf{c}}_{12})\hat{\mathcal{A}}_2). \quad (4.58)$$

Upon introducing the alternative Lax matrix

$$\check{\mathcal{A}} = \mathcal{H}^{-1} \hat{\mathcal{A}} \mathcal{H} = \mathcal{H}^{-1} \mathcal{A}, \quad (4.59)$$

the combination of the formulae (4.48), (4.50), (4.56) and (4.58), together with the explicit expressions (3.105) and (3.107), leads to the following result.

Theorem 13. *With the aid of the \mathfrak{g} -valued function*

$$\begin{aligned} \hat{\mathcal{B}} = & \frac{1}{2} \sum_{\alpha, \epsilon} \frac{\text{tr}(X_{\alpha}^{+, \epsilon}(\hat{\mathcal{A}} - \check{\mathcal{A}}))X_{\alpha}^{-, \epsilon} - \text{tr}(X_{\alpha}^{-, \epsilon}(\hat{\mathcal{A}} + \check{\mathcal{A}}))X_{\alpha}^{+, \epsilon}}{\alpha(\lambda)} \\ & + \frac{1}{2} \sum_{c=1}^n \text{tr}(S_c(\hat{\mathcal{A}} + \check{\mathcal{A}}) + T_c(\hat{\mathcal{A}} - \check{\mathcal{A}}))D_c^+ - \frac{\kappa}{2} \sum_{c=1}^n \frac{\text{tr}(X_{2\epsilon_c}^{-, i}\check{\mathcal{A}})}{\lambda_c \sqrt{\lambda_c^2 + \kappa^2}} D_c^- \end{aligned} \quad (4.60)$$

the derivative of the Lax matrix $\hat{\mathcal{A}}$ (4.44) along the Hamiltonian vector field \mathbf{X}_{HR} takes the Lax form $\mathbf{X}_{HR}[\hat{\mathcal{A}}] = [\hat{\mathcal{B}}, \hat{\mathcal{A}}]$. In other words, $\hat{\mathcal{B}}$ provides a Lax pair for $\hat{\mathcal{A}}$.

5 Discussion

One of the most important objects in the algebraic formulation of the theory of classical integrable systems is undoubtedly the r -matrix structure encoding the tensorial Poisson bracket of the Lax matrix. In the context of the A_n -type CMS and RSvD models the underlying dynamical r -matrix structure is under complete control, even in the elliptic case (see e.g. [36, 37, 38, 39]). In sharp contrast, for the models associated with the non- A_n -type root systems the theory is far less developed. By generalizing the ideas of Avan, Babelon and Talon [31], in our earlier paper [32] we constructed a dynamical r -matrix structure for the most general hyperbolic BC_n Sutherland system with three independent coupling constants. However, for the elliptic case only partial results are available [40]. For the non- A_n -type RSvD systems the situation is even more delicate. Prior to our present paper, the r -matrix structure of the BC_n RSvD systems was studied only in [41], based on the special one-parameter family of Lax matrices coming from \mathbb{Z}_2 -folding of the A_{2n} root system. Nevertheless, in the present paper we succeeded in constructing a quadratic r -matrix structure for the rational BC_n RSvD systems with the maximal number of three coupling parameters, as formulated in Theorems 11 and 12. It is also clear that by applying a standard analytic continuation argument on our formulae, one can easily derive a dynamical r -matrix structure for the rational RSvD system appearing in [42].

Regarding the hyperbolic, trigonometric and elliptic variants of the non- A_n -type RSvD systems we also face many interesting questions. Indeed, except from some very special cases [24, 43, 44], even the construction of Lax matrices for these models is a wide open problem. However, let us note that in the last couple of years many results for the A_n -type models have been reinterpreted in a more geometric context using advanced techniques from the theory of reductions (see e.g. [45, 46]). Relatedly, it would be of considerable interest to see whether the underlying classical r -matrix structures can be explored from these geometric pictures along the line of our present paper. We also expect that the various reduction approaches eventually may lead to a progress in the rigorous geometric theory of the non- A_n -type trigonometric, hyperbolic and elliptic RSvD systems as well. As a starting point, it is worth mentioning the recent paper [47], in which a Hamiltonian reduction approach based on the Heisenberg double of $SU(n, n)$ gives rise to a new integrable particle system, that in the cotangent bundle limit gives back the familiar hyperbolic BC_n Sutherland model with three independent coupling parameters.

Turning back to our quadratic r -matrix algebra (4.47), let us observe that the structure matrices $\hat{\mathbf{a}}_{12}$, $\hat{\mathbf{b}}_{12}$, $\hat{\mathbf{c}}_{12}$ and $\hat{\mathbf{d}}_{12}$ are fully dynamical, i.e. they depend on all variables of the phase space \mathcal{P}^R (1.2) in an essential way. It is in contrast with the CMS models, where the naturally appearing dynamical r -matrices usually depend only on the configuration space variables. Moreover, in many variants of the CMS models the r -matrices can be related to the dynamical Yang–Baxter equation, as first realized in [48]. However, in the A_n case Suris [49] observed that in some special choice of gauge the CMS and the RSvD models can be characterized by the same dynamical r -matrices. Working in this gauge, Nagy, Avan and Rollet proved that the quadratic structure matrices of the hyperbolic A_n RSvD system do obey certain dynamical quadratic Yang–Baxter equations (see Proposition 1 in [50], and relatedly also [51]). As a natural next step, we find it an important question whether such claims can be made about the quadratic algebra relation (4.47) in an appropriate gauge. Also, it would be of considerable interest to investigate whether the non- A_n -type RSvD models can be characterized by numerical, i.e. non-dynamical r -matrices. In the A_n case the answer is in the affirmative (see [52]), but in the BC_n case the analogous tasks seem to be quite challenging even for the rational models. Nevertheless, we wish to come back to these problems in later publications.

Acknowledgments. Our work was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. This work was also supported by a Lendület Grant; we wish to thank Z. Bajnok for hospitality in the MTA Lendület Holographic QFT Group.

References

- [1] S.N.M. Ruijsenaars and H. Schneider, A new class of integrable models and its relation to solitons, *Ann. Phys. (N.Y.)* **170** (1986) 370-405.
- [2] J.F. van Diejen, Deformations of Calogero–Moser systems and finite Toda chains, *Theor. Math. Phys.* **99** (1994) 549-554.
- [3] S.N.M. Ruijsenaars, Action-angle maps and scattering theory for some finite dimensional integrable systems I. The pure soliton case, *Commun. Math. Phys.* **115** (1988) 127-165.
- [4] O. Babelon and D. Bernard, The sine-Gordon solitons as a N -body problem, *Phys. Lett. B* **317** (1993) 363-368.
- [5] S.N.M. Ruijsenaars, Action-angle maps and scattering theory for some finite dimensional integrable systems II. Solitons, antisolitons and their bound states, *Publ. RIMS* **30** (1994) 865-1008.
- [6] S.N.M. Ruijsenaars, Action-angle maps and scattering theory for some finite dimensional integrable systems III. Sutherland type systems and their duals, *Publ. RIMS* **31** (1995) 247-353.
- [7] A. Gorsky and N. Nekrasov, Relativistic Calogero–Moser model as gauged WZW theory, *Nucl. Phys. B* **436** (1995) 582-608.

- [8] N. Nekrasov, Infinite-dimensional algebras, many-body systems and gauge theories, in *Moscow Seminar in Mathematical Physics*, eds. A.Yu. Morozov and M.A. Olshanetsky, AMS Transl. Ser. 2 (American Mathematical Society, 1999) pp. 263-299.
- [9] H.W. Braden, A. Marshakov, A. Mironov and A. Morozov, The Ruijsenaars–Schneider Model in the Context of Seiberg–Witten Theory, *Nucl. Phys. B* **558** (1999) 371-390.
- [10] N.A. Nekrasov and S.L. Shatavili, Quantization of integrable systems and four dimensional gauge theories, [arXiv:0908.4052](https://arxiv.org/abs/0908.4052) [[hep-th](https://arxiv.org/abs/0908.4052)].
- [11] A. Gadde, S. Gukov and P. Putrov, Walls, Lines, and Spectral Dualities in 3d Gauge Theories, *JHEP* **05** (2014) 047.
- [12] F. Calogero, Solution of the one-dimensional N -body problem with quadratic and/or inversely quadratic pair potentials, *J. Math. Phys.* **12** (1971) 419-436.
- [13] B. Sutherland, Exact results for a quantum many body problem in one dimension, *Phys. Rev. A* **4** (1971) 2019-2021.
- [14] J. Moser, Three integrable Hamiltonian systems connected with isospectral deformations, *Adv. Math.* **16** (1975) 197-220.
- [15] E. Bogomolny, O. Giraud and C. Schmit, Random Matrix Ensembles Associated with Lax Matrices, *Phys. Rev. Lett.* **103** (2009) 054103.
- [16] E. Bogomolny, O. Giraud and C. Schmit, Integrable random matrix ensembles, *Nonlinearity* **24** (2011) 3179-3213.
- [17] Y.V. Fyodorov and O. Giraud, High values of disorder-generated multifractals and logarithmically correlated processes, *Chaos, Solitons & Fractals* **74** (2015) 15-26.
- [18] E. Mukhin, V. Tarasov and A. Varchenko, Gaudin Hamiltonians generate the Bethe algebra of a tensor power of the vector representation of \mathfrak{gl}_N , *St. Petersburg Math. J.* **22** (2011) 463-472.
- [19] A. Alexandrov, S. Leurent, Z. Tsuboi and A. Zabrodin, The master T -operator for the Gaudin model and the KP hierarchy, *Nucl. Phys. B* **883** (2014) 173-223.
- [20] A. Gorsky, A. Zabrodin and A. Zotov, Spectrum of Quantum Transfer Matrices via Classical Many-Body Systems, *JHEP* **01** (2014) 070.
- [21] Z. Tsuboi, A. Zabrodin and A. Zotov, Supersymmetric quantum spin chains and classical integrable systems, *JHEP* **05** (2015) 086.
- [22] G. Aminov, S. Arthamonov, A. Smirnov and A. Zotov, Rational Top and its Classical R -matrix, *J. Phys. A: Math. Theor.* **47** (2014) 305207.

- [23] A. Levin, M. Olshanetsky and A. Zotov, Relativistic Classical Integrable Tops and Quantum R -matrices, *JHEP* **07** (2014) 012.
- [24] K. Chen, B.Y. Hou and W.L. Yang, Integrability of the C_n and BC_n Ruijsenaars–Schneider models, *J. Math. Phys.* **41** (2000) 8132-8147.
- [25] B.G. Puztai, Action-angle duality between the C_n -type hyperbolic Sutherland and the rational Ruijsenaars–Schneider–van Diejen models, *Nucl. Phys.* **B 853** (2011) 139-173.
- [26] B.G. Puztai, The hyperbolic BC_n Sutherland and the rational BC_n Ruijsenaars–Schneider–van Diejen models: Lax matrices and duality, *Nucl. Phys.* **B 856** (2012) 528-551.
- [27] B.G. Puztai, Scattering theory of the hyperbolic BC_n Sutherland and the rational BC_n Ruijsenaars–Schneider–van Diejen models, *Nucl. Phys.* **B 874** (2013) 647-662.
- [28] A.W. Knap, *Lie groups beyond an introduction*, Progress in Mathematics, vol. 140, Birkhäuser, Boston, MA, 2002.
- [29] L. Fehér and B.G. Puztai, Spin Calogero models associated with Riemannian symmetric spaces of negative curvature, *Nucl. Phys.* **B 751** (2006) 436-458.
- [30] L. Fehér and B.G. Puztai, A class of Calogero type reductions of free motion on a simple Lie group, *Lett. Math. Phys.* **79** (2007) 263-277.
- [31] J. Avan, O. Babelon and M. Talon, Construction of the classical r -matrices for the Toda and the Calogero models, *St. Petersburg Math. J.* **6** (1995) 255-274.
- [32] B.G. Puztai, On the r -matrix structure of the hyperbolic BC_n Sutherland model, *J. Math. Phys.* **53** (2012) 123528.
- [33] H.W. Braden, V.A. Dolgushev, M.A. Olshanetsky and A.V. Zotov, Classical r -matrices and the Feigin–Odesskii algebra via Hamiltonian and Poisson reductions, *J. Phys. A: Math. Gen.* **36** (2003) 6979-7000.
- [34] O. Babelon, D. Bernard and M. Talon, *Introduction to Classical Integrable Systems*, Cambridge University Press, 2003.
- [35] O. Babelon and C.-M. Viallet, Hamiltonian structures and Lax equations, *Phys. Lett.* **B 237** (1990) 411-416.
- [36] J. Avan and M. Talon, Classical R -matrix structure for the Calogero model, *Phys. Lett.* **B 303** (1993) 33-37.
- [37] E.K. Sklyanin, Dynamical r -matrices for the elliptic Calogero–Moser model, *St. Petersburg Math. J.* **6** (1995) 397-406.
- [38] H.W. Braden and T. Suzuki, R -matrices for elliptic Calogero–Moser models, *Lett. Math. Phys.* **30** (1994) 147-158.

- [39] F.W. Nijhoff, V.B. Kuznetsov, E.K. Sklyanin and O. Ragnisco, Dynamical r -matrix for the elliptic Ruijsenaars–Schneider system, *J. Phys. A: Math. Gen.* **29** (1996) L333-L340.
- [40] M. Forger and A. Winterhalder, Dynamical R -matrices for Calogero models, *Nucl. Phys.* **B 621** (2002) 523-570.
- [41] J. Avan and G. Rollet, Structures in BC_N Ruijsenaars–Schneider models, *J. Math. Phys.* **43** (2002) 403-416.
- [42] L. Fehér and T.F. Görbe, Duality between the trigonometric BC_n Sutherland system and a completed rational Ruijsenaars–Schneider–van Diejen system, *J. Math. Phys.* **55** (2014) 102704.
- [43] K. Chen, B.Y. Hou and W.L. Yang, The Lax pairs for elliptic C_n and BC_n Ruijsenaars–Schneider models and their spectral curves, *J. Math. Phys.* **42** (2001) 4894-4914.
- [44] K. Chen and B.Y. Hou, The D_n Ruijsenaars–Schneider model, *J. Phys. A: Math. Gen.* **34** (2001) 7579-7589.
- [45] L. Fehér and C. Klimčík, Poisson–Lie interpretation of trigonometric Ruijsenaars duality, *Commun. Math. Phys.* **301** (2011) 55-104.
- [46] L. Fehér and C. Klimčík, Self-duality of the compactified Ruijsenaars–Schneider system from quasi-Hamiltonian reduction, *Nucl. Phys.* **B 860** (2012) 464-515.
- [47] I. Marshall, A New Model in the Calogero–Ruijsenaars Family, *Commun. Math. Phys.* **338** (2015) 563-587.
- [48] J. Avan, O. Babelon and E. Billey, The Gervais–Neveu–Felder equation and the quantum Calogero–Moser systems, *Commun. Math. Phys.* **178** (1996) 281-299.
- [49] Y.B. Suris, Why is the Ruijsenaars–Schneider hierarchy governed by the same R -operator as the Calogero–Moser one?, *Phys. Lett.* **A 225** (1997) 253-262.
- [50] Z. Nagy, J. Avan and G. Rollet, Construction of dynamical quadratic algebras, *Lett. Math. Phys.* **67** (2004) 1-11.
- [51] G.E. Arutyunov, L.O. Chekhov and S.A. Frolov, R -Matrix Quantization of the Elliptic Ruijsenaars–Schneider Model, *Commun. Math. Phys.* **192** (1998) 405-432.
- [52] B.Y. Hou and W.L. Yang, The dynamical twisting and nondynamical r -matrix structure of the elliptic Ruijsenaars–Schneider model, *J. Math. Phys.* **41** (2000) 357-369.