

Oral administration of fumonisin B₁ and T-2 individually and in combination affects hepatic total and mitochondrial membrane lipid profile of rabbits

Running title: Hepatotoxicity of T-2 and FB₁

A Szabó^{1,2*}, J Szabó-Fodor², H Fébel³, M Mézes⁴, G Bajzik¹, M Kovács²

¹ Kaposvár University, Institute of Diagnostic Imaging and Radiation Oncology, 7400, Kaposvár, Guba S. u. 40., Hungary;

² Hungarian Academy of Sciences - Kaposvár University “MTA-KE Mycotoxins in the Food Chain” Research Group, 7400, Kaposvár, Guba S. u. 40. Hungary;

³ Research Institute for Animal Breeding, Nutrition and Meat Science, National Agricultural Research Center, Gesztenyés u. 1., 2053, Herceghalom, Hungary;

⁴ Szent István University, Faculty of Agricultural and Environmental Sciences, Department of Nutrition, 2013, Gödöllő, Páter K. u. 1., Hungary.

*corresponding author, Email: szanl125@freemail.hu

Tel.: +36 82 502 000/6020.ext.

Fax.: +36 82 502 020

fodor.judit@ke.hu

hedvig.febel@atk.hu

mezes.miklos@mkk.szie.hu

bajzik.gabor@sic.ke.hu

kovacs.melinda@ke.hu

Abstract

Weaned rabbits were fed diets contaminated with 2 mg/kg diet T-2 toxin alone, or 10 mg/kg diet fumonisin B₁ (FB₁) alone, and both toxins in combination (2+10 mg / kg, resp.), as compared to a toxin free control diet. Samplings were performed after 4 weeks (blood and liver). Bodyweight of T-2-fed group was lower after 4 weeks; the liver weight was increased dramatically (3-fold of control). Liver total phospholipids (PL) provided slight alterations in the fatty acid (FA) composition; all three toxin treated groups showed decrease in palmitoleic acid (C16:1 n7) proportion. In the liver mitochondrial PL FA composition margaric acid (C17:0) proportion decreased in the separated toxin treatments, compared to the combined setting. Oleic acid (C18:1 n9) proportion was increased and arachidonic acid (C20:4 n6) was decreased in the by FB₁-treated group, while docosapentaenoic acid (C22:5 n3) was decreased

in the separated treatments. The total monounsaturations were significantly higher in the FB₁ group's mitochondrial PL FA profile. After 4 weeks, all toxin treatments decreased the blood plasma reduced glutathione and glutathione peroxidase activity, and FB₁ increased the plasma sphinganine/sphingosine ratio. Both mycotoxins seem to cross the hepatocellular and the hepatic mitochondrial membrane, without drastic membrane disruption, as assessed from the PL FA composition, but inducing detectable lipid peroxidation.

Keywords: rabbit, T-2, FB₁, membrane lipids, liver, mitochondria, lipid peroxidation

Introduction

Mycotoxins are secondary metabolites of fungal (mold) metabolism with various biological activities, e.g. protein (T-2) or ceramide (Fumonisin B₁, FB₁) synthesis inhibition, but T-2 and FB₁ in high concentrations and prolonged consumption are as well carcinogens (9,11, 13). Since most fungal strains produce single or multiple mycotoxins, their combined effects are nowadays in focus (12).

T-2 toxin is a trichothecene, produced by *Fusarium* species, occurring mostly in corn and further cereals. T-2 toxin is metabolized (hydrolyzed to HT-2) after ingestion, yielding multiple metabolites with known or less known toxicity and membrane association capability (9). Their mode of toxic action is rather divergent, as the inhibition of protein synthesis (specifically in eukaryotes) and inhibition of DNA synthesis (9), or the disruption of the mitochondrial electron transport system (26), augmented lipid peroxidation, affecting ultimately cell membrane integrity. According to Ngampongsa et al. (26), the ATP-linked oxygen consumption rate of cardiomyocyte mitochondria is compromised by T-2 toxin, which is underpinned with our recent results, in which rabbit erythrocyte Na⁺/K⁺ ATPase activity (the rate of ATP breakdown) was markedly lowered by T-2 toxin (33). It was established that in the rat liver T-2 toxin inhibits mitochondrial electron transport system, concluding that mitochondria are a possible hepatic targets of T-2 toxin action (27). It is important to add that even in embryonic stem cells (being pluripotent!), T-2 toxin induced apoptosis via a reactive oxygen species mediated mitochondrial pathway (14). In summary, it is proven that T-2 toxin crosses the hepatic cellular and subcellular membranes and disrupts ATP production in mitochondria (28). An essential addition is that hepatic subcellular distribution of labelled [³H]T-2 toxin revealed immediate association to the plasma membrane and a concomitant association to the membranes of other cellular compartments, including mitochondria (28).

Anyhow, T-2 toxin is less known considering its membrane disruptor effect exerted during its membrane transitory movement, on which the present study was focused. The question is even more intriguing taking the solubility of pure T-2 toxin into account, namely it is only weakly water soluble (3).

Molecular - compositional changes underlying hepatic membrane lipid alterations (alteration of the lipid profile of phospholipid classes in the microsomal, the nuclear and the cellular membranes) caused by FB₁ were partly elucidated when feeding rats a drastically toxic, cancer promoting toxin level (250 mg/kg feed) (6). In the same study FB₁ altered lipid constituent proportions, in particular phosphatidylethanolamines (PE) were significantly increased in the mitochondrial and even in plasma membrane fractions. The fatty acid profile of the main polar lipid fractions (i.e. P-choline and PE) showed a decrease in polyunsaturated (especially C20:4 n6, arachidonic acid)/saturated (P/S) FA ratio, suggesting a less fluid membrane state. The decreased polyunsaturated fatty acid levels in PC along with a marked elevation of oleic (C18:1 n9) and linoleic (C18:2 n6) acids were indicative of an impaired Δ 6 desaturase activity. Authors concluded that FB₁ disrupts sphingolipid, phospholipid (the polar lipid fractions interaction with a polar toxin molecule), cholesterol and FA metabolism in the liver, in a manner to decrease membrane fluidity (ultimately leading to apoptosis). Arachidonic acid has a profound role in this, namely the outer plasma membrane lipid leaflet is enriched with this acid, being the precursor of inflammatory eicosanoids.

Since both T-2 and FB₁ are unequivocal membrane disruptors even at the subcellular level, their detailed study is reasoned.

The co-occurrence of mycotoxins makes the cytotoxicity assessment more complex, meanwhile Clarke et al. (9) used the EU regulation concentration limits for FB₁, ochratoxin and aflatoxin B₁ in a tertiary combination (9). In our recent study it was found that at levels of 2 mg/kg diet T-2 toxin alone, or 10 mg/kg diet fumonisin B₁ alone, and both toxins in combination exert hepatotoxic effect and act antagonistically onto the red cell sodium pump ATP breakdown capacity, T-2 markedly lowering its activity (33). However, red cell membrane fatty acyl chain composition was only slightly modified in the cells of the systemic circulation, supposing that both toxins act at the level of polychromatic erythroblasts. Here we tested the effects of the same toxin exposure onto the hepatic membrane and hepatic mitochondrial membrane composition and investigated possibly emerging lipid peroxidation.

Materials and Methods

Experimental design

Pannon white littermate male rabbits weaned at the age of 35 days were caged as pairs, and 4 groups were formed, each containing 10 animals. The animals were born, housed and treated at the rabbitry of the Experimental Farm of the Kaposvár University. One group was fed a rabbit diet complemented with 2 mg/kg T-2 toxin, another the same basal diet complemented with 10 mg/kg fumonisin B₁ (FB₁), while a further was given T-2 and FB₁ in a combined form (2+10 mg/kg diet). The control group was fed a toxin free diet containing the same ingredients as the experimental ones, with the exception of fungal cultures. Feed and water were *ad libitum* offered. Bodyweight and feed intake was measured weekly, individually, and the entire feeding period lasted for 4 weeks. All data were recorded on individually marked animals, from 6 rabbits (chosen out randomly of the total of 10) in each group.

Production of toxins and determination of mycotoxin content in fungal cultures and feeds

Fumonisin B₁ and T-2 toxin were produced with the application of *Fusarium verticillioides* strain MRC 826 and *Fusarium sporotrichioides* strain NRRL 3299, respectively (15). The homogenized fungal cultures contained FB₁ at a concentration of 3440 mg/kg, while T-2 toxin concentration was 1338 mg/kg (HT-2 toxin 285 mg/kg), respectively. These fungal cultures were mixed into the basal feed of the experimental animals, so as to provide contaminated feeds with 2 mg/kg feed T-2 toxin and/or 10 mg/kg feed FB₁.

The mycotoxin concentration of the control and the experimental feeds was determined with LC-MS (Shimadzu, Kyoto, Japan). The limit of detection (LOD) for FB₁ was 3 µg/kg, while for T-2 toxin it was 10 µg/kg. The diet fed to the control group did not contain detectable amounts of T-2 and FB₁.

Blood and liver tissue sampling

Fresh venous blood was sampled into heparinized (20 IU/ml whole blood) tubes and was centrifuged for 10 min at 1000 g, at the end of the 4 weeks. Plasma was removed and stored at -70 °C, for sphinganine and sphingosine analysis (*section 2.6.*).

After exsanguination whole livers were immediately dissected and weighed. Tissue samples of ca. 5 g were cut with scissors in a medium according to Brand et al. (5). Liver samples were rinsed and resuspended in 10 volumes of ice-cold medium, containing 250 mM sucrose, 5 mM Tris/HCl and 2 mM EGTA (pH 7.4 at 4 °C), then homogenized using six

passes of a motorized Teflon/glass Potter-Elvehjem homogenizer. All chemicals were purchased from Sigma-Aldrich (Schnelldorf, Germany). The homogenate was centrifuged at 1000 g for 3 minutes, and the supernatant was centrifuged at 12000 g for further 10 minutes at 4 °C. The pellet was then suspended, respun at 12000 g / 10 minutes (MLW K24 Janetzky, Engelsdorf, Germany), resuspended in isolation medium and kept at -70 °C.

Lipid analysis

Whole tissue samples, as well as isolated mitochondria were homogenized (IKA T25 Digital Ultra Turrax, Staufen, Germany) in a 20-fold volume of chloroform:methanol (2:1 vol:vol) and total lipid content was extracted (16). Solvents were ultrapure-grade (Sigma-Aldrich, Schnelldorf, Germany) and 0.01 % w:v butylated hydroxytoluene was added to prevent fatty acid oxidation. In the frame of lipid fractionation, extracted total lipids were transferred to glass chromatographic columns, containing 300 mg silica gel (230-400 mesh) for 10 mg of lipids (24). Neutral lipids were eluted with 10 mL chloroform for the above fat amount, then 15 mL acetone:methanol (9:1 vol:vol) was added, while 10 mL pure methanol eluted the total phospholipids. This latter fraction was evaporated to dryness under a nitrogen stream and was transmethylated with a base-catalysed NaOCH₃ method (8). Fatty acid methyl-esters were extracted into 300 µL ultrapure n-hexane for gas chromatography. Latter was performed on a Shimadzu 2010 apparatus (AOC 20i automatic injector), equipped with a Phenomenex Zebron ZB-WAX Capillary GC column (30 m x 0.25 mm ID, 0.25 micrometer film, Phenomenex Inc., Torrance, CA, USA) and flame ionisation detector (FID 2×10⁻¹¹). Characteristic operating conditions were: injector temperature: 270 °C, detector temperature: 300 °C, helium flow: 28 cm/sec. The oven temperature was graded: from 80 to 205 °C: 2.5 °C/min, 5 min at 205 °C, from 205 to 250 °C 10 °C/min and 5 min at 210 °C. The makeup gas was nitrogen. To identify individual FA, an authentic external FA standard (37 Component FAME Mix, Sigma-Aldrich, Cat No: CRM47885) was used. Fatty acid results were expressed as weight % of total fatty acid methyl esters.

Unsaturation index (UI) was defined as the number of double bonds in 100 fatty acyl chains. From the FA results unsaturation index (UI) was calculated as: $UI = [(1 \times \Sigma \text{ monoenoic FA}) + (2 \times \Sigma \text{ dienoic FA}) + (3 \times \Sigma \text{ trienoic FA}) + (4 \times \Sigma \text{ tetraenoic FA}) + (5 \times \Sigma \text{ pentaenoic FA}) + (6 \times \Sigma \text{ hexaenoic FA})]$ (7). The average FA chain length was calculated from the multiplication of the chain length values and the respective proportion of each fatty acid.

Assessment of antioxidant defence capacity

Lipid peroxidation was assessed in blood plasma with the determination of malondialdehyde (MDA) levels with 2-thiobarbituric acid method (30). The concentration of plasma reduced glutathione (GSH) was measured with photometry (31) and the activity of plasma glutathione peroxidase (GSHPx) according to Lawrence and Burk (22).

Determination of the plasma sphinganine (Sa) to sphingosine (So) ratio

Determination of Sa/So was carried out using acetonitrile (900 µl) addition to the plasma (100 µl) and the mixture was centrifuged at 10000 rpm for 5 min and filtered through a 0.45 µm filter. Sphinganine and sphingosine levels were determined by LC-MS. Chromatographic separation was carried out using a Waters Symmetry C₁₈ column (150 x 2.1 mm ID 5 µm; Waters, Mildford, MA, USA). Quantitative measurement was based on linear calibration to authentic standards.

Statistical evaluation

For the comparison of group means (bodyweight, liver weight fatty acid profile) multivariate analysis of variance was used (evaluating as well the interactions), with the Tukey „*post hoc*” test for the inter-group differences, preceded with the analysis of normal distribution (Kolmogorov-Smirnov test) (19).

Ethical issues

The experiment was carried out according to the regulations of the Hungarian Animal Protection Act. The allowance number for this study was 656-1/2012.

Results

Bodyweight and liver weight

After 4 weeks of feeding, the T-2 and the T-2+FB₁ groups bodyweight was significantly lower, as compared to the control or even to the pure FB₁ data, although the feed intake was not different among the groups (data not shown). The liver weight at the end of the experiment in the group fed T-2 fed toxin was significantly higher compared to the other treatment groups and reached ca. 3 times higher weight than the control (**Table 1**). However, T-2+FB₁ in combination or FB₁ alone did not significantly alter liver weight.

Hepatic total phospholipids

In **Table 2**, showing the total hepatic PL FA composition it is visible that all three toxin treatments decreased the proportion of palmitoleic acid (C16:1 n7). Arachidic acid (C20:0) proportion in the liver total PL fraction was increased by the combined toxin treatment (T-2+FB₁), as compared to the control. Interestingly, the unsaturation index (UI) was the highest in the combined toxin (T-2+FB₁) exposed animals, differing significantly only from the control, but not from the two individual toxin treatments (T-2 and FB₁ separately). A totally identical alteration was observed in the average fatty acyl chain length (ACL). Altogether, neither the separated nor the combined mycotoxin treatments induced drastic fatty acid compositional alteration in the hepatic total phospholipids.

Liver mitochondrial phospholipids

In the isolated mitochondria, the membrane lipids revealed to be more prone towards the mycotoxin exposure. **Table 3** summarizes the results concerning the alterations of the mitochondrial membrane PL FA composition. Margaric acid (C17:0) proportion was decreased in both separated toxin treatments, as compared to the combined setting. Oleic acid (C18:1 n9) (a major membrane component of the PL fraction) proportion was significantly increased by FB₁. Arachidonic acid (C20:4 n6) was decreased in the FB₁ group, as compared to the control and the binary mixture treatment. Docosapentaenoic acid (C22:5 n3) proportion was decreased significantly in both separate treatments, as compared to the combined group, but not differing from the control.

In the calculated FA data, total monounsaturated (MUFA) was significantly higher in the FB₁ group, as compared to the control. A fully opposite result was found for the total polyunsaturation, while the total n3 FA proportion was increased (on the expense of the total n6 FAs).

Antioxidant defense capacity

In the blood plasma the concentration of reduced glutathione was the highest in the control, significantly differing from all toxin treated groups. Identical finding was detected by glutathione-peroxidase, but no alteration was detected in the plasma malondialdehyde concentration (Figure 1).

Plasma sphinganine (Sa) to sphingosine (So) ratio

After 4 weeks of mycotoxin exposure, plasma sphinganine/sphingosine ratio was increased by FB₁ alone, but to a lesser extent, even in combination with T-2 toxin, while T-2 toxin alone did not influence the plasma Sa/So ratio (Figure 2).

Discussion

Bodyweight and liver weight

The toxin exposure lasted for 28 days until the animals reached the approximate Hungarian conventional slaughter weight. T-2 alone or in combination (with FB₁) appeared as a growth depressing factor. Meanwhile fumonisin B₁ was administered in relatively high dose (10 mg / kg feed), as a single toxin it had no detectable effect either on bodyweight or on liver weight, as compared to the control.

In contrast, T-2 toxin led to a marked growth depression, independently of the administration type (single toxin or co-occurring with FB₁). In a recent study (34) the <0.02 mg/kg bodyweight/day T-2 concentration value was defined as the no observed adverse effect level (NOAEL) for rabbit bucks. This present treatment was exceeding this value (ca. 0.96 mg/kg bodyweight/day), leading to well established compromised growth. The three-fold increased liver weight (as in this study) is a well-known consequence even in acute T-2 toxicosis in rabbits, accompanied by centrolobular dystrophy (34). (The most remarkable pathological findings included enlarged, friable, pale, yellowish liver, with spot-like necrotic areas (unpublished observations). The results clearly refer to the hypothesis that T-2 toxin, as a type A trichothecene with its characteristic hydroxyl and ester groups (thus being amphipathic (27)) exerts a direct hepatotoxic effect, unequivocally supporting the hypothesis that it – due to its molecular characteristics – effectively crosses the hepatocellular membrane (i.e. the amphipathic PLs) (9).

Fatty acid composition

Hepatic total phospholipids

In spite of the dramatically enlarged liver, T-2 toxin alone was less affecting hepatic total phospholipid fatty acyl chain composition. There was no single fatty acid of which T-2 toxin clearly altered the proportion, as compared to any of the other (combined or single) toxin

treatments. Altogether one monounsaturated (C16:1 n7, palmitoleic acid), one odd chain (C17:0, margaric acid) fatty acid showed altered proportion in the liver total PLs of the T-2 toxin group, as compared to the control. It was interesting that even on the calculated fatty acid parameters, T-2 toxin had only a slight effect, not differing significantly from the control, even not in the unsaturation index value (UI).

The only hint that FB₁ alters hepatic (microsomal) membrane lipid palmitoleic acid proportion has been published by Gelderblom et al. (17). These authors described a disturbed microsomal lipid metabolism with increased delta-9 desaturase activity (not proven in the hepatic total PLs in our study), but it has to be added that in the cited study 100 mg/kg feed FB₁ was fed to rats (carcinogenic level). Interestingly, the inhibition of ceramide synthase by FB₁ was not reflected in the microsomal membrane lipids (17), even at this high-dose treatment.

As the first level interpretation, for FB₁ identical results were found to that of T-2 toxin in the liver total PLs. However, as compared to the rat microsomal membranes, FB₁ alone was not a very potent modulator/disruptor of hepatic total PL FA composition, neither inducing liver weight alteration.

In contrast, the combined application (T-2+FB₁) of the two mycotoxins, as compared to the control or even the individual toxin exposures was “effective”, in that this treatment reduced the C16:1 n7 proportion (but did not perturb delta-9 desaturation significantly!) and slight, but significant increase was found for arachidic acid, the unsaturation index (UI) and the average fatty acyl chain length (ACL). In the multivariate statistical model, there was no interaction for ACL between the treatments (i.e. T-2 and FB₁), but for C16:1 n7, C20:0 and UI the interaction (i.e. T-2 x FB₁) of the two toxins was significant at the level of P<0.001. We thus suppose a synergism in these cases, the two fusariotoxins augmenting their common effects upon the latter FAs and unsaturation.

Comparing our results to literature data, it must be emphasized that total hepatic phospholipids have not yet been analyzed as the targets of combined T-2+FB₁ toxins. It thus seems that at the moderately high levels neither T-2 toxin, nor FB₁ are hepatocellular membrane disruptors in the leporine model, when the exposure is relatively short (4 weeks), but T-2 toxin provides a rather effective translocation across the hepatocellular membrane with major parenchymal consequences.

Liver mitochondrial phospholipids

In isolated mitochondria, the PLs were more strongly modified by the mycotoxin exposure. Margaric acid (C17:0) proportion was decreased by both individual toxin treatments (T-2 and FB₁) as compared to the combined setting, while the combined treatment increased its proportion (but was not different from the control). Their statistical interaction in the multivariate model was significant ($P < 0.001$), suggesting a synergistic effect. In rabbit tissue membranes, margaric acid abundance is clearly depending on two factors. Once, odd chain FAs are exclusively synthesized by the gut microflora, and as second, odd chain saturated FAs are peripherally (position *sn*-1) located in phosphatidylcholine and are primary targets of beta-oxidation (being quasi equivalent with paired chain saturated FAs in this respect), when hydrolyzed by the most abundant PLase A₂ (23). The enzymatic cleavage of an odd chain saturated FA at position *sn*-1 may be a part of the Lands mechanism (10, 21), a means to regulate membrane fluidity via the modification of the glycerolipid FA side chain combinations.

Oleic acid (C18:1 n9) proportion was significantly increased only in the FB₁ group, and in the combined treatment the difference was not significant among individual treatments and control. In our earlier study (33), in the erythrocyte membrane, oleic acid provided the highest proportion in the combined treatment. This partly agrees with the results of Gelderblom et al. (17), who reported that FB₁ at 100 mg/kg feed induced delta-9 desaturase activation and increased membrane monounsaturations. It is important to add that we only fed 4 weeks of this concentration, but it is a necessary addition that rabbits perform coecotrophy (33), while rats coprophagy (17); this means that the exact toxin (or mycotoxin metabolite) concentration at the site of action (liver) is with the highest probability higher than that in the feed. The reason why FB₁ seems to be a more potent modulator of delta-9 desaturation may be due to the lower stability of T-2 toxin, which is very quickly, enzymatically hydrolyzed to HT-2 (9). Meanwhile FB₁ is a more stable molecule and there are further factors to be considered in rabbits. In monogastric animals (piglets), the accumulative absorption of fumonisin B₁ is only ca. 4% (15). However, as a result of coecotrophy in rabbit, the accumulative absorption may increase, while the hidden proportion, occurring in a matrix associated form may be as well liberated by the digestive enzymes (during the multiplied uptake via coecotrophy), leading ultimately to a higher toxin exposure than determined analytically from the food (34).

The biological background to decrease polyunsaturation in parallel to increase monounsaturations is a means to maintain membrane fluidity by decreasing the sensitivity

towards lipid peroxidation (29). Although it was not directly measured from the liver, in the blood plasma the concentration of reduced glutathione was the highest in the control, significantly differing from all other toxin treatments, referring to an effective scavenger role against the emerging reactive oxygen substances (Figure 1). The same alteration was found for glutathione-peroxidase, but no alteration was detected in the plasma malondialdehyde concentration. MDA is formed in the terminal phase of lipid peroxidation from FAs having at least three double bonds, and is strongly cytotoxic (23). In this study the extent of lipid peroxidation, as assessed from the blood plasma was maintained at a relatively low level.

It thus seems that the cross-membrane transport has a more widespread effect, even detectable in the systemic circulation (4), as published for T-2 (broiler chickens) and for FB₁ in pigs (20). The role of GSHPx is to “up-oxidate” the reduced glutathione moiety. Indeed, our results supported this fact, namely GSHPx was likewise in a more saturated enzymatic phase, as compared to the control.

Mitochondrial membrane arachidonic acid (C20:4 n6) proportion decreased only in the FB₁ group, as compared to the control and the binary mixture treatment. In the biological membranes, primarily in phospholipids, arachidonic acid is the root component of the eicosanoids, liberated via the activity of phospholipase A₂. Although different cell organelles and lipid fractions react to FB₁ induced stimuli divergently, Burger et al. (6) reported supportive results in that FB₁ was liberated from the PC fraction, augmenting prostaglandin E₂ series synthesis, inducing sustaining cell proliferation (2). The further fact to take into account is that impaired arachidonic acid metabolism, in particular its proportional decrease in the PC fraction has been implied as a growth stimulus for hepatocellular nodules (1).

Docosapentaenoic acid (C22:5 n3) proportion was decreased significantly in both separate treatments, as compared to the combined group, but not differing from the control. According to Abel et al. (2001), the proportional decrease of C20-C22 FAs is a factor to control/improve antioxidant properties. This is underpinned with the fact that meanwhile the pentaenoic FA proportion and the monounsaturations were increased and there is a ca. 4-fold sensitivity difference in the peroxidation sensitivity between oleic and docosapentaenoic acids (18). In the calculated FA data total monounsaturations (MUFA) was significantly higher in the FB₁ group, as compared to the control. A fully opposite result was found for the total polyunsaturations, while the total n3 FA proportion increased.

Summarizing the above findings in the mitochondrial fatty acid composition, Burger et al. (6) reported that the modified fatty acid profile may be coupled with an altered growth pattern of FB₁ intoxicated hepatocytes, including as well the perturbation of the ceramide synthesis. Indeed, we did not measure hepatocellular, but only plasma sphinganine/sphingosine ratio, it was found that FB₁ alone, but to a lesser extent, even in combination with T-2 is able to markedly increase this ratio, while T-2 did not influence the plasma Sa/So ratio (Figure 2).

In conclusion, it was found that at the pre-defined mycotoxin concentration values young rabbits are more prone towards T-2 toxin administration, which slightly alters hepatic plasma and mitochondrial membrane composition. When T-2 toxin was fed in combination with FB₁, synergistic effects was proven for saturated (C16:0, C17:0, C20:0) and monounsaturated fatty acids. All toxin treatments induced lipid peroxidation and activated the antioxidant defense (GSHPx). Fumonisin was moreover proven to perturb membrane ceramide homeostasis, even at moderate concentration and a relatively short exposition time.

Acknowledgement

The research was supported by the Hungarian Academy of Sciences (within the frame of the MTA-KE “Mycotoxins in the food chain” Research Group), by the National Scientific Research Fund (OTKA K 100810) to M.K., and János Bolyai Research Grants of the Hungarian Academy of Sciences (BO/499/13) to J.Sz.F.

REFERENCES

1. Abel S, Smuts CM, de Villiers C, Gelderblom W: Changes in essential fatty acid patterns associated with normal liver regeneration and the progression of hepatocyte nodules in rat hepatocarcinogenesis. *Carcinogenesis* 22, 795–804 (2001)
2. Bagga D, Wang L, Farias-Eisner R, Glaspy JA, Reddy ST: Differential effects of prostaglandin derived from ω -6 and ω -3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion. *Proc. Natl. Acad. Sci.* 100, 1751–1756 (2003)
3. Black JA, Foarde KK, Menetrez MY: Solvent comparison in the isolation, solubilization, and toxicity of *Stachybotrys chartarum* spore trichothecene mycotoxins in an established in vitro luminescence protein translation inhibition assay. *J. Microbiol. Methods* 66, 354–361 (2006)

4. Bócsai A, Pelyhe C, Zándoki E, Ancsin Z, Szabó-Fodor J, Erdélyi M, Mézes M, Balogh K: Short-term effects of T-2 toxin exposure on some lipid peroxide and glutathione redox parameters of broiler chickens. *J. Anim. Physiol. Anim. Nutr. in press* (Berl). (2015)
5. Brand MD, Turner N, Ocloo A, Else PL, Hulbert AJ: Proton conductance and fatty acyl composition of liver mitochondria correlates with body mass in birds. *Biochem. J.* 376, 741–748 (2003)
6. Burger HM, Abel S, Snijman PW, Swanevelder S, Gelderblom WCA: Altered lipid parameters in hepatic subcellular membrane fractions induced by fumonisin B₁. *Lipids* 42, 249–261 (2007)
7. Buttemer WA, Battam H, Hulbert AJ: Fowl play and the price of petrel: long-living Procellariiformes have peroxidation-resistant membrane composition compared with short-living Galliformes. *Biol. Lett.* 4, 351–354 (2006)
8. Christie WW: A simple procedure for rapid transmethylolation of glycerolipids and cholesteryl esters. *J. Lipid Res.* 23, 1072–1075 (1982)
9. Clarke R, Connolly L, Frizzell C, Elliott CT: Cytotoxic assessment of the regulated, co-existing mycotoxins aflatoxin B₁, fumonisin B₁ and ochratoxin, in single, binary and tertiary mixtures. *Toxicon* 90, 70–81 (2014)
10. Cevc G: *Phospholipids Handbook*. Marcel Dekker, New York, USA. (1993)
11. Dutton MF: Fumonisin, mycotoxins of increasing importance: their nature and their effects. *Pharmacol. Ther.* 70, 137–61. (1996)
12. EFSA (European Food Safety Authority): Scientific opinion on the risks for animal and public health related to the presence of T-2 and HT-2 toxin in food and feed. *EFSA* 9, 2481-2668 (2011)
13. Escrivá L, Font G, Manyes L: In vivo toxicity studies of fusarium mycotoxins in the last decade: a review. *Food Chem. Toxicol.* 78, 185–206 (2015)
14. Fang H, Wu Y, Guo J, Rong J, Ma L, Zhao Z, Zuo D, Peng S: T-2 toxin induces apoptosis in differentiated murine embryonic stem cells through reactive oxygen species-mediated mitochondrial pathway. *Apoptosis* 17, 895–907 (2012)
15. Fodor J, Balogh K, Weber M, Miklós M, Kametler L, Pósa R, Mamet R, Bauer J, Horn P, Kovács F, Kovács M: Absorption, distribution and elimination of fumonisin B(1) metabolites in weaned piglets. *Food Addit. Contam. Part A.* 25, 88–96 (2008)
16. Folch JM, Lees M, Sloane-Stanley GH: A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226, 495-509 (1957)

17. Gelderblom WCA, Moritz W, Swanevelder S, Smuts CM, Abel S: Lipids and delta6-desaturase activity alterations in rat liver microsomal membranes induced by fumonisin B₁. *Lipids*. 37, 869–877 (2002)
18. Gunstone FD, Harwood JL, Padley FB: *The Lipid Handbook*. Chapman and Hall, London, UK (1994)
19. IBM SPSS 20 for Windows. SPSS Inc., Chicago, IL. USA (2012)
20. Klarić MS, Pepeljnjak S, Domijan AM, Petrik J: Lipid peroxidation and glutathione levels in porcine kidney PK15 cells after individual and combined treatment with fumonisin B(1), beauvericin and ochratoxin A. *Basic Clin. Pharmacol. Toxicol.* 100, 157–164 (2007)
21. Lands WE: Metabolism of glycerolipids. 2. The enzymatic acylation of lysolecithin. *J. Biol. Chem.* 235, 2233–2237 (1960)
22. Lawrence RA, Burk RF: Species, tissue and subcellular distribution of non Se-dependent glutathione peroxidase activity. *J. Nutr.* 108, 211–215 (1978)
23. Mead JF, Alfin-Slater RB, Howton DR, Popják, G: *Lipids: Chemistry, biochemistry and nutrition*. Plenum Press, N.Y., USA (1986)
24. Leray C, Andriamampandry M, Gutbier G, Cavadenti J, Klein-Soyer C, Gachet C, Cazenave JP: Quantitative analysis of vitamin E, cholesterol and phospholipid fatty acids in a single aliquot of human platelets and cultured endothelial cells. *J. Chromatogr. B. Biomed. Sci. Appl.* 696, 33–42 (1997)
25. Moon WS, Kim JH, Kang MJ, Lee DG: Early ultrastructural changes of apoptosis induced by fumonisin B₁ in rat liver. *Yonsei Med. J.* 41, 195–204 (2000)
26. Ngampongsa S, Hanafusa M, Ando K, Ito K, Kuwahara M, Yamamoto Y, Yamashita M, Tsuru Y, Tsubone H: Toxic effects of T-2 toxin and deoxynivalenol on the mitochondrial electron transport system of cardiomyocytes in rats. *J. Toxicol. Sci.* 38, 495–502 (2013)
27. Pace JG: Effect of T-2 mycotoxin on rat liver mitochondria electron transport system. *Toxicon* 21, 675–680 (1983)
28. Pace JG, Watts MR: Hepatic subcellular distribution of [³H]T-2 toxin. *Toxicon*. 27, 1307–1311 (1989)
29. Pamplona R, Portero-Otín M, Ruiz C, Gredilla R, Herrero A, Barja G: Double bond content of phospholipids and lipid peroxidation negatively correlate with maximum longevity in the heart of mammals. *Mech. Ageing Dev.* 112, 169–183 (2000)
30. Placer ZA, Cushman LL, Johnson BC: Estimation of product of lipid peroxidation (malonyl dialdehyde) in biochemical systems. *Anal. Biochem.* 16, 359–164 (1966)

31. Sedlak J, Lindsay RH: Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal. Biochem.* 25, 192–205 (1968)
32. Sharma D, Asrani RK, Ledoux DR, Rottinghaus GE, Gupta VK: Toxic interaction between fumonisin B1 and moniliformin for cardiac lesions in Japanese quail. *Avian Dis.* 56, 545–554 (2012)
33. Szabó A, Szabó-Fodor J, Fébel H, Romvári R, Kovács M: Individual and combined haematotoxic effects of fumonisin B(1) and T-2 mycotoxins in rabbits. *Food Chem. Toxicol.* 72, 257–264 (2014)
34. Szabó-Fodor J, Dall'Asta C, Falavigna C, Kachlek M, Szécsi Á, Szabó A, Kovács M: Determination of the amount of bioaccessible fumonisin B₁ in different matrices after in vitro digestion. *World Mycotoxin J.* 8, 261–267 (2015)

Table 1. Liver and bodyweight of the four experimental groups (means \pm SD)

group	liver weight (g)			bodyweight (g; day 60)				
	mean	\pm	SD	mean	\pm	SD		
T-2	219.2	\pm	3.37	b	2280	\pm	82.6	A
FB ₁	75.8	\pm	4.35	ab	2695	\pm	84.1	B
T-2+FB ₁	72.8	\pm	4.35	a	2221	\pm	81.7	A
Control	91	\pm	8.39	ab	2653	\pm	69	B

a,b; A, B: difference between treatments at day 60

Table 2. The total hepatic phospholipid fatty acid composition of the control and the toxin treated groups (all data are means \pm SD of 6 animals)

group	T-2		FB ₁		T-2+FB ₁		Control	
liver PL	Mean	\pm SD	Mean	\pm SD	Mean	\pm SD	Mean	\pm SD
C14:0	0.06	\pm 0.01	0.08	\pm 0.02	0.07	\pm 0.01	0.12	\pm 0.09
C15:0	0.19	\pm 0.03	0.17	\pm 0.04	0.20	\pm 0.02	0.17	\pm 0.04
C16:0	19.1	\pm 0.97	19.2	\pm 1.33	17.5	\pm 1.73	20.3	\pm 2.81
C16:1 n7	0.39	\pm 0.13	0.45	\pm 0.05	0.31	\pm 0.14	0.66	\pm 0.22
C17:0	0.85	\pm 0.12	0.72	\pm 0.14	0.91	\pm 0.07	0.63	\pm 0.11
C18:0	24.4	\pm 0.96	23.1	\pm 1.08	24.1	\pm 1.88	22.5	\pm 1.64
C18:1 n9c	13.5	\pm 2.20	15.2	\pm 0.91	13.8	\pm 1.13	15.3	\pm 1.43
C18:2 n6c	29.2	\pm 2.47	29.1	\pm 0.74	29.1	\pm 0.07	28.4	\pm 2.38
C18:3 n6c	0.09	\pm 0.09	0.04	\pm 0.01	0.05	\pm 0.01	0.09	\pm 0.08
C18:3 n3	0.50	\pm 0.06	0.55	\pm 0.09	0.51	\pm 0.03	0.49	\pm 0.14
C20:0	0.04	\pm 0.00	0.04	\pm 0.00	0.05	\pm 0.01	0.04	\pm 0.01
C20:1 n9	0.27	\pm 0.05	0.32	\pm 0.16	0.35	\pm 0.15	0.31	\pm 0.12
C20:2 n6	0.68	\pm 0.14	0.89	\pm 0.17	0.90	\pm 0.29	0.75	\pm 0.26
C22:0	0.42	\pm 0.35	0.27	\pm 0.06	0.34	\pm 0.16	0.38	\pm 0.22
C20:3 n6	0.93	\pm 0.27	0.88	\pm 0.11	1.07	\pm 0.21	0.95	\pm 0.30
C20:3 n3	0.03	\pm 0.01	0.03	\pm 0.01	0.04	\pm 0.03	0.03	\pm 0.01
C20:4 n6	7.96	\pm 0.48	7.40	\pm 0.98	9.06	\pm 1.66	7.41	\pm 0.55
C22:5 n3	0.25	\pm 0.04	0.23	\pm 0.06	0.31	\pm 0.07	0.24	\pm 0.05
C22:6 n3	0.17	\pm 0.04	0.15	\pm 0.05	0.19	\pm 0.06	0.17	\pm 0.05
saturated	45.0	\pm 1.43	43.6	\pm 1.05	43.2	\pm 1.11	44.1	\pm 1.48
monoenoic	14.1	\pm 2.31	16.0	\pm 1.02	14.5	\pm 1.42	16.3	\pm 1.66
PUFA	39.8	\pm 2.67	39.3	\pm 1.05	41.2	\pm 1.72	38.6	\pm 2.73
n3	0.95	\pm 0.13	0.96	\pm 0.17	1.05	\pm 0.11	0.92	\pm 0.17
n6	38.9	\pm 2.60	38.3	\pm 0.93	40.2	\pm 1.63	37.6	\pm 2.70
n6/n3	41.2	\pm 4.26	40.8	\pm 6.17	38.7	\pm 3.41	42.0	\pm 6.58
UI	112.7	\pm 3.90	112.22	\pm 3.40	118.39	\pm 6.09	111.1	\pm 4.88
ACL	17.66	\pm 0.04	17.64	\pm 0.05	17.70	\pm 0.10	17.62	\pm 0.06

UI: unsaturation index; ACL: average chain length; a, b: significant ($P < 0.05$) difference between groups

Table 3. The hepatic mitochondrial phospholipid fatty acid composition of the control and the toxin treated groups (all data are means \pm SD of 6 animals)

group	T-2	FB ₁	T-2+FB ₁	Control
liver mit. PL	Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD
C14:0	0.21 \pm 0.19	0.28 \pm 0.12	0.15 \pm 0.10	0.22 \pm 0.16
C15:0	0.20 \pm 0.10	0.21 \pm 0.08	0.27 \pm 0.04	0.24 \pm 0.06
C16:0	20.6 \pm 2.03	21.0 \pm 3.73	18.2 \pm 1.24	18.4 \pm 1.78
C16:1 n7	1.27 \pm 0.46	1.20 \pm 0.17	0.75 \pm 0.23	1.13 \pm 0.43
C17:0	0.61 \pm 0.18 ^a	0.60 \pm 0.16 ^a	0.92 \pm 0.15 ^b	0.79 \pm 0.17 ^{ab}
C18:0	17.3 \pm 1.05	16.5 \pm 1.00	18.3 \pm 4.39	16.7 \pm 4.60
C18:1 n9c	23.4 \pm 4.68 ^{ab}	26.8 \pm 1.56 ^b	20.3 \pm 3.63 ^{ab}	19.1 \pm 4.99 ^a
C18:2 n6c	25.6 \pm 5.63	22.0 \pm 6.05	28.7 \pm 4.24	30.3 \pm 3.78
C18:3n6	0.19 \pm 0.06	0.38 \pm 0.17	0.09 \pm 0.05	0.19 \pm 0.24
C18:3n3	1.61 \pm 0.80	2.04 \pm 0.60	1.23 \pm 0.40	1.92 \pm 1.21
C20:0	0.08 \pm 0.02	0.09 \pm 0.02	0.08 \pm 0.01	0.08 \pm 0.03
C20:1	0.70 \pm 0.37	0.86 \pm 0.59	0.51 \pm 0.18	0.46 \pm 0.12
C20:3n6	0.90 \pm 0.44	0.80 \pm 0.37	1.39 \pm 0.31	1.28 \pm 0.45
C20:3n3	0.23 \pm 0.04 ^{ab}	0.27 \pm 0.20 ^b	0.07 \pm 0.03 ^{ab}	0.23 \pm 0.11 ^{ab}
C20:4n6	4.83 \pm 1.05 ^{ab}	4.04 \pm 0.34 ^a	6.39 \pm 1.58 ^b	6.43 \pm 1.32 ^b
C22:5n3	0.19 \pm 0.04 ^a	0.18 \pm 0.02 ^a	0.28 \pm 0.05 ^b	0.25 \pm 0.06 ^{ab}
C22:6n3	0.83 \pm 0.34	1.41 \pm 1.15	0.63 \pm 0.53	0.55 \pm 0.24
saturated	39.0 \pm 1.57	38.6 \pm 4.12	37.9 \pm 4.50	36.4 \pm 4.23
monoenoic	23.5 \pm 4.34 ^{ab}	28.7 \pm 1.37 ^b	21.7 \pm 3.66 ^{ab}	20.8 \pm 4.97 ^a
PUFA	34.4 \pm 5.69 ^{ab}	31.1 \pm 5.43 ^a	38.7 \pm 4.87 ^{ab}	41.2 \pm 4.55 ^b
n3	2.85 \pm 1.03 ^{ab}	3.90 \pm 0.82 ^b	2.20 \pm 0.44 ^a	2.95 \pm 1.55 ^a
n6	31.6 \pm 6.55 ^{ab}	27.2 \pm 6.04 ^a	36.5 \pm 5.15 ^{ab}	38.2 \pm 4.87 ^b
n6/n3	12.8 \pm 6.71	7.37 \pm 2.53	17.6 \pm 6.09	16.9 \pm 11.5
UI	102.9 \pm 20.0	108.7 \pm 5.44	118.0 \pm 9.68	122.5 \pm 9.39
ACL	17.5 \pm 0.14	17.5 \pm 0.17	17.5 \pm 0.13	17.5 \pm 0.10

UI: unsaturation index; ACL: average chain length; a,b: significant ($P < 0.05$) difference between groups




Figure 1. The extent of antioxidant capacity (GSH, GSHPx) and the concentration of the ultimate lipid peroxidation product, malondialdehyde, in the blood plasma of the rabbits (means \pm SD)




Figure 2. The plasma sphinganine to sphingosine ratio after four weeks of feeding FB₁, T-2 toxin, and T-2+FB₁ as compared to the control (means \pm SD)