



Árpád Baricz

Zeros of a cross-product of the Coulomb wave and Tricomi hypergeometric functions

Proceedings of the American Mathematical Society

DOI: 10.1090/proc/13331

Accepted Manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by AMS Production staff. Once the accepted manuscript has been copyedited, proofread, and finalized by AMS Production staff, the article will be published in electronic form as a “Recently Published Article” before being placed in an issue. That electronically published article will become the Version of Record.

This preliminary version is available to AMS members prior to publication of the Version of Record, and in limited cases it is also made accessible to everyone one year after the publication date of the Version of Record.

The Version of Record is accessible to everyone five years after publication in an issue.

ZEROS OF A CROSS-PRODUCT OF THE COULOMB WAVE AND TRICOMI HYPERGEOMETRIC FUNCTIONS

ÁRPÁD BARICZ

Dedicated to Prof. Péter T. Nagy on the occasion of his 70th birthday

ABSTRACT. Motivated by a problem on conditions for the existence of clines in genetics, we show that the positive zeros of a cross-product of the regular Coulomb wave function and the Tricomi hypergeometric function are increasing with respect to one of the parameters. In particular, this implies that the eigenvalues of a certain boundary value problem are increasing with the dimension.

1. INTRODUCTION

In his study about the existence of clines in genetics, Nagylaki [5] considered a partial differential equation in space and time satisfied by the gene frequency in a monoecious population distributed continuously over an arbitrary habitat. He showed that this partial differential equation reduces to the simplest multidimensional generalization of the classical Fisher-Haldane cline model, and investigated the efficacy of migration and selection in maintaining genetic variability at equilibrium in this model by deducing conditions for the existence of clines in various circumstances. The boundary value problem considered by Nagylaki reads as follows

$$(1.1) \quad \Delta p + \lambda^2 g(r)p = 0,$$

File: zeros-revised.tex, printed: 2016-06-9, 19.22

2010 *Mathematics Subject Classification.* 34B09, 34B30, 33C15, 33C10.

Key words and phrases. Coulomb wave function; Tricomi hypergeometric function; boundary value problem; zeros of a cross-product; eigenvalues; eigenfunctions; Bessel and modified Bessel functions; monotonicity of the zeros.

★The research of Á. Baricz was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. The author is very grateful to Prof. Mourad E.H. Ismail for suggesting the investigation of the zeros discussed in this paper of the cross-product of the regular Coulomb wave function and Tricomi hypergeometric function, and also for his kind hospitality during the author's visit at Department of Mathematics of the City University of Hong Kong in September 2011. The author is also very grateful to his friend and colleague Prof. Alexandru Kristály for the discussions and suggestions about the boundary value problems considered in this paper, as well as to the referee for his/her useful suggestions, which enhanced this paper.

where $p'(0) = 0$ and $p(\infty) < \infty$, Δp is the n -dimensional Laplacian, r is the distance from the origin of an n -dimensional vector x , and

$$g(r) = \begin{cases} 1, & r \in [0, 1], \\ -\alpha^2, & r > 1. \end{cases}$$

Nagylaki [5] conjectured that for each fixed $\alpha > 0$ the smallest positive eigenvalues of the above boundary value problem increase with the dimension. Motivated by Nagylaki's investigation, Ismail and Muldoon [4] considered the radial part of the boundary value problem (1.1), that is,

$$(1.2) \quad -(ry'(r))' + \nu^2 r^{-1} y(r) = \lambda^2 r g(r) y(r),$$

where $p'(0) = 0$, $p(\infty) < \infty$, $y(r) = r^\nu p(r)$ and $\nu = n/2 - 1$, and they showed that the positive eigenvalues of (1.2) are the positive zeros of fixed rank of the cross-product

$$J_{\nu+1}(r)K_\nu(\alpha r) - \alpha K_{\nu+1}(\alpha r)J_\nu(r),$$

where J_ν is the Bessel function of the first kind, while K_ν stands for the modified Bessel function of the second kind. Moreover, motivated by Askey's claim, Ismail and Muldoon [4] proved that the positive zeros of the cross-product

$$J_{\nu+\beta}(r)K_\nu(\alpha r) - \alpha^\beta K_{\nu+\beta}(\alpha r)J_\nu(r)$$

are increasing with respect to ν on $[-\beta/2, \infty)$, where $\beta \in (0, 1]$. Thus, it is clear that Nagylaki's conjecture follows from the case $\beta = 1$ of the above result. In [4], the authors actually stated more: they showed that the expression αr in the above affirmation can be changed to any strictly increasing differentiable function on $(0, \infty)$ and α^β can be replaced by an arbitrary positive constant. Motivated by the importance of the boundary value problem (1.1) and its radial part (1.2) in the existence of clines, and following the suggestion of Ismail, our aim is to show that Nagylaki's claim about the positive eigenvalues will be also true in a more general setting, we replace the Bessel function of the first kind by the regular Coulomb wave function, and the modified Bessel function of the second kind by the Tricomi hypergeometric function of the second kind. This is actually a generalization of the problem considered by Nagylaki. For more details on the special functions appearing in this paper we refer to [3, Chapter 6] and [6, Chapter 10].

2. THE EIGENVALUE PROBLEM RELATED TO COULOMB AND TRICOMI FUNCTIONS

In order to extend Nagylaki's problem we consider the boundary value problem

$$(2.1) \quad \Delta p + \varphi_\lambda(r)r^{-2}p = 0,$$

where $p'(0) = 0$ and $p(\infty) < \infty$,

$$\varphi_\lambda(r) = L(L-1) + \lambda^2 r^2 g(r) - 2\eta\lambda r h(r),$$

$L = (n - 1)/2$, η is a real parameter, and

$$h(r) = \begin{cases} 1, & r \in [0, 1], \\ \alpha, & r > 1. \end{cases}$$

It can be shown that the radial part of the above boundary value problem (2.1) is

$$(2.2) \quad r^2 y''(r) + (\lambda^2 r^2 g(r) - 2\eta\lambda r h(r) - L(L + 1)) y(r) = 0,$$

where $p'(0) = 0$, $p(\infty) < \infty$ and $y(r) = r^L p(r)$. If we suppose that $r \in (0, 1]$, then we obtain

$$r^2 y''(r) + (\lambda^2 r^2 - 2\eta\lambda r - L(L + 1)) y(r) = 0.$$

By using the change of variable $u = \lambda r$ (and taking $y(r) = z(u)$), this equation becomes the Coulomb wave equation

$$u^2 z''(u) + (u^2 - 2\eta u - L(L + 1)) z(u) = 0.$$

Moreover, when $r > 1$, (2.2) becomes

$$r^2 y''(r) - (\alpha^2 \lambda^2 r^2 + 2\eta\alpha\lambda r - L(L + 1)) y(r) = 0,$$

which after the change of variable $v = \alpha\lambda r$ (and taking $y(r) = q(v)$) becomes a transformation of the Kummer confluent hypergeometric differential equation

$$v^2 q''(v) - (v^2 + 2\eta v + L(L + 1)) q(v) = 0.$$

Thus, when $r \in (0, 1]$ the differential equation (2.2) has as a particular solution the regular Coulomb wave function

$$y(r) = F_L(\eta, u),$$

while for $r > 1$ the equation (2.2) has the particular solution a transformation of the Tricomi hypergeometric function

$$y(r) = v^{L+1} e^{-v} \psi(L + \eta + 1, 2L + 2, 2v).$$

When $\eta = 0$ the above particular solutions reduce to

$$y(r) = \sqrt{\frac{\pi}{2u}} J_{L+\frac{1}{2}}(u) \quad \text{and} \quad y(r) = 2^{-L} \sqrt{\frac{2v}{\pi}} K_{L+\frac{1}{2}}(v),$$

which show that the boundary value problem (2.1) is a natural extension of (1.1), while (2.2) is a natural extension of (1.2).

Now, we are ready to state the main result of this paper.

Theorem 1. **a.** *The boundary value problem (2.2) has for its eigenvalues the zeros of the cross-product of the regular Coulomb wave and Tricomi hypergeometric functions*

$$F_L'(\eta, r) Q_L(\eta, \alpha r) - \alpha Q_L'(\eta, \alpha r) F_L(\eta, r)$$

and corresponding eigenfunctions

$$r \mapsto \Theta_L(\eta, r) = \begin{cases} Q_L(\eta, \alpha\lambda) \cdot F_L(\eta, \lambda r), & r \in (0, 1], \\ F_L(\eta, \lambda) \cdot Q_L(\eta, \alpha\lambda r), & r > 1, \end{cases}$$

where

$$Q_L(\eta, r) = r^{L+1} e^{-r} \psi(L + \eta + 1, 2L + 2, 2r).$$

- b.** For fixed $\alpha > 0$, $\eta \in \mathbb{R}$ such that $L + \eta > 0$, and $L > -3/2$, $L \neq -1$ if $\eta \neq 0$ and $L > -3/2$ if $\eta = 0$, the equation

$$(2.3) \quad F'_L(\eta, r)/F_L(\eta, r) = \alpha Q'_L(\eta, \alpha r)/Q_L(\eta, \alpha r)$$

has infinitely many positive roots, which we denote in increasing order by $\lambda_{L,\eta,\alpha,n}$, $n \in \mathbb{N}$. These zeros satisfy

$$x_{L,\eta,n-1} < \lambda_{L,\eta,\alpha,n} < x_{L,\eta,n},$$

$n \in \{2, 3, \dots\}$, where $x_{L,\eta,n}$ is the n th positive zero of the Coulomb wave function $\rho \mapsto F_L(\eta, \rho)$. Moreover, if $\alpha > 0$, $\eta \in \mathbb{R}$ and $L > -1/2$, then we have $\lambda_{L,\eta,\alpha,1} < x_{L,\eta,1}$.

- c.** For fixed $\alpha > 0$, $\eta \geq 0$ and $n \in \mathbb{N}$ the zeros $\lambda_{L,\eta,\alpha,n}$ increase with L on $[0, \infty)$.

We note that since the boundary value problem (2.1) is an extension of (1.1), while (2.2) is an extension of (1.2), if we take $\eta = 0$ in the above theorem, we obtain some of the main results of [4] for the case $\beta = 1$.

Proof of Theorem 1. **a.** Subject to the stated boundary condition the differential equation in (2.2) has solution

$$y(r) = \begin{cases} A \cdot F_L(\eta, \lambda r), & r \in (0, 1], \\ B \cdot Q_L(\eta, \alpha \lambda r), & r > 1. \end{cases}$$

Since y and y' are to be continuous at $r = 1$ we must have

$$A \cdot F_L(\eta, \lambda) = B \cdot Q_L(\eta, \alpha \lambda)$$

and

$$A \cdot \lambda F'_L(\eta, \lambda) = B \cdot \alpha \lambda Q'_L(\eta, \alpha \lambda),$$

and there will be a nontrivial solution of this system if and only if

$$F'_L(\eta, \lambda) Q_L(\eta, \alpha \lambda) = \alpha Q'_L(\eta, \alpha \lambda) F_L(\eta, \lambda).$$

Hence we may take $A = Q_L(\eta, \alpha \lambda)$ and $B = F_L(\eta, \lambda)$. Thus, indeed the boundary value problem (2.2) has for its eigenvalues the zeros of the cross-product of the regular Coulomb wave and Tricomi hypergeometric functions, that is,

$$F'_L(\eta, r) Q_L(\eta, \alpha r) - \alpha Q'_L(\eta, \alpha r) F_L(\eta, r),$$

and corresponding eigenfunctions $r \mapsto \Theta_L(\eta, r)$.

- b.** The equation (2.3) is equivalent to

$$(2.4) \quad \frac{F'_L(\eta, r)}{F_L(\eta, r)} - \frac{L+1}{r} = \alpha + \frac{2\alpha \psi'(L + \eta + 1, 2l + 2, 2\alpha r)}{\psi(L + \eta + 1, 2L + 2, 2\alpha r)}.$$

Now, we shall use the Mittag-Leffler expansion of regular Coulomb wave function (obtained from its infinite product representation, see [1, 7, 8] for

more details). Namely, since for $L > -3/2$, $L \neq -1$ if $\eta \neq 0$ and $L > -3/2$ if $\eta = 0$ we have [1, Lemma 1]

$$\frac{F'_L(\eta, r)}{F_L(\eta, r)} - \frac{L+1}{r} = \frac{\eta}{L+1} - \sum_{n \geq 1} \left(\frac{r}{x_{L,\eta,n}(x_{L,\eta,n} - r)} + \frac{r}{y_{L,\eta,n}(y_{L,\eta,n} - r)} \right),$$

the left-hand side of the equation (2.4) is decreasing on $(0, x_{L,\eta,1})$ and also on each interval $(x_{L,\eta,n}, x_{L,\eta,n+1})$, $n \in \mathbb{N}$. Here $y_{L,\eta,n}$ stands for the n th negative zero of the regular Coulomb wave function $r \mapsto F_L(\eta, r)$. When $r \searrow 0$ the left-hand side of (2.4) tends to $\eta/(L+1)$, when $r \nearrow x_{L,\eta,n}$, $n \in \mathbb{N}$ it tends to $-\infty$ and when $r \searrow x_{L,\eta,n}$, $n \in \mathbb{N}$ it tends to $+\infty$. On the other hand, according to [2, Remark 3] we know that for $a > 1$ and $c \in \mathbb{R}$ the function $r \mapsto \psi'(a, c, r)/\psi(a, c, r)$ is increasing on $(0, \infty)$. Note that this monotonicity result is in fact equivalent to a Turán-type inequality for Tricomi hypergeometric functions. Moreover, by using the recurrence relation

$$\psi'(a, c, r) = -a\psi(a+1, c+1, r),$$

the fact that $\psi(a, c, r)$ is positive for $a, c, r \in \mathbb{R}$, and the asymptotic expansion

$$\psi(a, c, r) = r^{-a}(1 + \mathcal{O}(r^{-1})) \quad \text{as } r \rightarrow \infty,$$

it follows that $r \mapsto \psi'(a, c, r)/\psi(a, c, r)$ maps $(0, \infty)$ into $(-\infty, 0)$. Thus, the right-hand side of (2.4) is increasing on $(0, \infty)$ for $\alpha > 0$ and $L + \eta > 0$ and maps the interval $(0, \infty)$ into $(-\infty, \alpha)$. These show that the equation (2.4) has infinitely many positive roots, and starting from the second positive root they are certainly located between the positive zeros of the regular Coulomb wave function. Now, by using the asymptotic relation

$$\psi(a, c, r) \sim \Gamma(c-1)r^{1-c}/\Gamma(a) \quad \text{as } r \rightarrow 0 \quad \text{and } c > 1,$$

it follows that $\psi'(a, c, r)/\psi(a, c, r) \sim (1-c)/r$ as $r \rightarrow 0$ and $c > 1$, and thus the right-hand side of (2.4) tends to $-\infty$ as $r \rightarrow 0$ and $L > -1/2$. This shows that if $\alpha > 0$, $\eta \in \mathbb{R}$ and $L > -1/2$, then we have $\lambda_{L,\eta,\alpha,1} < x_{L,\eta,1}$.

c. We shall follow the approach considered in [4], namely, the Sturmian-type arguments and the approach of the Hellman-Feynman theorem of quantum chemistry. Since $r \mapsto \Theta_L(\eta, r)$ are eigenfunctions of the boundary value problem (2.2), we have

$$-\Theta''_L(\eta, r)\Theta_L(\eta, r) + 2\eta\lambda\frac{1}{r}h(r)\Theta_L^2(\eta, r) + L(L+1)\frac{1}{r^2}\Theta_L^2(\eta, r) = \lambda^2g(r)\Theta_L^2(\eta, r).$$

Integrating from zero to infinity we get

$$\begin{aligned} & \int_0^\infty \left(\lambda^2g(r) - 2\eta\lambda\frac{1}{r}h(r) \right) \Theta_L^2(\eta, r) dr \\ &= L(L+1) \int_0^\infty \frac{1}{r^2} \Theta_L^2(\eta, r) dr + \int_0^\infty (\Theta'_L(\eta, r))^2 dr, \end{aligned}$$

where we used integration by parts in the last integral. Since the right-hand side of the above relation is positive for $L \geq 0$, it follows that

$$(2.5) \quad \lambda^2 \int_0^\infty g(r) \Theta_L^2(\eta, r) dr \geq \lambda \int_0^\infty 2\eta \frac{1}{r} h(r) \Theta_L^2(\eta, r) dr.$$

Now, writing λ_L instead of $\lambda_{L, \eta, \alpha, n}$, multiplying the equations

$$-\Theta_L''(\eta, r) + 2\eta \lambda_L \frac{1}{r} h(r) \Theta_L(\eta, r) + L(L+1) \frac{1}{r^2} \Theta_L(\eta, r) = \lambda_L^2 g(r) \Theta_L(\eta, r),$$

$$-\Theta_M''(\eta, r) + 2\eta \lambda_M \frac{1}{r} h(r) \Theta_M(\eta, r) + M(M+1) \frac{1}{r^2} \Theta_M(\eta, r) = \lambda_M^2 g(r) \Theta_M(\eta, r)$$

by $\Theta_M(\eta, r)$, $\Theta_L(\eta, r)$ respectively, subtracting and integrating between 0 and ∞ we get

$$\begin{aligned} & (\Theta_M'(\eta, r) \Theta_L(\eta, r) - \Theta_L'(\eta, r) \Theta_M(\eta, r)) \Big|_0^\infty \\ & + (\lambda_L - \lambda_M) \int_0^\infty 2\eta \frac{1}{r} h(r) \Theta_L(\eta, r) \Theta_M(\eta, r) dr \\ & + [L(L+1) - M(M+1)] \int_0^\infty \frac{1}{r^2} \Theta_L(\eta, r) \Theta_M(\eta, r) dr \\ & = (\lambda_L^2 - \lambda_M^2) \int_0^\infty g(r) \Theta_L(\eta, r) \Theta_M(\eta, r) dr. \end{aligned}$$

Note that the integrated term vanishes at 0 and ∞ for $L, M > 0$, and consequently dividing both parts of the above equation by $L - M$ and taking the limit $M \rightarrow L$ we obtain

$$\begin{aligned} & \frac{d\lambda_L}{dL} \int_0^\infty 2\eta \frac{1}{r} h(r) \Theta_L^2(\eta, r) dr + (2L+1) \int_0^\infty \frac{1}{r^2} \Theta_L^2(\eta, r) dr \\ & = \frac{d\lambda_L^2}{dL} \int_0^\infty g(r) \Theta_L^2(\eta, r) dr \end{aligned}$$

or equivalently

$$\begin{aligned} & \frac{d\lambda_L}{dL} \int_0^\infty 2\eta \frac{1}{r} h(r) \Theta_L^2(\eta, r) dr + (2L+1) \int_0^\infty \frac{1}{r^2} \Theta_L^2(\eta, r) dr \\ & = 2\lambda_L \frac{d\lambda_L}{dL} \int_0^\infty g(r) \Theta_L^2(\eta, r) dr. \end{aligned}$$

This implies that

$$\begin{aligned} & \frac{d\lambda_L}{dL} \left(2\lambda_L \int_0^\infty g(r) \Theta_L^2(\eta, r) dr - \int_0^\infty 2\eta \frac{1}{r} h(r) \Theta_L^2(\eta, r) dr \right) \\ & = (2L+1) \int_0^\infty \frac{1}{r^2} \Theta_L^2(\eta, r) dr > 0, \end{aligned}$$

which in view of (2.5) and the fact that λ_L is positive according to part **b**, yields $d\lambda_L/dL > 0$, so for fixed $\alpha > 0$ and $\eta \geq 0$ the zero λ_L is increasing with respect to L on $[0, \infty)$. \square

REFERENCES

- [1] Á. BARICZ, Turán type inequalities for regular Coulomb wave functions, *J. Math. Anal. Appl.* 430(1) (2015) 166–180.
- [2] Á. BARICZ, M.E.H. ISMAIL, Turán type inequalities for Tricomi confluent hypergeometric functions, *Constr. Approx.* 37(2) (2013) 195–221.
- [3] A. ERDÉLYI ET AL., *Higher Transcendental Functions*, vol. 1, McGraw-Hill, New York, 1955.
- [4] M.E.H. ISMAIL, M.E. MULDOON, Monotonicity of the zeros of a cross-product of Bessel functions, *SIAM J. Math. Anal.* 9(4) (1978) 759–767.
- [5] T. NAGYLAKI, Conditions for the existence of clines, *Genetics* 80 (1975) 595–615.
- [6] F.W.J. OLVER, D.W. LOZIER, R.F. BOISVERT, C.W. CLARK (EDS.), *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press, Cambridge, 2010.
- [7] F. ŠTAMPACH, P. ŠŤOVÍČEK, Orthogonal polynomials associated with Coulomb wave functions, *J. Math. Anal. Appl.* 419(1) (2014) 231–254.
- [8] J. WIMP, Some explicit Padé approximants for the function Φ'/Φ and a related quadrature formula involving Bessel functions, *SIAM J. Math. Anal.* 16(4) (1985) 887–895.

INSTITUTE OF APPLIED MATHEMATICS, ÓBUDA UNIVERSITY, BUDAPEST, HUNGARY

DEPARTMENT OF ECONOMICS, BABEȘ-BOLYAI UNIVERSITY, CLUJ-NAPOCA, ROMANIA
E-mail address: bariczocsi@yahoo.com