

REVIEW

Septic acute kidney injury: molecular mechanisms and the importance of stratification and targeting therapy

Eric D Morrell¹, John A Kellum^{1,2,3,4}, Núria M Pastor-Soler^{1,2,5*} and Kenneth R Hallows^{1,2,5}

Abstract

The most common cause of acute kidney injury (AKI) in hospitalized patients is sepsis. However, the molecular pathways and mechanisms that mediate septic AKI are not well defined. Experiments performed over the past 20 years suggest that there are profound differences in the pathogenesis between septic and ischemic AKI. Septic AKI often occurs independently of hypoperfusion, and is mediated by a concomitant pro- and anti-inflammatory state that is activated in response to various pathogen-associated molecular patterns, such as endotoxin, as well as damage-associated molecular patterns. These molecular patterns are recognized by Toll-like receptors (TLRs) found in the kidney, and effectuate downstream inflammatory pathways. Additionally, apoptosis has been proposed to play a role in the pathogenesis of septic AKI. However, targeted therapies designed to mitigate the above aspects of the inflammatory state, TLR-related pathways, and apoptosis have failed to show significant clinical benefit. This failure is likely due to the protean nature of septic AKI, whereby different patients present at different points along the immunologic spectrum. While one patient may benefit from targeted therapy at one end of the spectrum, another patient at the other end may be harmed by the same therapy. We propose that a next important step in septic AKI research will be to identify where patients lie on the immunologic spectrum in order to appropriately target therapies at the inflammatory cascade, TLRs, and possibly apoptosis.

Introduction

Acute kidney injury (AKI) is a very common and especially formidable clinical problem in the ICU, where mortality rates approach 25% and soar to 50 to 60% when severe enough to require renal replacement therapy [1]. These statistics have not significantly improved over the past 50 years, though patients today are generally older and have more comorbidities [1]. The most common cause of AKI in hospitalized patients is sepsis [2], and AKI occurs with regularity even in non-severe sepsis where clinically evident hemodynamic changes are not readily apparent [3]. The combination of sepsis and AKI portends a dire clinical situation that is associated with a hospital mortality rate as high as 70% [4]. However, the basic pathophysiologic mechanisms undergirding the

association between sepsis and the clinical manifestations of AKI are not completely understood.

In this review, we will first summarize the findings of published human studies that support the concept that the pathogenesis of septic AKI in humans should not be exclusively viewed in the context of distributive shock-associated ischemia, but rather also within the context of a dysregulated and ill-defined inflammatory response to septic stimuli. We then discuss why previous attempts at modulating the inflammatory response in septic AKI may have failed, and highlight more recent trials and therapies aimed at targeted treatment based on where patients lie along the immunologic spectrum. We also discuss the role Toll-like receptors (TLRs) have been shown to play in endotoxemic models of AKI, and again focus on why agents used to block TLR-related pathways may have failed to show significant clinical benefit. Finally, we review the data supporting and refuting a central role for apoptosis in septic AKI, and argue that much of the clinical

* Correspondence: pastorn@pitt.edu

¹Renal-Electrolyte Division, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA

²The Center for Critical Care Nephrology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA

Full list of author information is available at the end of the article

manifestations in septic AKI occur prior to significant cell death secondary to apoptosis or necrosis.

Septic acute kidney injury: is it hyperemic kidney injury?

The morphologic changes that are seen in classic ischemic and toxic acute tubular necrosis are often lacking [5,6] or are microscopically distinct in septic AKI [7,8]. While the use of renal biopsy in the management of septic AKI is exceedingly rare in many western nations, most human studies have found no consistent histopathologic changes in septic AKI, and acute tubular necrosis is infrequently observed [5,6]. In a systematic review of six studies describing histopathologic changes seen in septic AKI, Langenberg and colleagues [6] found that tubular epithelial cell necrosis was seen in only 22% of human samples taken after septic injury. A more recent study by Takasu and colleagues [8] examined post-mortem kidneys taken from 67 patients who died with sepsis, and while they found focal tubular injury in 78% of septic kidneys, the majority of tubular cells were normal.

Perhaps an even greater distinction between non-septic AKI and septic AKI might be made when examining the role of overall vascular hemodynamics. Traditionally, septic AKI has been thought to be secondary to a demand and perfusion mismatch, whereby renal blood flow was diminished in the midst of increased metabolic demand [9,10]. Furthering this idea are the FINNAKI and SEPSIS-PAM studies, two recently published trials that suggest septic patients with target mean arterial pressure higher than 65 mmHg might have less progression to AKI [11] and less need for renal-replacement therapy if they suffer from chronic hypertension [12].

Nevertheless, the data directly measuring overall kidney blood flow during sepsis suggest that flow is increased in the septic physiologic state. Despite a plethora of conflicting animal studies - some of which show overall increased kidney blood flow while others show overall decreased kidney blood flow [10,13] - Langenberg and associates [13] found only three human studies that have directly measured renal blood flow in septic AKI by invasive hemodynamic measurements. All three studies showed that septic AKI occurs in the midst of renal arterial vasodilation and preserved overall renal blood flow [14-16]. In another study with a small cohort of 10 patients with sepsis, renal blood flow was either preserved or increased [17]. Bellomo and associates [18] have thus proposed a new concept for understanding septic AKI hemodynamics, so-called hyperemic AKI.

Despite the above human studies that suggest an increase in overall renal blood flow in septic AKI, the complex interplay between regional microcirculation and hemodynamics may play a much more important role in overall kidney function during sepsis. As Langenberg and colleagues [13] and Molitoris [19] point out,

glomerular filtration rate (GFR) and cellular perfusion can decrease even if overall renal blood flow is increased due to disproportionate vascular resistance between the afferent and efferent arterioles, regional microvascular flow rates, or renal venous congestion. There are limited human data regarding microcirculatory function in septic AKI. However, using implanted laser Doppler flow probes, Di Giandomasso and colleagues [20] showed that, in sheep hyperdynamic septic models using *Escherichia coli* infusion as a source of infection, regional redistribution of blood flow from the medulla to the cortex was absent during sepsis. Local inhibition of endothelial nitric oxide synthase by liposaccharide (LPS) may play a role in the decreased GFR observed in endotoxemic models of AKI, at least in rats [21]. In a hyperdynamic endotoxemic pig model, Cohen and associates [22] likewise observed a significant increase in overall kidney blood flow after LPS administration; however, almost all of the increased blood flow was shunted to the medulla, with no increase in cortical blood flow. In a sepsis pig model using peritonitis as a source of infection, Chvojka and colleagues [23] observed that while overall renal blood flow increased in septic pigs with AKI, renal perfusion pressure significantly decreased as a result of increased renal venous pressure. In parallel to this, there was a significant decrease in renal cortex microvascular perfusion as measured by laser Doppler blood flow, as well as increased renal venous acidosis and lactate concentrations [23]. These septic physiologic observations related to regional kidney blood flow have been found in humans as well. In a retrospective observational study, it was shown that elevated central venous pressure was associated with progression to AKI in septic patients (odds ratio = 1.22, 95% confidence interval (CI) 1.08 to 1.39), suggesting that venous congestion may contribute to septic AKI in humans [24].

An in-depth discussion about renal microcirculation in septic AKI is beyond the scope of this review. As mentioned above, there is a dearth of human data on this important topic and most of the animal data are conflicting and difficult to interpret, given the different models used and measurement modalities employed. While it is clear from above that ischemic and microvascular dysfunction may contribute to septic AKI, septic AKI can occur independently of overall renal hypoperfusion or ischemia [25]. Indeed, AKI occurs in up to one-third of non-critically ill patients with pneumonia, of which less than 10% received any vasopressor therapy [3]. Additionally, there have been many studies published over the past decade showing that plasma and urinary biomarkers are often distinct in septic and non-septic AKI (reviewed in [26]). This suggests a fundamental mechanistic distinction that may involve primarily a dysregulated immune response and possibly apoptosis as opposed to ischemic factors related to distributive shock [25].

Sepsis-induced inflammation and its effects on the kidney

For many years, organ damage and other deleterious sequelae from sepsis were thought to result from a hyper-inflammatory 'cytokine storm,' triggered in response to either an infectious pathogen or major physiologic insult [27]. Thus, a significant amount of sepsis research in the 1990s focused on identifying specific cytokines that were associated with the stereotypical sepsis syndrome. However, while cytokines such as TNF- α and IL-1 β were found in very high concentrations in septic patients, large clinical trials testing specific cytokine inhibition failed to show any clinical benefit [28,29].

As a consequence of these failed trials, new ideas regarding the pathogenesis and pathophysiology of sepsis emerged [30]. Utilizing pathologic data showing impaired immune function in some septic patients, Hotchkiss and Remick have both hypothesized that the host's dysregulated and ill-defined inflammatory response to stimuli is associated with both pro- and anti-inflammatory states [31-33]. Many septic patients are chronically ill, and their development of sepsis is not necessarily associated with a hyper-inflammatory response [31]. One current goal in sepsis research is to identify where a septic patient lies on the immunologic spectrum, and identify which patients will benefit from specific therapies.

Indeed, the early trials focusing on inhibiting TNF- α or IL-1 β did not risk-stratify their patients in terms of their individual inflammatory status or genetic predilections towards certain types of phenotypic responses [28,29]. It seems likely that the next step in sepsis and septic AKI treatment will be targeted therapy based on stratification of patients that have a higher likelihood of responding to various treatments. Panacek and colleagues first attempted this strategy with a randomized controlled trial to study afelimomab, an anti-TNF F(ab')₂ monoclonal antibody fragment, in septic patients with and without elevated serum levels of IL-6 [34]. In this study, for those patients who had serum IL-6 levels >1,000 pg/mL, they demonstrated a 4% absolute risk reduction in mortality at 28 days after treatment in patients treated with the agent than in those not treated. Although rates of kidney failure were not reported in this trial, there was a statistically significant improvement in the overall Sequential Organ Failure Assessment (SOFA) score starting at 48 hours after drug administration.

As an alternative to targeted treatment, some researchers have proposed implementing treatment strategies that are able to broadly regulate the overall inflammatory response in sepsis by using extra-corporeal blood purification. Multiple *in vitro* and rat *in vivo* studies have demonstrated that the broad reduction of inflammatory cytokines by hemoabsorption can protect against renal and liver injury [35], improve bacterial clearance [36] and mean arterial pressure [37], decrease apoptosis and tubular cell

damage [38], and improve overall survival [35,37]. However, its efficacy in humans is still unknown. The Early Use of Polymyxin B Hemoperfusion in Abdominal Sepsis (EUPHAS) trial showed improved hemodynamics, organ function, and 28-day mortality in patients suffering from severe sepsis or septic shock from intra-abdominal Gram-negative infections who were treated with two sessions of polymyxin B hemoperfusion compared with controls [39]. Renal SOFA scores improved in the treatment group (-0.3, 95% CI -0.7 to 0.1 versus 0.6, 95% CI 0.1 to 1.1) versus control, as did overall cardiovascular function. However, there were several methodological flaws in this small trial, including a failure to measure endotoxin levels in the treatment and control groups and the use of 28-day mortality as a secondary endpoint. While these findings suggest the need for larger clinical trials to be performed before widespread implementation can be recommended, these preliminary results appear promising.

Although the *in vivo* animal studies cited above have shown that broad regulation of cytokine levels is possible with the use of cytokine-reducing resins, modulation of cytokine levels by continuous hemodialysis in humans has not been clearly demonstrated. Despite the very short half-life of many inflammatory cytokines, small initial studies suggested that high-volume hemofiltration might be able to modulate overall cytokine profiles in humans [40,41]. However, larger randomized controlled trials failed to show that continuous hemofiltration modulates cytokine levels [42], nor has any clinical benefit been demonstrated with the use of high-volume hemofiltration in the treatment of sepsis [42-44]. While the large VA/NIH Acute Renal Failure Trial Network study showed no mortality difference in the septic subgroup of patients who underwent more intense renal replacement, subsequent analysis has shown higher rates of non-recovery of kidney function and increased mortality in patients with higher circulating concentrations of pro- and anti-inflammatory cytokines and apoptosis markers [45]. This result again suggests that targeting certain subgroups along the immunologic spectrum might be beneficial not only in targeted cytokine therapy, but also in the broad modulation of inflammatory cytokines via hemofiltration and absorption.

Finally, the use of agents to augment immune function in patients suffering from sepsis-associated immunosuppression has shown clinical promise. As mentioned above, sepsis represents a complex inflammatory state with both pro- and anti-inflammatory characteristics. Thus, identifying and treating septic patients who are on the more immunosuppressed spectrum of sepsis with immune augmentation may improve outcomes. Meisel and associates [46] administered granulocyte-macrophage colony-stimulating factors (GM-CSFs) to septic patients who had decreased monocyte HLA-DR expression and found that monocyte immunocompetence could be restored in this

subgroup of patients. While there was no change in the clinical outcomes that were measured (ventilator days and hospital length of stay), their data showed that biomarker-guided GM-CSF therapy in sepsis is safe and could hold promise in larger randomized controlled trials.

Toll-like receptor-4 and the effects of sepsis on renal epithelial cells

Initial molecular research on sepsis suggested that the generalized inflammatory state described above is initiated when the innate immune system recognizes highly conserved pathogen-associated molecular patterns (PAMPs), such as LPS [47]. However, in the late 1990s Polly Matzinger [48] proposed the 'danger model', which argues that the immune system is more concerned with 'danger signals' than with foreign signals. Subsequent research has found that there are numerous 'damage-associated molecular patterns' (DAMPs) that can elicit immune responses similar to PAMPs [49]. The extent to which both PAMPs (a subgroup of DAMPs) and more broadly DAMPs activate the immune system and contribute to sepsis and septic AKI is not fully understood, but recent research has provided a significant amount of insight into the process.

TLRs have provided a link between host and pathogen [50,51], as well as host and DAMPs [52]. In the 1990s, TLR-4 was found to be the main receptor that binds LPS, leading to the release of TNF- α and IL-1 and activation of numerous downstream intracellular signaling pathways via the NF- κ B transcription factor [50,53] (Figure 1). Additionally, in sterile models of inflammation, TLR-4 has been shown to respond to DAMPs, such as high mobility group-B1 [54-56]. Indeed, the capacity of TLRs to respond to PAMPs as well as other DAMPs provides a potential link between systemic inflammation and kidney damage that is independent of ischemia and hypotension. In the kidney, TLR-4 is constitutively expressed in mouse primary renal tubular epithelial cells [57] and in mouse cells harvested from *in vivo* experiments [58], in renal tubular cells harvested from rats [59,60], and in human kidney tubular epithelial cells taken from surgical patients [61]. TLR-4 expression in the glomeruli is controversial [59,60], but its expression in tubular epithelial cells has been shown to modulate septic AKI and inflammation (Figure 1) in multiple models of injury [60,62,63]. The final downstream mechanisms that produce AKI after TLR-4 activation are complex and incompletely understood. Of note, recent work suggests that one important mechanism underpinning septic AKI is that this inflammatory pathway induces renal tubular transport dysfunction, which enhances NaCl delivery to the macula densa and increases tubuloglomerular feedback, thus decreasing GFR [64].


NF- κ B is a highly expressed transcription factor that is often activated in response to cellular stresses and induces the synthesis and release of TNF- α , IL-1, IL-6, and IL-8 [65]. NF- κ B-mediated processes occur in isolated primary rat proximal tubular epithelial cells [66], but the overall contribution of inflammatory cytokine levels in the kidney from local versus systemic production of those cytokines is unknown. Experiments performed on TLR-4 knockout mice transplanted with TLR-4 expressing kidneys suggest systemic production may be more important [62]. Direct inhibition of NF- κ B with agents such as methyl-2-acetamidoacrylate (an ethyl pyruvate analogue) [67], ethyl pyruvate [68], and nicotinic acetylcholine [69] can lower cytokine levels in mice (TNF- α , IL-6, IFN- γ , and IL-10) and have been associated with improved clinical characteristics in mice suffering from septic AKI [67-69].

However, a recent randomized controlled trial examining the effect of an MD2-TLR-4 antagonist failed to show any mortality benefit in human patients suffering from severe sepsis [70]. In this study, none of the patients enrolled were stratified in terms of what type of infection they were suffering from (polymicrobial, Gram-positive, or Gram-negative). Moreover, levels of endotoxin were not measured, which leaves open the possibility that some patients may have benefited depending on where they stood on the immunologic spectrum. Nonetheless, this trial possibly highlights some of the pitfalls of associating LPS endotoxemia with human sepsis that have manifested themselves over the past 20 years in sepsis research.

Indeed, multiple studies have shown that in polymicrobial and cecal ligation and puncture models the mechanism of damage induced by sepsis may be very different than in LPS-induced models [71], might be independent of TLRs altogether [72-75], and instead could be mediated by direct MyD88 regulation (Figure 1) [72,75]. Interestingly, the presence of a specific TLR-4 mutation (D299G) in humans, which impairs LPS signaling, has not been shown to influence TNF- α , IL-6, and IL-10 levels in sepsis [73]. Nor has this particular allele been associated with any change in outcome or development of sepsis in patients undergoing major visceral surgery [73] or in patients with meningococcal disease [76]. The effect of this specific mutation has not been examined in kidneys directly, and its study may provide significant insight into the mechanisms of septic AKI in humans. Of note, the use of various mouse models to study human inflammation and sepsis has recently been questioned due to the significant differences in the two species' genomic responses to stress [77].


Does apoptosis have a role in septic acute kidney injury?

The traditional model of apoptosis in kidney and other cells is similar [78]. Processes involving the mitochondria



and cell respiration are labeled 'intrinsic,' while binding of cell surface proteins leading to subsequent intracellular signaling events are termed 'extrinsic' processes (Figure 2) [78]. However, the mechanisms whereby cells in the septic kidney succumb (or do not succumb) to apoptosis are still under investigation. One important route may be via the tumor necrosis factor receptor-1 (TNFR-1; Figure 2). TNFR-1 knockout mice were resistant to LPS-induced kidney failure while they also manifested much less apoptosis and neutrophil accumulation, suggesting a direct role for both necrosis and apoptosis in septic AKI [79]. Downstream of TNFR-1, Jo and colleagues [80] have shown that apoptosis in LPS, TNF-α, and IL-1 exposed HK-2 cells is

likely dependent in part on Fas and caspase expression. Caspase inhibition has also been shown to prevent apoptotic proximal, distal, and peri-tubular cell death in LPS models of septic AKI [81]. Cantaluppi and associates [38] have demonstrated a role for the Fas and caspase-mediated pathway in septic AKI as well. Using primary human proximal tubule epithelial cells exposed to septic human plasma, they showed that: 1) apoptosis as detected through DNA TUNEL assays was significantly increased in cells exposed to septic plasma; 2) levels of Fas and caspase-3, -8, and -9 were all significantly increased in response to septic plasma; and 3) levels of those effector signals and apoptosis can be decreased by removal of



inflammatory mediators via a resin adsorption method to filter septic plasma [38]. Indeed, initiation of apoptosis was observed in cultured kidney tubular cells exposed to plasma taken from patients suffering from severe burns and septic AKI [82].

However, in most of the *in vitro* and *in vivo* studies examining the presence or role of apoptosis in septic AKI, there were major methodological shortcomings. LPS was used as a model for sepsis in most of the studies cited above [79-81], as well as many experiments evaluating kidney tubular apoptosis in primary mouse distal convoluted tubule cells [83] and mouse MCT proximal tubule cells [84]. As mentioned in the previous section, LPS endotoxemia is not a model of sepsis. Perhaps even more confounding to the argument that apoptosis is a major factor in septic AKI is the lack of histological evidence in patients with septic AKI. Histologic sections of septic human kidneys generally demonstrate very little necrosis or cell damage [5-8]. Moreover, a significant amount of data taken from 'warm' autopsies [5,8] has also

failed to show substantial apoptosis, at least as measured by conventional light microscopy [5], electron microscopy [8], and DNA agarose gel electrophoresis for apoptotic laddering [5]. In the seminal septic autopsy study performed by Hotchkiss and colleagues [5], they failed to observe any evidence of kidney apoptosis despite the fact that 65% of their samples were taken from patients with clinically manifest sepsis-associated kidney dysfunction. While apoptosis likely plays a major role in lymphocytes and other cells within the human body during sepsis [30], its role in septic AKI is unclear. Further study using more physiologically relevant models of sepsis (as opposed to LPS) are needed to help reconcile much of the conflicting data that currently exist.

Perhaps one of the reasons for the discrepancy between histologic data and functional performance in septic AKI is that most of the insults occur prior to significant cell death from apoptosis and necrosis. Indeed, apoptosis and necrosis are end-stage manifestations of cellular insults, and significant impairment of cellular function can occur

before the histologic and genomic changes that are measured in most apoptosis and necrosis assays. Despite a measured decrease in kidney perfusion in their septic mouse model, Tran and associates found no elevation in hypoxia-inducible factor-1 α , a transcription factor that is significantly upregulated [85] in hypoxic and ischemic rat kidneys [86]. They hypothesized that this was due to significantly reduced oxygen consumption by kidney proximal tubule cells during sepsis [86]. Further study showed that this process was likely mediated by PPAR γ coactivator-1 α (PGC-1 α), a major regulator of mitochondrial function. PGC-1 α was significantly downregulated in septic rats exposed to LPS and TNF- α , whereas increased levels of PGC-1 α were associated with restored oxygen consumption and improved kidney function [86]. While increased expression of hypoxia-inducible factor-1 α was not seen in mice exposed to LPS [86], sepsis has been associated with significant intra-renal oxidative and nitrosative stress that is mediated by inflammatory cytokines rather than local hypoxia or ischemia [87,88]. The idea of 'cellular stunning', whereby a cell that is living within a septic environment is trying to defend itself from death by decreasing its metabolic activity, has been proposed as a mechanism not only in sepsis by Hotchkiss and Karl [30], but also in ischemic models in the heart [89].

Conclusion

While the relative contributions that ischemia, TLR-4-mediated pathways, apoptosis, and other subcellular processes have towards overall organ dysfunction are currently unclear, the abundance of research in the area has shed a significant amount of light on the basic mechanisms that drive septic AKI. Perhaps the most important finding is that therapeutic approaches targeted at a single mediator for all patients may not succeed given the protean nature of the disease. Indeed, targeted inhibition of cytokines such as TNF- α and IL-1 as well as TLR antagonists have failed to show consistent and significant clinical benefit, most likely because only certain subsets of septic patients fit an immunologic profile that would benefit from those therapies. An important next step in sepsis research should be to identify immunologic markers in patients that can help tailor specific therapies for patients based on where they lie along the immunologic spectrum.

Abbreviations

AKI: Acute kidney injury; CI: Confidence interval; DAMP: Damage-associated molecular pattern; GFR: Glomerular filtration rate; GM-CSF: Granulocyte-macrophage colony-stimulating factor; IFN: Interferon; IL: Interleukin; LPS: Lipopolysaccharide; NF: Nuclear factor; PAMP: Pathogen-associated molecular pattern; PGC-1 α : PPAR γ coactivator-1 α ; SOFA: Sequential Organ Failure Assessment; TLR: Toll-like receptor; TNF: Tumor necrosis factor; TNFR: Tumor necrosis factor receptor.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

Work performed in our laboratories and the preparation of this review were supported by grants from the National Institutes of Health: P30 DK079307 to NMPS and KRH, R01 DK084184 to NMPS, the University of Pittsburgh, Department of Medicine Junior Scholar Awards to NMPS, R01 DK075048 to KRH, R01 HL080926 to JAK, and R01 DK070910 to JAK.

Author details

¹Renal-Electrolyte Division, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA. ²The Center for Critical Care Nephrology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA. ³CRISMA (Clinical Research Systems Modeling of Acute Illness) Center, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA. ⁴Department of Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA. ⁵Department of Cell Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA.

Published online: 02 September 2014

References

1. Ympa YP, Sakr Y, Reinhart K, Vincent JL: **Has mortality from acute renal failure decreased? A systematic review of the literature.** *Am J Med* 2005, **118**:827–832.
2. Uchino S, Kellum JA, Bellomo R, Doig GS, Morimatsu H, Morgera S, Schetz M, Tan I, Bouman C, Macedo E, Gibney N, Tolwani A, Ronco C, Beginning and Ending Supportive Therapy for the Kidney (BEST Kidney) Investigators: **Acute renal failure in critically ill patients: a multinational, multicenter study.** *JAMA* 2005, **294**:813–818.
3. Murugan R, Karajala-Subramanyam V, Lee M, Yende S, Kong L, Carter M, Angus DC, Kellum JA: **Acute kidney injury in non-severe pneumonia is associated with an increased immune response and lower survival.** *Kidney Int* 2010, **77**:527–535.
4. Bagshaw SM, Uchino S, Bellomo R, Morimatsu H, Morgera S, Schetz M, Tan I, Bouman C, Macedo E, Gibney N, Tolwani A, Oudemans-van Straaten HM, Ronco C, Kellum JA, Beginning and Ending Supportive Therapy for the Kidney (BEST Kidney) Investigators: **Septic acute kidney injury in critically ill patients: clinical characteristics and outcomes.** *Clin J Am Soc Nephrol* 2007, **2**:431–439.
5. Hotchkiss RS, Swanson PE, Freeman BD, Tinsley KW, Cobb JP, Matuschak GM, Buchman TG, Karl IE: **Apoptotic cell death in patients with sepsis, shock, and multiple organ dysfunction.** *Crit Care Med* 1999, **27**:1230–1251.
6. Langenberg C, Bagshaw SM, May CN, Bellomo R: **The histopathology of septic acute kidney injury: a systematic review.** *Crit Care* 2008, **12**:R38.
7. Lerolle N, Nochy D, Guerot E, Bruneval P, Fagon JY, Diehl JL, Hill G: **Histopathology of septic shock induced acute kidney injury: apoptosis and leukocytic infiltration.** *Intensive Care Med* 2010, **36**:471–478.
8. Takasu O, Gaut JP, Watanabe E, To K, Fagley RE, Sato B, Jarman S, Efimov IR, Janks DL, Srivastava A, Bhayani SB, Drewry A, Swanson PE, Hotchkiss RS: **Mechanisms of cardiac and renal dysfunction in patients dying of sepsis.** *Am J Respir Crit Care Med* 2013, **187**:509–517.
9. Schrier RW, Wang W, Poole B, Mitra A: **Acute renal failure: definitions, diagnosis, pathogenesis, and therapy.** *J Clin Invest* 2004, **114**:5–14.
10. Bougle A, Duranteau J: **Pathophysiology of sepsis-induced acute kidney injury: the role of global renal blood flow and renal vascular resistance.** *Contrib Nephrol* 2011, **174**:89–97.
11. Poukkanen M, Wilkman E, Vaara ST, Pettilä V, Kaukonen KM, Korhonen AM, Uusaro A, Hovilehto S, Inkinen O, Laru-Sompa R, Hautamäki R, Kuitunen A, Karlsson S, the FINNAKI Study Group: **Hemodynamic variables and progression of acute kidney injury in critically ill patients with severe sepsis: data from the prospective observational FINNAKI study.** *Crit Care* 2013, **17**:R295.
12. Asfar P, Meziani F, Hamel JF, Grelon F, Megarbane B, Anguel N, Mira JP, Dequin PF, Gergaud S, Weiss N, Legay F, Le Tulzo Y, Conrad M, Robert R, Gonzalez F, Guittou C, Tamion F, Tonnelier JM, Guezennec P, Van Der Linden T, Vieillard-Baron A, Mariotte E, Pradel G, Lesieur O, Ricard JD, Hervé F, du Cheyron D, Guerin C, Mercat A, Teboul JL, et al: **High versus low blood-pressure target in patients with septic shock.** *N Engl J Med* 2014, **370**:1583–1593.
13. Langenberg C, Bellomo R, May C, Wan L, Egi M, Morgera S: **Renal blood flow in sepsis.** *Crit Care* 2005, **9**:R363–R374.

14. Brenner M, Schaefer GL, Mallory DL, Suffredini AF, Parrillo JE: **Detection of renal blood flow abnormalities in septic and critically ill patients using a newly designed indwelling thermodilution renal vein catheter.** *Chest* 1990, **98**:170-179.
15. Lucas CE, Rector FE, Werner M, Rosenberg IK: **Altered renal homeostasis with acute sepsis. Clinical significance.** *Arch Surg* 1973, **106**:444-449.
16. Rector F, Goyal S, Rosenberg IK, Lucas CE: **Sepsis: a mechanism for vasodilatation in the kidney.** *Ann Surg* 1973, **178**:222-226.
17. Bradley VE, Shier MR, Lucas CE, Rosenberg IK: **Renal hemodynamic response to furosemide in septic and injured patients.** *Surgery* 1976, **79**:549-554.
18. Wan L, Bagshaw SM, Langenberg C, Saotome T, May C, Bellomo R: **Pathophysiology of septic acute kidney injury: what do we really know?** *Crit Care Med* 2008, **36**:S198-S203.
19. Molitoris BA: **Renal blood flow in sepsis: a complex issue.** *Crit Care* 2005, **9**:327-328.
20. Di Giandomasso D, Morimatsu H, May CN, Bellomo R: **Intrarenal blood flow distribution in hyperdynamic septic shock: effect of norepinephrine.** *Crit Care Med* 2003, **31**:2509-2513.
21. Schwartz D, Mendonca M, Schwartz I, Xia Y, Satriano J, Wilson CB, Blantz RC: **Inhibition of constitutive nitric oxide synthase (NOS) by nitric oxide generated by inducible NOS after lipopolysaccharide administration provokes renal dysfunction in rats.** *J Clin Invest* 1997, **100**:439-448.
22. Cohen RI, Hassell AM, Marzouk K, Marini C, Liu SF, Scharf SM: **Renal effects of nitric oxide in endotoxemia.** *Am J Respir Crit Care Med* 2001, **164**:1890-1895.
23. Chvojka J, Sykora R, Krouzicky A, Radej J, Varnerova V, Karvunidis T, Hes O, Novak I, Radermacher P, Matejovic M: **Renal haemodynamic, microcirculatory, metabolic and histopathological responses to peritonitis-induced septic shock in pigs.** *Crit Care* 2008, **12**:R164.
24. Legrand M, Dupuis C, Simon C, Gayat E, Mateo J, Lukaszewicz AC, Payen D: **Association between systemic hemodynamics and septic acute kidney injury in critically ill patients: a retrospective observational study.** *Crit Care* 2013, **17**:R278.
25. Kellum JA: **Impaired renal blood flow and the 'spicy food' hypothesis of acute kidney injury.** *Crit Care Med* 2011, **39**:901-903.
26. Zarjou A, Agarwal A: **Sepsis and acute kidney injury.** *J Am Soc Nephrol* 2011, **22**:999-1006.
27. Thomas L: **Germes.** *N Engl J Med* 1972, **287**:553-555.
28. Fisher CJ Jr, Agosti JM, Opal SM, Lowry SF, Balk RA, Sadoff JC, Abraham E, Schein RM, Benjamin E: **Treatment of septic shock with the tumor necrosis factor receptor:Fc fusion protein, The Soluble TNF Receptor Sepsis Study Group.** *N Engl J Med* 1996, **334**:1697-1702.
29. Fisher CJ Jr, Dhainaut JF, Opal SM, Pribble JP, Balk RA, Slotman GJ, Iberti TJ, Rackow EC, Shapiro MJ, Greenman RL, Reines HD, Shelly MP, Thompson BW, LaBrecque JF, Catalano MA, Knaus WA, Sadoff JC, Astiz M, Carpati C, Bone RC, Friedman B, Mure AJ, Brathwaite C, Shapiro E, Melhorn L, Taylor R, Keegan M, O'Brien J, Schein R, Pena M, et al: **Recombinant human interleukin 1 receptor antagonist in the treatment of patients with sepsis syndrome. Results from a randomized, double-blind, placebo-controlled trial. Phase III rhlL-1ra Sepsis Syndrome Study Group.** *JAMA* 1994, **271**:1836-1843.
30. Hotchkiss RS, Karl IE: **The pathophysiology and treatment of sepsis.** *N Engl J Med* 2003, **348**:138-150.
31. Boomer JS, To K, Chang KC, Takasu O, Osborne DF, Walton AH, Bricker TL, Jarman SD 2nd, Kreisel D, Krupnick AS, Srivastava A, Swanson PE, Green JM, Hotchkiss RS: **Immunosuppression in patients who die of sepsis and multiple organ failure.** *JAMA* 2011, **306**:2594-2605.
32. Osuchowski MF, Welch K, Siddiqui J, Remick DG: **Circulating cytokine/inhibitor profiles reshape the understanding of the SIRS/CARS continuum in sepsis and predict mortality.** *J Immunol* 2006, **177**:1967-1974.
33. Hotchkiss RS, Monneret G, Payen D: **Immunosuppression in sepsis: a novel understanding of the disorder and a new therapeutic approach.** *Lancet Infect Dis* 2013, **13**:260-268.
34. Panacek EA, Marshall JC, Albertson TE, Johnson DH, Johnson S, MacArthur RD, Miller M, Barchuk WT, Fischkoff S, Kaul M, Teoh L, Van Meter L, Daum L, Lemeshow S, Hicklin G, Doig C, Monoclonal Anti-TNF- α Randomized Controlled Sepsis Study Investigators: **Efficacy and safety of the monoclonal anti-tumor necrosis factor antibody F(ab')₂ fragment afelimomab in patients with severe sepsis and elevated interleukin-6 levels.** *Crit Care Med* 2004, **32**:2173-2182.
35. Peng ZY, Wang HZ, Carter MJ, Dileo MV, Bishop JV, Zhou FH, Wen XY, Rimmelé T, Singbartl K, Federspiel WJ, Clermont G, Kellum JA: **Acute removal of common sepsis mediators does not explain the effects of extracorporeal blood purification in experimental sepsis.** *Kidney Int* 2012, **81**:363-369.
36. Namas RA, Namas R, Lagoa C, Barclay D, Mi Q, Zamora R, Peng Z, Wen X, Fedorchak MV, Valenti IE, Federspiel WJ, Kellum JA, Vodovotz Y: **Hemoadsorption reprograms inflammation in experimental gram-negative septic peritonitis: insights from in vivo and in silico studies.** *Mol Med* 2012, **18**:1366-1374.
37. Peng ZY, Carter MJ, Kellum JA: **Effects of hemoadsorption on cytokine removal and short-term survival in septic rats.** *Crit Care Med* 2008, **36**:1573-1577.
38. Cantaluppi V, Weber V, Lauritano C, Figliolini F, Beltramo S, Biancone L, De Cal M, Cruz D, Ronco C, Segolini GP, Tetta C, Camussi G: **Protective effect of resin adsorption on septic plasma-induced tubular injury.** *Crit Care* 2010, **14**:R4.
39. Cruz DN1, Antonelli M, Fumagalli R, Foltran F, Brienza N, Donati A, Malcangi V, Petrini F, Volta G, Bobbio Pallavicini FM, Rottoli F, Giunta F, Ronco C: **Early use of polymyxin B hemoperfusion in abdominal septic shock: the EUPHAS randomized controlled trial.** *JAMA* 2009, **301**:2445-2452.
40. De Vriese AS, Colardyn FA, Philippe JJ, Vanholder RC, De Sutter JH, Lameire NH: **Cytokine removal during continuous hemofiltration in septic patients.** *J Am Soc Nephrol* 1999, **10**:846-853.
41. Cole L, Bellomo R, Journois D, Davenport P, Baldwin I, Tipping P: **High-volume haemofiltration in human septic shock.** *Intensive Care Med* 2001, **27**:978-986.
42. Payen D, Mateo J, Cavaillon JM, Fraise F, Floriot C, Vicaut E: **Impact of continuous venovenous hemofiltration on organ failure during the early phase of severe sepsis: a randomized controlled trial.** *Crit Care Med* 2009, **37**:803-810.
43. Joannes-Boyau O, Honoré PM, Perez P, Bagshaw SM, Grand H, Canivet JL, Dewitte A, Flamens C, Pujol W, Grandoulier AS, Fleureau C, Jacobs R, Broux C, Floch H, Branchard O, Franck S, Rozé H, Collin V, Boer W, Calderon J, Gauche B, Spapen HD, Janvier G, Ouattara A: **High-volume versus standard-volume haemofiltration for septic shock patients with acute kidney injury (IVOIRE study): a multicentre randomized controlled trial.** *Intensive Care Med* 2013, **39**:1535-1546.
44. VA/NIH Acute Renal Failure Trial Network, Palevsky PM, Zhang JH, O'Connor TZ, Chertow GM, Crowley ST, Choudhury D, Finkel K, Kellum JA, Paganini E, Schein RM, Smith MW, Swanson KM, Thompson BT, Vijayan A, Watnick S, Star RA, Pедуzzi P: **Intensity of renal support in critically ill patients with acute kidney injury.** *N Engl J Med* 2008, **359**:7-20.
45. Wen X, Murugan R, Kong L, Shah N, Lee MJ, Carter MJ, Elder MM, Palevsky PM, Unruh ML, Kellum JA: **Higher concentrations of inflammatory and apoptotic biomarkers are associated with mortality and nonrecovery of kidney function.** *J Am Soc Nephrol* 2011, **22**:83A.
46. Meisel C, Schefold JC, Pischowski R, Baumann T, Hetzger K, Gregor J, Weber-Carstens S, Hasper D, Keh D, Zuckermann H, Reinke P, Volk HD: **Granulocyte-macrophage colony-stimulating factor to reverse sepsis-associated immunosuppression: a double-blind, randomized, placebo-controlled multicenter trial.** *Am J Respir Crit Care Med* 2009, **180**:640-648.
47. Cohen J: **The immunopathogenesis of sepsis.** *Nature* 2002, **420**:885-891.
48. Matzinger P: **The danger model: a renewed sense of self.** *Science* 2002, **296**:301-305.
49. Zhang Q, Raoof M, Chen Y, Sumi Y, Sursal T, Junger W, Brohi K, Itagaki K, Hauser CJ: **Circulating mitochondrial DAMPs cause inflammatory responses to injury.** *Nature* 2010, **464**:104-107.
50. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Riccardi-Castagnoli P, Layton B, Beutler B: **Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene.** *Science* 1998, **282**:2085-2088.
51. Medzhitov R, Preston-Hurlburt P, Janeway CA Jr: **A human homologue of the Drosophila Toll protein signals activation of adaptive immunity.** *Nature* 1997, **388**:394-397.
52. Klune JR, Dhupar R, Cardinal J, Billiar TR, Tsung A: **HMGB1: endogenous danger signaling.** *Mol Med* 2008, **14**:476-484.

53. Chow JC, Young DW, Golenbock DT, Christ WJ, Gusovsky F: **Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction.** *J Biol Chem* 1999, **274**:10689–10692.
54. Park JS, Arcaroli J, Yum HK, Yang H, Wang H, Yang KY, Choe KH, Strassheim D, Pitts TM, Tracey KJ, Abraham E: **Activation of gene expression in human neutrophils by high mobility group box 1 protein.** *Am J Physiol Cell Physiol* 2003, **284**:C870–C879.
55. Park JS, Svetkauskaite D, He Q, Kim JY, Strassheim D, Ishizaka A, Abraham E: **Involvement of toll-like receptors 2 and 4 in cellular activation by high mobility group box 1 protein.** *J Biol Chem* 2004, **279**:7370–7377.
56. Park JS, Gamboni-Robertson F, He Q, Svetkauskaite D, Kim JY, Strassheim D, Sohn JW, Yamada S, Maruyama I, Banerjee A, Ishizaka A, Abraham E: **High mobility group box 1 protein interacts with multiple Toll-like receptors.** *Am J Physiol Cell Physiol* 2006, **290**:C917–C924.
57. Tsuboi N, Yoshikai Y, Matsuo S, Kikuchi T, Iwami K, Nagai Y, Takeuchi O, Akira S, Matsuguchi T: **Roles of toll-like receptors in C-C chemokine production by renal tubular epithelial cells.** *J Immunol* 2002, **169**:2026–2033.
58. Wolfs TG, Buurman WA, van Schadewijk A, de Vries B, Daemen MA, Hiemstra PS, van Veer C: **In vivo expression of Toll-like receptor 2 and 4 by renal epithelial cells: IFN-gamma and TNF-alpha mediated up-regulation during inflammation.** *J Immunol* 2002, **168**:1286–1293.
59. Laestadius A, Soderblom T, Aperia A, Richter-Dahlfors A: **Developmental aspects of Escherichia coli-induced innate responses in rat renal epithelial cells.** *Pediatric Res* 2003, **54**:536–541.
60. El-Achkar TM, Huang X, Plotkin Z, Sandoval RM, Rhodes GJ, Dagher PC: **Sepsis induces changes in the expression and distribution of Toll-like receptor 4 in the rat kidney.** *Am J Physiol Renal Physiol* 2006, **290**:F1034–F1043.
61. Samuelsson P, Hang L, Wullt B, Irfala H, Svanborg C: **Toll-like receptor 4 expression and cytokine responses in the human urinary tract mucosa.** *Infect Immun* 2004, **72**:3179–3186.
62. Cunningham PN, Wang Y, Guo R, He G, Quigg RJ: **Role of Toll-like receptor 4 in endotoxin-induced acute renal failure.** *J Immunol* 2004, **172**:2629–2635.
63. Patole PS, Schubert S, Hildinger K, Khandoga S, Khandoga A, Segerer S, Henger A, Kretzler M, Werner M, Krombach F, Schlöndorff D, Anders HJ: **Toll-like receptor-4: renal cells and bone marrow cells signal for neutrophil recruitment during pyelonephritis.** *Kidney Int* 2005, **68**:2582–2587.
64. Morrell ED, Kellum JA, Hallows KR, Pastor-Soler NM: **Epithelial transport during septic acute kidney injury.** *Nephrol Dial Transplant* 2014, **29**:1312–1319.
65. Senftleben U, Karin M: **The IKK/NF-kappa B pathway.** *Crit Care Med* 2002, **30**:S18–S26.
66. Wang Y, Rangan GK, Goodwin B, Tay YC, Harris DC: **Lipopolysaccharide-induced MCP-1 gene expression in rat tubular epithelial cells is nuclear factor-kappaB dependent.** *Kidney Int* 2000, **57**:2011–2022.
67. Leelahavanichkul A, Yasuda H, Doi K, Hu X, Zhou H, Yuen PS, Star RA: **Methyl-2-acetamidoacrylate, an ethyl pyruvate analog, decreases sepsis-induced acute kidney injury in mice.** *Am J Physiol Renal Physiol* 2008, **295**:F1825–F1835.
68. Miyaji T, Hu X, Yuen PS, Muramatsu Y, Iyer S, Hewitt SM, Star RA: **Ethyl pyruvate decreases sepsis-induced acute renal failure and multiple organ damage in aged mice.** *Kidney Int* 2003, **64**:1620–1631.
69. Chatterjee PK, Yeboah MM, Dowling O, Xue X, Powell SR, Al-Abed Y, Metz CN: **Nicotinic acetylcholine receptor agonists attenuate septic acute kidney injury in mice by suppressing inflammation and proteasome activity.** *PLoS One* 2012, **7**:e35361.
70. Opal SM, Laterre PF, Francois B, LaRosa SP, Angus DC, Mira JP, Wittebole X, Dugernier T, Perrotin D, Tidswell M, Jauregui L, Krell K, Pachel J, Takahashi T, Peckelsen C, Cordasco E, Chang CS, Oeyen S, Aikawa N, Maruyama T, Schein R, Kalil AC, Van Nuffelen M, Lynn M, Rossignol DP, Gogate J, Roberts MB, Wheeler JL, Vincent JL, ACCESS Study Group: **Effect of eritoran, an antagonist of MD2-TLR4, on mortality in patients with severe sepsis: the ACCESS randomized trial.** *JAMA* 2013, **309**:1154–1162.
71. Remick DG, Newcomb DE, Bolgos GL, Call DR: **Comparison of the mortality and inflammatory response of two models of sepsis: lipopolysaccharide versus cecal ligation and puncture.** *Shock* 2000, **13**:110–116.
72. Dear JW, Yasuda H, Hu X, Hieny S, Yuen PS, Hewitt SM, Sher A, Star RA: **Sepsis-induced organ failure is mediated by different pathways in the kidney and liver: acute renal failure is dependent on MyD88 but not renal cell apoptosis.** *Kidney Int* 2006, **69**:832–836.
73. Feterowski C, Emmanuilidis K, Miethke T, Gerauer K, Rump M, Ulm K, Holzmann B, Weighardt H: **Effects of functional Toll-like receptor-4 mutations on the immune response to human and experimental sepsis.** *Immunology* 2003, **109**:426–431.
74. McMasters KM, Peyton JC, Hadjiminas DJ, Cheadle WG: **Endotoxin and tumour necrosis factor do not cause mortality from caecal ligation and puncture.** *Cytokine* 1994, **6**:530–536.
75. Weighardt H, Kaiser-Moore S, Vabulas RM, Kirschning CJ, Wagner H, Holzmann B: **Cutting edge: myeloid differentiation factor 88 deficiency improves resistance against sepsis caused by polymicrobial infection.** *J Immunol* 2002, **169**:2823–2827.
76. Read RC, Pullin J, Gregory S, Borrow R, Kaczmarski EB, di Giovine FS, Dower SK, Cannings C, Wilson AG: **A functional polymorphism of toll-like receptor 4 is not associated with likelihood or severity of meningococcal disease.** *J Infect Dis* 2001, **184**:640–642.
77. Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, Richards DR, McDonald-Smith GP, Gao H, Hennessy L, Finnerty CC, López CM, Honari S, Moore EE, Minei JP, Cuschieri J, Bankey PE, Johnson JL, Sperry J, Nathens AB, Billiar TR, West MA, Jeschke MG, Klein MB, Gamelli RL, Gibran NS, Brownstein BH, Miller-Graziano C, Calvano SE, Mason PH, et al: **Genomic responses in mouse models poorly mimic human inflammatory diseases.** *Proc Natl Acad Sci U S A* 2013, **110**:3507–3512.
78. Havasi A, Borkan SC: **Apoptosis and acute kidney injury.** *Kidney Int* 2011, **80**:29–40.
79. Cunningham PN, Dyanov HM, Park P, Wang J, Newell KA, Quigg RJ: **Acute renal failure in endotoxemia is caused by TNF acting directly on TNF receptor-1 in kidney.** *J Immunol* 2002, **168**:5817–5823.
80. Jo SK, Cha DR, Cho WY, Kim HK, Chang KH, Yun SY, Won NH: **Inflammatory cytokines and lipopolysaccharide induce Fas-mediated apoptosis in renal tubular cells.** *Nephron* 2002, **91**:406–415.
81. Guo R, Wang Y, Minto AW, Quigg RJ, Cunningham PN: **Acute renal failure in endotoxemia is dependent on caspase activation.** *J Am Soc Nephrol* 2004, **15**:3093–3102.
82. Mariano F, Cantaluppi V, Stella M, Romanazzi GM, Assenzio B, Cairo M, Biancone L, Triolo G, Ranieri VM, Camussi G: **Circulating plasma factors induce tubular and glomerular alterations in septic burns patients.** *Crit Care* 2008, **12**:R42.
83. Duranton C, Rubera I, L'Hoste S, Coughon M, Poujeol P, Barhanin J, Tauc M: **KCNQ1 K+ channels are involved in lipopolysaccharide-induced apoptosis of distal kidney cells.** *Cell Physiol Biochem* 2010, **25**:367–378.
84. Ortiz-Arduan A, Danoff TM, Kalluri R, Gonzalez-Cuadrado S, Karp SL, Elkon K, Egido J, Neilson EG: **Regulation of Fas and Fas ligand expression in cultured murine renal cells and in the kidney during endotoxemia.** *Am J Physiol* 1996, **271**:F1193–F1201.
85. Rosenberger C, Mandriota S, Jurgensen JS, Wiesener MS, Horstrup JH, Frei U, Ratcliffe PJ, Maxwell PH, Bachmann S, Eckardt KU: **Expression of hypoxia-inducible factor-1alpha and -2alpha in hypoxic and ischemic rat kidneys.** *J Am Soc Nephrol* 2002, **13**:1721–1732.
86. Tran M, Tam D, Bardia A, Bhasin M, Rowe GC, Kher A, Zsengeller ZK, Akhavan-Sharif MR, Khankin EV, Saintgeniez M, David S, Burstein D, Karumanchi SA, Stillman IE, Arany Z, Parikh SM: **PGC-1alpha promotes recovery after acute kidney injury during systemic inflammation in mice.** *J Clin Invest* 2011, **121**:4003–4014.
87. Huet O, Chereau C, Nicco C, Dupic L, Conti M, Borderie D, Pene F, Vicaut E, Benhamou D, Mira JP, Duranteau J, Batteux F: **Pivotal role of glutathione depletion in plasma-induced endothelial oxidative stress during sepsis.** *Crit Care Med* 2008, **36**:2328–2334.
88. Heyman SN, Rosen S, Rosenberger C: **A role for oxidative stress.** *Contrib Nephrol* 2011, **174**:138–148.
89. Sawyer DB, Loscalzo J: **Myocardial hibernation: restorative or preterminal sleep?** *Circulation* 2002, **105**:1517–1519.
90. Wright SD, Ramos RA, Tobias PS, Ulevitch RJ, Mathison JC: **CD14, a receptor for complexes of lipopolysaccharide (LPS) and LPS binding protein.** *Science* 1990, **249**:1431–1433.
91. Bussolati B, David S, Cambi V, Tobias PS, Camussi G: **Urinary soluble CD14 mediates human proximal tubular epithelial cell injury induced by LPS.** *Int J Mol Med* 2002, **10**:441–449.

92. Fearn C, Loskutoff DJ: **Role of tumor necrosis factor alpha in induction of murine CD14 gene expression by lipopolysaccharide.** *Infect Immun* 1997, **65**:4822–4831.
93. El-Achkar TM, Dagher PC: **Renal Toll-like receptors: recent advances and implications for disease.** *Nat Clin Pract Nephrol* 2006, **2**:568–581.
94. Medzhitov R, Preston-Hurlburt P, Kopp E, Stadlen A, Chen C, Ghosh S, Janeway CA Jr: **MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways.** *Mol Cell* 1998, **2**:253–258.
95. Taal MW, Brenner BM, Rector FC: *Brenner & Rector's the Kidney*. 9th edition. Philadelphia, PA: Elsevier/Saunders; 2012.
96. Green DR, Kroemer G: **The pathophysiology of mitochondrial cell death.** *Science* 2004, **305**:626–629.

doi:10.1186/s13054-014-0501-5

Cite this article as: Morrell *et al.*: Septic acute kidney injury: molecular mechanisms and the importance of stratification and targeting therapy. *Critical Care* 2014 **18**:501.