

POSTER PRESENTATION

Open Access

Dopamine-NMDA interactions and relevance to gamma band synchrony in schizophrenia

Kübra Kömek^{1*}, George B Ermentrout^{1,2}, Raymond Y Cho^{1,3}

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

An increasing body of behavioral and physiological evidence shows that cortical activity operates optimally within a limited range of dopamine (DA) transmission within a broader inverted-U shaped relationship [1]. However, the exact mechanism through which DA leads to such non-monotonic modulation of network activity needs further investigation. Our recent work suggests that DA's modulation of potassium currents in the fast-spiking interneurons could be a potential mechanism underlying this inverted-U relationship [2]. Here we propose that another potential mechanism could contribute in the form of interactions between DA D1 receptors and NMDA conductance in the cortex. This has been an area of interest for many reasons including its relevance to the etiology of schizophrenia with deficits in both NMDA-mediated glutamate and DA transmission. Studies examining the relationship between DA and NMDA conductance in the cortex revealed that DA D1 receptors enhance NMDA responses [3-5].

In the current study, we incorporated DA's modulation of NMDA conductance at a network level and analyzed whether DA's modulation of NMDA currents could lead to an inverted-U shaped relationship with cortical activity. With this motivation, we simulated a neural network with 200 excitatory and 40 inhibitory quadratic integrate-and-fire neurons coupled with biologically realistic probabilities. In the simulated network, the effects of DA were implemented through varying the NMDA conductances in various combinations of synaptic types including excitatory-excitatory and excitatory-inhibitory. The network synchronization was analyzed through examining gamma band power in the local field potential. Parametrically varying the NMDA conductance revealed an inverted-U

shaped relationship, with lower gamma band power at both low and high conductance levels and optimal synchronization occurring at intermediate conductance levels. Our findings reveal that DA's modulation of NMDA currents at the single cell level gives rise to a non-monotonic relationship between cortical gamma band synchrony and DA levels. These results together with those of [2] suggest that DA's modulation of potassium currents as well as NMDA-mediated currents may be complementary or synergistic mechanisms that give rise to the inverted-U shaped pattern in cortical activity across dopamine levels.

Author details

¹Center for the Neural Basis of Cognition, Carnegie Mellon University, Pittsburgh, PA, 15213, USA. ²Department of Mathematics, University of Pittsburgh, Pittsburgh, PA, 15213, USA. ³Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, 15213, USA.

Published: 8 July 2013

References

1. Williams GV, Castner SA: **Under the curve: critical issues for elucidating D1 receptor function in working memory.** *Neuroscience* 2006, **139**(1):263-276.
2. Kömek K, Ermentrout GB, Walker CP, Cho RY: **Dopamine and gamma band synchrony in schizophrenia—insights from computational and empirical studies.** *Eur J Neurosci* 2012, **36**:2146-2155.
3. Seamans JK, Durstewitz D, Christie BR, Stevens CF, Sejnowski TJ: **Dopamine D1/D5 receptor modulation of excitatory synaptic inputs to layer V prefrontal cortex neurons.** *PNAS* 2001, **98**(1):301-306.
4. Wang J, O'Donnell P: **D(1) dopamine receptors potentiate NMDA-mediated excitability increase in layer V prefrontal cortical pyramidal neurons.** *Cereb Cortex* 2001, **11**(5):452-462.
5. Chen G, Greengard P, Yan Z: **Potentiation of NMDA receptor currents by dopamine D1 receptors in prefrontal cortex.** *PNAS* 2004, **101**(8):2596-2600.

doi:10.1186/1471-2202-14-S1-P218

Cite this article as: Kömek et al.: Dopamine-NMDA interactions and relevance to gamma band synchrony in schizophrenia. *BMC Neuroscience* 2013 **14**(Suppl 1):P218.

* Correspondence: kubrakomek@gmail.com

¹Center for the Neural Basis of Cognition, Carnegie Mellon University, Pittsburgh, PA, 15213, USA

Full list of author information is available at the end of the article