



MEETING ABSTRACT

Open Access

Positron emission tomography imaging of amyloid-beta plaque deposition: a decade of translation

Julie C Price^{1*}, Chester A Mathis¹, William E Klunk²

From 2012 Sino-American Symposium on Clinical and Translational Medicine (SAS-CTM) Shanghai, China. 27-29 June 2012

Positron emission tomography (PET) radiotracer developments over the past decade have enabled *in vivo* measurement of amyloid-beta (A β) plaque deposition, a key neuropathological hallmark of Alzheimer's disease (AD). This presentation will provide an overview of the translational research path for one of the most widely used PET A β imaging agents, ¹¹C-labeled Pittsburgh compound-B (¹¹C]PiB). Early preclinical *ex-vivo* characterization revealed the capacity of ¹¹C]PiB to bind to fibrillar A β plaques [1]. Real-time *in vivo* multiphoton microscopy demonstrated that ¹¹C]PiB labeled individual A β plaques in transgenic mouse models of AD [2]. Human proof-of-concept studies then showed nearly 2-fold greater uptake of ¹¹C]PiB in AD patients relative to controls in areas of brain known to contain amyloid in AD (frontal cortex, $p < 0.0001$), while retention was equivalent for both groups in areas known to be relatively unaffected by amyloid deposition (subcortical white matter, pons, cerebellum; $p > 0.2$) [3]. These semi-quantitative studies were followed by fully quantitative arterial-based kinetic modeling PET studies that supported the validity of simplified (non-arterial) ¹¹C]PiB PET retention outcomes that exhibited good test-retest reliability (5-10%) needed for improved study feasibility for clinical application on cross-sectional and longitudinal bases [4] and for large collaborative multi-site studies, such as the Alzheimer's disease neuroimaging initiative (ADNI).

Early translational findings include cross-sectional evidence of amyloid deposition (i.e., ¹¹C]PiB retention) in 20-30% of cognitively normal elderly controls, variable retention in subjects with mild cognitive impairment

(MCI) that ranged from negligible to AD-like levels and additional findings in those at risk for the development of AD [[5], for review]. Emerging longitudinal results indicate small significant increases in ¹¹C]PiB retention in AD and MCI groups, and in controls who had high baseline retention [6]. Efforts are ongoing to establish relationships between *in vivo* imaging measures and post-mortem measures of A β deposition in those scanned with ¹¹C]PiB PET imaging before death. Anti-amyloid therapies have shown promise based on ¹¹C]PiB PET imaging results in early treatment trials. Lastly, the development of ¹⁸F-labeled A β imaging agents (several-fold longer radioactive half-life than ¹¹C) has allowed greater distribution and translational capability, with one such agent recently being approved by the U.S. Food and Drug Administration for use in the rejection of AD diagnosis in patients under evaluation for AD. A decade of translation has further clarified the need for sensitive early detection of AD pathophysiology in order to identify those who might benefit most from future therapeutic intervention.

Author details

¹Departments of Radiology, University of Pittsburgh, Pittsburgh PA, 15213, USA. ²Departments of Psychiatry, University of Pittsburgh, Pittsburgh PA, 15213, USA.

Published: 17 October 2012

References

1. Mathis CA, Wang Y, Holt DP, et al: Synthesis and evaluation of ¹¹C-labeled 6 substituted 2-arylbenzothiazoles as amyloid imaging agents. *J Med Chem* 2003, **46**:2740-54.
2. Backskai BJ, Hickey GA, Skoch J, et al: Four-dimensional multiphoton imaging of brain entry, amyloid binding, and clearance of an amyloid-beta ligand in transgenic mice. *Proc Natl Acad Sci* 2003, **100**:12462-7.
3. Klunk WE, Engler H, Nordberg A, et al: Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol* 2004, **55**:306-19.

* Correspondence: pricjc@upmc.edu

¹Departments of Radiology, University of Pittsburgh, Pittsburgh PA, 15213, USA

Full list of author information is available at the end of the article

4. Price JC, Klunk WE, Lopresti BJ, *et al*: Kinetic modeling of amyloid binding in humans using PET imaging and Pittsburgh Compound-B. *J Cereb Blood Flow Metab* 2005, **25**:1528-47.
5. Wolk DA, Klunk W: Update on amyloid imaging: from healthy aging to Alzheimer's disease. *Curr Neurol Neurosci Rep* 2009, **9**:345-52.
6. Villemagne VL, Pike KE, Chetelat G, *et al*: Longitudinal assessment of Abeta and cognition in aging and Alzheimer disease. *Ann Neurol* 2011, **69**:181-92.

doi:10.1186/1479-5876-10-S2-A31

Cite this article as: Price *et al.*: Positron emission tomography imaging of amyloid-beta plaque deposition: a decade of translation. *Journal of Translational Medicine* 2012, **10**(Suppl 2):A31.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

