

The presence of a high- K_m hexokinase activity in dog, but not in boar, sperm

Josep M. Fernández-Novell^a, Joan Ballester^b, Antonio Medrano^b, Pedro J. Otaegui^c,
Teresa Rigau^b, Joan J. Guinovart^a, Joan E. Rodríguez-Gil^{b,*}

^aDepartment of Biochemistry and Molecular Biology and IRBB, Barcelona Science Park, University of Barcelona, E-08028 Barcelona, Spain

^bUnit of Animal Reproduction, Department of Animal Medicine and Surgery, School of Veterinary Medicine, Autonomous University of Barcelona, E-08193 Bellaterra, Spain

^cDepartment of Biochemistry and Molecular Biology, School of Veterinary Medicine, Autonomous University of Barcelona, E-08193 Bellaterra, Spain

Received 2 March 2004; revised 31 May 2004; accepted 8 June 2004

Available online

Edited by Judit Ovádi

Abstract The presence of a high- K_m hexokinase activity was tested in both dog and boar spermatozoa. Hexokinase kinetics from dog extracts showed the presence of a specific activity (dog-sperm glucokinase-like protein, DSGLP), in the range of glucose concentrations of 4–10 mM, whereas boar sperm did not show any DSGLP activity. Furthermore, dog-sperm cells, but not those of boar, showed the presence of a protein which specifically reacted against a rat-liver anti-glucokinase antibody. This protein also had a molecular weight equal to that observed in rat-liver extracts, suggesting a close similarity between both the proteins. This glucokinase-like protein was distributed in the peri- and post-acrosomal zones of the head, and the midpiece and principal piece of tail of dog spermatozoa. These results indicate that dog spermatozoa have functional high- K_m hexokinase activity, which could contribute to a very fine regulation of their hexose metabolism. This strict regulation could ultimately be very important in optimizing dog-sperm function along its life-time.

© 2004 Published by Elsevier B.V. on behalf of the Federation of European Biochemical Societies.

Keywords: High- K_m hexokinase; Dog-sperm; Glucokinase-regulatory protein

1. Introduction

Carbohydrates are, probably, the most important substrates for the maintenance of energy levels in mammalian sperm from fresh ejaculates. Thus, sugars such as glucose, fructose and mannose are utilized as energy sources by sperm cells from species like bull, dog and boar [1]. However, recent findings support the hypothesis that sugars can play another role, at least in some mammalian species, and not only as mere energy substrates. This is especially evident in dog, where the incubation of sperm cells from fresh ejaculates with either glucose or fructose induced hexose-specific changes in functional parameters such as motility [2] or tyrosine phosphorylation

patterns [3]. These glucose- or fructose-specific effects were related to specific actions on the majority of the evaluated metabolic parameters, such as intracellular levels of glucose 6-phosphate and glycogen or production of L-lactate and CO₂ [3]. Glucose and fructose also showed separate effects on hexokinase activity [3], and they even induced separate effects, not only on glycogen synthase activity [4], but also on the intracellular, specific location of this enzyme in dog sperm [5]. All of these results indicate that dog sperm would have very sophisticated mechanisms to specifically identify sugars that they are consuming in order to direct them to induce the sugar-linked, functional effects. These mechanisms might be related to the intake of sugars, since the fructose-specific transporter, GLUT5, and the more glucose-specific transporter, GLUT3, are located in separate zones, not only in dog sperm [3], but also in other mammalian species, such as bull, mice and human [6]. However, there are probably other systems that allow sperm to optimize these hexose-differentiating mechanisms, at least in dog.

Vertebrate glucokinase (hexokinase type IV) is a member of the hexokinase protein family which shows some remarkable characteristics that clearly differentiate it from the other mammalian hexokinases. In fact, glucokinase does not have a strict specificity for substrate, since it can phosphorylate not only glucose, but also fructose or mannose [7]. Nevertheless, glucokinase's elevated K_m for glucose, together with its specific expression in the liver and pancreas, allows it to be a sensitive and efficient control step for the maintenance of mammalian glucose metabolism [7]. The existence of a similar, high- K_m hexokinase activity in mammalian sperm could be an efficient system to control that described above, i.e., hexose-specific functional changes observed, at least in dog. Taking this all into consideration, the main aim of this work is to test the presence of a high- K_m hexokinase activity in mammalian sperm, which could act similarly to hepatic glucokinase in the control of sperm's hexose metabolism. For this purpose, sperm cells from dog and boar were used, since they are species which show very different functional characteristics, from their motion parameters (dog cells are fast and linear, whereas boar cells are much slower, see [2,8]) to their life-span after ejaculation (dog spermatozoa last about one week inside the female genital tract, whereas boar cells last only about 48 h, see [9]). In these cells the total hexokinase activity kinetics was deter-

* Corresponding author. Fax: +34-935812006.

E-mail address: juanenrique.rodruiguez@uab.es (J.E. Rodríguez-Gil).

Abbreviations: DSGLP, dog-sperm glucokinase-like protein; PBS, phosphate-buffered saline; SDS, sodium dodecyl sulfate; TRITC, tetramethylrhodamine isothiocyanate

91 mined, and then the presence of proteins with immunological
92 properties like rat-liver glucokinase and glucokinase regula-
93 tory protein was tested, in order to find some glucokinase-like,
94 or at least some high- K_m hexokinase, activity in the cells. Our
95 results indicate that dog spermatozoa, but not those of boar,
96 showed a clear high- K_m hexokinase activity, as well as a pro-
97 tein which specifically reacts to an anti-rat-liver glucokinase
98 antibody. This high- K_m hexokinase activity could be related to
99 the dog-sperm's ability to specifically react in front of glucose
100 or fructose.

101 2. Materials and methods

103 2.1. Animals and sample collection

104 Canine semen was obtained from 11 purebred Beagle dogs ranging
105 from 2 to 7 years of age. The care of the dogs followed the guidelines
106 indicated in the Catalan Animal Welfare Law (Generalitat de Cata-
107 lunya, Spain). Semen was collected once or twice weekly by manual
108 stimulation. Only the sperm-rich fraction of the ejaculates was used.

109 Porcine semen was obtained from 10 healthy boars ranging from 2 to
110 5 years of age that belonged to a commercial herd (Servicios Genéticos
111 Porcinos, S.L.; Roda de Ter, Spain). Ejaculates were manually col-
112 lected and only the rich-sperm fractions were used.

113 2.2. Total hexokinase activity kinetics

114 The kinetics of the total hexokinase activity in dog and boar sper-
115 matozoa was determined as in [3]. For this, both boar and dog samples
116 were centrifuged at $600 \times g$ for 10 min and were then resuspended in 50
117 mL of a Krebs–Ringer–Henseleit solution without sugars at 15 °C
118 (KRH– medium; pH 7.4). Sperm were again centrifuged at $600 \times g$ for
119 10 min and sedimented cells were then resuspended in an additional 50
120 mL of the KRH– medium. The centrifugation–resuspension step was
121 repeated once more, in order to completely eliminate all substances
122 included in seminal plasma that could affect kinetic and immunological
123 properties of sperm hexokinases. The final, 50-mL-sperm suspension
124 was again centrifuged at $600 \times g$ for 10 min. After this, the supernatant
125 was discarded and the resultant pellets were immediately homogenized
126 by sonication in 250 μ L of an ice-cold buffer (pH 7.4) containing 500
127 mM glycylglycine, 2 M KCl, 100 mM dithiothreitol, 300 IU/mL
128 aprotinin and 100 mM phenylmethylsulfonyl fluoride (hexokinase
129 buffer). Homogenized samples were centrifuged at $10\,000 \times g$ for 15
130 min at 4 °C and hexokinase activity was measured both in the resultant
131 supernatants and in the pellets. For this purpose, the pellets were
132 washed once in 500 μ L of hexokinase buffer and were further resus-
133 pended in 250 μ L of hexokinase buffer. Hexokinase activity was
134 measured as in [10] with the addition of increasing concentrations of
135 glucose in the reaction buffer after adaptation of the technique to a
136 Cobas Bio autoanalyzer (Roche Biomedical, Basel, Switzerland).

137 2.3. Immunological techniques

138 For this experiment, semen samples were pooled both from two
139 (boar semen) or four ejaculates (dog semen). Samples were initially
140 treated through the three times centrifugation/KRH– resuspension
141 washing step described above. After this, both dog and boar sperm
142 cells were suspended in a final 5-mL KRH– medium at 37 °C. Aliquots
143 of the suspension were placed in open vials and incubated with con-
144 tinuous shaking at 37 °C, with the addition to the medium of either
145 glucose or fructose at a final concentration of 10 mM in both cases.
146 Concentrations of sperm cells in the final suspension were of 3.5 –
147 4×10^5 spermatozoa/mL in dog samples and 3 – 6×10^6 spermatozoa/
148 mL in those of boar. Finally, aliquots were taken at the indicated times
149 and processed for immunological detection.

150 Western blot analyses were performed in samples homogenized by
151 sonication in ice-cold 10-mM Tris–HCl buffer (pH 7.4) containing 1%
152 (w/v) sodium dodecyl sulfate (SDS) and 1 mM Na_2VO_4 (homogeniza-
153 tion buffer, proportion 1:5, v/v). The samples were briefly boiled and
154 were then centrifuged at $10\,000 \times g$ for 14 min at 4 °C. Mammalian sperm
155 has very low amounts of cytoplasm and a very compartmentalized
156 structure [11]. These particularities led us to consider the presence of the

tested proteins in either the supernatant or the pellet obtained after
homogenization, centrifugation and the boiling of the samples, since
proteins could be in a free form or they could be linked to the internal
sperm structures. For this purpose, the obtained pellets were resus-
pended in 20–30 μ L of the homogenization buffer, and Western blot
analyses were performed in both supernatants and resuspended pellets
obtained in all of this process.

Western blot was based on SDS gel electrophoresis [12], followed by
transfer to nitrocellulose [13]. The transferred samples were tested with
the antibodies at a dilution of 1:1000 (v/v). Immunoreactive proteins
were tested using peroxidase-conjugated goat, anti-rabbit second
antibody (dilution 1:200, v/v) and the reaction was developed with an
ECL-Plus detection system (Amersham, Buckinghamshire, England).

Immunocytochemistry was carried out with spermatozoa seeded onto
glass coverslips, which were washed with phosphate-buffered saline
(PBS; pH 7.4) and were fixed for 30 min in PBS containing 4% (w/v)
paraformaldehyde. The fixed samples were incubated with 1 mg/mL
 NaBH_4 to eliminate autofluorescence, and blocked in 3% (w/v) bovine
serum albumin in PBS. Spermatozoa were further incubated with the
anti-glucokinase antibody (dilution 1:200, v/v) for 2 h at 15–17 °C, washed
with PBS and treated with a tetramethylrhodamine isothiocyanate
(TRITC)-conjugated swine anti-rabbit immunoglobulin (Dako, Glost-
rup, Denmark). Finally, fluorescent images were obtained by a Leica
TCS 4D confocal scanning laser microscope (Leica Lasertechnik, Hei-
delberg, Germany), adapted to an inverted Leitz DMIRBE microscope
and a $63\times$ (NA 1.4 oil) Leitz Plan-Apo Lens (Leitz, Stuttgart, Germany).
The light source was an argon/krypton laser (75 mW).

2.4. Suppliers

Anti-rat-liver glucokinase and anti-rat-liver glucokinase regulatory
protein were produced and tested in the laboratory of Dr. Guinovart
(IRBB, Barcelona Science Park, University of Barcelona; see [14]). All
of the reactives used were of analytical grade.

3. Results

3.1. Kinetics of dog and boar total hexokinase activity

Supernatants obtained from homogenates of dog sperm
showed an increase in total hexokinase activity when it was
determined in the presence of glucose in a range from 0.05 mM
(1.4 ± 0.1 IU/mg protein) to 2 mM (3.7 ± 0.4 IU/mg protein,
see Fig. 1A). A further, and noticeable increase of total
hexokinase activity was then determined in a range of glucose
concentration from 4 mM (4.2 ± 0.4 IU/mg protein) to 10 mM
(8.4 ± 0.8 IU/mg protein, see Fig. 1A). No further increase in
total hexokinase activity was detected at glucose concentra-
tions above 10 mM. On the other hand, total hexokinase ac-
tivity from pellets obtained after homogenization of dog sperm
from fresh ejaculates also showed an increase in total hexo-
kinase activity in the glucose concentration range from 0.05
mM (0.60 ± 0.1 IU/mg protein) to 4 mM (4.0 ± 0.2 IU/mg pro-
tein). Again, a further and noticeable increase of total hexo-
kinase activity was observed between 4 mM glucose (4.0 ± 0.2
IU/mg protein) and 10 mM glucose (6.2 ± 0.3 IU/mg protein,
see Fig. 1B). These results were reflected in the Lineweaver–
Burke representation. Thus, as shown in Fig. 1C, hexokinase
kinetics of supernatants from dog-sperm extracts described a
biphasic diagram, with two separate lines, the first in the glu-
cose range from 0.05 to 6 mM, and the second in the glucose
range from 6 to 50 mM. A theoretical, approximate calculus of
the K_m of both lines resulted in values of 8.5 and 0.08 mM,
assuming that these values are only approximate. These results
were compatible with the presence of a glucokinase-like ac-
tivity in dog-sperm supernatants. On the other hand, the Li-
neweaver–Burke representation of pellets from dog-sperm
homogenates also showed the presence of two separate lines




Fig. 1. Kinetics of total hexokinase activity in dog- and boar-sperm extracts. (A and B) Relationship between total hexokinase activity and glucose concentration in supernatants (A) and resuspended pellets (B) from dog (◆) and boar (■) sperm extracts. (C and D) Semi-logarithmic representation between inverse values of hexokinase-specific activity and glucose concentration in supernatants (A) and resuspended pellets (B) from dog (◆) and boar (■) sperm extracts. Lines show the apparent hexokinase activity types from dog (continuous lines) and boar samples (broken lines) revealed by this representation. Results are means \pm S.E.M. for eight separate semen samples.

220 with very different slopes, the first in the glucose range from
221 0.05 to 6 mM, and the second once again from 6 to 50 mM
222 (Fig. 1D). In this case, the approximate, theoretical values of
223 the K_m of both lines were of 6.9 and 0.09 mM.

224 Total hexokinase kinetics of boar sperm showed totally
225 different results. Thus, supernatants from homogenates of
226 these cells did not show an appreciable increase, from 0.05 mM
227 glucose (1.9 ± 0.2 IU/mg protein) to 50 mM glucose (2.0 ± 0.3
228 IU/mg protein, see Fig. 1A), thus indicating the presence of
229 only one hexokinase activity that was very sensitive to the
230 presence of glucose in the medium. Slightly different results
231 were observed in pellets from these homogenates, since in this
232 case there was a progressive increase in hexokinase activity in
233 the glucose range from 0.05 mM (3.9 ± 0.7 IU/mg protein) to
234 0.5 mM (4.9 ± 0.8 mM, see Fig. 1B), and further increases of
235 hexokinase activity were not noticeable (7.2 ± 0.7 IU/mg pro-
236 tein in the presence of 50 mM glucose). These results were
237 reflected in the Lineweaver–Burke representation, where both
238 supernatants and pellets from boar sperm showed only one
239 line from 0.05 mM glucose to 50 mM glucose (Figs. 1C and D).
240 The calculated, approximate K_m value of this line was of 0.03
241 mM in supernatants and 0.06 mM in resuspended pellets.

242 3.2. Presence of an immunoreactive protein against 243 anti-glucokinase antibody

244 The Western blot using an anti-rat-liver glucokinase anti-
245 body revealed the presence of a protein which specifically re-

246 acted against this antibody in dog sperm (Fig. 2). This protein
247 was clear in the pellets, with a molecular weight of about 50
248 kDa (Fig. 2A). On the other hand, the Western blot from
249 supernatants showed two fainter bands of a molecular weight
250 of about 45–50 kDa (Fig. 2B) The intensity and the molecular
251 weight of these bands were not modified after incubation in the




Fig. 2. Western blot against sperm glucokinase. The figure shows two representative Western blots obtained from supernatants (A) and resuspended pellets (B) from dog and boar (P) sperm homogenates. Dog sperm was analyzed from fresh ejaculates (D0) or subjected to incubation with 10 mM glucose for 10 min (DG) before being subjected to Western blot analysis. M: Molecular weight markers. L: Rat-liver extracts. The total number of independent replicates for these Western blots was 5.

252 presence of 10 mM glucose after up to 30 min (Fig. 2, and data
253 not shown). It is noteworthy that liver extracts used as positive
254 controls showed a positive band for glucokinase of about 50
255 kDa, which match to those observed in both supernatants and
256 pellets from dog-sperm homogenates (Fig. 2). Finally, boar
257 homogenates did not show any 50-kDa band which could
258 correspond to the result observed in dog extracts, although
259 they revealed a positive signal of about 45 kDa, similar to
260 another non-specific reactivity band detected in liver extracts
261 (Fig. 2).

262 Confocal images showed the presence of a specific, reactive
263 protein against the anti-rat-liver glucokinase antibody in both
264 the head and the tail of dog spermatozoa from fresh ejaculates
265 (Fig. 3). Tail marking was located at both the midpiece and the
266 main piece, whereas head location was established in both the
267 peri-acrosomal and the post-acrosomal zones. These locations
268 were not significantly modified after incubation in the presence
269 of 10 mM glucose after up to 30 min (Fig. 3C, and data not
270 shown). No positive reaction in front of the anti-rat-liver
271 glucokinase antibody was observed in boar spermatozoa (data
272 not shown).

3.3. Detection of sperm glucokinase regulatory protein

273 Western blot from dog-sperm extracts did not demonstrate
274 the presence of a specific reactivity against an anti-rat-liver
275 glucokinase regulatory protein antibody, neither in superna-
276 tants nor in resuspended pellets obtained after homogenization
277 of samples (Fig. 4, and data not shown). On the other hand,
278 supernatants, but not resuspended pellets, obtained after ho-
279 mogenization of boar spermatozoa showed a clear, specific 70-
280 kDa protein, which was equal to that obtained in rat-liver
281 extracts (Fig. 4A, and data not shown). Molecular weight and
282 density of this band in boar sperm were not modified after
283 incubation with 10 mM glucose and 10 mM fructose after up
284 to 30 min (data not shown). 285

4. Discussion and conclusions

286 Our results indicate the presence of a high- K_m hexokinase
287 activity with a similarity to glucokinase in dog sperm, but not
288 in boar cells. This can be sustained by the following facts: 289




Fig. 3. Immunocytochemistry of dog-sperm glucokinase. The figure shows representative images of dog spermatozoa immunolocalized against glucokinase. A: Negative control. B: Cells from fresh, control samples. C: Cells from samples incubated with 10 mM glucose for 5 min. The total number of independent replicates for these experiments was 5. Images have a multiplication factor of 75. Arrows indicate sperm tails, whereas asterisks indicate sperm heads.




Fig. 4. Western blot against sperm glucokinase regulatory protein. The figure shows a representative Western blot obtained from supernatants obtained after homogenization of either dog or boar (P) sperm homogenates, following the technique described in Section 2. Dog sperm was from fresh ejaculates (D0) or it was subjected to incubation with 10 mM glucose for 10 min (DG) before being subjected to Western blot analysis. M: Molecular weight markers. L: Rat-liver extracts. The total number of independent replicates for these Western blots was 5.

290 (i) The presence, in dog spermatozoa, of a specific hexokinase
 291 activity which shows a high K_m for glucose, which was in
 292 the range described for mammalian glucokinase (5–10
 293 mM, see [7,10]).
 294 (ii) The presence of a dog-sperm protein which specifically re-
 295 acts against an anti-rat-liver glucokinase antibody and,
 296 moreover, shows a molecular weight similar to mamma-
 297 lian glucokinase (about 50 kDa, see [7]).
 298 Thus, immunological and kinetic data point towards this
 299 interpretation. Of course, we cannot affirm that this protein is
 300 truly glucokinase. Moreover, other proteins different from
 301 glucokinase have been described with high- K_m characteristics,
 302 such as *N*-acetylglucosamine kinase [15] or the high- K_m
 303 hexokinase present in the mhAT3F hepatoma cell line
 304 (mhAT3F-HK, see [16]). However, kinetic and immunological
 305 characteristics indicate that the dog-sperm, high- K_m hexoki-
 306 nase (DSGLP) is more similar to glucokinase than these pro-
 307 teins. Thus, *N*-acetylglucosamine kinase shows a K_m for
 308 glucose (370 mM, see [15]) far higher than that of DSGLP (7–9
 309 mM, see Section 3). Furthermore, the comparison of the
 310 structural sequences between glucokinase (reference number
 311 NP-034422.2, see [17]) and *N*-acetylglucosamine kinase (ref-
 312 erence number Q9QZ08, see [17]) revealed a percentage of
 313 overall structural affinity of 11.6%. On the other hand, the
 314 structural comparison between the epitope utilized to develop
 315 the anti-glucokinase antibody [14] and the whole sequence of
 316 the *N*-acetylglucosamine kinase rendered the following result:
 317 Epytope of liver glucokinase:
 318 414-KLHPSFKERFASVR-428
 319 Sequence of *N*-acetylglucosamine kinase which rendered the
 320 higher homology:
 321 274-KSWELLKEGFLALT-288
 322 As shown in this comparison, in the best case only four amino-
 323 acids, which were not linear (highlighted letters), are the same
 324 in both sequences. These data seem to indicate that our antibody
 325 has a low degree of affinity for *N*-acetylglucosamine kinase.
 326 Furthermore, the theoretical molecular weight of *N*-acetylglu-
 327 cosamine kinase, calculated from its aminoacidic composition
 328 [17], is about 37–38 kDa. Our Western blot analysis did not
 329 detect any band with could be included in the 35–40 kDa mo-
 330 lecular-weight range. Summing up all of these data, we can
 331 conclude that *N*-acetylglucosamine kinase does not seem to be
 332 DSGLP. On the other hand, mhAT3F-HK also shows a K_m for
 333 glucose (40 mM, see [16]) higher than DSGLP. Moreover, al-
 334 though this protein recognized an anti-glucokinase antibody, its
 335 molecular weight was about 30 kDa [16]. Our results did not

336 reveal the presence of any protein with this molecular weight.
 337 The sum of all of these results, of course, does not preclude the
 338 presence of *N*-acetylglucosamine kinase and mhAT3F-HK in
 339 dog sperm. However, our results also indicate that DSGLP with
 340 immunological properties similar to rat-liver glucokinase, re-
 341 gardless of the existence of other high- K_m hexokinases.
 342 Nevertheless, some differences between glucokinase and
 343 DSGLP exist. Thus, glucokinase activity is mainly regulated in
 344 two ways, first through controlling its expression [7]. However,
 345 mammalian sperm does not have the ability to express its
 346 DNA to synthesize proteins [11], so, in this manner, dog sperm
 347 cannot control DSGLP activity through this way. The second
 348 way is through changes in its spatial position inside the cell
 349 [14]. This latter control is achieved through a glucose-modu-
 350 lated linking between glucokinase and the glucokinase-regu-
 351 latory protein, which displaces glucokinase in the hepatocyte
 352 to areas with or without glucose, thus controlling its ability to
 353 phosphorylate the monosaccharide [14]. On the other hand,
 354 displacements of glucokinase-regulatory protein through the
 355 hepatocyte to achieve its control of glucokinase activity are
 356 possible due to the hepatocyte being a cell with an active and
 357 more-or-less relaxed nuclear structure as well as with a great
 358 amount of cytoplasm. Spermatozoa have neither requirement,
 359 since their amount of cytoplasm is very scarce and their nu-
 360 clear structure is condensed and very inactive, totally different
 361 from that observed in hepatocytes [18]. Moreover, the presence
 362 of a clear immunoreactivity of glucokinase-regulatory protein
 363 in boar sperm, which did not show DSGLP activity, suggests
 364 the lack of a clear regulatory role for the glucokinase-regula-
 365 tory protein in dog sperm. Thus, DSGLP activity regulation
 366 would have probably been achieved by other ways, like
 367 phospho-dephosphorylation mechanisms that are also opera-
 368 tive on glucokinase [19].
 369 The functional role of DSGLP could be explained as a regu-
 370 latory mechanism of dog-sperm hexose metabolism. Previous
 371 reports have shown that dog-sperm cells have a very complex
 372 hexose metabolism, which includes the presence of anabolic
 373 paths like glycogen metabolism, and differentiated, functional
 374 roles for separate hexoses [2,4,5]. The elevated K_m for glucose
 375 shown by DSGLP could be an efficient system to control glu-
 376 cose-specific mechanisms of regulation of dog-sperm function-
 377 ality, in a similar way that glucokinase controls glucose-induced
 378 changes in hepatocytes and pancreatic β -cells [7,10].
 379 It is noteworthy that DSGLP is present in dog spermatozoa,
 380 but not in boar cells. This would mark great differences in
 381 hexose metabolism and, hence, in the energy status manage-
 382 ment of both species. In fact, hexose metabolism of sperma-
 383 tozoa from several mammals like boar or bull are basically
 384 glycolytic, with elevated L-lactate formation rates, low glucose
 385 6-phosphate levels, low activity in anabolic pathways like
 386 glycogen synthesis and small differences, if any, in the selective
 387 utilization of hexoses such as glucose or fructose as energy
 388 substrates [20,21]. In these spermatozoa, the presence of
 389 DSGLP makes no sense, since they utilize different monosac-
 390 charides in the same way for the same ultimate necessity, the
 391 attainment of energy. Nevertheless, as commented above, en-
 392 ergy management of dog spermatozoa is very different, and the
 393 selective, functional utilization that these cells have of glucose
 394 and fructose can easily explain the necessity of DSGLP. Thus,
 395 our results indicate that there is not a single, simple mechanism
 396 that explains the regulation of mammalian sperm function

397 under the energetic point of view. This has to be taken into
398 consideration when trying to apply new strategies in the con-
399 servation of sperm, since it would vary depending on the
400 specific metabolic and hexose-related functional profiles of
401 each species.

402 5. Conclusions

403 The presence of DSGLP in dog spermatozoa would play an
404 important role in the control of both the energy management
405 pathways and the hexose-related functional mechanisms that
406 dog spermatozoa show from fresh ejaculates. Notwithstand-
407 ing, at this moment we have no real knowledge of the exact
408 physiological role and control mechanisms of DSGLP, and
409 experiments regarding substrate specificity, besides purification,
410 sequencing and cloning of DSGLP, will be needed to
411 clarify these important points.

412 *Acknowledgements:* We thank Mr. Chuck Simmons for his valuable
413 assistance in preparing the English version of the manuscript, Mrs.
414 Raquel García-Olivas (Serveis Científico-Técnicos, Barcelona Science
415 Park, University of Barcelona) for her assistance in obtaining confocal
416 images and Mr. Daniel Cifuentes (Barcelona Science Park, University
417 of Barcelona) for his kind collaboration in the comparison of protein
418 sequences. J.B. was a recipient of a fellowship from the “Generalitat de
419 Catalunya” (Spain). The manuscript has been partially financed by
420 Grant AGL2001-2568 from the “Ministerio de Ciencia y Tecnología”
421 (Spain).

422 References

- 423 [1] Mann, A. (1975) in: Handbook of Physiology (Greep, R.O. and
424 Astwood, E.B., Eds.), pp. 321–347, American Physiology Society,
425 Washington, DC.
426 [2] Rigau, T., Farré, M., Ballester, J., Peña, A. and Rodríguez-Gil,
427 J.E. (2001) *Theriogenology* 56, 801–815.
428 [3] Rigau, T., Rivera, M., Palomo, M.J., Mogas, T., Ballester, J.,
429 Peña, A., Otaegui, P.J., Guinovart, J.J. and Rodríguez-Gil, J.E.R.
430 (2002) *Reproduction* 123, 579–591.

- [4] Ballester, J., Fernández-Novell, J.M., Rutllant, J., García-Rocha,
M., Palomo, M.J., Mogas, T., Peña, A., Rigau, T., Guinovart, J.J.
and Rodríguez-Gil, J.E. (2000) *Mol. Reprod. Dev.* 56, 207–
219. 431
[5] Palomo, M.J., Fernández-Novell, J.M., Peña, A., Guinovart, J.J.,
Rigau, T. and Rodríguez-Gil, J.E. (2003) *Mol. Reprod. Dev.* 64,
349–359. 432
[6] Angulo, C., Rauch, M.C., Droppelmann, A., Reyes, A.M., Slebe,
J.C., Delgado-López, F., Guaiquil, V.H., Vera, J.C. and Concha,
I.L. (1998) *J. Cell. Biochem.* 71, 189–203. 433
[7] Cárdenas, M.L., Cornish-Bowden, A. and Ureta, T. (1998)
Biochim. Biophys. Acta. 434
[8] Quintero-Moreno, A., Rigau, T. and Rodríguez-Gil, J.E. (2003)
Theriogenology, in press. 435
[9] Feldman, E.C. and Nelson, R.W. (1987) in: *Canine and Feline*
Endocrinology and Reproduction (Pedersen, E., Ed.), pp. 420–
421, WB Saunders, Philadelphia. 436
[10] Otaegui, P.J., Ferré, T., Pujol, A., Riu, E., Jiménez, R. and Bosch,
F. (2000) *Hum. Gene Therapy* 11, 15443–15552. 437
[11] Bedford, J.M. and Hoskins, D.D. (1990) in: *Marshall's Physiology*
of Reproduction (Lamming, G.E., Ed.), pp. 379–568, Churchill
Livingstone, Edinburgh. 438
[12] Laemmli, U.K. (1970) *Nature* 227, 680–685. 439
[13] Burnet, W.N. (1981) *J. Anal. Biochem.* 112, 195–203. 440
[14] De la Iglesia, N., Veiga-da-Cunha, M., Van Schaftingen, E.,
Guinovart, J.J. and Ferrer, J.C. (1999) *FEBS Lett.* 456, 332–
338. 441
[15] Davagnino, J. and Ureta, T. (1980) *J. Biol. Chem.* 255, 2633–
2636. 442
[16] Rencurel, F., Muñoz-Alonso, M.J., Girard, J. and Leturque, A.
(1998) *J. Biol. Chem.* 273, 26187–26193. 443
[17] National Center for Biotechnology Information Database. Beth-
esda, MD, 2004. 444
[18] Eddy, E.M. (1988) in: *The Physiology of Reproduction* (Knobil,
E. and Neil, J.D., Eds.), pp. 27–68, Raven Press, New York. 445
[19] Grimsby, J., Sarabu, R., Corbett, W.L., Haynes, N.-E., Bizarro,
F.T., Coffey, J.W., Guertin, K.R., Hiliard, D.W., Kester, R.F.,
Mahaney, P.E., Marcus, L., Qi, L., Spence, C.L., Tnegi, J.,
Magnuson, M.A., Chu, C.A., Dvoroziak, M.T., Natschinsky,
F.M. and Grippo, J.F. (2003) *Science* 301, 370–373. 446
[20] Hammersted, R.H. and Lardy, H.A. (1983) *J. Biol. Chem.* 258,
8759–8768. 447
[21] Marín, S., Chiang, K., Bassilian, S., Lee, W.N.P., Boros, L.G.,
Fernández-Novell, J.M., Centelles, J.J., Medrano, A., Rodríguez-
Gil, J.E. and Cascante, M. (2003) *FEBS Lett.* 554, 342–346. 448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475