

On two and three periodic Lyness difference equations*

Anna Cima⁽¹⁾, Armengol Gasull⁽¹⁾ and Víctor Mañosa⁽²⁾

⁽¹⁾ *Dept. de Matemàtiques, Facultat de Ciències,
Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
cima@mat.uab.cat, gasull@mat.uab.cat*

⁽²⁾ *Dept. de Matemàtica Aplicada III (MA3),
Control, Dynamics and Applications Group (CoDALab)
Universitat Politècnica de Catalunya (UPC)
Colom 1, 08222 Terrassa, Spain
victor.manosa@upc.edu*

Abstract

We describe the sequences $\{x_n\}_n$ given by the non-autonomous second order Lyness difference equations $x_{n+2} = (a_n + x_{n+1})/x_n$, where $\{a_n\}_n$ is either a 2-periodic or a 3-periodic sequence of positive values and the initial conditions x_1, x_2 are as well positive. We also show an interesting phenomenon of the discrete dynamical systems associated to some of these difference equations: the existence of one oscillation of their associated rotation number functions. This behavior does not appear for the autonomous Lyness difference equations.

2000 Mathematics Subject Classification: 39A20, 39A11

Keywords: Difference equations with periodic coefficients, circle maps, rotation number.

1 Introduction and main result

This paper fully describes the sequences given by the non-autonomous second order Lyness difference equations

$$x_{n+2} = \frac{a_n + x_{n+1}}{x_n}, \quad (1)$$

where $\{a_n\}_n$ is a k -periodic sequence taking positive values, $k = 2, 3$, and the initial conditions x_1, x_2 are as well positive. This question is proposed in [5, Sec. 5.43]. Recall

***Acknowledgements.** GSD-UAB and CoDALab Groups are supported by the Government of Catalonia through the SGR program. They are also supported by MCYT through grants MTM2008-03437 (first and second authors) and DPI2008-06699-C02-02 (third author).

that non-autonomous recurrences appear for instance as population models with a variable structure affected by some seasonality [11, 12], where k is the number of seasons. Some dynamical issues of similar type of equations have been studied in several recent papers [1, 9, 10, 13, 15, 17, 18].

Recall that when $k = 1$, that is $a_n = a > 0$, for all $n \in \mathbb{N}$, then (1) is the famous Lyness recurrence which is well understood, see for instance [2, 19]. The cases $k = 2, 3$ have been already studied and some partial results are established. For both cases it is known that the solutions are persistent near a given k -periodic solution, which is stable. This is proved by using some known invariants, see [15, 17, 18]. Recall that in our context it is said that a solution $\{x_n\}_n$ is persistent if there exist two real positive constants c and C , which depend on the initial conditions, such that for all $n \geq 1$, $0 < c < x_n < C < \infty$. We prove:

Theorem 1. *Let $\{x_n\}_n$ be any sequence defined by (1) and $k \in \{2, 3\}$. Then it is persistent. Furthermore, either*

- (a) *the sequence $\{x_n\}_n$ is periodic, with period a multiple of k ; or*
- (b) *the sequence $\{x_n\}_n$ densely fills one or two (resp. one, two or three) disjoint intervals of \mathbb{R}^+ when $\{a_n\}_n$ is 2-periodic (resp. 3-periodic). Moreover it is possible by algebraic tools to distinguish which is the situation.*

Our approach to describe the sequences $\{x_n\}_n$ is based on the study of the natural dynamical system associated to (1) and on the results of [7]. The main tool that allows to distinguish the number of intervals for the adherence of the sequences $\{x_n\}_n$ is the computation of several resultants, see Section 4.

It is worth to comment that Theorem 1 is an extension of what happens in the classical case $k = 1$. There, the same result holds but in statement (b) only appears one interval. Our second main result will prove that there are other more significative differences between the case $k = 1$ and the cases $k = 2, 3$. These differences are related with the lack of monotonicity of certain rotation number functions associated to the dynamical systems given by the Lyness recurrences, see Theorem 3. The behaviors of these rotation number functions are important for the understanding of the recurrences, because they give the possible periods for them, see [2, 4, 19].

On the other hand in [10, 18] it is proved that, at least for some values of $\{a_n\}_n$, the behaviour of $\{x_n\}_n$ for the case $k = 5$ is totally different. In particular unbounded positive solutions appear. In the forthcoming paper [8] we explore in more detail the differences between the cases $k = 1, 2, 3$ and $k \geq 4$.

This paper is organized as follows: Section 2 presents the difference equations that we are studying as discrete dynamical systems and we state our main results on them, see

Theorems 2 and 3. Section 3 is devoted to the proof of Theorem 2. By using it, in Section 4, we prove Theorem 1 and we give some examples of how to apply it to determine the number of closed intervals of the adherence of $\{x_n\}_n$. In Section 5 we demonstrate Theorem 3 and we also present some examples where we study in more detail the rotation number function of the dynamical systems associated to (1).

2 Main results from the dynamical systems point of view

In this section we reduce the study of the sequence $\{x_n\}_n$ to the study of some discrete dynamical systems and we state our main results on them.

First we introduce some notations. When $k = 2$, set

$$a_n = \begin{cases} a & \text{for } n = 2\ell + 1, \\ b & \text{for } n = 2\ell, \end{cases} \quad (2)$$

and when $k = 3$, set

$$a_n = \begin{cases} a & \text{for } n = 3\ell + 1, \\ b & \text{for } n = 3\ell + 2, \\ c & \text{for } n = 3\ell, \end{cases} \quad (3)$$

where $\ell \in \mathbb{N}$ and $a > 0, b > 0$ and $c > 0$.

We also consider the maps $F_\alpha(x, y)$, with $\alpha \in \{a, b, c\}$, as

$$F_\alpha(x, y) = \left(y, \frac{\alpha + y}{x} \right),$$

defined on the open invariant set $Q^+ := \{(x, y) : x > 0, y > 0\} \subset \mathbb{R}^2$.

Consider for instance $k = 2$. The sequence given by (1),

$$x_1, x_2, x_3, x_4, x_5, x_6, x_7, \dots, \quad (4)$$

can be seen as

$$(x_1, x_2) \xrightarrow{F_a} (x_2, x_3) \xrightarrow{F_b} (x_3, x_4) \xrightarrow{F_a} (x_4, x_5) \xrightarrow{F_b} (x_5, x_6) \xrightarrow{F_a} \dots$$

Hence the behavior of (4) can be obtained from the study of the dynamical system defined in Q^+ by the map:

$$F_{b,a}(x, y) := F_b \circ F_a(x, y) = \left(\frac{a + y}{x}, \frac{a + bx + y}{xy} \right).$$

Similarly, for $k = 3$ we can consider the map:

$$F_{c,b,a}(x, y) := F_c \circ F_b \circ F_a(x, y) = \left(\frac{a + bx + y}{xy}, \frac{a + bx + y + cxy}{y(a + y)} \right).$$

It can be proved that both maps have an only fixed point in Q^+ , which depends on a, b (and c), that for short we denote by \mathbf{p} .

It is easy to interpret the invariants for (1) and $k = 2, 3$, given in [13, 15], in terms of first integrals of the above maps, see also Lemma 6. We have that

$$V_{b,a}(x, y) := \frac{ax^2y + bxy^2 + bx^2 + ay^2 + (b^2 + a)x + (b + a^2)y + ab}{xy},$$

is a first integral for $F_{b,a}$ and

$$V_{c,b,a}(x, y) := \frac{cx^2y + axy^2 + bx^2 + by^2 + (a + bc)x + (c + ab)y + ac}{xy},$$

is a first integral for $F_{c,b,a}$. Notice that

$$abV_{b,a}(x, y) + a^2b^2 + a^3 + b^3 = \frac{(bx + a)(ay + b)(ax + by + ab)}{xy}$$

and

$$abcV_{c,b,a}(x, y) + a^2b^2 + a^2c^2 + b^2c^2 = \frac{(bcx + aby + ac)(acxy + abx + bcy + abc)}{xy}$$

are as well first integrals of $F_{b,a}$ and $F_{c,b,a}$, respectively. They remain the well-known expression

$$V_a(x, y) = \frac{(x + 1)(y + 1)(x + y + a)}{xy}$$

of the first integral of the Lyness' map F_a .

The topology of the level sets of $V_{b,a}$ and $V_{c,b,a}$ in Q^+ as well as the dynamics of the maps restricted to them is described by the following result, that will be proved in Section 3.

Theorem 2. (i) *Each level set of $V_{b,a}$ (resp. $V_{c,b,a}$) in $Q^+ \setminus \{\mathbf{p}\}$ is diffeomorphic to a circle surrounding \mathbf{p} , which is the unique fixed point of $F_{b,a}$ (resp. $F_{c,b,a}$).*

(ii) *The action of $F_{b,a}$ (resp. $F_{c,b,a}$) on each level set of $V_{b,a}$ (resp. $V_{c,b,a}$) contained in $Q^+ \setminus \{\mathbf{p}\}$ is conjugated to a rotation of the circle.*

Once a result like the above one is established the study of the possible periods of the sequences $\{x_n\}_n$ given by (1) is quite standard. It suffices, first to get the rotation interval, which is the open interval formed by all the rotation numbers given by the above theorem, varying the level sets of the first integrals. Afterwards, it suffices to find which are the denominators of all the irreducible rational numbers that belong to the corresponding interval, see [4, 6, 19].

The study of the rotation number of these kind of rational maps is not an easy task, see again [2, 4, 6, 19]. In particular, in [4] was proved that the rotation number function

parameterized by the energy levels of the Lyness map $F_a, a \neq 1$, is always monotonous, solving a conjecture of Zeeman given in [19], see also [16]. As far as we know, in this paper we give the first simple example for which this rotation number function is neither constant nor monotonous. We prove:

Theorem 3. *There are positive values of a and b , such that the rotation number function $\rho_{b,a}(h)$ of $F_{b,a}$ associated to the closed ovals of $\{V_{b,a} = h\} \subset Q^+$ has a local maximum.*

Hence, apart from the known behaviors for the autonomous Lyness maps, that is global periodicity or monotonicity of the rotation number function, which trivially holds for $F_{b,a}$, taking for instance $a = b = 1$ or $a = b \neq 1$, respectively, there appear more complicated behaviors for the rotation number function.

Our proof of this result relies on the study of lower and upper bounds for the rotation number of $F_{b,a}$ on a given oval of a level set of $V_{b,a}$ given for some $(a, b) \in Q^+$ and $\{V_{b,a}(x, y) = V_{b,a}(x_0, y_0)\}$, for $(x_0, y_0) \in Q^+$. This can be done because the map on this oval is conjugated to a rotation and it is possible to use an algebraic manipulator to follow and to order a finite number iterates on it, which are also given by points with rational coordinates. So, only exact arithmetic is used. A similar study could be done for $F_{c,b,a}$.

3 Proof of Theorem 2

Proof of (i) of Theorem 2. Case $k = 2$. The orbits of $F_{b,a}$ lie on the level sets of $V_{b,a}$. First we prove that if $h > 0$ the sets $\{(x, y) \in Q^+ : V_{b,a}(x, y) \leq h\}$ are compact. From the inequality

$$V_{b,a}(x, y) = ax + by + b\frac{x}{y} + a\frac{y}{x} + \frac{b^2 + a}{y} + \frac{b + a^2}{x} + \frac{ab}{xy} \leq h$$

we get that

$$\frac{b + a^2}{h} \leq x \leq \frac{h}{a} \quad \text{and} \quad \frac{a + b^2}{h} \leq y \leq \frac{h}{b}.$$

Notice that this result in particular already implies the persistence of the sequences given by (1).

Next, let us show that in Q^+ , the set of fixed points of $F_{b,a}$ and the set of singular points of $V_{b,a}$ coincide and they only contain a point \mathbf{p} . Indeed, the fixed points of $F_{b,a}$ are given by $x^2 = a + y$ and $y^2 = b + x$. Since $f(x) := a + \sqrt{b + x} - x^2$ is concave, $f(0) > 0$, $f'(0) > 0$ and $\lim_{x \rightarrow +\infty} f(x) = -\infty$, there is a unique positive zero of f , and therefore there is a unique fixed point \mathbf{p} in Q^+ .

Now, notice that

$$\frac{\partial V_{b,a}(x, y)}{\partial x} = \frac{ay + b}{x^2 y} (x^2 - y - a) \quad \text{and} \quad \frac{\partial V_{b,a}(x, y)}{\partial y} = \frac{bx + a}{xy^2} (y^2 - x - b).$$

So in Q^+ the only fixed point of $F_{b,a}$ coincides with the only singular point of $V_{b,a}$, which necessarily is the point where it attains its absolute minimum.

The above facts prove that each level set of $V_{b,a}$ in Q^+ is, for $h > h_c := V_{b,a}(\mathbf{p})$, diffeomorphic to a circle: it is compact, a differentiable closed curve without singular points and the boundary of the connected open set $\{V_{b,a}(x, y) < h\}$ from statement (5) of [3, Prop. 2.1].

Case $k = 3$. By using the same kind of arguments as in the previous case the result follows. We will only give the details of the proof that the fixed point of $F_{c,b,a}$ in Q^+ is unique and coincides with the unique singular point of $V_{c,b,a}$ in Q^+ .

Clearly a fixed point in Q^+ satisfies

$$y = g(x) := \frac{bx + a}{x^2 - 1} \quad \text{and} \quad y(y + a) = x(x + c). \quad (5)$$

So the fixed point in Q^+ is given by the unique intersection point in $\{x > 1\}$ of the graphs of the decreasing function $y = g(x)$ and the increasing one $y = (-a + \sqrt{a^2 + 4x(x + c)})/2$.

The point (x, y) is a singular point of $V_{c,b,a}$ if and only if

$$x^2 = \frac{(by + c)(y + a)}{cy + b} \quad \text{and} \quad y^2 = \frac{(bx + a)(x + c)}{ax + b}. \quad (6)$$

Consider $\Delta := x(x + c) - y(y + a)$. By taking $x + c$ from the right equation in (6) and $y + a$ from the left one we obtain that

$$\Delta = \frac{y(ax + b)(by + c) - x(bx + a)(cy + b)}{(bx + a)(by + c)} xy.$$

Some computations, by using once more equations in (6), show that

$$\begin{aligned} & y(ax + b)(by + c) - x(bx + a)(cy + b) \\ &= b(axy^2 + by^2 + cy - cx^2y - bx^2 - ax) = b(y^2(ax + b) + cy - x^2(cy + b) - ax) \\ &= b[(bx + a)(x + c) + cy - (by + c)(y + a) - ax] = b^2 [x(x + c) - y(y + a)]. \end{aligned}$$

Therefore

$$\Delta = \frac{b^2 xy}{(bx + a)(by + c)} \Delta.$$

If $\Delta \neq 0$ we obtain that $aby + cbx + ac = 0$, which is impossible. So $\Delta = 0$, which means that any singular point verifies the second equation of the fixed point.

Combining the right equations of (5) and (6), we get

$$ay = (x + c) \left(x - \frac{bx + a}{ax + b} \right) \quad \text{or equivalently} \quad \frac{y}{x + c} = \frac{x^2 - 1}{ax + b}.$$

By using again the right equation of (6) we get that $y = (bx + a)/(x^2 - 1)$ as we wanted to prove. ■

3.1 Proof of (ii) of Theorem 2

In [7] it is proved a result that characterizes the dynamics of integrable diffeomorphisms having a *Lie Symmetry*, that is a vector field X such that $X(F(p)) = (DF(p))X(p)$. Next theorem states it, particularized to the case we are interested.

Theorem 4 ([7]). *Let $\mathcal{U} \subset \mathbb{R}^2$ be an open set and let $\Phi : \mathcal{U} \rightarrow \mathcal{U}$ be a diffeomorphism such that:*

(a) *It has a smooth regular first integral $V : \mathcal{U} \rightarrow \mathbb{R}$, having its level sets $\Gamma_h := \{z = (x, y) \in \mathcal{U} : V(z) = h\}$ as simple closed curves.*

(b) *There exists a smooth function $\mu : \mathcal{U} \rightarrow \mathbb{R}^+$ such that for any $z \in \mathcal{U}$,*

$$\mu(\Phi(z)) = \det(D\Phi(z)) \mu(z).$$

Then the map Φ restricted to each Γ_h is conjugated to a rotation with rotation number $\tau(h)/T(h)$, where $T(h)$ is the period of Γ_h as a periodic orbit of the planar differential equation

$$\dot{z} = \mu(z) \left(-\frac{\partial V(z)}{\partial y}, \frac{\partial V(z)}{\partial x} \right)$$

and $\tau(h)$ is the time needed by the flow of this equation for going from any $w \in \Gamma_h$ to $\Phi(w) \in \Gamma_h$.

Next lemma is one of the key points for finding a Lie symmetry for families of periodic maps, like the 2 and 3-periodic Lyness maps.

Lemma 5. *Let $\{G_a\}_{a \in A}$ be a family of diffeomorphisms of $\mathcal{U} \subset \mathbb{R}^2$. Suppose that there exists a smooth map $\mu : \mathcal{U} \rightarrow \mathbb{R}$ such that for any $a \in A$ and any $z \in \mathcal{U}$, the equation $\mu(G_a(z)) = \det(DG_a(z)) \mu(z)$ is satisfied. Then, for every choice $a_1, \dots, a_k \in A$, we have*

$$\mu(G_{[k]}(z)) = \det(DG_{[k]}(z)) \mu(z),$$

where $G_{[k]} = G_{a_k} \circ \dots \circ G_{a_2} \circ G_{a_1}$.

Proof. It is only necessary to prove the result for $k = 2$ because the general case follows easily by induction. Consider $a_1, a_2 \in A$ then

$$\begin{aligned} \mu(G_{a_2, a_1}(z)) &= \mu(G_{a_2} \circ G_{a_1}(z)) = \det(DG_{a_2}(G_{a_1}(z))) \mu(G_{a_1}(z)) = \\ &= \det(DG_{a_2}(G_{a_1}(z))) \det(DG_{a_1}(z)) \mu(z) = \det(D(G_{a_2} \circ G_{a_1}(z))) \mu(z) = \\ &= \det(DG_{a_2, a_1}(z)) \mu(z), \end{aligned}$$

and the lemma follows. ■

Proof of (ii) of Theorem 2. From part (i) of the theorem we know that the level sets of $V_{b,a}$ and $V_{c,b,a}$ in $Q^+ \setminus \{\mathbf{p}\}$ are diffeomorphic to circles. Moreover these functions are first integrals of $F_{b,a}$ and $F_{c,b,a}$, respectively. Notice also that for any a , the Lyness map $F_a(x, y) = (y, \frac{a+y}{x})$ satisfies

$$\mu(F_a(x, y)) = \det(DF_a(x, y))\mu(x, y),$$

with $\mu(x, y) = xy$. Hence, by Lemma 5,

$$\mu(F_{b,a}(x, y)) = \det(DF_{b,a}(x, y))\mu(x, y) \quad \text{and} \quad \mu(F_{c,b,a}(x, y)) = \det(DF_{c,b,a}(x, y))\mu(x, y).$$

Thus, from Theorem 4, the result follows. ■

It is worth to comment that once part (i) of the theorem is proved it is also possible to prove that the dynamics of $F_{b,a}$ (resp. $F_{c,b,a}$) restricted to the level sets of $V_{b,a}$ (resp. $V_{c,b,a}$) is conjugated to a rotation by using that they are given by cubic curves and that the map is birational, see [14]. We prefer our approach because it provides a dynamical interpretation of the rotation number together with its analytic characterization.

4 Proof of Theorem 1

In order to prove Theorem 1 we need a preliminary result. Consider the maps $F_{b,a}$ and $F_{a,b}$, jointly with their corresponding first integrals $V_{b,a}$ and $V_{a,b}$. In a similar way consider $F_{c,b,a}$, $F_{a,c,b}$ and $F_{b,a,c}$ with $V_{c,b,a}$, $V_{a,c,b}$ and $V_{b,a,c}$. Some simple computations prove the following elementary but useful lemma. Notice that it can be interpreted as the relation between the first integrals and the non-autonomous invariants.

Lemma 6. *With the above notations:*

$$(i) \quad V_{b,a}(x, y) = V_{a,b}(F_a(x, y)).$$

$$(ii) \quad V_{c,b,a}(x, y) = V_{a,c,b}(F_a(x, y)) = V_{b,a,c}(F_b(F_a(x, y))).$$

Proof of Theorem 1. We split the proof in two steps. For $k = 2, 3$ we first prove that there are only two types of behaviors for $\{x_n\}_n$, either this set of points is formed by kp points for some positive integer p , or it has infinitely many points whose adherence is given by at most k intervals. Secondly, in this later case, we provide an algebraic way for studying the actual number of intervals.

First step: We start with the case $k = 2$. With the notation introduced in (2), it holds that

$$F_{b,a}(x_{2n-1}, x_{2n}) = (x_{2n+1}, x_{2n+2}), \quad F_{a,b}(x_{2n}, x_{2n+1}) = (x_{2n+2}, x_{2n+3}),$$

where $(x_1, x_2) \in Q^+$ and $n \geq 1$. So the odd terms of the sequence $\{x_n\}_n$ are contained in the projection on the x -axis of the oval of $\{V_{b,a}(x, y) = V_{b,a}(x_1, x_2) = h\}$ and the even terms in the corresponding projection of $\{V_{a,b}(x, y) = V_{a,b}(F_a(x_1, x_2)) = V_{b,a}(x_1, x_2) = h\}$, where notice that we have used Lemma 6.

Recall that the ovals of $V_{b,a}$ are invariant by $F_{b,a}$ and the ovals of $V_{a,b}$ are invariant by $F_{a,b}$. Notice also that the trivial equality $F_a \circ F_{b,a} = F_{a,b} \circ F_a$ implies that the action of $F_{b,a}$ on $\{V_{b,a}(x, y) = h\}$ is conjugated to the action of $F_{a,b}$ on $\{V_{a,b}(x, y) = h\}$ via F_a .

From Theorem 2 we know that $F_{b,a}$ on the corresponding oval is conjugated to a rotation of the circle. Hence, if the corresponding rotation number is rational, then the orbit starting at (x_1, x_2) is periodic, of period say q , then the sequence $\{x_n\}_n$ is $2q$ -periodic. On the other hand if the rotation number is irrational, then the orbit of (x_1, x_2) generated by $F_{b,a}$ fulfills densely the oval of $\{V_{b,a}(x, y) = h\}$ in Q^+ and hence the subsequence of odd terms also fulfills densely the projection of $\{V_{b,a}(x, y) = h\}$ in the x -axis. Clearly, the sequence of even terms do the same with the projection of the oval of $\{V_{a,b}(x, y) = h\}$.

Similarly when $k = 3$ the equalities

$$\begin{aligned} F_{c,b,a}(x_{3n-2}, x_{3n-1}) &= (x_{3n+1}, x_{3n+2}), \\ F_{a,c,b}(x_{3n-1}, x_{3n}) &= (x_{3n+2}, x_{3n+3}), \\ F_{b,a,c}(x_{3n}, x_{3n+1}) &= (x_{3n+3}, x_{3n+4}), \end{aligned}$$

where $n \geq 1$, allow to conclude that each term x_m , of the sequence $\{x_n\}_n$ where we use the notation (3), is contained in one of the projections on the x -axis of the ovals $\{V_{c,b,a}(x, y) = V_{c,b,a}(x_1, x_2) =: h\}$ and $\{V_{a,c,b}(x, y) = h\}$ and $\{V_{b,a,c}(x, y) = h\}$, according with the remainder of m after dividing it by 3. The rest of the proof in this case follows as in the case $k = 2$. So the first step is done.

Second step: From the above results it is clear that the problem of knowing the number of connected components of the adherence of $\{x_n\}_n$ is equivalent to the control of the projections of several invariant ovals on the x -axis. The strategy for $k = 3$, and analogously for $k = 2$, is the following. Consider the ovals contained in the level sets given by $\{V_{c,b,a}(x, y) = h\}$, $\{V_{a,c,b}(x, y) = h\}$ and $\{V_{b,a,c}(x, y) = h\}$ and denote by $I = I(a, b, c, h)$, $J = J(a, b, c, h)$ and $K = K(a, b, c, h)$ the corresponding closed intervals of $(0, \infty)$ given by their projections on the x -axis.

We want to detect the values of h for which two of the intervals, among I , J and K , have exactly one common point. First we seek their boundaries. Since the level sets are given by cubic curves, that are quadratic with respect the y -variable, these points will correspond with values of x for which the discriminant of the quadratic equation with respect to y is

zero. So, we compute

$$R_1(x, h, a, b, c) := \text{dis}(xyV_{c,b,a}(x, y) - hxy, y) = 0,$$

$$R_2(x, h, a, b, c) := \text{dis}(xyV_{a,c,b}(x, y) - hxy, y) = 0,$$

$$R_3(x, h, a, b, c) := \text{dis}(xyV_{b,a,c}(x, y) - hxy, y) = 0.$$

Now we have to search for relations among a, b, c and h for which two of these three functions have some common solution, x . These relations can be obtained by computing some suitable resultants.

Taking the resultants of R_1 and R_2 ; R_2 and R_3 ; and R_1 and R_3 with respect x we obtain three polynomial equations $R_4(h, a, b, c) = 0$, $R_5(h, a, b, c) = 0$ and $R_6(h, a, b, c) = 0$. In short, once a, b and c are fixed we have obtained three polynomials in h such that a subset of their zeroes give the bifurcation values which separate the number of intervals of the adherence of $\{x_n\}_n$. See the results of Proposition 7 and Example 8 for concrete applications of the method.

Before ending the proof we want to comment that for most values of a, b and c , varying h there appear the three possibilities, namely 1, 2 or 3 different intervals. The last case appears for values of h near $h_c := V_{c,b,a}(\mathbf{p})$, because the first coordinates of the three points \mathbf{p} , $F_a(\mathbf{p})$ and $F_b(F_a(\mathbf{p}))$ almost never coincide. The other situations can be obtained by increasing h . ■

Proposition 7. *Consider the recurrence (1) with $k = 2$ and $\{a_n\}_n$ as in (2) taking the values $a = 3$ and $b = 1/2$. Define $h_c = (12z^3 - 33z + 7)/(2(z^2 - 3)) \simeq 17.0394$, where $z \simeq 2.1513$ is the biggest positive real root of $2z^4 - 12z^2 - 2z + 17$, and $h^* \simeq 17.1198$, is the smallest positive root of*

$$p_4(h) := 112900h^4 - 2548088h^3 - 48390204h^2 + 564028596h + 7613699255.$$

Then,

(i) *The initial condition $(x_1, x_2) = (z, z^2 - 3)$ gives a two periodic recurrence $\{x_n\}_n$. Moreover $V_{1/2,3}(z, 3 - z^2) = h_c$.*

(ii) *Let (x_1, x_2) be any positive initial conditions, different from $(z, z^2 - 3)$, and set $h = V_{1/2,3}(x_1, x_2)$. Let $\rho(h)$ denote the rotation number of $F_{1/2,3}$ restricted to the oval of $\{V_{1/2,3}(x, y) = h\}$. Then*

(I) *If $\rho(h) = p/q \in \mathbb{Q}$, with $\text{gcd}(p, q) = 1$, then the sequence $\{x_n\}_n$ is $2q$ -periodic.*

(II) *If $\rho(h) \notin \mathbb{Q}$ and $h \in (h_c, h^*)$ then the adherence of the sequence $\{x_n\}_n$ is formed by two disjoint closed intervals.*

(III) If $\rho(h) \notin \mathbb{Q}$ and $h \in [h^*, \infty)$ then the adherence of the sequence $\{x_n\}_n$ is one closed interval.

We want to remark that, from a computational point of view, the case (I) almost never is detected. Indeed, taking a and b rational numbers and starting with rational initial conditions, by using Mazur's theorem it can be seen that almost never the rotation number will be rational, see the proof of [2, Prop. 1]. Therefore, in numeric simulations only situations (II) and (III) appear, and the value $h = h^*$ gives the boundary between them. In general, for $k = 2$, the value h^* is always the root of a polynomial of degree four, which is constructed from the values of a and b .

Proof of Proposition 7. Clearly $(z, 3-z^2)$ is the fixed point of $F_{b,a}$ in Q^+ . Some computations give the compact expression of $h_c := V_{a,b}(z, z^2 - 3)$. To obtain the values h^* we proceed as in the proof of Theorem 1. In general,

$$\begin{aligned} R_1(x, h, a, b) &:= \text{dis}(xyV_{b,a}(x, y) - hxy, y) \\ &= (ax^2 - hx + a^2 + b)^2 - 4(bx + a)(bx^2 + b^2x + ax + ab), \\ R_2(x, h, a, b) &:= \text{dis}(xyV_{a,b}(x, y) - hxy, y) \\ &= (bx^2 - hx + a + b^2)^2 - 4(ax + b)(ax^2 + a^2x + bx + ab). \end{aligned}$$

Then we have to compute the resultant of the above polynomials with respect to x . It always decomposes as the product of two quartic polynomials in h . Its expression is very large, so we only give it when $a = 3$ and $b = 1/2$. It writes as

$$\frac{625}{65536} (4h^4 - 1176h^3 + 308h^2 + 287380h + 1816975) p_4(h).$$

It has four real roots, two for each polynomial. Some further work proves that the one that interests us is the smallest one of p_4 . ■

We also give an example when $k = 3$, but skipping all the details.

Example 8. Consider the recurrence (1) with $k = 3$ and $\{a_n\}_n$ as in (3) taking the values $a = 1/2, b = 2$ and $c = 3$. Then for any positive initial conditions x_1 and x_2 , $V_{c,b,a}(x_1, x_2) = h \geq V_{c,b,a}(\mathbf{p}) = h_c \simeq 15.9283$. Moreover if the rotation number of $F_{c,b,a}$ associated to the oval $\{V_{c,b,a}(x, y) = h\}$ is irrational then the adherence of $\{x_n\}_n$ is given by:

- Three intervals when $h \in (h_c, h^*)$, where $h^* \simeq 15.9614$;
- Two intervals when $h \in [h^*, h^{**})$, where $h^{**} \simeq 16.0015$;
- One interval when $h \in [h^{**}, \infty)$.

The values h^* and h^{**} are roots of two polynomials of degree 8 with integer coefficients that can be explicitly given.

Finally, for $k = 2$, we give an easy sufficient condition to ensure that the projections of the level sets of $\{V_{b,a}(x, y) = h\}$ and $\{V_{a,b}(x, y) = h\}$ on the x -axis are not disjoint.

Remark 9. Consider the recurrence (1) with $k = 2$ and $\{a_n\}_n$ as in (2). Let $\mathbf{p}_{b,a}$ and $\mathbf{p}_{a,b}$ be the fixed points of $F_{b,a}$ and $F_{a,b}$ in Q^+ respectively. Then, if either $V_{b,a}(\mathbf{p}_{a,b}) < V_{b,a}(x_1, x_2)$ or $V_{a,b}(\mathbf{p}_{b,a}) < V_{a,b}(x_2, x_3)$, the sequence $\{x_n\}$ is periodic or densely fills exactly one interval of \mathbb{R}^+ .

5 Some properties of the rotation number function

From Theorem 2 it is natural to introduce the *rotation number function* for $F_{b,a}$:

$$\rho_{b,a} : [h_c, \infty) \longrightarrow (0, 1),$$

where $h_c := V_{b,a}(\mathbf{p})$, as the map that associates to each invariant oval $\{V_{b,a}(x, y) = h\}$, the rotation number $\rho_{b,a}(h)$ of the function $F_{b,a}$ restricted to it. The following properties hold:

(i) The function $\rho_{b,a}(h)$ is analytic for $a > 0, b > 0, h > h_c$ and it is continuous at $h = h_c$. This can be proved from the tools introduced in [6, Sec. 4].

(ii) The value $\rho_{b,a}(h_c)$ is given by the argument over 2π of the eigenvalues (which have modulus one due to the integrability of $F_{b,a}$) of the differential of $F_{b,a}$ at \mathbf{p} .

(iii) $\rho_{b,a}(h) = \rho_{a,b}(h)$.

(iv) $\rho_{a,a}(h) = 2\rho_a(h) \pmod{1}$, where ρ_a is the rotation number* function associated to the classical Lyness map. Then, from the results of [4] we know that $\rho_{1,1}(h) \equiv 3/5$, that for $a \neq 1$, positive, $\rho_{a,a}(h)$ is monotonous and $\lim_{h \rightarrow \infty} \rho_{a,a}(h) = 3/5$.

Note that item (iii) follows because $F_{a,b}$ is conjugated with $F_{b,a}$ via $\psi = F_a$ which is a diffeomorphism of Q^+ , because $\psi^{-1}F_{a,b}\psi = F_a^{-1}F_aF_bF_a = F_bF_a = F_{b,a}$. Since ψ preserves the orientation, the rotation number functions of $F_{a,b}$ and $F_{b,a}$ restricted to the corresponding ovals must coincide.

Similar results to the ones given above hold for $F_{c,b,a}$ and its corresponding rotation number function. Some obvious differences are:

$$\begin{aligned} \rho_{c,b,a}(h) &= \rho_{b,a,c}(h) = \rho_{a,c,b}(h) & \rho_{a,a,a}(h) &= 3\rho_a(h) \pmod{1}, \\ \rho_{1,1,1}(h) &= 2/5, & \lim_{h \rightarrow \infty} \rho_{a,a,a}(h) &= 2/5. \end{aligned}$$

*Notice that given a map of the circle there is an ambiguity between ρ and $1 - \rho$ when one considers its rotation number. So, while for us the rotation number of the classical Lyness map for $a = 1$ is $4/5$, in other papers it is computed as $1/5$.

We are convinced that when $a > 0$ and $b > 0$,

$$\lim_{h \rightarrow \infty} \rho_{b,a}(h) = 3/5 \quad \text{and} \quad \lim_{h \rightarrow \infty} \rho_{c,b,a}(h) = 2/5,$$

but we have not been able to prove these equalities. If they were true, by combining them with the values of the rotation number function at $h = h_c$ this would give a very useful information to decide if, apart from the trivial cases $a = b = 1$ ($c = 1$), there are other cases for which the rotation number function is constant. Notice that in these situations the maps $F_{b,a}$ or $F_{c,b,a}$ would be globally periodic in Q^+ . This information, together with the values at h_c , also would be useful to know the regions where the corresponding functions could be increasing or decreasing. Finally notice that this rotation number at infinity is not continuous when we approach to $a = 0$ or $b = 0$, where the recurrence and the first integral are also well defined on Q^+ . For instance $\rho_{0,0}(h) \equiv 2/3$ and the numerical experiments of next subsection seem to indicate that for $a > 0$ or $b > 0$,

$$\lim_{h \rightarrow \infty} \rho_{0,a}(h) = \lim_{h \rightarrow \infty} \rho_{b,0}(h) = 5/8.$$

Before proving Theorem 3 we introduce with an example the algorithm that we will use along this section to compute lower and upper bounds for the rotation number. We have implemented it in an algebraic manipulator. Notice also that when we apply it taking rational values of a and b and rational initial conditions, it can be used as a method to achieve proofs, see next example or the proof of Theorem 3.

Fix $a = 3$, $b = 2$ and $(x_0, y_0) = (1, 1)$. Then $h = V_{2,3}(1, 1) = 34$. Compute for instance the 27 points of the orbit starting at $(1, 1)$,

$$(x_1, y_1) = (4, 6), \quad (x_2, y_2) = \left(\frac{9}{4}, \frac{17}{24}\right), \quad (x_3, y_3) = \left(\frac{89}{54}, \frac{788}{153}\right), \dots$$

and consider them as points on the oval $\{V_{2,3}(x, y) = 34\}$, see Figure 1.

We already know that the restriction of $F_{2,3}$ to the given oval is conjugated to a rotation, with rotation number $\rho := \rho_{2,3}(34)$ that we want to estimate. This can be done by counting the number of turns that give the points (x_j, y_j) , after fixing some orientation in the closed curve. We orientate the curve with the counterclockwise sense. So, for instance we know that the second point (x_2, y_2) has given more than one turn and less than two, giving that $1 < 2\rho < 2$, and hence that $\rho \in (1/2, 1)$. Doing the same reasoning with all the points




Figure 1: Oval of $\{V_{2,3}(x, y) = 34\}$ with 27 iterates of $F_{2,3}$. The label 0 indicates the initial condition $(1, 1)$, and the label $k, k = 1, \dots, 26$, corresponds with the k -th point of the orbit.

computed we obtain,

$$\begin{aligned}
 4 < 7\rho < 5 & \Rightarrow \rho \in \left(\frac{4}{7}, \frac{5}{7}\right), \\
 8 < 14\rho < 9 & \Rightarrow \rho \in \left(\frac{8}{14}, \frac{9}{14}\right), \\
 10 < 19\rho < 11 & \Rightarrow \rho \in \left(\frac{10}{19}, \frac{11}{19}\right), \\
 14 < 26\rho < 15 & \Rightarrow \rho \in \left(\frac{14}{26}, \frac{15}{26}\right),
 \end{aligned}$$

where we have only written the more relevant information obtained, which are given by the points of the orbit closer to the initial condition. So, we have shown that

$$0.5714 \simeq \frac{4}{7} < \rho_{2,3}(34) < \frac{15}{26} \simeq 0.5769.$$

In Figure 2 we represent several successive lower and upper approximations obtained while the orbit is turning around the oval. We plot around six hundred steps, after skipping the first fifty ones. By taking 1000 points we get

$$0.5761246 \simeq \frac{338}{578} < \rho_{2,3}(34) < \frac{473}{821} \simeq 0.5761267,$$

and after 3000 points,

$$0.57612457 \simeq \frac{338}{578} < \rho_{2,3}(34) < \frac{1472}{2555} \simeq 0.57612524.$$




Figure 2: Lower and upper bounds for $\rho_{2,3}(34)$ obtained after following some points of the orbit starting at $(1, 1)$.

In fact when we say that $\rho_{2,3}(34) \in (\rho_{\text{low}}, \rho_{\text{upp}})$, the value ρ_{low} is the upper lower bound obtained by following all the considered points of the orbit, and ρ_{upp} is the lowest upper bound. Notice that taking 1000 or 3000 points we have obtained the same lower bound for $\rho_{2,3}(34)$.

Let us prove Theorem 3 by using the above approach.

Proof of Theorem 3. Consider $a = 1/2$, $b = 3/2$ and the three points

$$\mathbf{p}^1 = \left(\frac{149}{100}, \frac{173}{100} \right), \quad \mathbf{p}^2 = \left(\frac{3}{40}, \frac{173}{100} \right), \quad \mathbf{p}^3 = \left(\frac{1}{1000}, \frac{173}{100} \right).$$

Notice that

$$\begin{aligned} h_1 &:= V_{3/2, 1/2}(\mathbf{p}^1) = \frac{10655559}{1288850} \simeq 8.27, \\ h_2 &:= V_{3/2, 1/2}(\mathbf{p}^2) = \frac{9328327}{207600} \simeq 44.93, \\ h_3 &:= V_{3/2, 1/2}(\mathbf{p}^3) = \frac{1056238343}{346000} \simeq 3052.71. \end{aligned}$$

Hence $h_c < h_1 < h_2 < h_3$. By applying the algorithm described above, using 100 points of each orbit starting at each \mathbf{p}^j , $j = 1, 2, 3$, we obtain that

$$\rho_{3/2, 1/2}(h_1), \rho_{3/2, 1/2}(h_3) \in \left(\frac{3}{5}, \frac{59}{98} \right) \quad \text{and} \quad \rho_{3/2, 1/2}(h_2) \in \left(\frac{56}{93}, \frac{53}{88} \right).$$

Since $59/98 < 56/93$ we have proved that the function $\rho_{3/2, 1/2}(h)$ has at least a local maximum in (h_1, h_3) . From the continuity of the rotation number function, with respect a, b and h , we notice that this result also holds for all values of a and b in a neighborhood of $a = 1/2, b = 3/2$. ■

We believe that with the same method it can be proved that a similar result to the one given in Theorem 3 holds for some maps $F_{c,b,a}$, but we have decided to not perform this study.

5.1 Some numerical explorations for $k = 2$.

We start by studying with more detail the rotation number function $\rho_{3/2,1/2}(h)$, that we have considered to prove Theorem 3. In this case the fixed point is $\mathbf{p} \simeq (1.493363282, 1.730133891)$ and $h_c = V_{b,a}(\mathbf{p}) \simeq 8.267483381$. Moreover $\rho_{b,a}(h_c) \simeq 0.6006847931$. By applying our algorithm for approximating the rotation number, with 5000 points on each orbit, we obtain the results presented in Table 1. In Figure 3 we also plot the upper and lower bounds of $\rho_{3/2,1/2}(h)$ that we have obtained by using a wide range of values of h .

Init. cond. (x, \bar{y})	Energy level h	$\rho_{\text{low}}(h)$	$\rho_{\text{upp}}(h)$
\bar{x}	$h_c \simeq 8.2675$	$\simeq 0.6006848$	$\simeq 0.6006848$
1.3	8.3068	$\frac{173}{288} \simeq 0.6006944$	$\frac{2938}{4891} \simeq 0.6006951$
0.75	9.2747	$\frac{1435}{2388} \simeq 0.6009213$	$\frac{2087}{3473} \simeq 0.6009214$
0.3	14.7566	$\frac{1548}{2573} \simeq 0.6016323$	$\frac{2285}{3798} \simeq 0.6016324$
0.075	44.9347	$\frac{657}{1091} \simeq 0.6021998$	$\frac{2354}{3909} \simeq 0.6022001$
0.001	3052.75	$\frac{2927}{4867} \simeq 0.6013972$	$\frac{86}{143} \simeq 0.6013986$
$5 \cdot 10^{-6}$	609716.07	$\frac{1832}{3049} \simeq 0.6008527$	$\frac{1409}{2345} \simeq 0.6008529$
$5 \cdot 10^{-256}$	$6.097 \cdot 10^{255}$	$\frac{3}{5} = 0.6$	$\frac{2999}{4998} \simeq 0.6000400$

Table 1: Lower and upper bounds of the rotation number $\rho_{3/2,1/2}(h)$, for some orbits of $F_{3/2,1/2}$ starting at (x, \bar{y}) , where $\mathbf{p} = (\bar{x}, \bar{y})$.

For other values of a and b we obtain different behaviors. All the experiments are performed by starting at the fixed point $\mathbf{p} = (\bar{x}, \bar{y})$, and increasing the energy level by taking initial conditions of the form (x, \bar{y}) , by decreasing x to 0. With this process we take orbits approaching to the boundary of Q^+ , that is lying on level sets of $V_{b,a}$ with increasing energy. The step in the decrease of x (and therefore in the increase of h) is not uniform, and it has been manually tuned making it smaller in those regions where a possible non monotonous behavior could appear.

Consider the set of parameters $\Gamma = \{(a, b), \in [0, \infty)^2\}$, where notice that we also consider the boundaries $a = 0$ or $b = 0$, where the map $F_{b,a}$ is well defined. We already know that the rotation number function behaves equal at (a, b) and (b, a) . Moreover we know perfectly its behavior on the diagonal (a, a) (when $a < 1$ it is monotonous decreasing and when $a > 1$ it




Figure 3: Lower and upper bounds for $\rho_{3/2, 1/2}(h)$. On the horizontal axis we represent $-\log_{10}(h)$ and, on the vertical axis, the value of the rotation number. Notice that for values of $-\log_{10}(h)$ smaller than 70 both values are indistinguishable in the Figure.

is monotonous increasing) and that $\rho_{1,1}(h) \equiv 4/5$ and $\rho_{0,0}(h) \equiv 2/3$. Hence a good strategy for a numerical exploration can be to produce sequences of experiments using our algorithm by fixing some $a \geq 0$ and varying b . For instance we obtain:

- Case $a = 1/2$. For all the values of $b > 0$ considered, the rotation number function seems to tend to $3/5$ when h goes to infinity. Moreover it seems
 - monotone decreasing for $b \in \{1/4, 1\}$;
 - to have a unique maximum when $b \in \{7/5, 3/2\}$;
 - monotone increasing for $b \in \{2, 3\}$.
- Case $a = 0$. For all the values of $b > 0$ considered, the rotation number function seems to tend to $5/8$ when h goes to infinity. Moreover it seems
 - monotone decreasing for $b \in \{1/10, 3/10, 1/2\}$;
 - to have a unique maximum when $b \in \{7/10, 3/4\}$;
 - monotone decreasing for $b \in \{1, 5\}$.

The above results, together with some other experiments for other values of a and b , not detailed in this paper, indicate the existence of a subset of positive measure in Γ where the

corresponding rotation number functions seem to present a unique maximum. This subset probably separates two other subsets of Γ , one where $\rho_{b,a}(h)$ is monotonically decreasing to $3/5$, and another one where $\rho_{b,a}(h)$ increases monotonically to the same value. The “oscillatory subset” seems to shrink to $(a, b) = (1, 1)$ when it approaches to the line $a = b$ and seems to finish in one interval on each of the borders $\{a = 0\}$ and $\{b = 0\}$. Further analysis must be done in this direction in order to have a more accurate knowledge of the bifurcation diagram associated to the behavior of $\rho_{b,a}$ on Γ .

Acknowledgement. The authors are sincerely grateful to the referee for her/his valuable comments that helped to simplify several points of the work.

References

- [1] W.J. Beyn, T. Hüls, M.Ch. Samtenschneider. *On r -periodic orbits of k -periodic maps*, J. Difference Equ. Appl. 14 (2008), 865–887.
- [2] G. Bastien, M. Rogalski. *Global behavior of the solutions of Lyness’ difference equation $u_{n+2}u_n = u_{n+1} + a$* , J. Difference Equ. Appl. 10 (2004), 977–1003.
- [3] G. Bastien, M. Rogalski. *On the algebraic difference equations $u_{n+2}u_n = \psi(u_{n+1})$ in \mathbb{R}_*^+ , related to a family of elliptic quartics in the plane*, Adv. Difference Equ. 2005:3 (2005), 227–261.
- [4] F. Beukers, R. Cushman. *Zeeman’s monotonicity conjecture*, J. Differential Equations 143 (1998), 191–200.
- [5] E. Camouzis, G. Ladas. *Dynamics of third-order rational difference equations with open problems and conjectures*. Advances in discrete mathematics and applications vol. 5. Chapman & Hall, Boca Raton FL, 2008.
- [6] A. Cima, A. Gasull, V. Mañosa. *Dynamics of the third order Lyness’ difference Equation*, J. Difference Equ. Appl. 13 (2007), 855-884.
- [7] A. Cima, A. Gasull and V. Mañosa. *Studying discrete dynamical systems through differential equations*, J. Differential Equations 244 (2008), 630–648.
- [8] A. Cima, A. Gasull and V. Mañosa. *On k -periodic Lyness difference equations*. In preparation 2010.
- [9] C.A. Clark, E.J. Janowski, M.R.S. Kulenović. *Stability of the Gumowski–Mira equation with period–two coefficient*, Int. J. Bifurcations & Chaos 17 (2007), 143–152.

- [10] V. de Angelis. *Notes on the non-autonomous Lyness equation*, J. Math. Anal and Appl. 307, 292–304 (2005).
- [11] S. Elaydi, R.J. Sacker. *Global stability of periodic orbits of non-autonomous difference equations and population biology*, J. Differential Equations 208, 258–273 (2005).
- [12] S. Elaydi, R.J. Sacker. *Periodic difference equations, population biology and Cushing–Henson conjectures*, Math. Biosciences 201, 195–207 (2006).
- [13] E.J. Janowski, M.R.S. Kulenović, Z. Nurkanović. *Stability of the k th order Lyness’ equation with period- k coefficient*, Int. J. Bifurcations & Chaos 17, 143–152 (2007).
- [14] D. Jogia, J. A. G. Roberts, F. Vivaldi, *An algebraic geometric approach to integrable maps of the plane*, J. Phys. A 39, 1133–1149 (2006).
- [15] M.R.S. Kulenović, Z. Nurkanović. *Stability of Lyness’ equation with period-three coefficient*, Radovi Matematički 12, 153–161 (2004).
- [16] J. Esch and T.D. Rogers, *The screensaver map: dynamics on elliptic curves arising from polygonal folding*, Discrete Comput. Geom. 25, 477–502 (2001).
- [17] J. Feuer, E.J. Janowski, G. Ladas. *Invariants for some rational recursive sequence with periodic coefficients*, J. Difference Equations and Appl. 2, 167–174 (1996).
- [18] E.A. Grove, C.M. Kent and G. Ladas. *Boundedness and persistence of nonautonomous Lyness and Max equations*, J. Difference Equations and Appl. 3, 241–258 (1998).
- [19] E. C. Zeeman. *Geometric unfolding of a difference equation*, Hertford College, Oxford (1996). Unpublished.