

# Heteroclinic, homoclinic and closed orbits in the Chen system

G. Tigan, J. Llibre

the date of receipt and acceptance should be inserted later

**Abstract** Bounded orbits such as closed, homoclinic and heteroclinic orbits are discussed in this work for a Lorenz-like 3D nonlinear system. For a large spectrum of the parameters the system has neither closed nor homoclinic orbits but has exactly two heteroclinic orbits, while under other constraints the system has symmetrical homoclinic orbits.

**Keywords** ODE systems · bifurcations · homoclinic and heteroclinic orbits

## 1 Introduction

In this paper we consider a Lorenz-like three-dimensional system [13], namely the Chen system. Interesting results reported recently on this system concerning closed, homoclinic and heteroclinic orbits are found in [7]. More results on the Chen system are given in [3], [8], [9], [10], [11], [14], [17] and in some references therein.

In this paper we first refine some results reported in [7] and present a different proof of these results. Secondly, we consider an important case not treated in [7] and show the system has homoclinic orbits for a large spectrum of the parameters, by transforming the system into a new form and using results reported in [1]. Proving existence of homoclinic or heteroclinic orbits in nonlinear ode systems is in general a difficult task [2], [5], [6], [12], [15], [16].

The Chen system is given by:

$$\dot{x} = a(y - x), \dot{y} = (c - a)x - xz + cy, \dot{z} = xy - bz, \quad (1)$$

where  $a > 0$ ,  $b > 0$  and  $c > 0$  are positive real parameters.

G. Tigan

Department of Mathematics, Politehnica University of Timisoara, Romania, E-mail: gheorghe.tigan@upt.ro,

J. Llibre

Department of Mathematics, University Autònoma of Barcelona, Catalonia, Spain, E-mail: jllibre@mat.uab.cat

Recall that, if  $x_0$  is an equilibrium hyperbolic point such that the stable manifold  $W^s(x_0)$  intersected with the unstable manifold  $W^u(x_0)$  is not empty, then the orbits belonging to  $W^s(x_0) \cap W^u(x_0) \neq \Phi$  are called homoclinic orbits. They are doubly asymptotic to an equilibrium point. Similarly, if  $x^1, x^2$  are two hyperbolic equilibrium points such there exists an orbit  $\Gamma \subset W^s(x^1) \cap W^u(x^2)$  or inversely  $\Gamma \subset W^u(x^1) \cap W^s(x^2)$ , then  $\Gamma$  is called a hyperbolic orbit.

The system (1) has the origin  $O$  as an equilibrium point for any  $a, b, c > 0$  and it has two more equilibrium points

$$S_1 = (\sqrt{b(2c - a)}, \sqrt{b(2c - a)}, 2c - a)$$

and

$$S_2 = (-\sqrt{b(2c - a)}, -\sqrt{b(2c - a)}, 2c - a),$$

for  $2c > a$ . Assume also further  $a > c$ . As the transformation  $(x, y, z) \rightarrow (-x, -y, z)$  leaves the system unchanged, the orbits of the system are symmetrical with respect to the  $z$ -axis.

## 2 Existence of heteroclinic orbits

The matrix of the linearized system to (1) at the origin has the eigenvalues and the corresponding eigenvectors given by:

$$d_1 = \frac{1}{2}c - \frac{1}{2}a + \frac{1}{2}\sqrt{6ac - 3a^2 + c^2},$$

$$u_1 = \left( \frac{1}{2a-2c} \left( a + c - \sqrt{6ac - 3a^2 + c^2} \right) \ 1 \ 0 \right)^T;$$

$$d_2 = \frac{1}{2}c - \frac{1}{2}a - \frac{1}{2}\sqrt{6ac - 3a^2 + c^2},$$

$$u_2 = \left( \frac{1}{2a-2c} \left( a + c + \sqrt{6ac - 3a^2 + c^2} \right) \ 1 \ 0 \right)^T;$$

$$d_3 = -b,$$

$$u_3 = (0 \ 0 \ 1)^T.$$

Considering  $2c > a > c > 0$ , it follows that  $d_1 > 0$  and  $d_{2,3} < 0$ , that is, the origin is a saddle point having a one-dimensional unstable manifold  $W_0^u$  and a two-dimensional stable manifold  $W_0^s$ . The tangent unstable subspace  $TW_0^u$  is

given by

$$TW_0^u = \left\{ 2(a-c)y = \left( a+c - \sqrt{6ac - 3a^2 + c^2} \right) x, z = 0 \right\}.$$

The unstable manifold  $W_0^u$  contains  $O(0,0,0)$  and is tangent to  $TW_0^u$  at the origin.

Using the method of undetermined coefficients in a small neighborhood of the origin we get

$$W_0^u = \left\{ (x, y, z) \in \mathbb{R}^3 \mid \begin{array}{l} y = a_1 x + O(x^2) \\ z = \frac{a_1}{-2a+b+2aa_1} x^2 + O(x^3) \end{array} \right\},$$

where  $|x| \ll 1$  and  $a_1 = \frac{1}{2(a-c)} \left( a+c - \sqrt{6ac - 3a^2 + c^2} \right)$ .

Note that  $W_0^u$  is indeed tangent to  $TW_0^u$  since  $z'(0) = 0$ ,  $y'(0) = a_1$  and the vector  $(1, y'(0), z'(0))^T$  is collinear to the direction vector  $u_1$  of the line  $TW_0^u$ . Note also that the  $z$ -axis is included in the stable manifold  $W_0^s$ .

Denote in the following by  $\phi_t u_0 = (x(t, u_0), y(t, u_0), z(t, u_0))$  a solution of the system (1) through the initial point  $u_0 = (x_0, y_0, z_0)$  and by  $W_+^u$  ( $W_-^u$ ) the positive (negative) branch of the unstable manifold  $W_0^u$  corresponding to  $x > 0$  ( $x < 0$ ).

Define a Lyapunov-like function

$$U(x, y, z) = A(y-x)^2 + B(z-x^2/b)^2 + C(x^2 - b(2c-a))^2.$$

Choosing

$$A = b - 2a \geq 0, B = b > 0 \text{ and } C = \frac{1}{2ab} (b - 2a) \geq 0,$$

it follows that

$$\frac{dU}{dt} = -2(b-2a)(a-c)(x-y)^2 - 2(bz-x^2)^2 \leq 0. \quad (2)$$

*Remark 1* Different from [7], the Lyapunov function  $U$  is defined both for  $b > 2a$  and  $b = 2a$ .

**PROPOSITION 1** *If  $2c > a > c > 0$ ,  $b \geq 2a$  the following assertions are true:*

a) *If there exist  $t_1$  and  $t_2$  such that  $t_1 < t_2$  and  $U$  satisfies  $U(\phi_{t_1} u_0) = U(\phi_{t_2} u_0)$ , then either*

- $u_0$  is an equilibrium point of system (1), or
- $b = 2a$  and the orbit  $\phi_t u_0$  is contained in the parabolic cylinder  $bz = x^2$ .

b) *Assume  $b > 2a$ . If  $\phi_t u_0 \rightarrow O$  as  $t \rightarrow -\infty$  and  $x(t, u_0) > 0$  for some  $t$ , then*

$$U(O) > U(\phi_t u_0) \text{ and } x(t, u_0) > 0, \text{ for all } t \in \mathbb{R}.$$

Consequently  $u_0 \in W_+^u$ .

*Proof* a) From (2) and from the hypothesis of a), one gets  $\frac{dU}{dt}(\phi_t u_0) \equiv 0$  for all  $t \in (t_1, t_2)$ , which implies either

$$\dot{x}(\phi_t u_0) = \dot{y}(\phi_t u_0) = \dot{z}(\phi_t u_0) \equiv 0 \quad (3)$$

for all  $t \in (t_1, t_2)$ , i.e.  $u_0$  is one of the equilibria of (1), or  $b > 2a$  and the orbit  $\phi_t u_0$  is contained in the intersection of the plane  $x = y$  with the parabolic cylinder  $bz = x^2$ . But this latter case leads again to (3), i.e.  $u_0$  is one of the equilibria of system (1), because from  $\phi_t u_0 \in \{x = y\} \cap \{bz = x^2\}$  for all  $t$ , we get  $\dot{x}(t, u_0) = 0$ ,  $\dot{z}(t, u_0) = 0$ . Hence  $x(t) = x_0$ , but  $y(t) = x(t)$  for all  $t$ , i.e.  $\dot{y}(t, u_0) = 0$ . We notice that the three

equilibrium points lie on the non-invariant curve  $\{x = y\} \cap \{bz = x^2\}$ .

Finally, (2) and the hypothesis of a) lead also to  $b = 2a$  and the orbit  $\phi_t u_0$  is contained in the parabolic cylinder  $bz = x^2$  (for these values of the parameters this cylinder is invariant by the flow of system (1), i.e. if an orbit has a point in it the whole orbit is contained in the cylinder).

b) We prove first  $U(O) > U(\phi_t u_0)$  for all  $t \in \mathbb{R}$ ;  $U(O) > 0$ . To this end, assume by contrary that there exists a  $t_0 \in \mathbb{R}$  such that  $0 < U(O) \leq U(\phi_{t_0} u_0)$ . From  $\phi_t u_0 \rightarrow O$  as  $t \rightarrow -\infty$  and  $U$  continuous on  $t$ , it follows that there exists a sequence  $t_n \rightarrow -\infty$  and an integer positive number  $n_1$  such that  $|U(\phi_{t_n} u_0) - U(O)| < \varepsilon$  for all  $\varepsilon > 0$  and  $n > n_1$ . Since  $t_n \rightarrow -\infty$  and  $t_0 \in \mathbb{R}$ , there is an integer positive number  $n_2$  such that  $t_n < t_0$  for all  $n > n_2$ . Denote further by  $n_0 = \max\{n_1, n_2\}$  and take  $\varepsilon = \frac{1}{2}(U(\phi_{t_0} u_0) - U(O))$ . It is clear that  $\varepsilon \geq 0$ . Then  $U(\phi_{t_n} u_0) - U(\phi_{t_0} u_0) = U(\phi_{t_n} u_0) - U(O) + U(O) - U(\phi_{t_0} u_0) < \varepsilon + U(O) - U(\phi_{t_0} u_0) = -\varepsilon \leq 0$ . On the other hand  $U(t)$  is decreasing with respect to  $t$ , which, by definition, leads to  $U(\phi_{t_n} u_0) \geq U(\phi_{t_0} u_0)$  for all  $t_n < t_0$  and  $n > n_0$ . Therefore,  $U(\phi_{t_n} u_0) = U(\phi_{t_0} u_0)$  and by virtue of a) we get that  $u_0$  is an equilibrium point of system (1). Since  $\phi_t u_0 \rightarrow O$  we get  $u_0 \equiv O$  and  $x(t, u_0) = 0$  for all  $t$ . But this contradicts the hypothesis  $x(t, u_0) > 0$  for some  $t$ . Hence,  $U(O) > U(\phi_t u_0)$  for all  $t \in \mathbb{R}$ .

Let us prove now that  $x(t, u_0) > 0$  for all  $t \in \mathbb{R}$ . Assuming that there exists a  $t' \in \mathbb{R}$  such that  $x(t', u_0) \leq 0$  and using  $x(t'', u_0) > 0$  for some  $t'' \in \mathbb{R}$  from the hypothesis of b), one gets that there exists a  $\tau \in \mathbb{R}$  such that  $x(\tau, u_0) = 0$ . As  $U(O) > U(\phi_t u_0)$  for all  $t \in \mathbb{R}$ , it follows that  $\phi_\tau u_0 \in \Omega \cap P$ , where  $\Omega = \{(x, y, z) : U(O) > U(x, y, z)\}$  and  $P$  is the plane  $x = 0$ . On the other hand,  $\Omega \cap P$  is given by  $Ay^2 + Bz^2 + Cb^2(2c-a)^2 < Cb^2(2c-a)^2$ , i.e.  $Ay^2 + Bz^2 < 0$  with  $A, B \geq 0$ . It leads to  $\Omega \cap P = \emptyset$  which is a contradiction. Therefore  $x(t, u_0) > 0$  for all  $t \in \mathbb{R}$ . This completes the proof of the proposition.

**Theorem 1** *Consider  $2c > a > c > 0$ ,  $b > 2a$  and the above function  $U$ . Then the following assertions are true:*

a) *The  $\omega$ -limit of any trajectory of system (1) is an equilibrium point. In particular system (1) has no closed trajectories.*

b) *System (1) has no homoclinic trajectories.*

c) *System (1) has exactly two heteroclinic trajectories.*

*Proof* While the function  $U$  is different, the proof is similar to the one presented in [7]. However, we choose to present it here as it may be a useful handy exercise for some readers.

a) If  $a > c > 0$ , the function  $U$  is decreasing along trajectories of the system both for  $b > 2a$  and  $b = 2a$ , except perhaps for the orbits on the cylinder  $bz = x^2$  if  $b = 2a$ , and the orbits contained in  $\{x = y\} \cap \{bz = x^2\}$  if  $b > 2a$ , where the function  $U$  is constant and equal to zero. This implies that

for all  $t \geq 0$

$$0 \leq U(\phi_t u_0) \leq U(u_0), \quad (4)$$

where  $\phi_t u_0$  is a trajectory of the system through the initial point  $u_0$ . It implies that, the limit  $\lim_{t \rightarrow \infty} U(\phi_t u_0)$  exists. Denote it by  $U^*(u_0)$ . From (4) one gets that  $U(\phi_t u_0)$  is bounded for  $t \geq 0$ , which implies further that  $x(t, u_0), y(t, u_0), z(t, u_0)$  are bounded for  $t \geq 0$ , i.e.  $\phi_t u_0$  is bounded for  $t \geq 0$ . Denote by  $\Omega(u_0)$  the  $\omega$ -limit set of the orbit  $\phi_t u_0$ . It is known that, if  $u \in \Omega(u_0)$ , then all points of the orbit through  $u$  belong to  $\Omega(u_0)$ , i.e.  $\phi_t u \in \Omega(u_0)$ . Therefore, for any point  $\phi_t u, t \geq 0$ , there exists a sequence  $t_n \rightarrow \infty$  for  $n \rightarrow \infty$  such that  $\lim_{n \rightarrow \infty} \phi_{t_n} u_0 = \phi_t u$  which leads to

$$U(\phi_t u) = \lim_{n \rightarrow \infty} U(\phi_{t_n} u_0) = U^*(u_0) = \text{const},$$

for all  $t \geq 0$ . So there exists  $t_1 < t_2$  such that  $U(\phi_{t_1} u) = U(\phi_{t_2} u)$  for all  $t \in (t_1, t_2)$ , and by Proposition 1 either  $u$  is one of the equilibria of the system, or  $b = 2a$  and the point  $u$  is contained in the invariant cylinder  $bz = x^2$ , or  $b > 2a$  and the point  $u$  is contained in the curve  $\{x = y\} \cap \{bz = x^2\}$ .

Assume  $b = 2a$  and the point  $u$  is contained in the invariant cylinder  $bz = x^2$ . Then the function  $U$  takes the value zero on this cylinder, and eventually the system can have periodic orbits on the cylinder.

Assume  $b > 2a$  and  $u \in \{x = y\} \cap \{bz = x^2\}$ . Since in the connected curve  $\{x = y\} \cap \{bz = x^2\}$  there are the three equilibria of the system, and  $u$  is an  $\omega$ -limit set,  $u$  is one of the equilibria of the system by the Bendixson-Poincaré Theorem (see for instance Corollary 1.30 of [4]).

b) Assume that the system has a homoclinic orbit  $\gamma(t)$  at one of the equilibria  $O, S_1$  or  $S_2$ , that is,  $\lim_{t \rightarrow \pm\infty} \gamma(t) = q$  where  $q \in \{O, S_1, S_2\}$ . Since  $U$  is decreasing along the trajectories of the system, it follows that

$$U(q) \leq U(\gamma(t)) \leq U(q), \quad (5)$$

c) By statement a) every one-dimensional branch of the unstable manifold  $W^u$  has  $\omega$ -limit an equilibrium point  $p$ .


Assume  $b > 2a$ . Since  $U(O) > U(S_i) = 0$  for  $i = 1, 2$ , the equilibrium point  $p$  must be either  $S_1$  or  $S_2$ , and by the symmetry of the system with respect to the  $z$ -axis, one of the two branches of  $W^u$  must go to  $S_1$  and the other to  $S_2$ , obtaining in this way two heteroclinic orbits. A numerical case with two heteroclinic orbits is illustrated in Fig.1 a). This completes the proof of the theorem.

The case  $b = 2a$  has been studied in [7] using a Lyapunov-like function in the form

$$V(x, y, z) = a^2 (y - x)^2 + \frac{1}{4} (x^2 - 2a(2c - a))^2.$$

In particular, the following results have been reported:

a) If the negative orbit from a point  $u_0$  is bounded, then the solution  $\phi_t u_0$  approaches one of the equilibria of the system as  $t \rightarrow -\infty$ . Consequently, the system has no closed orbits.



**Fig. 1** a) Two symmetrical heteroclinic orbits for  $a = 1, b = 2.1, c = 0.8$ . (up); b) Two symmetrical homoclinic orbits for  $a = 1, b = 1.17, c = 0.8$  (down).

b) The system (1) has no homoclinic orbits.

c) The system (1) has exactly two heteroclinic orbits: one linking  $O$  to  $S_1$  and the other  $O$  to  $S_2$ .

### 3 Existence of homoclinic orbits

Consider in the following the case  $2c > a > c > 0, 2a > b > 0$  and make the nonsingular transformation:

$$u = \alpha x, \quad v = \beta(y - x), \quad w = \gamma \left( z - \frac{x^2}{2a} \right),$$

and the rescaling  $\tau = rt$ . Then the system (1) becomes:

$$\begin{aligned} \dot{u} &= \frac{a\alpha}{r\beta} v, \\ \dot{v} &= \frac{\beta}{2ar\alpha^3} \left( \frac{2a}{\gamma} \alpha^2 - \frac{2a\alpha^2}{\gamma} w - u^2 \right) u + \frac{c-a}{r} v, \\ \dot{w} &= \frac{u^2}{\alpha^2} \left( 1 - \frac{b}{2a} \right) \frac{\gamma}{r} - \frac{b}{r} w. \end{aligned} \quad (6)$$

Choosing

$$\alpha = \sqrt{\frac{1}{2a(2c-a)}} > 0, \quad \beta = \frac{\sqrt{2}}{2(2c-a)} > 0, \quad \gamma = \frac{1}{2c-a} > 0,$$

and  $r = \sqrt{a(2c-a)} > 0$ , system (6) reads:

$$\begin{aligned} \dot{u} &= v, \\ \dot{v} &= -(u^2 + w - 1)u - \lambda v, \\ \dot{w} &= -\sigma w + \delta u^2, \end{aligned} \quad (7)$$

where  $\lambda = \frac{a-c}{r}$ ,  $\sigma = \frac{b}{\sqrt{a(2c-a)}}$ ,  $\delta = (2a-b)\sqrt{\frac{1}{a(2c-a)}}$ .

Notice that  $\beta = 2ar\alpha^3 = \frac{a}{r}\alpha$ ,  $\gamma = 2a\alpha^2$  and  $r^2 = \frac{1}{2\alpha^2}$ . But system (7) is treated in [1] and after a laborious study and using a method based on comparison systems, the following result is reported:

**Theorem 2** *For each  $\sigma > 0$ , there is, in the region of positive parameters  $\delta$  and  $\lambda$ , a bifurcation curve  $\{\rho(\sigma, \delta, \lambda) = 0\}$ , beginning at  $(0, 0)$  and going to infinity for  $\delta \rightarrow \infty$ , corresponding to a homoclinic orbit to the saddle point  $O(0, 0, 0)$  of system (7).*

It implies that for any  $a, b, c$  with  $2c > a > c > 0$  and  $2a > b > 0$ , system (1) has two symmetrical homoclinic orbits to the equilibrium point  $O$ . A particular numerical case in this regard is presented in Fig.1 b) where the homoclinic orbits are depicted.

## 4 Conclusions

In this paper we have investigated closed, homoclinic and heteroclinic orbits in a three-dimensional autonomous system, known as the Chen system. Using a convenient Lyapunov-like function, we proved that the system under some constraints of its parameters has no homoclinic orbits and no closed orbits but it has exactly two heteroclinic orbits, symmetrically with respect to the  $z$ -axis. Moreover, transforming the system to a new form and using some known results from [1], we obtained results on cases not covered in [7] concerning homoclinic orbits. More exactly, we have proved that the system has two symmetrical homoclinic orbits to the equilibrium point  $O$ .

**Acknowledgements** This research was supported by FP7-PEOPLE-2012-IRSES-316338.

## References

1. V.N. Belykh, Bifurcations of separatrices of a saddle point of the Lorenz system, *Differential Equations*, 20(10) (1984), 1184-1191.
2. Y.Y. Cao, K.W. Chung, J. Xu, A novel construction of homoclinic and heteroclinic orbits in nonlinear oscillators by a perturbation-incremental method, *Nonlinear Dynamics*, 64(3) (2011), 221-236.
3. Y. Chang, G. Chen, Complex dynamics in Chen's system, *Chaos, Solitons and Fractals*, 27(2006), 75-86.
4. F. Dumortier, J. Llibre, J.C. Artés, *Qualitative theory of planar differential systems*, UniversiText, Springer-Verlag, New York, 2006.
5. S.V. Gonchenko, I.I. Ovsyannikov, D. Turaev, On the effect of invisibility of stable periodic orbits at homoclinic bifurcations, *Physica D*, 241(13), 2012, 1115-1122.
6. V. Grines et al., On heteroclinic separators of magnetic fields in electrically conducting fluids, *Physica D*, 294(15)(2015), 1-5.
7. T. Li, G. Chen, G. Chen, On homoclinic and heteroclinic orbits of the Chen's system, *IJBC*, 16(10)(2006), 3035-3041.
8. X-F. Li et al., Nonlinear dynamics and circuit realization of a new chaotic flow: A variant of Lorenz, Chen and Lü, *Nonlinear Analysis: Real World Applications*, 10(4)(2009), 2357-2368.
9. X. Li, P. Wang, Hopf bifurcation and heteroclinic orbit in a 3D autonomous chaotic system, *Nonlinear Dynamics*, 73(1-2)(2013), 621-632.
10. J. Llibre, M. Messias, P.R. da Silva, Global dynamics in the Poincaré ball of the Chen system having invariant algebraic surfaces, *IJBC*, 22(2012).
11. J. Llibre, A. Rodrigues, On the Dynamics of the Unified Chaotic System Between Lorenz and Chen Systems, *IJBC*, 25(2015).
12. A. Lohse, Stability of heteroclinic cycles in transverse bifurcations, *Physica D*, 310(2015), 95-103.
13. E.N. Lorenz, Deterministic nonperiodic flow, *J. Atmos. Sci.*, 20(1963), 130-141.
14. T. Lu, X. Zhang, Darboux polynomials and algebraic integrability of the Chen system, *IJBC*, 17(08)(2007), 2739-2748.
15. G. Tigan, D. Turaev, Analytical search for homoclinic bifurcations in the Shimizu-Morioka model, *Physica D*, 240(12)(2011), 985-989.
16. K. Yagasaki, T. Wagenknecht, Detection of symmetric homoclinic orbits to saddle-centres in reversible systems, *Physica D*, 214(2)(2006), 169-181.
17. T. Zhou, Y. Tang, G. Chen, Chen's attractor exists, *IJBC*, 14(9)(2004), 3167-3177.