

Rapid Molecular Characterization of *Acinetobacter baumannii* Clones with rep-PCR and Evaluation of Carbapenemase Genes by New Multiplex PCR in Hospital District of Helsinki and Uusimaa

Tanja Pasanen*, Suvi Koskela, Sointu Mero, Eveliina Tarkka, Päivi Tissari, Martti Vaara, Juha Kirveskari

Division of Clinical Microbiology, Helsinki University Hospital, Helsinki, Finland

Abstract

Multidrug-resistant *Acinetobacter baumannii* (MDRAB) is an increasing problem worldwide. Prevalence of carbapenem resistance in *Acinetobacter* spp. due to acquired carbapenemase genes is not known in Finland. The purpose of this study was to examine prevalence and clonal spread of multiresistant *A. baumannii* group species, and their carbapenemase genes. A total of 55 *Acinetobacter* isolates were evaluated with repetitive PCR (DiversiLab) to analyse clonality of isolates, in conjunction with antimicrobial susceptibility profile for ampicillin/sulbactam, colistin, imipenem, meropenem, rifampicin and tigecycline. In addition, a new real-time PCR assay, detecting most clinically important carbapenemase genes just in two multiplex reactions, was developed. The assay detects genes for KPC, VIM, IMP, GES-1/-10, OXA-48, NDM, GIM-1, SPM-1, IMI/NMC-A, SME, CMY-10, SFC-1, SIM-1, OXA-23-like, OXA-24/40-like, OXA-58 and ISAbal-OXA-51-like junction, and allows confident detection of isolates harbouring acquired carbapenemase genes. There was a time-dependent, clonal spread of multiresistant *A. baumannii* strongly correlating with carbapenemase gene profile, at least in this geographically restricted study material. The new carbapenemase screening assay was able to detect all the genes correctly suggesting it might be suitable for epidemiologic screening purposes in clinical laboratories.

Citation: Pasanen T, Koskela S, Mero S, Tarkka E, Tissari P, et al. (2014) Rapid Molecular Characterization of *Acinetobacter baumannii* Clones with rep-PCR and Evaluation of Carbapenemase Genes by New Multiplex PCR in Hospital District of Helsinki and Uusimaa. PLoS ONE 9(1): e85854. doi:10.1371/journal.pone.0085854

Editor: Igor Mokrousov, St. Petersburg Pasteur Institute, Russian Federation

Received: September 29, 2013; **Accepted:** December 2, 2013; **Published:** January 21, 2014

Copyright: © 2014 Pasanen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: As a preliminary notice, however, HUSLAB has informed us that they may not be able to commit to all publication costs unlike previously agreed due to bad economical situation. Therefore, another fundign source, including the authors' new employee Mobidiag Ltd must be considered. (No agreement from their side yet). If they decide to grant some portion of the total amount, it will be a personal hand-out to this academic study, and in such case it is fair to state: The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. For the sake of clarity, JK's affiliation has changed after submission of the original manuscript version. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: JK has filed a patent application related to test described in the paper. Mobidiag is a privately owned company developing and selling molecular diagnostics tests, including antibiotic resistance content. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: tanja.holma@hus.fi

Introduction

Acinetobacter baumannii is a hospital-acquired pathogen which commonly causes pneumonia, bloodstream infections, meningitis, wound infections and urinary tract infections, especially in patients with impaired host defences. *A. baumannii* isolates are resistant to many antimicrobial classes: fluoroquinolones, tetracyclines, cephalosporines and aminoglycosides [1]. However, today carbapenem resistance is more frequently encountered [1–3]. In *A. baumannii* carbapenem resistance is usually conferred by carbapenem-hydrolyzing class D oxacillinases (CHDLs), including OXA-23-like (*bla*_{OXA-23-like}), OXA-40-like (*bla*_{OXA-40-like}), OXA-58-like (*bla*_{OXA-58-like}), and OXA-143-like (*bla*_{OXA-143-like}) oxacillinases. Additionally *A. baumannii* has the intrinsic OXA-51-like (*bla*_{OXA-51-like}) oxacillinase [4,5]. Although CHDLs exhibit weak carbapenem hydrolysis, they can confer resistance when overexpressed. This resistance is mediated through a combination of naturally low permeability to β -lactams, efflux pumps and IS*Aba* elements located upstream of the gene, providing a strong promoter activity [6]. In addition, *A. baumannii* may harbour many other carbapenemases

more commonly found among *Enterobacteriaceae* and *Pseudomonas* species [7].

To determine genetic and epidemiological relatedness, genomic fingerprinting of clinical isolates is required. One of the most effective method is the repetitive extragenic palindromic sequence-based polymerase chain reaction (rep-PCR), which is commercially available known as the DiversiLab microbial typing system (bioMérieux, Marcy L'Etoile, France) [8]. This system has been proven useful in the typing of *A. baumannii* and has demonstrated good discriminatory ability, comparable with pulsed-field gel electrophoresis (PFGE) and multilocus sequence typing (MLST) [9,10]. Recently this rep-PCR typing system, DiversiLab, has identified eight carbapenem-resistant *A. baumannii* clonal lineages (WW1 to WW8) that are distributed worldwide [4]. DiversiLab fingerprints between laboratories were recently tested and clustering was found to be conserved [11].

The carbapenem resistance has recently attracted new interest as a subset among tens of gene families has spread to *Enterobacteriaceae* [12–14], despite a much longer history among *Pseudomonas* and *Acinetobacter* species. *A. baumannii* may harbour

most of the acquired carbapenemase genes within *Enterobacteriaceae*, and *Pseudomonas* in addition to their characteristics CDHL genes [7].

Recently, new molecular assays have been described to detect most prevalent carbapenemase genes [15], or a subset of *A. baumannii* selective carbapenemase genes. Due to limited gene set, or technical limitations, most new tests are not suitable for clinical routine monitoring in low prevalence settings [16]. In addition, combinations of other resistance mechanisms, such as reduced permeability due porin mutations, or defect, and efflux pumps in conjunction with ampC β -lactamases are the most common cause of carbapenem resistance in low prevalence areas [14]. Therefore, an imipenem hydrolysis test or dedicated MALDI-TOF [17] and more extensive screening of resistance mechanisms in a reference laboratory are often needed to reliably exclude carbapenemase genes.

The aim of this study was to investigate the carbapenemase genes of *A. baumannii* and the correlation between these genes and clonal lineages. The feasibility of a new real-time PCR assay was tested for screening of most important carbapenemase genes detected among *A. baumannii*, *Enterobacteriaceae*, and *Pseudomonas* species.

Materials and Methods

Bacterial strains and culture conditions

A total of 55 *Acinetobacter* isolates from 44 patients were detected. 51 isolates with reduced susceptibility to carbapenem from HUSLAB (Laboratory of Helsinki University Central Hospital) between Jun 18th 1993 and Jan 18th 2008 were collected and four *Acinetobacter* isolates susceptible to carbapenems were included as controls. Helsinki University Hospital is responsible for the secondary and tertiary care of app. 1.5 million people. The culture samples from this area received by HUSLAB are both from these hospitals as well as from outpatients of this geographical area, the Helsinki and Uusimaa district in southern Finland. The culture samples in this study were from patients treated in nine different hospitals (Table S1).

Acinetobacter isolates were cultured in aerobic atmosphere on chocolate and cysteine lactose electrolyte deficient (CLED) agar and incubated at 35°C for 18 h. Colonies with typical morphology and biochemistry were identified as *A. baumannii* complex. Identification with the VITEK 2 (bioMérieux, Marcy L'Etoile, France) system with GN card was performed, as well. 16S rRNA gene sequencing was performed when biochemical identification was equivocal. In addition a house-keeping OXA-51-like (*bla*_{OXA-51-like}) gene was detected separately within all the clinical isolates with reduced susceptibility to carbapenems, whereas carbapenem susceptible control strains did not harbour OXA-51-like genes.

Antimicrobial susceptibility testing was performed by the disk diffusion method according to the CLSI guidelines (<http://www.clsi.org>). MICs for ampicillin/sulbactam, colistin, imipenem, meropenem, rifampicin and tigecycline by E-test (AB BIODISC, Solna, Sweden) were determined on Mueller-Hinton agar according to manufacturer's instructions.

Design of multiplex Real-Time carbapenemase gene screening assay

The assay was designed to detect most clinically relevant carbapenemase genes described within *A. baumannii*, *Pseudomonas aeruginosa*, and *Enterobacteriaceae* species. The design was performed using AlleleID software (<http://www.premierbiosoft.com>), taking into account all the globally known sub-variants in NCBI data

Table 1. Description of validation isolates.

Target	Species	Isolation site	Travel history
GES-1	<i>P. aeruginosa</i>	wound	no
GES-14	<i>K. pneumoniae</i>	trachea	n/a
GES-5	<i>P. aeruginosa</i>	incision wound	no
GES-5	<i>P. aeruginosa</i>	incision wound	no
IMI-1	<i>E. cloacae</i>	stool	Thailand
IMI-2	<i>E. cloacae</i>	wound	no
IMP-15	<i>P. aeruginosa</i>	blood	no
IMP-15	<i>P. aeruginosa</i>	wound	no
IMP-15	<i>P. aeruginosa</i>	incision wound	no
IMP-15	<i>P. aeruginosa</i>	urine	no
IMP-15	<i>P. aeruginosa</i>	urine	n/a
IMP-15	<i>P. aeruginosa</i>	incision wound	n/a
IMP-15	<i>P. aeruginosa</i>	urine	n/a
ISAbal-OXA-51	<i>A. baumannii</i>	stool	Spain
ISAbal-OXA-51	<i>A. baumannii</i>	stool	no
ISAbal-OXA-51	<i>A. baumannii</i>	trachea	no
KPC	<i>K. pneumoniae</i>	stool	US
KPC-2	<i>K. pneumoniae</i>	stool	Greece
KPC-2	<i>K. pneumoniae</i>	wound	Italy
KPC-2	<i>K. pneumoniae</i>	urine	no
KPC-2	<i>K. pneumoniae</i>	blood	Mexico/US
KPC-2	<i>K. pneumoniae</i>	urine	no
NDM-1	<i>K. pneumoniae</i>	stool	n/a
OXA-23	<i>A. baumannii</i>	blood	n/a
OXA-23	<i>A. baumannii</i>	wound	no
OXA-23	<i>A. baumannii</i>	trachea	Thailand
OXA-48	<i>E. coli</i>	stool	Syria
OXA-48	<i>K. pneumoniae</i>	stool	Turkey
OXA-48	<i>A. baumannii</i>	stool	n/a
OXA-58	<i>A. baumannii</i>	stool	Tunis
OXA-58	<i>A. baumannii</i>	wound	no
OXA-58	<i>A. baumannii</i>	wound	no
OXA-58	<i>A. baumannii</i>	stool	Greece
OXA-58	<i>A. baumannii</i>	incision wound	n/a
OXA-58	<i>A. baumannii</i>	urine	no
SFC-1	<i>S. fonticola</i>	control strain	Portugal
SIM-1	<i>A. baumannii</i>	control strain	South-Korea
SME	<i>S. marcescens</i>	control strain	n/a
VIM	<i>P. aeruginosa</i>	stool	Thailand
VIM	<i>K. pneumoniae</i>	stool	Spain
VIM	<i>K. pneumoniae</i>	stool	Greece
VIM	<i>K. pneumoniae</i>	CV catheter	n/a
VIM-1	<i>K. pneumoniae</i>	blood	Greece
VIM-2	<i>P. aeruginosa</i>	trachea	Russia
VIM-2	<i>P. aeruginosa</i>	trachea	Russia

doi:10.1371/journal.pone.0085854.t001

base. For practical purposes, the assay was divided in two multiplex reactions consisting of nine and eight gene families, respectively. The assay was validated *in vitro* using 43 positive

Table 2. Primers used for amplification of resistance genes by polymerase chain reaction (PCR).

Primer	Sequence 5' - 3'	Reference	Oligomix
F_ges_001	ACACCTGGCGACC-TCAGAGATAC	This study	1
R_ges_001	ACTTGACCGACA-GAGGCAACTAATTC	This study	1
F_gim_001	CGAATGGGTTGGTAG-TTCTGGATAATAATC	This study	1
R_gim_001	ATGTGTATGTAGG-AATTGACTTTGA-ATTTAGC	This study	1
F_imi1_001	AAACAAGGGAA-TGGGTGGAGACTG	This study	1
R_imi1_001	AAGGTATGCTTT-GAATTTGCGTTG	This study	1
F_imp_10	AATAATGACGCCT-ATCTAATTGACACTCC	This study	1
R_imp_10	ATTCACCCGTA-CTGTCCGATG	This study	1
F_imp_11	TGACGCCTATCTG-ATTGACACTCC	This study	1
R_imp_11	GCTGTCGCTATGG-AAATGTGAGG	This study	1
F_kpc_001	CAGCGGCAGCAG-TTTGTTGATTG	This study	1
R_kpc_001	CCAGACGACGGC-ATAGTCATTTG	This study	1
F_oxa48_003	TTACTGAACATAA-ATCACAGGCGTAG	This study	1
R_oxa48_003	ATTATTCGTAATC-CTTGCTGCTTATTCTC	This study	1
F_sme_006	CAGATGAGCGGT-CCCTTTATGC	This study	1
R_sme_006	CAGAAGCCATA-TCACCTAATGTCATACC	This study	1
F_spm_001	CCTACAATCTAA-CGGCGACCAAG	This study	1
R_spm_001	AACGGCGAAGA-GACAATGACAAC	This study	1
F_vim_03	GTGTTTGGTCGCA-TATCGCAAC	This study	1
R_vim_03	GCTGTATCAATC-AAAAGCAACTCATC	This study	1
F_cmy_01	CAGGTGCTCTTC-AACAAG	This study	2
R_cmy_01	CGCCCTCTTTTCAAC	This study	2
F_IS51_01	GTCATAGTATT-CGTCGTTAGA	This study	2
R_IS51_01	GTAAGAGTGCTTTAATG-TTCATA	This study	2
F_ndm_01	CGATCAAACCGTTGGAAG	This study	2
R_ndm_01	AAGGAAAACCTTGATG-GAATTG	This study	2
F_oxa24_02	ACTTTAGGTGAGGCAATG	This study	2
R_oxa24_02	TAACCTCTTGTA-CTGGTGTA	This study	2
F_oxa27_001	ATATTTACTTGCTATG-TGGTTGCTTCTC	This study	2
R_oxa27_001	TCTCCAATCCGATC-AGGGCATTC	This study	2

Table 2. Cont.

Primer	Sequence 5' - 3'	Reference	Oligomix
F_oxa58_02	GACAATTACACCTA-TACAAGAAG	This study	2
R_oxa58_02	CGCTCTACATAC-AACATCTC	This study	2
F_sfc_01	CCTGGTGATGAT-AGAGATAC	This study	2
R_sfc_01	ATAATCGTTGGCTGTACC	This study	2
F_sim_01	CTGCTGGGATAGAGTGGCTTAATACThis study2R_sim_01TCAATAGTATGCGTCTCCGATTCThis study2		

doi:10.1371/journal.pone.0085854.t002

control strains (Table 1), which were confirmed at National Institute for Health and Welfare, Turku, Finland [14]. Since the target primer regions were fully conserved in silico, it was considered adequate to demonstrate PCR performance with one or more control species representing all the gene variants. In addition, synthetic gene constructs for SFC, CMY-1/10, SIM, SME, OXA-25, and OXA-58 genes containing a partial, non-functional resistance gene in *E. coli* plasmid (pIDTsmart), including the amplicon and app. 20 bp upstream and downstream sequence (Integrated DNA Technologies Inc, CA, USA). The plasmid was then transfected into the TOP10 strain according to manufacturer's instructions. The construct was ordered from IDT using pSMART plasmid, blunt-ended, containing a kanamycin resistance gene. The SFC, and SIM the control strains were obtained later (as a kind gift from Dr. Correia and Dr. Yunsop Chong and Kyungwon Lee, consequently). All the gene products were confirmed by sequencing with reference primers, or the gene

Table 3. Control strains.

Gene	Bacterium	Ct (50 ng/μl)	T(m)	PCR reaction
GES-1	<i>K. pneumoniae</i>	23	84	PCR1
GIM-1	<i>P. aeruginosa</i>	16	80	PCR1
IMI-2	<i>E. cloacae</i>	14	78	PCR1
IMP-15	<i>P. aeruginosa</i>	15	77	PCR1
KPC-2	<i>K. pneumoniae</i>	17	87	PCR1
OXA-48	<i>E. coli</i>	15	75	PCR1
SME	<i>S. marcescens</i>	11	77	PCR1
SPM-1	<i>P. aeruginosa</i>	16	80	PCR1
VIM-1	<i>K. pneumoniae</i>	17	81	PCR1
CMY-1/10	<i>E.coli</i> *	16	88	PCR2
ISaba1-OXA-51- family	<i>Acinetob. spp</i>	19	72	PCR2
NDM-1	<i>K. pneumoniae</i>	18	87	PCR2
OXA-23- family	<i>Acinetob. spp</i>	22	78	PCR2
OXA-24/40- family	<i>Acinetob. spp</i>	17	79	PCR2
OXA-58	<i>E.coli</i> *	15	76	PCR2
SFC-1	<i>E.coli</i> *	16	81	PCR2
SIM-1	<i>Acinetob. spp</i>	21	80	PCR2

* = gene construct containing the partial, non-functional resistance gene in *E. coli* plasmid.

doi:10.1371/journal.pone.0085854.t003

Table 4. Primers used for sequencing of resistance genes by polymerase chain reaction (PCR).

Gene	Primer	Sequence (5'- 3')	Size (bp)	T (m)	Reference
CMY	F_cmy_s1	TAAGATACTTCGGATGAGGAG	695	60	
	R_cmy_s1	GCATCTTCTCGGATGAATC			This study
	GES-C	GTTTTGCAATGTGCTCAACG	371	60	
GES	GES-D	TGCCATAGCAATAGGCGTAG			[25]
	GIM-1F	AGAACCTTGACCGAACGACG	748	60	
GIM	GIM-1R	ACTCATGACTCCTCACGAGG			[25]
	IMI-A	ATAGCCATCCTGTTTAGCTC	818	55	
IMI	IMI-B	TCTGCGATTACTTTATCTCTC			[25]
	F_IMP-1	TGAGCAAGTTATCTGTATTC	740	55	
IMP	R_IMP-1	TTAGTTGCTTGGTTTTGATG			[25]
	F_IMP-2	GGCAGTCGCCCTAAAACAAA	737	55	
IMP	R_IMP-2	TAGTTACTTGGCTGTGATGG			[25]
ISaba1/OXA-51	F_IS51_01	GTCATAGTATTCGTCGTTAGA	301	60	
	R_oxa51_001	GCACGAGCAAGATCATTACCATAGC			This study
	F_KPC	ATGTCACTGTATCGCCGTCT	893	55	
KPC	R_KPC	TTTTCAGAGCCTTACTGCCC			[25]
NDM	F_ndm_s1	GACAACGCATTGGCATAAG	447	60	
	R_ndm_s1	AAAGGAAAACCTGTATGGAATTG			This study
OXA-23 family	F_oxa23_s1	GTGTCATAGTATTCGTCGTTAG	592	60	
	R_oxa23_s1	TATCAACCTGCTGTCCAAT			This study
OXA-24 family	F_oxa25_s1	ATTAGGGCTTGAGTGAAAA	521	60	
	R_oxa25_s1	TTGTATGATTGTCAACTGCTAT			This study
	OXA-48A	TTGGTGGCATCGATTATCGG	744	62	
OXA-48	OXA-48B	GAGCACTCTTTTGTGATGGC			[25]
SFC	F_sfc_s1	CTCATTCTCCTGTGACTGA	351	60	
	R_sfc_s1	TTGCTCCTCTGTGTATT			This study
	SIM1-F	TACAAGGGATTGGCATCG	571	60	
SIM	SIM1-R	TAATGGCCTGTTCCCATGTG			[25]
	F_sme_s1	AAGGCTCAGGTATGACATT	410	60	
SME	R_sme_s1	GGCATAATCATTGCGAGTA			This study
	SPM-1F	CCTACAATCTAACGGCGACC	650	55	
SPM	SPM-1R	TCGCCGTGTCAGGTATAAC			[25]
	F_VIM-1	TTATGGAGCAGCAACCGATGT	920	60	
VIM	R_VIM-1	CAAAAGTCCCGCTCCAACGA			[25]
	F_VIM-2	AAAGTTATGCCGCACTCACC	865	60	
VIM	R_VIM-2	TGCAACTTCATGTTATGCCG			[25]

doi:10.1371/journal.pone.0085854.t004

specific primers alone, when published reference primers were not available. For additional species identification, OXA-51 gene (*bla*_{OXA-51-like}), with or without *ISAbal*, was detected separately, using F_oxa51_001 AATTTATTTAACGAAGCACACACATACGG, and R_oxa51_001 GCACGAGCAAGATCATTACCATAGC primers and the PCR program shown below.

The specificity was tested with 58 carbapenem susceptible *Enterobacteriaceae* isolates (Table S2) [18], and 710 isolates with putative reduced susceptibility *A. baumannii*, *P. aeruginosa* and *Enterobacteriaceae* isolated from clinical samples during 2008–2011. These isolates were selected among samples growing on CHROMagar ESB, or CHROMagar KPC plates (bioMérieux, Marcy L'Etoile, France), or from other culture isolates with disk

diffusion diameter <25 mm for ertapenem, or <22 mm for meropenem, or MIC>0,5 mg/l for ertapenem and meropenem.

Validation of multiplex Real-Time PCR assay

Template DNA was extracted from a single colony on CLED plate grown overnight, and re-suspended in 100 µl TE-buffer (0,5 McF) and boiled 15 min. Each 20 µl real time PCR-reaction included 10 µl Maxima SYBR Green qPCR Master Mix (2X) (Scientific Fermentas, Schwerte, Germany), 6 µl Oligomix 1 or 2 (Table 2), IDT (Integrated DNA Technologies, Inc.), 3 µl H₂O, and 1 µl DNA template. Amplification was performed as follows: 95°C 10 min initial denaturation, 30 cycles with 95°C 20 sec denaturation, 58°C 30 sec annealing and extension, final extension 58°C 1 min and final denaturation 95°C 30 sec (MxPro

3005P, Stratagene, La Jolla, CA, USA). Melting curve was determined between temperatures 58–95°C. Control strains are presented in Table 3.

The PCR was run as a preformed oligonucleotide mixture with master mixture and template to avoid quality variations between the runs. A new oligonucleotide mixture was always tested with all the panel targets with set expected 19–25 Cq range in qPCR depending on the target (Table 3). The oligonucleotide mixture was stored in stock concentrations in small aliquots, and a working dilution was formed for short term usage only. In addition, each PCR run including a representative negative and positive control for the given multiplex: KPC for multiplex 1 and NDM for multiplex 2. An acceptance range for positive controls (target +/- 3 Cq) was implemented to accept test series.

All positive isolates were confirmed by further analysing by an independent, conventional PCR and by sequencing the carbapenemase gene. Primers used in sequencing are presented in Table 4. Reaction included 2,5 mM dNTP 1,6 µl, HotStarTaq polymerase (Qjagen, Helsinki, Finland), 0,1 µl, Polymerase Buffer 10x2 µl, primer F and R 1 µl each, H₂O 13,3 µl and 1 µl template making a total of 20 µl reaction volume. Amplification was performed as follows: initial denaturation 95°C 15 min, 35 cycles with denaturation 94°C 30 sec, variable annealing temperature 55/60/62°C 30 sec depending on the carbapenemase gene to be amplified, extension 72°C 10 min, final extension 72°C 10 min (DNA Engine Tetrad 2, Peltier Thermal Cycler, BioRad, CA, USA).

Rep-PCR

DNA was extracted from colonies on GLED plates using the UltraClean microbial DNA isolation kit (Mo Bio Laboratories, Solona Beach, CA, USA) and diluted to 35 ng/µl. The DNA was amplified using the DiversiLab *Acinetobacter* kit (Bacterial Barcodes, Inc. cat no DL-AB01, Athens, GA, USA) for DNA fingerprinting following the manufacturer’s instructions. PCR was run on preheated thermal cycler (DNA Engine Tetrad 2, Peltier Thermal Cycler BioRad, Hercules, CA, USA) using the parameters according to manufacturer’s recommendations. The kit specific positive and negative controls were run with each reaction set for the validation of amplification. The rep-PCR products were detected and the amplicons were separated using microfluidics lab-on-a-chip technology and analysed using the DiversiLab system (Bacterial Barcodes, Inc.). Further analysis was performed with the web-based DiversiLab software (version 3.4) using the band-based modified Kullback-Leibler distance for the calculation of percent similarities. The manufacturer provides guidelines for strain-level discrimination; similarity more than 97% is considered as indistinguishable (no differences in fingerprints), similarity more than 95% as similar (1-2 band difference in fingerprints) and similarity less than 95% as different. In this study optimal cut-off for clustering was 95%.

Ethics statement

The bacterial isolates analyzed in this study belong to the microbiological collections of HUSLAB (Laboratory of Helsinki University Central Hospital) and were obtained as part of routine clinical care in the past. Furthermore, all patient identifiers had been previously removed and data were analyzed anonymously. As the isolates were not clinical samples in the legal sense, no written or verbal consent was needed.




Figure 1. DiversiLab analysis. Dendrogram and computer-generated image of rep-PCR banding patterns showing clustering between oxacillinase genes; OXA-23-like, OXA-24-like and OXA-58. doi:10.1371/journal.pone.0085854.g001




Figure 2. Time-dependent distribution of acquired oxacillinase genes; OXA-23-like, OXA-24-like and OXA-58.
doi:10.1371/journal.pone.0085854.g002

Results

Characterization of carbapenemase genes with *A. baumannii*

All the strains were analysed for 17 carbapenemase gene groups using the new assay. Among these *A. baumannii* isolates the most prevalent gene was OXA-23-like (bla_{OXA-23} -like). In addition we also found eight OXA 58 (bla_{OXA-58}) genes and one OXA-24-like (bla_{OXA-24} -like) gene (Figure 1). No other carbapenemase genes, including genes for KPC, VIM, IMP, GES-1/-10, OXA-48, NDM, GIM-1, SPM-1, IMI/NMC-A, SME, CMY-10, SFC-1, and SIM-1, were detected. The IS*AbaI*-OXA-51-like junction PCR was negative in all strains, as well (data not shown).

Temporal variation of prevalent, endemic *A. baumannii* clones

A time dependent clonal variation among the analysed *A. baumannii* was observed. A predominant clone was detected during the follow-up period, typically lasting a few years, which was then substituted by a new clone (Figure 2). Briefly, first a few isolates, harbouring a mobile element with OXA-58 gene, appeared 1993–1996 and 2003–2006 (Clone 1, Figure 1), which was not detected in the following years, followed by a clone harbouring a mobile OXA-23-like gene (Clone 2, Figure 1). The results were consistent with DiversiLab typing, and characteristic antibiotic susceptibility profile associated with the OXA clones analyzed. Only five out of 55 species having OXA-23/-58 gene displayed a different rep-PCR profile. Based on rep-PCR analysis, two predominant clones were detected. One isolate having OXA-24-like gene was unique in DiversiLab analysis, as well. As expected, all the control isolates from patient with no known connection were unique in their rep-PCR profiles.

Association of antibiotic susceptibility with clonality and carbapenemase gene profile

In our study, OXA-58 isolates had lower MIC-values for to meropenem than OXA-23-like positive isolates that systematically had higher MIC-values (Table 5). The isolates with non-acquired OXA-gene, displayed a marked variation and they included also some carbapenem resistant isolates. The control isolates (Figure 1) consisted of *Acinetobacter* spp not harbouring any of the OXA genes analyzed. These isolates were all carbapenem susceptible (Table 5).

Discussion

The carbapenemase producing multi-resistant gram negative rods are probably the most important challenge for hospital hygiene at the moment [13,19]. The great variety of underlying mechanisms, in contrast to simple *mecA* or *mecC* in MRSA, possesses a significant challenge to clinical screening process. Phenotypes are highly variable and many overlapping other resistance mechanisms complicate any simple screening approach. A straight-forward, economical method suitable for routine clinical diagnostics has not been available yet. In this paper we demonstrate the good performance of a new multiplex real-time PCR assay, detecting most important carbapenemases based on melting curve analysis, by applying it to an epidemiologically important set of clinical *A. baumannii* isolates. In a striking contrast to carbapenemase producing *Enterobacteriaceae*, which were first detected in Finland 2008 [14], the carbapenem resistant *A. baumannii* were detected in Finland already three decades ago. This study highlights the emergence of carbapenem-resistant *A. baumannii* isolates carrying the bla_{OXA-23} -like gene (Clone 1), which replaced the bla_{OXA-58} gene (Clone 2) in three years (Figure 2). These major clones might have been endemic.

The new carbapenemase detection assay was initially developed to detect carbapenemase producing *Enterobacteriaceae* isolates, but it also appeared to be a useful tool for *P. aeruginosa* and *A. baumannii*. After three years of clinical use, it has been proved to be sensitive and highly specific screening assay among more than 700 hundred

Table 5. MIC distributions for 55 *Acinetobacter* isolates.

Drug	Cumulative percentage of isolates inhibited at MIC (mg/l) value of:										Isolate	
	≤0.5	≤1	≤2	≤4	≤8	≤16	≤32	≤64	≤128	≤256		
MP	0,0	0,0	12,5	75,0	75,0	87,5	100,0					OXA-58
IP	0,0	0,0	0,0	0,0	12,5	37,5	100,0					OXA-58
RI	0,0	0,0	0,0	75,0	75,0	75,0	100,0					OXA-58
AB	0,0	0,0	12,5	37,5	50,0	50,0	87,5	87,5	87,5	100,0		OXA-58
TGC	0,0	12,5	25,0	87,5	87,5	87,5	87,5	100,0				OXA-58
CO	100,0											OXA-58
MP	0,0	0,0	0,0	0,0	0,0	0,0	100,0					OXA-23
IP	0,0	0,0	0,0	0,0	0,0	3,0	100,0					OXA-23
RI	0,0	0,0	0,0	85,0	88,0	88,0	100,0					OXA-23
AB	0,0	0,0	0,0	6,0	12,0	74,0	89,0	95,0	95,0	100,0		OXA-23
TGC	0,0	6,0	12,0	94,0	97,0	100,0						OXA-23
CO	100,0											OXA-23
MP	0,0	0,0	0,0	0,0	0,0	0,0	100,0					OXA-24
IP	0,0	0,0	0,0	0,0	0,0	0,0	100,0					OXA-24
RI	0,0	0,0	0,0	100,0								OXA-24
AB	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	100,0		OXA-24
TGC	0,0	0,0	100,0									OXA-24
CO	100,0											OXA-24
MP	20,0	40,0	50,0	50,0	50,0	80,0	100,0					non OXA
IP	10,0	50,0	50,0	50,0	50,0	50,0	100,0					non OXA
RI	0,0	0,0	0,0	30,0	90,0	90,0	100,0					non OXA
AB	0,0	0,0	0,0	0,0	20,0	90,0	100,0					non OXA
TGC	0,0	0,0	0,0	70,0	100,0							non OXA
CO	100,0											non OXA
MP	75,0	100,0										Control
IP	100,0											Control
RI	0,0	25,0	50,0	50,0	75,0	100,0						Control
AB	0,0	75,0	75,0	75,0	75,0	75,0	75,0	75,0	100,0			Control
TGC	50,0	100,0										Control
CO	100,0											Control

MP, meropenem; IP, imipenem; RI, rifampicin; AB, ampicillin+sulbactam; TGC, tigecycline; CO, colistin.
doi:10.1371/journal.pone.0085854.t005

isolates with reduced carbapenem susceptibility analysed to date [14]. One of the major problems related to molecular detection of many antibiotic resistance genes is the appearance of new genomic variants. For example, the variable regions of *bla*_{OXA-181} are up to 9% different from *bla*_{OXA-48} [20]. The new variants may not be detectable with the existing systems. To minimize the risk for false negative results, the primers were designed at conserved gene regions to achieve optimal amplification of all the current and forthcoming sub-variants. The SYBR Green chemistry was preferred to avoid false negative results due to minor mutations in the probe sequence. The probe based assays are often sensitive to just 1–2 mutations in probe sequence, whereas primers are usually less sensitive to minor target mutations. These design features were considered relevant to achieve a high exclusion power of clinically relevant, acquired carbapenemase genes among carbapenem resistant strains.

A. baumannii is a nosocomial pathogen, and epidemiological tools are important to develop effective strategies for better monitoring of MDRAB clinical isolates [21]. In this study we used rep-PCR

because the method is suitable for comparison of isolate genetic profiles using standardized and automated format [22]. This method has previously demonstrated good discrimination ability of *A. baumannii* isolates [23,24]. We found two major clones with DiversiLab (Clone 1 and 2, Figure 1.) harbouring most of the isolates with *bla*_{OXA-23-like} and *bla*_{OXA-58} genes. There were only few exceptions. The cases were mostly from departments of treating patients with severe burn trauma, or intensive care units.

In this study, a good correlation between the carbapenemase gene and DiversiLab typing suggested that they both could be effectively applied for epidemiological screening of *A. baumannii* species. The new carbapenemase gene screening assay has been in clinical use for more than three years, and it has been a highly suitable method for rapid unequivocal identification of isolates harbouring acquired carbapenemase genes among *Acinetobacter*, *Pseudomonas aeruginosa*, and *Enterobacteriaceae* species. This study suggests that the new molecular methods could be successfully applied in clinical diagnostics to monitor acquired carbapenemase genes, provided that they are user-friendly and cost-effective as well.

Supporting Information

Table S1 Acinetobacter isolate description.
(DOCX)

Table S2 Species included in analytical specificity testing.
(DOCX)

References

- Dijkshoorn L, Nemec A, Seifert H (2007) An increasing threat in hospitals: multidrug-resistant *Acinetobacter baumannii*. *Nat Rev Microbiol* 5: 939–951.
- Kohlenberg A, Brummer S, Higgins PG, Sohr D, Piening BC, et al. (2009) Outbreak of carbapenem-resistant *Acinetobacter baumannii* carrying the carbapenemase OXA-23 in a German university medical centre. *J Med Microbiol* 58: 1499–1507.
- Peleg AY, Seifert H, Paterson DL (2008) *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 21: 538–582.
- Higgins PG, Dammhayn C, Hackel M, Seifert H (2010) Global spread of carbapenem-resistant *Acinetobacter baumannii*. *J Antimicrob Chemother* 65: 233–238.
- Poirel L, Naas T, Nordmann P (2010) Diversity, epidemiology, and genetics of class D beta-lactamases. *Antimicrob Agents Chemother* 54: 24–38.
- Turton JF, Ward ME, Woodford N, Kaufmann ME, Pike R, et al. (2006) The role of ISAbal1 in expression of OXA carbapenemase genes in *Acinetobacter baumannii*. *FEMS Microbiol Lett* 258: 72–77.
- Poirel L, Pitout JD, Nordmann P (2007) Carbapenemases: molecular diversity and clinical consequences. *Future Microbiol* 2: 501–512.
- Healy M, Huong J, Bittner T, Lising M, Frye S, et al. (2005) Microbial DNA typing by automated repetitive-sequence-based PCR. *J Clin Microbiol* 43: 199–207.
- Yan ZQ, Shen DX, Cao JR, Chen R, Wei X, et al. (2010) Susceptibility patterns and molecular epidemiology of multidrug-resistant *Acinetobacter baumannii* strains from three military hospitals in China. *Int J Antimicrob Agents* 35: 269–273.
- Higgins PG, Janssen K, Fresen MM, Wisplinghoff H, Seifert H (2012) Molecular epidemiology of *Acinetobacter baumannii* bloodstream isolates from the United States 1995–2004 using rep-PCR and multilocus sequence typing. *J Clin Microbiol* 15.
- Higgins PG, Hujer AM, Hujer KM, Bonomo RA, Seifert H (2012) Interlaboratory reproducibility of DiversiLab rep-PCR typing and clustering of *Acinetobacter baumannii* isolates. *J Med Microbiol* 61: 137–141.
- Miriagou V, Cornaglia G, Edelstein M, Galani I, Giske CG, et al. (2010) Acquired carbapenemases in Gram-negative bacterial pathogens: detection and surveillance issues. *Clin Microbiol Infect* 16: 112–122.
- Walsh TR (2010) Emerging carbapenemases: a global perspective. *Int J Antimicrob Agents* 36 (suppl): 8–14.
- Osterblad M, Kirveskari J, Hakanen AJ, Tissari P, Vaara M, et al. (2012) Carbapenemase-producing Enterobacteriaceae in Finland: the first years (2008–11). *J Antimicrob Chemother* 31.
- Poirel L, Walsh TR, Cuvillier V, Nordmann P (2011) Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis* 70: 119–123.
- Kaase M, Szabados F, Wassill L, Gatermann SG (2012) Detection of carbapenemases in enterobacteriaceae by a commercial multiplex PCR. *J Clin Microbiol* 50: 3115–3118.
- Kempf M, Bakour S, Flaudrops C, Berrazeg M, Brunel JM, et al. (2012) Rapid detection of carbapenem resistance in *Acinetobacter baumannii* using matrix-assisted laser desorption ionization-time of flight mass spectrometry. *PLoS One* 7: e31676.
- Antikainen J, Tarkka E, Haukka K, Siitonen A, Vaara M, et al. (2009) New 16-plex PCR method for rapid detection of diarrheagenic *Escherichia coli* directly from stool samples. *Eur J Clin Microbiol Infect Dis* 28: 899–908.
- Nordmann P, Cuzon G, Naas T (2009) The real threat of *Klebsiella pneumoniae* carbapenemase-producing bacteria. *Lancet Infect Dis* 9: 228–236.
- Potron A, Nordmann P, Lafeuille E, Al Maskari Z, Al Rashdi F, et al. (2011) Characterization of OXA-181, a carbapenem-hydrolyzing class D beta-lactamase from *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 55: 4896–4899.
- Runnegar N, Sidjabat H, Goh HM, Nimmo GR, Schembri MA, et al. (2010) Molecular epidemiology of multidrug-resistant *Acinetobacter baumannii* in a single institution over a 10-year period. *J Clin Microbiol* 48: 4051–4056.
- Carretto E, Barbarini D, Farina C, Grosini A, Nicoletti P, et al. (2008) Use of the DiversiLab semiautomated repetitive-sequence-based polymerase chain reaction for epidemiologic analysis on *Acinetobacter baumannii* isolates in different Italian hospitals. *Diagn Microbiol Infect Dis* 60: 1–7.
- Fontana C, Favaro M, Minelli S, Bossa MC, Testore GP, et al. (2008) *Acinetobacter baumannii* in intensive care unit: a novel system to study clonal relationship among the isolates. *BMC Infect Dis* 8: 79.
- Grisold AJ, Zarfel G, Strenger V, Feierl G, Leitner E, et al. (2010) Use of automated repetitive-sequence-based PCR for rapid laboratory confirmation of nosocomial outbreaks. *J Infect* 60: 44–51.
- Queenan AM, Bush K (2007) Carbapenemases: the versatile beta-lactamases. *Clin Microbiol Rev* 20: 440–58, table of contents.

Author Contributions

Conceived and designed the experiments: TP PT JK. Performed the experiments: TP SK SM JK. Analyzed the data: TP JK. Contributed reagents/materials/analysis tools: TP SK SM ET JK. Wrote the paper: TP SK SM ET PT MV JK.