

RESEARCH ARTICLE

Open Access



Healthy Food Intake Index (HFII) – Validity and reproducibility in a gestational-diabetes-risk population

Jelena Meinilä^{1*} , Anita Valkama^{1,2}, Salla B. Koivusalo³, Beata Stach-Lempinen⁴, Jaana Lindström⁵, Hannu Kautiainen^{1,6}, Johan G. Eriksson^{1,2} and Majjaliisa Erkkola⁷

Abstract

Background: The aim was to develop and validate a food-based diet quality index for measuring adherence to the Nordic Nutrition Recommendations (NNR) in a pregnant population with high risk of gestational diabetes (GDM).

Methods: This study is a part of the Finnish Gestational Diabetes Prevention Study (RADIEL), a lifestyle intervention conducted between 2008 and 2014. The 443 pregnant participants (61 % of those invited), were either obese or had a history of GDM. Food frequency questionnaires collected at 1st trimester served for composing the HFII; a sum of 11 food groups (available score range 0–17) with higher scores reflecting higher adherence to the NNR.

Results: The average HFII of the participants was 10.2 (SD 2.8, range 2–17). Factor analysis for the HFII component matrix revealed three factors that explained most of the distribution (59 %) of the HFII. As an evidence of the component relevance 9 out of 11 of the HFII components independently contributed to the total score (item-rest correlation coefficients <0.31). Saturated fatty acids, monounsaturated fatty acids, polyunsaturated fatty acids, sucrose, and fiber intakes (among other nutrients) showed linearity across the HFII categories ($P \leq 0.030$ for all nutrients tested); the higher the HFII, the closer the nutrient intake to the recommended intake level. Educational attainment ($P = 0.0045$), BMI ($P = 0.0098$), smoking ($P = 0.007$), and leisure time physical exercise ($P = 0.038$) showed linearity across the HFII categories. Intra-class correlation coefficient for the HFII was 0.85 (CI 0.79, 0.90).

Conclusions: The HFII components reflect the food guidelines of the NNR, intakes of relevant nutrients, and characteristics known to vary with diet quality. It largely ignores energy intake, its components have independent contribution to the HFII, and it exhibits reproducibility. The main shortcomings are absence of red and processed meat component, and the validation in a selected study population. It is suitable for ranking participants according to the adherence to the NNR in pregnant women at high risk of GDM.

Keywords: Diet quality index, Dietary pattern, Gestational diabetes, Nutrition and pregnancy, Validation, Nordic nutrition recommendations

* Correspondence: jelena.meinila@helsinki.fi

¹Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, P.O. Box 20Tukholmankatu 8 B, Biomedicum Helsinki, 00014 Helsinki, Finland

Full list of author information is available at the end of the article



Background

Dietary analysis through single nutrients or foods is often too constricted [1]. The effect of a single nutrient may be insufficient due to interactions and inter-correlations with other nutrients, foods, or dietary patterns [2], and moreover, it may be too small to be identified. In contrast the cumulative effect of several foods, measured by dietary index for instance, addressing interactions and inter-correlations may become observable [3]. Few indices include only foods [4], and those that do [5, 6] tend to be rather concise and not specific enough for measuring subtle yet possibly essential elements, such as the quality of fat or the quality of carbohydrates, as well as the consumption of energy-rich/nutrient-poor foods. A nutrient-based index is rather burdensome for both the participant and the evaluator as it employs the use of a detailed diet record, a nutrient-calculation software, and a nutrient composition database. Food-based indices are necessary and useful when the calculations for precise nutrient intakes are infeasible, when the interest is in dietary choices rather than in nutrient intakes, or for a quick screening of a patient's diet in health care settings.

Evidence based national and international nutrition recommendations provide a good basis for the construction of a diet quality index aimed to measure healthy eating [7]. Nordic Nutrition Recommendations (NNR) food-based guidelines are also applicable for pregnant women and adherence to these guidelines will be adequate to meet the nutritional needs of a healthy pregnant woman [8]. The NNR food guidelines [9] may also help in preventing gestational diabetes (GDM), since many of the food recommendations, namely high consumption of fruits, vegetables, whole-grains, fish and poultry, and low intakes of red and processed meat, and refined grains, have been associated with a lower risk of GDM [10–12]. Existing indices that are at least indirectly based on the NNR, such as The Baltic Sea Diet Score (BSDS) [13] and the Diet Quality Index, which is based on the Swedish Nutrition Recommendations 2005 (DQI-SNR) [14], are not solely food-based. The New Nordic Diet score (NND) [15], for example, takes into account issues beyond our aims, such as how environmentally friendly foods are and the locality of foods.

What led us to develop the Healthy Food Intake Index (HFII) was simply the lack of an appropriate food-based diet quality index for RADIEL and for other epidemiological studies where only food instead of nutrient intake data is available. This kind of index could be further adapted and validated for clinical use for diet quality screening. The aim of this study was to create a simple food-based diet quality index for measuring the adherence to the NNR food guidelines [9] and to validate the index among Finnish pregnant women at high risk of GDM.

Methods

The study participants were part of the Finnish Gestational Diabetes Prevention Study (RADIEL), a multicenter lifestyle intervention study conducted in two Southern Finnish districts, namely the Helsinki Metropolitan area and Lappeenranta, between 2008 and 2014. The 727 participants were Finnish women with an elevated risk for GDM due to obesity (BMI ≥ 30 kg/m²), or a history of GDM, who were less than 20 weeks pregnant ($n = 492$) or were planning pregnancy ($n = 235$). Another criterion for inclusion in the study was that the participants were at least 18 years old. Exclusion criteria included diabetes diagnosed before pregnancy, medication influencing glucose metabolism, multiple pregnancy, physical disability, current substance abuse, severe psychiatric disorder and substantial communication difficulties. The data presented here is from the first trimester of pregnancy and it was collected during the woman's first visit to the study nurse (between 2008 and 2014) during pregnancy. From the 235 women who were recruited before their pregnancy, 30 did not conceive within one year, 61 dropped out without getting pregnant or before the first visit to the study nurse during pregnancy, and 9 were excluded because they became pregnant with twins. From the 627 (86 %) participants who visited the study nurse in the first trimester of pregnancy, diet records were available for 485 (77 % of 627), but were inaccurately recorded for 4 women. The food frequency questionnaire (FFQ) was available for 624 participants, while for 586 participants it was complete enough to calculate their dietary score. The final number of participants that provided a HFII score and a food record was 443 (61 % of all) of whom 96 (22 %) were recruited before pregnancy and 347 (78 %) during pregnancy. For reproducibility analysis we used a subsample of control participants who did not have GDM diagnosis at the first trimester of pregnancy ($n = 122$). This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by The Ethics Committee of the Department of Obstetrics and Gynecology of Helsinki and Uusimaa Hospital District. Written informed consent was obtained from all participants.

Data collection

Food intake data for constructing the Healthy Food Intake Index (HFII)

The data for the HFII were derived from the non-validated semi-quantitative food frequency questionnaires (FFQ). The validation and reproducibility process of the HFII also compensates the lack of validity assessment of the underlying FFQ; reproducibility of the HFII also reflects reproducibility of the FFQ, and comparison between the HFII and nutrient intakes from food records provides information on the validity of the FFQ. The participants filled in the FFQs

at their first visit to the study nurse when they were between 6 and 18 weeks pregnant, with a mean gestational age of 12.5 weeks (SD 1.9). For the reproducibility analysis we also used the FFQs from weeks 22 to 30 (mean gestational age 26 (SD 1.9)), for calculating the HFII_{2trimester}. The FFQs included 48 foods with 7 frequency options ranging from “less than once a week or never” to “more than 4 times per day”. Twenty-one foods from the FFQ were used for the HFII. In addition, there were 9 qualitative questions in the FFQ, of which two on the quality of fats, namely fat spreads and cooking fat, and these were used to develop the HFII. The options for fat spreads were margarine, low-fat margarine, sterol margarine, butter-oil mix, or butter, and those for cooking fats were vegetable oil, margarine, liquid margarine, baking margarine, butter-oil mix, or butter.

Components of the Healthy Food Intake Index (HFII)

The components for the HFII were selected to reflect the content of food-based guidelines of the Nordic Nutrition Recommendations (NNR) [9]. The main food-based guidelines of the HFII are presented in the first column of Table 1. The HFII comprised 11 components and covered the following food groups; vegetables, fruits and berries, high-fiber grains, fish, low-fat milk, low-fat cheese, cooking fat, fat spread, snacks, sugar-sweetened beverages, and fast food. The HFII is a sum of the 11 components, and the scores range between 0 and 17.

The HFII components snacks, sugar-sweetened beverages, fast food, vegetables, and fruits and berries, included all the foods that the FFQ provided for those food groups. The foods not incorporated into the index, such as potato, soy products, internal organs, coffee or tea, were not relevant considering the content of the NNR. Within the main 12 NNR food groups, three groups were missing from the HFII, namely meat, alcohol, and nuts and seeds (Table 1). The different types of meats (red / white and processed), and nuts (salted / non-salted) were impossible to separate from each other, because of the way the questions had been laid out in the FFQ. Alcohol intake in the current population was marginal, and was therefore not included in the HFII. Liquid dairy products included only milk, because reliable categorization of yoghurts, based on their fat content, was not possible since this information had not been collected. The high-fiber grain-category did not include breakfast cereals as we could not classify them according to their sugar and fat content.

Scoring of the Healthy Food Intake Index (HFII)

The scoring of the HFII is presented in Table 1. The score was set to reflect the food-based guidelines of the NNR where feasible. Where NNR did not provide unequivocal or numerical recommendations, we applied Finnish Nutrition Recommendations (FNR; based on the NNR) [16]

and a consensus agreement in a panel of nutrition experts was applied in the cut-offs.

Scores for consumption of cheese and milk were assigned based on their fat content. Participant's choice between high SFA vs. low SFA content was applied for cooking fat and fat spread components. For high-fiber grains, vegetables, fruits and berries, and fish the cut-offs were set according to the FNR recommendation on the number of portions per day. For the rest of the components (fast food, snacks, and sugar-sweetened beverages) numerical recommendations were absent from the NNR and the FNR, leading to the cut-offs being set according to medians and tertiles of the frequency of use in the study population.

Weighting of the HFII components

Each score component was assigned a maximum score value of either 1 or 2, based on a priori assumption of the relative importance of the category for the overall diet quality. Based on findings in the latest national Findiet survey [17], fat used as spread was considered more important source of total and saturated/unsaturated fats compared to cooking fat. Similarly, type of milk was considered to contribute more to total dairy fat intake than cheese. Snacks component includes more food groups than sugar-sweetened beverages and fast food and was, therefore, assigned maximum of 2 points whereas maximum of one point was assigned for both sugar-sweetened beverages and fast foods.

Nutrient intake data

The three-day estimated food records were used for assessing criterion validity, i.e. as a reference method for the validity of the HFII. The food records were collected prior to the FFQ. The participants were asked to record all foods and beverages they consumed during three consecutive days (2 weekdays and 1 weekend day) using household measures or weights, where applicable, and submit the food record at their first visit to the study nurse. Two trained nutritionists assessed and entered the data into the nutrient-calculation software AivoDiet, version 2.0.1.5 (Aivo Finland Oy). The Finnish National Institute for Health and Welfare (www.fineli.fi) provided the food composition database used by the software. A more detailed description of the collection and handling of the food records have been presented in our previous paper [18].

Demographic characteristics and anthropometric measures

The participants filled in a questionnaire on their health, lifestyle habits, and history of pregnancies and pregnancy-related issues. The questionnaire also gathered information on age, basic education, highest level of education, smoking (yes/no), and time used for physical exercise. The number of years of education was calculated based on the

Table 1 Components and scoring of the Healthy Food Intake Index (HFII) validated among Finnish pregnant women at high risk of gestational diabetes

| Main guidelines in NNR | HFII component | Included foods | Intake freq. | Score | Principle for cut-off | |
|---|---------------------------|---|--------------|----------|-----------------------|----------|
| 1. Limit: | | | | | | |
| Beverages and foods with added sugar or salt | Snacks | candy, chocolate, pastries, chips, ice cream | ≤4 x / wk | 2 | tertiles | |
| | | | 5–6 x / wk | 1 | | |
| | | | ≥1 x / d | 0 | | |
| | Sugar-sweetened beverages | sugar-sweetened soft drink and sugar-sweetened juice | <1 x / wk | 1 | median | |
| | | | ≥1 x / wk | 0 | | |
| | Fast food | hamburgers ja pizza | <1 x / wk | 1 | median | |
| ≥1 x / wk | | | 0 | | | |
| Processed and red meat | - | | | | | |
| Alcohol | - | | | | | |
| 2. Exchange: | | | | | | |
| Refined cereals to → whole grain cereals | High-fiber grains | dark bread, brown rice and pasta, porridge | ≥3 x / d | 2 | FNR + CP | |
| | | | 1–2 x / d | 1 | | |
| | | | <1 x / d | | | |
| Butter to → vegetable oils | Cooking fat | vegetable oil/margarine /liquid margarine/no fat butter, butter-oil mix /baking margarine | | 1 | NNR | |
| | | | | 0 | | |
| | | | | | | |
| Butter based bread fat spreads to → oil-based fat spreads | Fat spread | margarine, low-fat margarine sterol margarine, or if options from more than one category chosen butter or butter-oil-mix/no spread /no spread | | 2 | NNR | |
| | | | | 1 | | |
| | | | | 0 | | |
| | | | | | | |
| High-fat dairy to → low-fat dairy | Low-fat cheese | fat percentage ≤17 % fat percentage >17 % / no cheese | | 1 | NNR | |
| | | | | 0 | | |
| | Low-fat milk | only low-fat milk (≤1 % fat) both low-fat and full fat milk full fat milk or no milk at all | | 2 | NNR | |
| | | | | 1 | | |
| | | | 0 | | | |
| 3. Increase: | | | | | | |
| Vegetables, fruits, and berries | Vegetables | vegetables, legumes | >2 x / d | 2 | FNR + CP | |
| | | | 1–2 x / d | 1 | | |
| | | | <1 x / d | 0 | | |
| | Fruits and berries | | | ≥1 x/d | 1 | FNR + CP |
| | | | | <1 x / d | 0 | |
| | | | | | | |
| Fish and seafood | Fish | | ≥1 x / wk | 2 | FNR + CP | |
| | | | <1 x / wk | 0 | | |
| Nuts and seeds | - | | | | | |
| Total HFII, maximum score | | | | 17 | | |

NNR Nordic Nutrition Recommendations 2012 [9], FNR Finnish Nutrition Recommendations 2014 [16], CP Consensus Panel decision

reported basic and highest education. Leisure time physical activity was queried as time per week used for physical exercise during the last month. Body mass index (BMI) was calculated from the weight and height measured at the woman's first visit to the study nurse during pregnancy. The validation, component-analysis and reproducibility-analysis protocol are presented in Table 2.

Statistical methods

Descriptive results are expressed as percentages, mean or median, standard deviation (SD), or interquartile range (IQR). Differences between the participants with available dietary data (who were included in the further analysis) and participants without available dietary data (excluded from the further analysis) were tested by Chi-square test, Student's *t*-test, Mann-Whitney *U*-test or Fisher's exact test, depending on the distribution of the variables. Construct validity was studied by iterated principal factor analysis with varimax rotation for the HFII component matrix of polychoric correlations. Item analysis of the HFII components was performed by analyzing item discriminating power (corrected item correlation) and item difficulty (item mean), depicted by explanatory data analysis. Corrected item correlation was estimated using polychoric or polyserial correlations. In order to test how the HFII was associated with nutrient intakes and with characteristics associated with a healthy diet, the HFII was divided into three categories by setting cut-off limits at \pm one deviation from the mean (from here onwards referred to as HFII categories). Statistical comparisons between the

categories were performed using bootstrap-type general linear models with the appropriate contrast. The bootstrap method was used when the theoretical distribution of the test statistics was unknown or when it violated the assumptions (e.g. non-normality). The normality of the variables was tested using the Shapiro-Wilk *W* test. Reproducibility of the HFII components was evaluated from the HFII_{1trimester} and the HFII_{2trimester} using the weighted Kappa coefficient, and that of the total HFII using intra-class correlation coefficient (ICC) with one-way random-effects model. Thresholds for the Kappa coefficients were considered according to Landis & Koch [19]: 0 = less than chance agreement; 0.01–0.20 = slight agreement; 0.21–0.40 = fair agreement; 0.41–0.60 = moderate agreement; 0.61–0.80 = substantial agreement; 0.81–0.99 = almost perfect agreement. Stata 13.1, Stata-Corp LP (College Station, TX, USA) statistical package was used for the analyses.

Results

The participants with available dietary data had less frequently a history of GDM (49 % vs. 59 %, $p = 0.03$), and higher educational attainment (14.4 vs. 14.1 years, $p = 0.04$) compared to participants without available dietary data. No other differences were observed (results not shown). The average age of the 443 participating pregnant women with a high risk of GDM was 32.4 years (SD 4.5). Of this number, 140 (31 %) were nulliparous, 85 (19 %) had no history of GDM, and 218 (49 %) had a history of GDM. The average number of years of education was

Table 2 Validation protocol of the Healthy Food Intake Index (HFII) among pregnant Finnish women at high risk of GDM: type of validity and HFII components, and adopted approach for evaluation

| | |
|--|--|
| Reproducibility | 1. Kappa coefficients between 1st and 2nd pregnancy trimesters' HFII components, intra-class correlation coefficient for 1st and 2nd trimesters' total HFII. |
| 1. Does the HFII ₁ measured at 1st trimester adequately agree with the HFII ₂ measured at 2nd trimester? | |
| Content validity | 2. Comparing the content of the HFII with NNR. |
| 2. Do the index components cover all the food groups of the underlying recommendations of healthy diet (NNR)? | |
| Construct validity | 3. Item analysis of the HFII components: corrected item correlation and item mean. |
| 3. Does the HFII create variation in the population? | 4. Energy intake from food records. Comparisons between the HFII categories: general linear models |
| 4. Is scoring independent from energy intake? | 5. Iterated principal factor analysis for the HFII components matrix. |
| 5. Does the HFII have multidimensional construct and what are the dimensions it measures? | |
| Components | 6. Corrected item correlation |
| 6. Do the components have independent roles within the HFII? | 7. Item mean |
| 7. Which components provide the highest and lowest scores? | |
| Criterion validity | 8. Statistical comparisons between the HFII categories: bootstrap-type general linear models with the appropriate contrast |
| 8. Does a linear trend in nutrient intake exist across the HFII categories, or index component categories? | 9. Same as 8 |
| 9. Does the HFII distinguish between groups with known differences in diet quality? ^a | |
| a. Age | |
| b. Education | |
| c. BMI | |
| d. Smoking | |
| e. Physical activity | |

14.4 (SD 4.7), and the mean BMI at the first study visit was 31.9 kg/m² (SD 5.7). The number of smokers among the participants was 22 (5 %) and the median time used for physical exercise was 60 min/week (IQR 30; 140).

The mean score (HFII) among pregnant women at high risk of GDM was 10.2 points (SD 2.8), with a range of 2 to 17. Three participants (0.7 %) had the highest possible score of 17. The distribution of the HFII did not differ from normality ($P = 0.41$) (Additional file 1) but was slightly negatively skewed ($P = 0.081$).

Content validity

The Healthy Food Intake Index (HFII) covered all the food groups addressed in the NNR food guidelines, apart from alcohol, processed and red meat, and nuts (Table 1).

Criterion validity

The energy-adjusted intakes of the energy-yielding nutrients, including saturated fatty acids (SFA), monounsaturated fatty acids (MUFA), and polyunsaturated fatty acids (PUFA), and the intakes of fiber, sucrose, vitamins A, D, E, and folate, showed linearity across the three total HFII categories (Table 3). The intake of energy was not statistically significantly associated with the three total HFII categories ($P = 0.24$), which is an indication of the HFII measuring diet quality instead of diet quantity. The linearity of the nutrient intakes in the HFII-component categories is presented in Fig. 1. The components that provided

Table 3 Intake of nutrients among pregnant Finnish women at high risk of gestational diabetes by Healthy Food Intake Index (HFII) categories^a

| | HFII | | | | | | P-value ^b |
|-----------------------|------|------|------|------|-------|------|----------------------|
| | 0–7 | | 8–12 | | 13–17 | | |
| | Mean | SD | Mean | SD | Mean | SD | |
| Energy | 1943 | 551 | 1918 | 405 | 1856 | 411 | 0.24 |
| Carbohydrate E % | 42.8 | 7.4 | 45.3 | 5.8 | 46.1 | 5.7 | 0.002 |
| Protein E % | 17.5 | 3.7 | 17.8 | 3.0 | 19.0 | 2.8 | 0.001 |
| Fat E % | 36.9 | 7.5 | 33.5 | 5.8 | 30.9 | 5.6 | <0.001 |
| SFA % from total fat | 38.9 | 5.3 | 36.6 | 5.4 | 33.6 | 5.4 | <0.001 |
| MUFA % from total fat | 33.5 | 3.0 | 34.4 | 2.7 | 34.9 | 3.0 | 0.0017 |
| PUFA % from total fat | 15.5 | 3.3 | 17.4 | 3.8 | 19.6 | 4.1 | <0.001 |
| Sucrose E % | 9.6 | 4.6 | 8.9 | 3.6 | 7.4 | 3.3 | <0.001 |
| Dietary fiber g/MJ | 2.5 | 0.9 | 2.9 | 0.9 | 3.5 | 0.8 | <0.001 |
| Vitamins: | | | | | | | |
| C mg/MJ | 14.9 | 10.7 | 17.7 | 7.9 | 20.4 | 8.5 | <0.001 |
| E mg/MJ | 1.3 | 0.3 | 1.3 | 0.3 | 1.4 | 0.3 | <0.001 |
| A µg/MJ | 89.6 | 45.0 | 92.9 | 44.1 | 103.3 | 40.2 | 0.03 |
| D µg/MJ | 0.8 | 0.4 | 0.9 | 0.5 | 1.1 | 0.8 | <0.001 |
| Folate µg/MJ | 32.4 | 9.4 | 35.6 | 8.4 | 40.0 | 6.9 | <0.001 |

^aCut-off limits \pm one deviation from the mean. ^bTested by bootstrap-type general linear models with linear contrast

the highest scores (fish, low-fat milk, fat spread, and snacks) as well as the component with the lowest score (low-fat cheese), all reflected the intake levels of SFA and PUFA.

Educational attainment (years of education) ($p = 0.0045$) and physical activity ($p = 0.038$) showed positive linearity across the three HFII categories. BMI ($p = 0.0098$) and smoking (0.0079) showed negative linearity across the three HFII categories whereas age ($p = 0.18$) did not show a statistically significant trend.

Construct validity


Factor analysis for the HFII component matrix revealed that within all the HFII components (Table 4), there were three distinct factors that explained most of the variation (59 %) within the score. According to the food group loadings, three major factors were identified. The first was characterized by high loadings for components cooking fat, fat spread, low-fat cheese, and low-fat milk (named as Fat factor). The second was characterized by high loadings for high-fiber grains, vegetables, fruits and berries, and fish (named as Healthy foods). The third factor was characterized by high loadings for snacks, sugar-sweetened beverages, and fast food (named Unhealthy foods). This provided information on correlational structure of the components and that the HFII measured meaningful dimensions of diet.

Components of the HFII

The components of the HFII that provided the highest scores were milk, fish, fat spread, and snacks (Fig. 2). The component that provided the lowest scores was low-fat cheese. All components had item-rest correlation coefficients less than 0.31 with the rest of the components, snacks having the lowest item-rest correlation coefficient (0.03). High-fiber grains and vegetables had item-rest correlation coefficients of similar quantity and item means close to each other, which prevents us from interpreting whether they act independently in the HFII. The component scores of high-fiber grains and vegetables, however, reflected nutrient intakes of different magnitude which suggests that they had at least some independency. All components independently contributed to the total score and created variation within the study population, but for high-fiber grains and vegetables independency may have been weak.

Reproducibility

Weighted Kappa coefficients between the HFII_{1trimester} and the HFII_{2trimester} components ranging from 0.41 to 0.69 (Table 5) suggest moderate to substantial agreement [19]. The ICC between the HFII_{1trimester} and the HFII_{2trimester} was 0.85 (95 % CI 0.79 to 0.90). The repeatability coefficient of the HFII was 3.7 (95 % CI 3.3–4.3) units,



meaning there was a 95 % chance that two measurements will differ by less than 3.7.

Discussion

The study evaluated validity and reliability of the Healthy Food Intake Index (HFII) in a population of pregnant Finnish women who had a high risk of GDM and were participating in the Finnish Gestational Diabetes Prevention Study RADIEL.

The HFII succeeded in reflecting the level of adherence to the recommended nutrient intake levels of the NNR. The study showed that when moving from a lower to a higher HFII category, the nutrient density, apart from fat was higher. This result was similar to the results of a study

of the Baltic Sea Diet Score (BSDS) [13], Diet Quality Index – Swedish Nutrient Recommendations 2005 (DQI-SNR) [14] and the New Nordic Diet score (NND) [15]. An advantage compared to the BSDS [13] was that the HFII additionally reflected the intake of sucrose. High intake of sucrose may contribute to the development of gestational diabetes [20] and overweight [21]. The highest category in the current study did not reflect sufficient intake of folate and vitamin D as per the recommended nutrition density of the NNR. This should be taken into account when using the HFII. Sufficient intake of folate is especially important in early pregnancy [22] and the recommended nutrient intake level of the NNR [16] is higher for pregnant women than for the general population. The intakes

Table 4 Factors^a among the components of the Healthy Food Intake Index (HFII) among pregnant Finnish women at high risk of gestational diabetes

| Component | “Fats” | “Healthy foods” | “Unhealthy foods” |
|---------------------------|--------|-----------------|-------------------|
| Low-fat cheese | 0.67 | | |
| Low-fat milk | 0.66 | | |
| Fat spread | 0.70 | | |
| Cooking fat | 0.79 | | |
| Fish | | 0.66 | |
| Vegetables | | 0.81 | |
| Fruit and berries | | 0.60 | |
| High-fiber grains | | 0.45 | |
| Snacks | | | 0.72 |
| Fast food | | | 0.80 |
| Sugar-sweetened beverages | | | 0.58 |

^aExplanatory factor analysis with varimax-rotated factor loadings. Factor loadings with values < 0.45 not shown

of folate and vitamin D in the Finnish population have been below the NNR’s recommended daily intakes [23, 24] and supplementation is recommended for all pregnant women. The finding that the HFII score was rather independent of energy intake, indicates that the HFII does, to a large extent, ignore the quantity of food intake. The unwanted consequence of providing higher scores for eating more food in general, was in part reduced by the components that measured intake frequencies and habitual choices rather than amounts of intake.

Table 5 Agreement of the HFII_{1trimester} and the HFII_{2trimester} components and Kappa coefficients between them

| | Observed agreement % (95 % CI) ^a | k (95 % CI) |
|---------------------------|---|-------------------|
| Snacks | 68 (58, 75) | 0.56 (0.42, 0.70) |
| Low-fat cheese | 79 (70, 84) | 0.56 (0.42, 0.71) |
| Fast food | 78 (69, 84) | 0.44 (0.26, 0.61) |
| Low-fat milk | 81 (75, 89) | 0.65 (0.48, 0.79) |
| High fiber grains | 67 (60, 75) | 0.60 (0.45, 0.72) |
| Fish | 75 (66, 81) | 0.43 (0.26, 0.60) |
| Fruits and berries | 76 (70, 84) | 0.48 (0.32, 0.64) |
| Cooking fat | 92 (87, 98) | 0.67 (0.49, 0.86) |
| Fat spread | 80 (74, 88) | 0.69 (0.55, 0.81) |
| Vegetables | 72 (65, 80) | 0.59 (0.45, 0.72) |
| Sugar-sweetened beverages | 70 (63, 80) | 0.41 (0.25, 0.57) |

^aConfidence intervals (95 % CI) were obtained by bias-corrected and accelerated bootstrapping (5000 replications)

Being able to avoid energy-adjustment contributes considerably to the HFII’s usability because it rules out the need for detailed dietary data. Further evidence for the capacity of the HFII to rank participants according to the healthiness of the diet was provided by the parallel association of the HFII with the characteristics known to associate with the healthiness of diet [25–29]. The reproducibility measures of the current study were similar to the few earlier reproducibility studies on food intake during pregnancy [30, 31]. In the light of studies of dietary changes through pre-pregnancy to pregnancy, assessing reproducibility during pregnancy seems reasonable. Cuco and her colleagues [32] as well as Crozier and her colleagues [33] both found little overall change in dietary patterns from pre-pregnancy through early to late pregnancy.

The factors identified as “fat”, “healthy”, and “unhealthy”, show that the HFII captured reasonable aspects of diet; such food choices have also occurred jointly in other studies [2, 34, 35]. The HFII-components nevertheless had independent roles within the total HFII, which was supported by the low item-rest correlation coefficients. A very low correlation with the rest of the index, however, may suggest a flaw within the component. One such component which we decided to retain in the current study because it linearly reflected the intakes of six important nutrients was snacks. Its low scores reflected the intakes further from and the high scores reflected intakes closer to the recommended nutrient intake levels of the NNR. Lower item-rest correlation of snacks may, however, indicate cut-off limits that are not comparable to the other components in reflecting a healthy diet. Re-setting the cut-off limits for snacks could improve the internal consistency of the HFII and should be considered.

The HFII distinguished whole-grain products from refined grains, and fatty milk from skimmed milk, as




Fig. 2 Item-analysis of the components of the Healthy Food Intake Index (HFII) among pregnant Finnish women at high risk of gestational diabetes. X-axis represents the mean score that the participants got from the component, and the Y-axis represents the correlation of a component to the rest of the components. The dashed line represents the mean score of all the components. 1 = Snacks, 2 = Low-fat cheese, 3 = Fish, 4 = Low-fat milk, 5 = Vegetables, 6 = Fruits and berries, 7 = Sugar-sweetened beverages, 8 = High-fiber grains, 9 = Fast food, 10 = Fat spread, 11 = Cooking fat

proposed by Waijers and co-workers [4]. Excluding red and processed meat may have impaired the accuracy of the HFII in reflecting all aspects of a healthy diet, because red and processed meat may increase the risk for cancer, cardiovascular disease, type 2 diabetes [20], and gestational diabetes [11]. In future studies, a component of red and processed meat could be added. Subjective interpretations in scoring, setting cut-off limits, and weighting, all common problems of indices [2], may have resulted in some suboptimal scorings, for example in the case of the snacks component. A more detailed FFQ could have provided more accurate cut-offs for whole grain foods, vegetables, fruits and berries, and fish. Population-based cut-offs may perform differently in other populations, and, therefore, scores between different populations may not be comparable with each other.

As a food-based index, the HFII takes into account the complexity of foods, and makes detailed nutrient-intake data and the use of major resources unnecessary [36]. Its components, nutrient intakes and characteristics in its categories compare well with the findings from the BSDS [13], the DQI-SNR [14], and the NND [15]. A major advantage of the HFII compared to many other indices is its independency of energy intake [13–15, 37]. Other notable advantages of the HFII are its simplicity and the approved multi-perspective evaluation including analysis of the relevance of its components and their interrelationships, something that the majority of index validation studies fail to recognize [4].

One weakness of the current study was that the FFQ underlying the HFII was not validated, which may have affected the accuracy of the scores. This, however, is at least partly compensated by testing validity and reliability of the HFII. Comparison of food record, that measures short term intake [38], with HFII, that measures habitual intake, may have attenuated the association between the HFII and the nutrient intakes [39]. A replicate food record could have improved the evaluation [40]. Obesity, and pregnancy are factors that result in under-reporting of dietary intake [41, 42] and therefore constitute a challenging group for dietary assessment. Thus, the scores may be over-estimates. However, because the HFII proved to be independent of energy intake, it may be less sensitive to under-reporting. Available dietary data seemed to add to the selection bias, which was already prevalent because of the study design. Obese pregnant women and women with GDM history do not represent general pregnant population, but cover a wide proportion of it; approximately one third of Finnish [43] women at childbearing age are either overweight or obese, the prevalence of GDM being approximately 13 % [44, 45]. For more generalizable results validation in a more general population, however, is required. Since nutrition recommendations in Western countries tend to have similar main

principles [46], the HFII could be used in other Western countries with minor adaptations in the food groups and with re-evaluation. Similar food guidelines apply for pregnant and non-pregnant populations [9] so the HFII could be also suitable for other adult populations. Based on the current study the HFII can be used among pregnant women in countries with similar food consumption and food guidelines to Finland, namely the Nordic countries. For further evaluation studies in different populations we have now provided a detailed scoring system and thorough validation protocol to be applied.

Conclusions

Despite the shortcomings of the HFII, it covered all relevant food groups mentioned in the food guidelines of the NNR, excluding red and processed meat. Secondly, the nutrient intakes came closer to the recommended intake of the NNR for all macronutrients and all vitamins and minerals measured when stepping towards the higher HFII categories. Thirdly, all components had a contribution to the HFII. Fourthly, demographic characteristics varied across the HFII categories meaningfully. Thus, the HFII can be used without detailed dietary data or energy-adjustment in studies for ranking the participants according to the level of adherence to the food-guidelines of the Nordic Nutrition Recommendations among overweight and obese pregnant women or pregnant women with a history of GDM. It has also great potential to be adapted in other adult populations in countries with similarities to Finland in dietary patterns.

Additional file

Additional file 1: Distribution of the scores of the Healthy Food Intake Index (HFII) among pregnant Finnish women at high risk for gestational diabetes. (DOCX 22 kb)

Abbreviations

BMI, body mass index; BSDS, The Baltic Sea Diet Score; DQI-SNR, Diet Quality Index – Swedish Nutrient Recommendations 2005; GDM, gestational diabetes; HFII, Healthy Food Intake Index; NND, new nordic diet score; NNR, nordic nutrition recommendations

Acknowledgements

We are grateful to the women who participated as subjects, the study nurses and research scientists who contributed to this study.

Funding

This work was supported by the Ahokas Foundation, Finnish Foundation for Cardiovascular Disease, Special state subsidy for health science research of Helsinki University Central Hospital, Samfundet Folkhälsan, The Finnish Diabetes Research Foundation, Foundation for Medical Research Liv och Hälsa, State Provincial Office of Southern Finland, The Social Insurance Institution of Finland.

Availability of data and materials

The statistical codes and the templates of the questionnaires are available from the corresponding author on request. The data set will not be shared at this stage because of planned future studies of the data.

Authors' contribution

JM designed, conducted research, performed statistical analysis, wrote the paper, had primary responsibility for final content, and was a member of the consensus panel of nutrition experts. AV designed and conducted research, wrote the paper, was a member of the consensus panel of nutrition experts, and read and approved the final manuscript, HK designed research, performed statistical analysis, critically revised the manuscript with expert advice and comments, read and approved the final manuscript, JL designed research, was a member of the consensus panel of nutrition experts, critically revised the manuscript with expert advice and comments, read and approved the final manuscript, JE designed and conducted research, critically revised the manuscript with expert advice and comments, read and approved the final manuscript. BS-L designed, conducted research, critically revised the manuscript with expert advice and comments, and read and approved the final manuscript. SK designed and conducted research, critically revised the manuscript with expert advice and comments, and read and approved the final manuscript. ME designed, performed statistical analysis, wrote the paper, had primary responsibility for final content, was a member of the consensus panel of nutrition experts, and read and approved the final manuscript.

Competing interests

The authors declared that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Ethics Committees of Helsinki University Central Hospital and South-Karelia Central Hospital. Written informed consent was obtained from all participants.

Author details

¹Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, P.O. Box 20Tukholmankatu 8 B, Biomedicum Helsinki, 00014 Helsinki, Finland. ²Folkhälsan Research Centre, University of Helsinki, Helsinki, Finland. ³Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland. ⁴Department of Obstetrics and Gynecology, South-Karelia Central Hospital, Lappeenranta, Finland. ⁵Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland. ⁶Department of General Practice and Primary Health Care, University of Eastern Finland, Kuopio, Finland. ⁷Department of Food and Environmental Sciences, University of Helsinki, Helsinki, Finland.

Received: 7 January 2016 Accepted: 13 July 2016

Published online: 30 July 2016

References

- Willett WC. *Nutritional epidemiology*. 3rd ed. Oxford: Oxford University Press; 2012.
- Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*. 2002;13(1):3–9.
- Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA, Vogt TM, Cutler JA, Windhauser MM, Lin PH, Karanja N. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med*. 1997;336(16):1117–24.
- Wajjers PM, Feskens EJ, Ocke MC. A critical review of predefined diet quality scores. *Br J Nutr*. 2007;97(2):219–31.
- Lowik MR, Hulshof KF, Brussaard JH. Food-based dietary guidelines: some assumptions tested for The Netherlands. *Br J Nutr*. 1999;81 Suppl 2:S143–9.
- Osler M, Heitmann BL, Gerdes LU, Jorgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. *Br J Nutr*. 2001;85(2):219–25.
- Schwingshackl L, Hoffmann G. Diet Quality as Assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension Score, and Health Outcomes: A Systematic Review and Meta-Analysis of Cohort Studies. *J Acad Nutr Diet*. 2015;115(5):780–800. e5.
- Hasunen K, Kalavainen M, Keinonen H, Lagström H, Lyytikäinen A, Nurttila A, Peltola T, Talvia S. The Child, Family and Food. Nutrition recommendations for infants and young children as well as pregnant and breastfeeding mothers. 2004.
- Nordic Council of Ministers 2014. *Nordic Nutrition Recommendations 2012 -Integrating nutrition and physical activity*. 5th ed. Copenhagen: Nordic Council of Ministers; 2014. <https://www.norden.org/en/theme/nordic-nutrition-recommendation/nordic-nutrition-recommendations-2012>. Accessed January 1st 2016.
- Tobias DK, Zhang C, Chavarro J, Bowers K, Rich-Edwards J, Rosner B, Mozaffarian D, Hu FB. Prepregnancy adherence to dietary patterns and lower risk of gestational diabetes mellitus. *Am J Clin Nutr*. 2012;96(2):289–95.
- Zhang C, Schulze MB, Solomon CG, Hu FB. A prospective study of dietary patterns, meat intake and the risk of gestational diabetes mellitus. *Diabetologia*. 2006;49(11):2604–13.
- Karamanos B, Thanopoulou A, Anastasiou E, Assaad-Khalil S, Albache N, Bachaoui M, Slama CB, El Ghomari H, Jotic A, Lalic N, Lapolla A, Saab C, Marre M, Vassallo J, Savona-Ventura C, MGS-DGM Study Group. Relation of the Mediterranean diet with the incidence of gestational diabetes. *Eur J Clin Nutr*. 2014;68(1):8–13.
- Kanerva N, Kaartinen NE, Schwab U, Lahti-Koski M, Mannisto S. The Baltic Sea Diet Score: a tool for assessing healthy eating in Nordic countries. *Public Health Nutr*. 2014;17(8):1697–705.
- Drake I, Gullberg B, Ericson U, Sonestedt E, Nilsson J, Wallstrom P, Hedblad B, Wirfalt E. Development of a diet quality index assessing adherence to the Swedish nutrition recommendations and dietary guidelines in the Malmo Diet and Cancer cohort. *Public Health Nutr*. 2011;14(5):835–45.
- Hillesund ER, Bere E, Haugen M, Overby NC. Development of a New Nordic Diet score and its association with gestational weight gain and fetal growth - a study performed in the Norwegian Mother and Child Cohort Study (MoBa). *Public Health Nutr*. 2014;17(9):1–11.
- National Nutrition Council. *Suomalaiset ravitsemussuosituukset – Terveysttä ruoasta* [Finnish nutrition recommendations – health from food]. Committee report. Tampere: Juvenes Print; 2014. In Finnish.
- Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M, Virtanen S. The National FINDIET 2012 Survey. National Institute for Health and Welfare (THL). Tampere: Juvenes Print; 2013.
- Meinilä J, Koivusalo SB, Valkama A, Rono K, Erkkola M, Kautiainen H, Stach-Lempinen B, Eriksson JG. Nutrient intake of pregnant women at high risk of gestational diabetes. *Food Nutr Res*. 2015;59:26676.
- Landis JR, Koch GG. The Measurement of Observer Agreement for Categorical Data. *Biometrics*. 1977;33(1):159–74.
- Zhang C, Liu S, Solomon CG, Hu FB. Dietary fiber intake, dietary glycemic load, and the risk for gestational diabetes mellitus. *Diabetes Care*. 2006;29(10):2223–30.
- Knudsen VK, Heitmann BL, Halldorsson TI, Sorensen TI, Olsen SF. Maternal dietary glycaemic load during pregnancy and gestational weight gain, birth weight and postpartum weight retention: a study within the Danish National Birth Cohort. *Br J Nutr*. 2013;109(8):1471–8.
- Ramakrishnan U, Grant F, Goldenberg T, Zongrone A, Martorell R. Effect of women's nutrition before and during early pregnancy on maternal and infant outcomes: a systematic review. *Paediatr Perinat Epidemiol*. 2012;26 Suppl 1:285–301.
- Helldán A, Helakorpi A, Virtanen S, Uutela A. Health Behaviour and Health among the Finnish Adult Population. 2013. 21/2013: http://www.julkari.fi/bitstream/handle/10024/126023/URN_ISBN_978-952-302-447-2.pdf?sequence=1. Accessed January 1st 2016.
- Pietinen P, Paturi M, Reinivuo H, Tapanainen H, Valsta LM. FINDIET 2007 Survey: energy and nutrient intakes. *Public Health Nutr*. 2010;13(6A):920–4.
- Hiza HAB, Casavale KO, Guenther PM, Davis CA. Diet Quality of Americans Differs by Age, Sex, Race/Ethnicity, Income, and Education Level. *J Acad Nutr Diet*. 2013;113(2):297–306.
- Quintiliani L, Allen J, Marino M, Kelly-Weeder S, Li Y. Multiple health behavior clusters among female college students. *Patient Education & Counseling*. 2010;79(1):134–7.
- Kent LM, Worsley A. Trends in BMI, diet and lifestyle between 1976 and 2005 in North Sydney. *Asia Pac J Clin Nutr*. 2009;18(3):453–61.
- Schroder H, Benitez Arciniega A, Soler C, Covas MI, Baena-Diez JM, Marrugat J, REGICOR investigators, HERMES investigators. Validity of two short screeners for diet quality in time-limited settings. *Public Health Nutr*. 2012;15(4):618–26.
- Schuit AJ, van Loon AJM, Tijhuis M, Ocké MC. Clustering of Lifestyle Risk Factors in a General Adult Population. *Prev Med*. 2002;35(3):219–24.

30. Erkkola M, Karppinen M, Javanainen J, Rasanen L, Knip M, Virtanen SM. Validity and reproducibility of a food frequency questionnaire for pregnant Finnish women. *Am J Epidemiol*. 2001;154(5):466–76.
31. Vioque J, Navarrete-Munoz EM, Gimenez-Monzo D, Garcia-de-la-Hera M, Granado F, Young IS, Ramon R, Ballester F, Murcia M, Rebagliato M, Iniguez C, INMA-Valencia Cohort Study. Reproducibility and validity of a food frequency questionnaire among pregnant women in a Mediterranean area. *Nutr J*. 2013;12:26. 2891–12-26.
32. Cuco G, Fernandez-Ballart J, Sala J, Viladrich C, Iranzo R, Vila J, Arija V. Dietary patterns and associated lifestyles in preconception, pregnancy and postpartum. *Eur J Clin Nutr*. 2006;60(3):364–71.
33. Crozier SR, Robinson SM, Godfrey KM, Cooper C, Inskip HM. Women's dietary patterns change little from before to during pregnancy. *J Nutr*. 2009;139(10):1956–63.
34. Uusitalo U, Arkkola T, Ovaskainen ML, Kronberg-Kippila C, Kenward MG, Veijola R, Simell O, Knip M, Virtanen SM. Unhealthy dietary patterns are associated with weight gain during pregnancy among Finnish women. *Public Health Nutr*. 2009;12(12):2392–9.
35. Gorst-Rasmussen A, Dahm CC, Dethlefsen C, Scheike T, Overvad K. Exploring dietary patterns by using the treelet transform. *Am J Epidemiol*. 2011;173(10):1097–104.
36. Kant AK. Indexes of overall diet quality: a review. *J Am Diet Assoc*. 1996;96(8):785–91.
37. Panagiotakos DB, Pitsavos C, Stefanadis C. Dietary patterns: A Mediterranean diet score and its relation to clinical and biological markers of cardiovascular disease risk. *Nutr Metab Cardiovasc Dis*. 2006;16(8):559–68.
38. Dodd KW, Guenther PM, Freedman LS, Subar AF, Kipnis V, Midthune D, Tooze JA, Krebs-Smith SM. Statistical Methods for Estimating Usual Intake of Nutrients and Foods: A Review of the Theory. *J Am Diet Assoc*. 2006; 106(10):1640–50.
39. Kipnis V, Subar AF, Midthune D, Freedman LS, Ballard-Barbash R, Troiano RP, Bingham S, Schoeller DA, Schatzkin A, Carroll RJ. Structure of dietary measurement error: results of the OPEN biomarker study. *Am J Epidemiol*. 2003;158(1):14–21. discussion 22–6.
40. Carroll RJ, Midthune D, Subar AF, Shumakovich M, Freedman LS, Thompson FE, Kipnis V. Taking advantage of the strengths of 2 different dietary assessment instruments to improve intake estimates for nutritional epidemiology. *Am J Epidemiol*. 2012;175(4):340–7.
41. Heitmann BL, Lissner L. Dietary underreporting by obese individuals—is it specific or non-specific? *BMJ*. 1995;311(7011):986–9.
42. McGowan CA, McAuliffe FM. Maternal nutrient intakes and levels of energy underreporting during early pregnancy. *Eur J Clin Nutr*. 2012;66(8):906–13.
43. National Institute for Health and Welfare. Perinatal statistics: parturients deliveries and newborns 2012. 2013. 24. Available from: <http://urn.fi/URN:NBN:fi-fe2014101345161>. Accessed January 1st 2016.
44. O'Sullivan EP, Avalos G, O'Reilly M, Dennedy MC, Gaffney G, Dunne F, Atlantic DIP collaborators. Atlantic Diabetes in Pregnancy (DIP): the prevalence and outcomes of gestational diabetes mellitus using new diagnostic criteria. *Diabetologia*. 2011;54(7):1670–5.
45. Lamberg S, Raitanen J, Rissanen P, Luoto R. Prevalence and regional differences of gestational diabetes mellitus and oral glucose tolerance tests in Finland. *Eur J Public Health*. 2012;22(2):278–80.
46. WHO Regional Office for Europe, Nutrition and Food Security programme. Food based dietary guidelines in the WHO European Region. 2003. EUR/03/5045414 E79832.
47. Schroder H, Fito M, Covas MI, REGICOR i. Association of fast food consumption with energy intake, diet quality, body mass index and the risk of obesity in a representative Mediterranean population. *Br J Nutr*. 2007;98(6):1274–80.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

