

Assessing climate impact indicators: Evaluation criteria and observed strengths and weaknesses

**Luis Costa, Mikael Hildén, Jürgen Kropp, Kristin Böttcher,
Stefan Fronzek, Rob Swart, Juliane Otto, Niall McCormick,
Milka Radojevic, Johannes Lückenkötter, Elke Keup-Thiel,
Kari Luojus, Tanya Singh, Juha Pöyry, Emilia Sanchez, Martin Jukes**



Assessing climate impact indicators: Evaluation criteria and observed strengths and weaknesses

**Luis Costa, Mikael Hildén, Jürgen Kropp, Kristin Böttcher,
Stefan Fronzek, Rob Swart, Juliane Otto, Niall McCormick,
Milka Radojevic, Johannes Lückenkötter, Elke Keup-Thiel,
Kari Luojus, Tanya Singh, Juha Pöyry, Emilia Sanchez, Martin Jukes**



S Y K E

REPORTS OF THE FINNISH ENVIRONMENT INSTITUTE 41 | 2016
Finnish Environment Institute

Author list: Luis Costa¹⁾, Mikael Hildén²⁾, Jürgen Kropp¹⁾, Kristin Böttcher²⁾, Stefan Fronzek²⁾, Rob Swart³⁾, Juliane Otto⁴⁾, Niall McCormick³⁾, Milka Radojevic⁶⁾, Johannes Lückenkötter⁷⁾, Elke Keup-Thiel⁴⁾, Kari Luojus⁸⁾, Tanya Singh³⁾, Juha Pöyry²⁾, Emilia Sanchez⁶⁾, Martin Juckes⁹⁾

Affiliation:

- ¹⁾ Potsdam Institute for Climate Impact Research, Germany
- ²⁾ Finnish Environment Institute, Finland
- ³⁾ Wageningen Environmental Research, Netherlands
- ⁴⁾ Helmholtz Zentrum Geesthacht, Climate Service Center Germany, Germany
- ⁵⁾ European Commission, Joint Research Centre, Italy
- ⁶⁾ Centre Européen de Recherche et de Formation Avancée en Calcul Scientifique, France
- ⁷⁾ Technische Universität Dortmund, Germany
- ⁸⁾ Finnish Meteorological Institute, Finland
- ⁹⁾ Science and Technology Facilities, Council, UK

Subject editor: Suvi Huttunen, SYKE

Publisher: Finnish Environment Institute (SYKE),
P.O. Box 140, FI-00251 Helsinki, Finland, Phone +358 295 251 000, syke.fi

Layout: Ritva Koskinen and Pirjo Lehtovaara

Cover photo: Image bank of the Environmental Administration / Riku Lumiaro

The publication is available in the internet (pdf): syke.fi/publications | helda.helsinki.fi/syke and in print: syke.juvenesprint.fi

ISBN 978-952-11-4650-3 (PDF)

ISSN 1796-1726 (online)

Year of issue: 2016

ABSTRACT

This report documents and reviews a selected set of climate change and impact indicators. They are documented according to reference criteria that were based on a literature study and later refinement in expert discussions. Methodological description, data requirements and availability, treatment of uncertainty, fitness for purpose of indicator time series, and seven other relevant criteria are documented for a total of 81 climate change and impact related indicators. The indicators were grouped into three tiers that reflect their main purpose of use, ranging from change in climate variables to the socio-economic consequences of climate change. A key observation is the limited availability of indicators that explicitly link climate change with socio-economic phenomena. This might be explained by the complexity of the system that hinders quantitative attribution of economic and multi-level societal development to climatic factors. The strengths and weaknesses of indicators are discussed at a general level and also outlined both on an indicator-by-indicator basis and with respect to their potential uses. The report presents a consistent set of criteria and approaches for the incorporation of indicator information into climate information portals. The collected information on climate change and impact indicators can support the development of the Copernicus Climate Services and the indicators that such services will promote.

Keywords: climate change, climate impact indicators, evaluation framework

TIIVISTELMÄ

Raportti on katsaus ilmastonmuutosta ja ilmastonmuutoksen vaikutuksia kuvaaviin indikaattoreihin. Indikaattorit on arvioitu kriteereillä, jotka perustuvat kirjallisuusselvitykseen ja joita on jalostettu edelleen asiantuntijakeskusteluissa. Menetelmällistä kuvausta, aineistovaatimuksia ja aineistojen saatavuutta, epävarmuuksien käsittelyä, aikasarjojen pituutta suhteessa indikaattorin käyttöön sekä seitsemää muuta kriteeriä on sovellettu yhteensä 81 ilmastonmuutosta ja ilmastonmuutoksen vaikutuksia kuvaavaan indikaattoriin. Indikaattorit ryhmiteltiin kolmeen tasoon, jotka kuvaavat niiden pääasiallista kohdetta, lähtien ilmaston fysikaalisesta muuttumisesta yhteiskunnallisiin seurauksiin. Raportti osoittaa, että on vain harvoja indikaattoreita, jotka kytkisivät ilmastonmuutoksen sosio-ekonomisiin ilmiöihin. Vaikeudet kytkeä riittävän yksiselitteisesti taloudellista ja muuta yhteiskunnallista kehitystä ilmastollisiin tekijöihin on keskeinen syy näiden indikaattoreiden puuttumiseen. Indikaattoreiden vahvuuksia ja heikkouksia on tarkasteltu yleisellä tasolla sekä arvioitu indikaattorikohtaisesti että indikaattorien mahdollisen käytön valossa. Raportissa on esitetty johdonmukainen kokoelma kriteerejä ja lähestymistapoja, joita voi käyttää kehitettäessä ilmastotiedon portaaleja. Indikaattoreista koottu tieto tukee Copernicus ilmastopalveluiden ja niihin liittyvien indikaattoreiden kehittämistä.

Asiasanat: ilmastonmuutos, ilmastoindikaattorit, arviointikehys

SAMMANDRAG

Rapporten presenterar en översikt över indikatorer som beskriver klimatförändringen och dess effekter. Indikatorerna har utvärderats med hjälp av kriterier som bygger på en litteraturöversikt or samt på expert diskussioner. Metodbeskrivningen, datakrav och datatillgänglighet, behandling av osäkerhet, tidsseriens längd i förhållande till användningen sam sju andra kriterier har tillämpats på sammanlagt 81 indikatorer som beskriver klimatförändringen och dess konsekvenser. Indikatorerna grupperades i tre nivåer som beskriver deras fokus från grundläggande fysikaliska förändringar i klimatet till samhällsliga konsekvenser. Rapporten visar att det finns få indikatorer som skulle koppla klimatförändringen till socio-ekonomiska fenomen. Svårigheterna att tillräckligt entydigt sammanbinda ekonomisk och övrig samhällsutveckling med klimatfaktorer förklarar bristen på dessa indikatorer. Indikatorernas styrka och svaghet har utvärderats på ett allmänt plan och indikator för indikator. Dessutom utvärderades indikatorerna i relation till möjlig användning. Rapporten presenterar en konsistent samling kriterier och angreppssätt som kan utnyttjas då man utvecklar indikatorer för klimatportaler. Den information som samlats in om indikatorerna stöder utvecklandet av Copernicus klimattjänster och de indikatorer som tjänsterna främjar.

Nyckelord: klimat förändring, klimat indikatorer, ramverk för utvärdering

ACKNOWLEDGMENTS

The report was prepared within the project “Climate Information Portal for Copernicus” (CLIPC), funded by the European Union’s 7th Framework Programme for research, technological development and demonstration (2013-2016, grant agreement no 607418). Within the project, a new web portal (www.clipc.eu) was developed providing access to climate information of direct relevance to a wide variety of users. The platform complements existing Copernicus pre-operational components, but focus on datasets which provide information on climate variability on decadal to centennial time scales from observed and projected climate change impacts in Europe, and provides a toolbox to generate, compare and rank key indicators. Within the project a catalogue of potential climate change and impact indicators to be made available via CLIPC was created. The criteria for the examination of indicators and the assessment of their strengths and weaknesses are included in this report that draws upon a project deliverable (Costa and Hildén 2015).

A joined CLIPC and European Environment Agency (EEA) expert workshop was organized at the EEA in May 2015 with the aim to discuss the criteria to be used for the evaluation and screening of climate change and impact indicators. The following experts participated in the workshop: L. Bärring (SMHI), T. Carter (SYKE), A. de Groot (Alterra), J. Fons-Esteve (UAB), H-M. Füßel (EEA), N. Gobron (JRC), A. Jol (EEA), B. Kurnick (EEA), J.-N. Thépaut (ECMWF). We acknowledge the contribution of all participants in the workshop whose input contributed especially to the understanding of the use of indicators.

Furthermore, the authors thank J. Attila, O.-P. Mattila, H. Pirtonen and A. Törhönen for providing information to the climate indicator collection.

EXECUTIVE SUMMARY

The report has two main objectives: 1) to elaborate a set of reference criteria for the selection and evaluation of impact indicators, and 2) to review available climate impact indicators, their strengths and weaknesses according to the proposed framework.

In this report, 81 climate change and impact indicators are examined using agreed consistent reference criteria. These include aspects such as the methodological description of indicators, the input data requirements and availability for indicator calculation, treatment of uncertainty or fitness for purpose of indicator time series.

To provide a structured collection of indicators, we decided to group them into *tiers*. Tier-1 indicators focus mainly on the state and changes in the climate system, Tier-2 indicators provide information on the impacts of climate change on bio-physical systems. Tier-3 indicators are mostly used to indicate how socio-economic systems are expected to be affected by climate change. The analysis of the scientific and technical strengths and weaknesses of indicators was feasible at an aggregated level. A particular strength of gathered indicators is the availability of easy-to-access input data for their calculation. This is mostly the case for Tier-1 and Tier-2 indicators, while half of the Tier-3 indicators are based on data with restricted access. Beside, the total number of Tier-3 indicators is relatively small in comparison with Tier-1 and 2.

Uncertainty analysis is identified as one key criterion for the objective assessment of the given indicators. According to the indicator documentation, some information on uncertainties was available for approximately 2/3 of the indicators. The detail of description varies, but in general information on uncertainty stemming from the method and the input data source were provided. An apparent weakness of the indicators documented is the lack of regular updating. This is particularly the case when the indicators have been developed and presented as the output of specific research projects and are not maintained by organizations responsible for monitoring or statistical data. An evaluation of strengths and weaknesses on an indicator-by-indicator basis has been proposed and conducted for particular cases but further detailed analysis is still required.

This report further explores the strengths and weaknesses of indicators in the context of user expectations as evaluated in the research project Climate Information Platform for Copernicus (CLIPC)¹. User consultation activities helped to identify general uses of indicators (e.g. production of risk and vulnerability assessments), but details on particular applications of indicator by users are missing. The evaluation of indicators from a user's perspective is limited due to incomplete knowledge of how much weight a particular user might attribute to particular strengths and weaknesses. The report gathered an extensive set of information on climate change and impact indicators and developed the approach for analysing strengths and weaknesses of impact indicators provided by the CLIPC portal. Consequently, this information will more generally support the development of the Copernicus Climate Services and the indicators that such services will promote.

The sample of indicators collected at the time of writing has been observed to match the user needs for using indicators as input for climate research and for the purposes of raising societal awareness. The indicators can support the elaboration of adaptation strategies and vulnerability studies. It is still preliminary to make definitive judgments on the usefulness of each individual indicator due to limited knowledge on how specific indicators are used.

¹ <http://www.clipc.eu>

CONTENTS

ABSTRACT	3
TIIIVISTELMÄ	3
SAMMANDRAG	4
ACKNOWLEDGMENTS	5
EXECUTIVE SUMMARY	6
ABBREVIATIONS	9
1 Introduction	10
2 Previous work on climate change and impact indicators	13
2.1 Climate change, impacts and vulnerability in Europe 2012: An indicator-based report	14
2.1.1 Objective, data, coverage and scenarios	14
2.1.2 Climate change and impact indicators	16
2.2 ESPON Climate: Climate change and territorial effects on regions and local economies in Europe.....	16
2.2.1 Objective, data, coverage and scenarios.....	16
2.2.2 Climate change and impact indicators	17
2.3 Urban Vulnerability Indicators and associated ETC scoping study.....	18
2.3.1 Objective, data, coverage and scenarios.....	18
2.3.2 Climate change and impact indicators	19
2.4 ENSEMBLES: Climate change and its impacts at seasonal, decadal and centennial timescales	20
2.4.1 Objective, data, coverage and scenarios.....	20
2.4.2 Climate change and impact indicators	20
2.5 ISIMIP: The Inter-Sectoral Impact Model Intercomparison Project	22
2.5.1 Objective, data, coverage and scenarios.....	22
2.5.2 Climate change and impact indicators	22
2.6 IMPACT 2C: Quantifying projected impacts under 2°C warming.....	23
2.6.1 Objective, data, coverage and scenarios.....	23
2.6.2 Climate change and impact indicators	23
2.7 PESETA I and II: Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up Analysis.....	24
2.7.1 Objective, data, coverage and scenarios	24
2.7.2 Climate change and impact indicators.....	24
2.8 Preliminary take-home messages	25
2.8.1 Challenges in specifying Tier-3 indicators and suggestions to improve the societal relevance of Tier-1 and Tier-2 indicators.....	26
2.8.2 Impact indicators and decision making	28
3 Criteria for examining climate impact indicators	29
3.1 General development of criteria	29
3.2 Scientific adequacy and feasibility	32
3.3 Usability, relevance and scope of use	33
4 Strengths and weaknesses of documented indicators.....	36
4.1 Indicator database.....	36
4.2 Scientific and technical evaluation of indicators	38
4.2.1 Scientifically documented relationship	39
4.2.2 Methodological transparency	40
4.2.3 Recognition of and ability to deal with uncertainty	40
4.2.4 Public availability of relevant data	42

4.2.5 Updating frequency of relevant data, length of time series and spatial resolution.....	42
4.2.6 Indicator-by-indicator evaluation	43
4.3 User needs	45
5 Conclusions	50
REFERENCES	52
ANNEX 1. Expert workshop at the EEA	54
ANNEX 2. Investigation of indicators according to the potential uses	63
ANNEX 3. Schematic illustration of SRES scenarios	66

ABBREVIATIONS

CDR	Climate Data Record
CLIPC	Climate Information Platform for Copernicus, EU FP7 research project (2013-2016)
CMIP5	Coupled Model Intercomparison Project Phase 5
DGVM	Dynamic Global Vegetation Model
ECV	Essential Climate Variable
EEA	European Environment Agency
ENSEMBLES	EU FP6 Integrated Project (2004-2009)
EPA	United States Environmental Protection Agency
ESGS	Earth System Grid Federation
ESPON Climate	Climate Change and Territorial Effects on Regions and Local Economies in Europe, project (2009-2011)
ETC	European Topic Centre
EU	European Union
FP6 and FP7	European Union 6 th and 7 th Framework Programme for Research and Technological Development
FWI	Forest fire Weather Index
GCM	Global Climate Model
GEOSS	Global Earth Observation System of Systems
ICCC	Indicators of Climate Change in California, report
ICDC	Integrated Climate Data Center
IMPACT2C	Quantifying projected impacts under 2°C warming, EU FP7 research project (2011-2015)
IPCC	Intergovernmental Panel on Climate Change
ISIMIP	Inter-Sectoral Impact Model Intercomparison Project
JRC	Joint Research Centre
NPP	Net Primary Production
NUTS	Nomenclature of territorial units for statistics
PESETA	Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up Analysis, project by JRC
RCM	Regional Climate Model
RCP	Representative Concentration Pathways
SEA	Strategic Environmental Assessment
SEBI	Streamlining European Biodiversity Indicators
SMHI-RCA	Swedish Meteorological and Hydrological Institute Rossby Centre regional atmospheric model
SRES	Special Report on Emission Scenarios, IPCC special report
SSP	Shared Socio-Economic Pathways
WSDI	Warm Spell Duration Index

1 Introduction

This report aims to develop a consistent framework for examining climate impact indicators, as well as delivering a state-of-the-art review on selected climate impact indicators. The concept of an impact indicator is likely to find many meanings, reflecting the different research perspectives that study the evolution of climate and its consequences for the environment and societies. The term has not been unambiguously defined in relation to climate change. The European Environment Agency (EEA 2012a, p.35) specifies indicators with reference to their purpose. If the purpose is “understanding the causes of impacts of climate change”, then the report refers to “climate change indicators”. One can assume that “understanding” involves some type of “description” and that “causes of impacts” refer to “changes in the climate system”. In broad terms an indicator can be defined as a measure of the state of a particular system that provides a way to track the evolution of more complex processes, such as different aspects of climate change. An indicator provides information about complex processes while maintaining a certain degree of simplicity.

There is also ambiguity with respect to the use of the term ‘index’. The Integrated Climate Data Center (ICDC) defines climate indices as a “calculated value that can be used to describe the state and the changes in the climate system”². “Climate indices” are usually measures that have been agreed on and are based on standardized calculation routines, while indicators are also used in a much wider sense. Sometimes an indicator which is constructed by combining two or more distinct metrics can also be called an index. An example is the Palmer Drought Index, which is a measurement of dryness based on recent precipitation and temperature (Palmer 1965). A second example would be the Forest fire Weather Index (FWI). But there are also climate indices that are determined by making use of a simple climatic variable, for example, the number of frost days, which is calculated by the sum of days in one year with daily minimum temperature below 0°C. In practice the distinction between indicators and indices appears not to be that important or clear cut. The distinctions reflect conventions and traditions and thus climate researchers commonly refer to indices based on air temperature, precipitation, air pressure and sea surface temperature.³

In the field of climate change, the essential climate variables (ECV) are specified as a particular group of indicators. An ECV is “a physical, chemical, or biological variable or a group of linked variables that critically contributes to the characterization of Earth’s climate” (Bojinski *et al.* 2014, p.1432). The calculation procedures for ECV’s tend to be fixed as for climate indices. It is therefore no surprise that whatever the semantic used to describe a climate change indicator, climate indices or ECV, resulting findings show quite similar attributes. For example, the indicator “European temperature” (EEA 2012a) provides essentially the same information as the index “Mean of daily temperature” (European Climate Assessment & Dataset project) and the essential climate variable “Air temperature” (Global Climate Observation System).

Indicators or related concepts are not restricted to the presentation of data in the form of graphs and charts. EEA (2012a) specifically stresses that indicators should help “understanding the consequences of climate change and determining vulnerability” and therefore the indicators of the EEA include a narrative component. Climate change indicators can be used, and often are, to deal with concepts of vulnerability such as exposure, hazard or intensity (Costa and Kropp 2013). Similarly, climate change indicators are used as input for discerning the consequences of climate change.

² <http://icdc.cen.uni-hamburg.de/1/daten/climate-indices.html>

³ Climate Indices http://icdc.zmaw.de/climate_indices.html?&L=1 [visited September 13 2016]

An example is the occurrence of storms (denoted by wind velocity) in causing economic damages (Prahl *et al.* 2015).

For the purposes of this work, an impact indicator is described as an observed or projected measure that indicates a ‘relevant’ environmental/human/economic impact, and whose causes can be linked to the interaction between changes in climate and the system it portrays. A meta-classification of impact indicators into three Tiers (see Figure 1) is proposed, from indicators mainly concerning natural systems to those reflecting changes in human systems. An additional distinction can be made based on the timeframe. Generally indicators have been developed based on historical observation-based data to infer trends, but climate change scenarios also play an important role. The obvious distinction is that indicators based on historical data are primarily driven by available observations whereas projections are based on model outputs. Validation of model output and bias correction methods provide links between the two types of indicators.

In this categorization, Tier-1 indicators are intended to give information on the past and future evolution of the climate system. For example, mean temperature change, ice cover extent or sea-level rise provide indications of the impact on the climate system that are caused by anthropogenic interference with the global energy balance. Tier-1 indicators are often the departing point to derive higher Tier indicators. Tier-2 indicators attempt to quantify the impacts of climate change in bio-physical systems. Flood risks, crop losses, soil erosion and changes in distributional ranges or phenology of organisms are examples of such variables that can be used as indicators. Tier-3 indicators primarily aim at providing information on the socio-economic systems affected by climate change. These indicators usually build on previous ones and make the bridge from a bio-physical change to social or economic loss/gain. For example, indicators based on the economic consequences of extreme weather events or morbidity during heat waves belongs to this group. It comes without saying that the classification is not free of inconsistencies as there are indicators that overlap the classes proposed. Nevertheless, this structuring of indicators is useful for the purposes




Figure 1. A framework for climate impact indicator classification.

of this report, since it establishes a reference for better communicating its outcomes to general audiences.

For the sake of clarity, we provide some of the working definitions for this report.

- **Climate impact indicator** - an observed or projected measure that indicates a 'relevant' environmental/human/economic impact that can be linked to changes in the climate.
- **Tier-1 climate impact indicator** - A climate impact indicator primarily intended to give information on the past and future evolution of the climate system. Changes in temperature and precipitation extremes, arctic ice coverage or sea-level changes are examples of such variables that belong to this indicator category.
- **Tier-2 climate impact indicator** - A climate impact indicator primarily intended to quantify the impacts of climate change in bio-physical systems. Flood hazards, crop losses, changes in distributional ranges or phenology of organisms or soil erosion are examples of such variables that belong to this indicator category.
- **Tier-3 climate impact indicator** - A climate impact indicator primarily intended to provide information on the socio-economic consequences entailed by the changes in Tier-1 and 2 indicators. Crop-value loss, human casualties and economic losses from floods or storm events are examples of such variables that belong to this indicator category. Several Tier-2 indicators can be converted into Tier-3 indicators provided that reliable estimates can be provided on the economic consequences of physical impacts.
- **Climate indices** - Calculated value that can be used to describe the state and the changes in the climate system. Indices are often used as synonyms for indicators.
- **Essential Climate Variable** - A physical, chemical, or biological variable or a group of linked variables that critically contributes to the characterization of Earth's climate.

2 Previous work on climate change and impact indicators

This section reviews a number of European initiatives (projects, reports, activities) that have produced climate and climate impact indicators. The search for these initiatives was done by making use of the Climate-ADAPT portal and it was restricted to the time period from the start of the FP6 program to the end of 2014. In addition to the European-funded initiatives, the newest developments on climate impacts from the Intra-Sectoral Impact Model Intercomparison (ISIMIP) are also addressed (see Table 1).

Table 1. Overview of investigated studies on climate change and impact indicators.

Project/report/activity	Source	Objectives	Topics addressed
Climate change, impacts and vulnerability in Europe 2012: An indicator-based report.	EEA (2012a)	Assess past and projected climate changes and their impacts, and the associated vulnerability to society and ecosystems in Europe.	<ul style="list-style-type: none"> • Climate system • Environmental systems • Socio-economic systems and health • Vulnerability
ESPON Climate: Climate Change and Territorial Effects on Regions and Local Economies.	Greiving et al. (2011)	Assess the degree of vulnerability of different European regions to climate change and the impact of climate change on the region's economic, social, and environmental dimensions of European regions.	<ul style="list-style-type: none"> • Physical • Environmental • Economic • Social • Cultural
Urban Vulnerability Indicators-A joint report of ETC-CCA and ETC-SIA and Urban regions: Vulnerabilities, Vulnerability Assessments by Indicators and Adaptation Options for Climate Change Impacts (a Scoping Study).	ETC-CCA and ETC-SIA (2012) ETC/ACC (2010)	Propose a system of urban vulnerability indicators, for assessing where European cities stand in terms of vulnerability and adaptation.	<ul style="list-style-type: none"> • Heat waves • Floods • Droughts/water scarcity • Forest fires
ENSEMBLES: Climate change and its impacts at seasonal, decadal and centennial timescales EU FP6 research project 2004 to 2009	Van der Linden and Mitchell (2009)	Formulation of very high resolution Regional Climate Model Ensembles for Europe. Global mitigation scenarios. Probabilistic projections of climate change. Impact analysis, both with RCM and probabilistic projections.	<ul style="list-style-type: none"> • Climate system
ISIMIP: The Inter-Sectoral Impact Model Intercomparison Project. Research activity at the Potsdam Institute for Climate Impact Research	Schellnhuber et al. (2014)	Quantitative estimate of impacts and uncertainties for different sectors and from multiple impact models.	<ul style="list-style-type: none"> • Agriculture • Biomes • Forestry • Energy • Health • Coastal infrastructure • Marine ecosystems • Water

Project/report/activity	Source	Objectives	Topics addressed
IMPACT2C: Quantifying projected impacts under 2°C warming FP7 EU research project 2011-2015	Vautard <i>et al.</i> (2014)	Identify and quantify the impacts and most appropriate response strategies of a 2°C global warming for Europe and three selected vulnerable regions in other parts of the world	<ul style="list-style-type: none"> ● Water ● Energy ● Infrastructure ● Tourism ● Agriculture ● Forestry ● Ecosystem services
PESETA I and II: Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up analysis, Project funded by JRC	Ciscar <i>et al.</i> (2014)	Consistent multi-sectoral assessment of economic impacts of climate change in Europe for the 2071-2100 time horizon.	<ul style="list-style-type: none"> ● Agriculture ● Coastal systems ● River floods ● Tourism ● Human health ● Energy* ● Transport infrastructure* ● Forest fires* ● Habitat suitability* <p>* included in PESETA II</p>

2.1 Climate change, impacts and vulnerability in Europe 2012: An indicator-based report

2.1.1 Objective, data, coverage and scenarios

In 2012 the European Environment Agency (EEA 2012a) compiled information on past and projected climate change, and related impacts in Europe, based on a range of indicators. The report aimed at providing a strong knowledge base for the development and implementation of adaptation strategies and actions at both national and EU levels. Furthermore the report updates and improves earlier indicator-based assessments of climate change impacts and vulnerability published by the EEA, namely in 2004 and 2008. The indicators gathered are made accessible via the web-based EEA indicator management system⁴ and the European Climate Adaptation Platform ClimateADAPT⁵. Approximately 40 indicators are included in the EEA 2012a report. The indicators have been organized in three broad topics. These topics are: *Changes in the climate system*; *Impacts on environmental systems* such as the coastal zones, soil or inland waters; and *Impacts on socio-economic systems and health* such as agricultural systems, energy or transport. The report contains a chapter dedicated to indicators of vulnerability to climate change, such as indicators of damage costs, as well as integrated approaches to operationalize the concept of vulnerability taken from the ESPON project. A tentative matching of the indicators provided in the EEA 2012a report to the Tier classification (Figure 2) shows the predominance of indicators associated with Tier-1 and 2, respectively 46 and 48% of the total indicator set. Tier-3 indicators comprise only about 6% of the indicator set reported in the EEA 2012a report.

The report included both observations and projections for the majority of the indicators. Indicators have been quantified using existing information; hence, the use of climate models, forcing scenarios, spatial resolutions and time frames for projections varied between indicators as the data were obtained from a large number of independent studies. Full harmonization of indicators with respect to models, climate scenarios, time frames or spatial coverage was therefore impossible. As an example, while the indicators under the topic “Changes in the climate system” are usually derived from large ensembles of Regional Climate Models (RCMs) and

⁴ <http://www.eea.europa.eu/data-and-maps/indicators>

⁵ <http://climate-adapt.eea.europa.eu>




Figure 2. Fraction of indicators provided in EEA (2012a) tentatively allocated according to the Tier classification.

Global Climate Models (GCMs), a substantial number of indicators belonging to the topic “Impacts on environmental systems” and “Climate impacts on socio-economic systems and health” are obtained from studies that have used a single or two climate models. This is the case for indicators providing information on agro-phenology, distribution and abundance of animal species, forest fires or water requirements. The use of socio-economic scenarios is also non-systematic; although in this case a considerable fraction of indicators are obtained for the Intergovernmental Panel on Climate Change (IPCC) Special Report Emission Scenarios (SRES) A1B (IPCC 2000), see Figure 3 for details. For the indicators with spatially-explicit projections (in the form of a map), about 40% had been derived using the A1B scenario. Indicators that have been calculated according to one of the other SRES scenarios constitute about 20% of the indicators. For approximately 40% of the indicators included in the EEA report, there was no spatial representation (maps) of future projections. For these indicators the report provided concise text with information on projections available in the literature; many of these also used SRES scenarios.

The time frame for which projections are available is also rather variable across and within the investigated topics. The time frames of 2021-2050 and 2071-2100 were most commonly used for the indicators presented in a spatially-explicit manner.




Figure 3. Fraction of indicators with spatial-explicit representation (maps) provided in EEA (2012a) according to the socio-economic scenario used (For a description of the socio-economic scenarios please refer to the Annex 3).

2.1.2 Climate change and impact indicators

The set of indicators compiled is based extensively on peer-reviewed as well as non-peer-reviewed work. Although it is mentioned in the report that the selection of indicators to be included adhered to criteria documented in an European Topic Centre (ETC) technical paper published in 2013 (Hildén and Marx 2013), the report falls short in making transparent how the selection process was carried out. Although both preceding reports in 2004 and 2008 provide explicit criteria for indicator selections, specific details remain unclear. Examples of such criteria are: the policy relevance of an indicator, its strength in establishing the causal links between climate change and observed impacts, its methodological soundness, issues of data quality, the availability of the indicator for long periods of observations, and information on robustness and uncertainty.

The assessment of past and projected climate change and impacts is reported mostly for indicators belonging to the topics *Changes in the climate system* and *Climate impacts on environmental systems*. The indicator set under these two topics ranges from basic (Tier-1) climate change indicators such as temperature averages and extremes, wind speeds or snow cover length; to more elaborated (Tier-2) indicators employing biophysical (e.g. river flow and flood return levels) or envelope models (e.g. distribution of plant and animal species). Regarding the investigation of *Climate impacts on socio-economic systems and health*, the report contains indicators that cover most of Europe's economic sectors; including agriculture, biodiversity, forestry, energy, transport, tourism, fisheries and human health. Also here the indicators range from simple climate indices constructed via the use of basic climate variables (e.g. heating degree days or flowering date of winter wheat) to metrics that imply the use of biophysical or statistical modelling, such as agricultural yield.

Indicators that could be mostly related to the Tier-3 classification, such as people affected, are included in the EEA 2012a section referring to indicators of *Vulnerability to climate change*. Under this topic indicators are largely dominated by human systems such as costs of flood damages or other natural disasters or projected economic costs of climate change. The establishment of systematic indicators has proved to be challenging due to lack of systematic data collection and analysis. The report specifically noted that there is a need for enhanced and sustained monitoring in Europe of "environmental systems, socio-economic systems and health, and of costs of damages of extreme weather events" (EEA 2012a, p.237).

2.2 ESPON Climate: Climate change and territorial effects on regions and local economies in Europe

2.2.1 Objective, data, coverage and scenarios

The ESPON Climate⁶ project had the objective of assessing the degree of vulnerability of different European regions to climate change. In this light, it was not an explicit objective of ESPON Climate to collect, or elaborate, impact indicators. Instead, the project strived to operationalize the concept of vulnerability for European regions by using an adapted version of the Füssel and Klein (2006) vulnerability framework. In order to determine the vulnerability of a system a number of intermediate steps have to be fulfilled, among those, the determination of the potential climate change impacts. Potential impacts are framed as a combination of climatic exposure (Tier-1

⁶ http://www.espon.eu/main/Menu_Projects/Menu_AppliedResearch/climate.html

indicators) and sensitivity. In this particular case sensitivity is assessed via socio-economic and bio-physical conditions of each system under analysis, for example, population, infrastructure or landscape. Thus, impact metrics in ESPON are, in theory, mainly related to Tier-2 and Tier-3 indicators as defined in this report. Individual impact metrics are divided into physical, environmental, social, cultural and economic dimensions. The project also devoted large efforts to the aggregation of impact metrics within and across each dimension.

Figure 4 illustrates the number of individual climate exposure and potential climate indicators, according to the five dimensions assessed, and available from ESPON Climate. Indicators of climatic exposure and those depicting the potential physical and environmental impact of climate change in European regions dominate the indicator set. Metrics informing on the potential social, cultural and economic impacts are the least represented in the indicator set, a characteristic also noted in the EEA 2012a




Figure 4. Number of individual climatic exposure and potential impact indicators in ESPON Climate. ESPON climate indicators are tentatively color-coded according to the Tier classification. Indicators informing on cultural impacts have not been coded and are shown in grey.

study. We have coded, in a tentative manner, the indicators available from ESPON climate according to the Tier classification in this report. Most of the indicators can primarily be classified as Tier-1 and 2 indicators. One of the most prominent characteristics of the ESPON Climate project has been to use, as far as possible, consistent socio-economic scenarios and time frames. The regional model COSMO-CLM⁷ was adopted for climate change runs with three realizations for the time period 1961-1990 and two realizations for each scenario for the time frame 2001-2100 based on the IPCC A1B scenario. Indicators of climate exposure always indicate the change of climate conditions from the reference time period (1961-1990) to those expected in the time period 2071-2100. Also consistent across the study is the homogenization of indicators to the same administrative level, in this case NUTS-3 regions.

2.2.2 Climate change and impact indicators

The indicators of climatic exposure in ESPON (Tier-1) were largely identical to those proposed in the EEA 2012a report. For example, *change in mean annual temperature, numbers of frost days, snow cover duration, mean precipitation and extreme precipitation* or

⁷ <http://www.clm-community.eu/index.php?menuid=198>

changes in the 100 year return flood level are included in both studies. The main difference is a stronger focus of EEA 2012a on the indicators indicating the state and evolution of the cryosphere. Only the exposure indicators of river and coastal flooding required additional processing of the output given by COSMO-CLM. For the case of river flooding, flood heights from the LISFLOOD⁸ hydrological model were used (also used in EEA 2012a to evaluate river flows). For coastal flooding ESPON made use of a tailor-made approach combining storm-surge heights from the Dynamic Interactive Vulnerability Analysis tool and a global digital elevation model.

Metrics on potential impacts for the physical, environmental, societal, economic and cultural dimensions (Tier-2) result from a deductive approach that is, using available scientific knowledge in form of frameworks, theories or models about the vulnerability of the system of interest in the selection and aggregation of indicating variables (Hinkel 2011). In ESPON Climate potential impacts were determined by combining climatic exposure indicators with the sensitivity of a system using in most of the cases previous knowledge from analogue work or specific case studies. As an illustrative example the metric depicting the *potential impact of climate change on airports and harbours* (due to floods) was determined by overlaying inundated areas (tailor-made approach, see above) and corresponding changes in inundation heights (LISFLOOD, see above) with a map of the infrastructure networks and facility locations. The logic in this case, and very much for all indicators of potential impact, is the following: if the same geographical region scores high in the intermediate indicators of exposure and sensitivity, then the potential impact is also expected to be high. The individual impact score for a region is normalized between 0 and 1 (although the ESPON project also provides the original scores) according to the maximum and minimum distribution of impact scores for the NUTS-3 regions. This normalization implies that all European regions are ranked between the lowest and highest absolute scores. An interesting feature of the ESPON project was the very high level of indicator aggregation of such normalized scores across the physical, environmental, social, economic and cultural impacts of climate change.

2.3 Urban Vulnerability Indicators and associated ETC scoping study

2.3.1 Objective, data, coverage and scenarios

The Urban Vulnerability Indicators study (ETC-CCA and ETC-SIA 2012) aims at proposing a system of urban vulnerability indicators, which would allow an assessment of European cities in terms of vulnerability and adaptation, and the areas where certain problems cluster. The study is a follow-up of the 2010 ETC/ACC⁹ scoping study on vulnerabilities to climate change hazards in urban regions. Therefore it is sensible to analyse both together for the purposes of this review. Both reports focus on assessing vulnerability indicators for the urban space. This is a new feature in this short review since until now we have been mostly evaluating work that dealt with a large number of economic sectors. At the core of both works sits the same vulnerability framework as in EEA 2012a and ESPON Climate (see above). Both urban studies are preparatory work that is currently followed up by implementation of a selected number of indicators by the European Topic Centers on Spatial Analysis and Information and on Climate Change Adaptation (ETC/SIA and ETC/CCA).

⁸ <https://ec.europa.eu/jrc/en/publication/eur-scientific-and-technical-research-reports/lisflood-distributed-water-balance-and-flood-simulation-model-revised-user-manual-2013>

⁹ http://acm.eionet.europa.eu/reports/ETCACC_TP_2010_12_Urban_CC_Vuln_Adapt

The Scoping study reviews a total of 26 vulnerability indicators for the urban space distributed across the topics of heat, decreased precipitation and drought, wildfires, fluvial flooding, intense precipitation, sea-level and coastal flooding. Vulnerability indicators are composed of exposure, sensitivity and, at times, adaptive capacity components. Figure 5 shows the distribution of climate exposure indicators across the investigated themes in the ETC Scoping study. Climate exposure indicators for heat, sea-level rise and coastal flooding dominate the (exposure) indicator set make up approximately 50%.

The Urban Vulnerability Indicator study presents climate exposure indicators for the themes of heat, floods (both fluvial and coastal), water scarcity/droughts and forest fires. With the exception of heat, for which two climate exposure indicators are considered, the remaining topics include a single climate exposure indicator. While most of the exposure indicators available can be related to the Tier-1 classification, some can be related to Tier-2. This is the case of indicators for fluvial and coastal flooding, which are, at times a, combination of a potential flood height and its propagation over the terrain.




Figure 5. Climate exposure indicators across the investigated themes of the ETC scoping study color-coded according the Tier classification.

2.3.2 Climate change and impact indicators

Examples of climate exposure indicators for the case of heat in the Scoping study are: Warm Spell Duration Index (WSDI), tropical nights, heat wave days, days with temperature above 30 °C or changes in average December, January and February maximum temperature by 2030. These indicators are primarily Tier-1 indicators, since they are selected or constructed from primary climate data without further impact modelling to capture a specific impact on human systems. Exposure indicators for pluvial flooding were found to be similar to those in EEA 2012a and ESPON Climate, namely, river flow and inundation depth, and coastal flooding, with emphasis on indicators such as inundated area and changes in storm surge height (Tier-2). Some of the exposure indicators use insights from vulnerability studies applied in urban regions outside Europe. As an illustrative case, the climate exposure metrics used in the case of wildfires refer specifically to those used in an Australian bush fire vulnerability indicator (see Preston *et al.* 2008).

The Urban Vulnerability indicator study narrows down the urban vulnerability indicators to a manageable number. The proposed set of indicators is derived from the

Scoping study. The final set of indicators to measure climatic exposure is: heat: combined number of hot days and warm nights, and effective temperature; Floods: area prone to flooding (both from fluvial and coastal flooding); Water scarcity/droughts: standard precipitation index; Forest fires: fire probability index. The report neither provides details on how this selection took place nor which indicator criteria (e.g., methodology or coverage) were used.

2.4 ENSEMBLES: Climate change and its impacts at seasonal, decadal and centennial timescales

2.4.1 Objective, data, coverage and scenarios

The ENSEMBLES project (Van der Linden and Mitchell 2009) aimed at providing researchers, decision makers, businesses and the public with climate information obtained through the use of the (at the time) latest climate modelling and analysis tools. The central feature of the project was the running of multiple climate models in order to improve the accuracy and reliability of results. The information was envisioned to help policy makers, at all levels, in determining future strategies to address climate change. From the many topics addressed in the ENSEMBLES project two are of particular interest: (i) the probabilistic estimate of uncertainty in future climate variables at seasonal to decadal and longer time-scales; and (ii) a linkage of outputs of the ensemble climate predictions to a range of sectoral impacts including agriculture, health, food security, energy, water resources, insurance and weather risk management.

Gridded observational datasets of daily precipitation and temperature have been developed using a European network of high-quality station series. The datasets cover the period from 1950 to 2008. A set of multi-model simulations was produced over the period 1860-2000 to simulate the long-term climate conditions. Subsequently, a multi-model set of coupled simulations over the 21st century was produced for the A2, A1B and B1 IPCC scenarios. ENSEMBLES made considerable efforts to construct probabilistic high-resolution regional climate scenarios and seasonal-decadal *hindcasts*. Results are available at 25 km resolution. For particular climate variables and regions, downscaling methods were applied to GCM output in order to obtain both climate change projections extended up to 2100 as well as seasonal to decadal *hindcasts*. The downscaled climatic variables all belong to Tier-1: daily temperature and precipitation, minimum and maximum temperature, marine surface wind, drought indices, river discharge, solar radiation, vapour pressure, wind speed and relative humidity.

2.4.2 Climate change and impact indicators

The ENSEMBLES project did not aim, as a core objective, to provide climate change or impact indicators. It did nevertheless support 1) the integration of process models of impacts on the natural and managed global environment into Earth System Models and 2) the modelling of the extreme weather events to evaluate impact risks. For example, the Dynamic Global Vegetation Model¹⁰ (DGVM) LPJmL was forced with the projected climatic patterns from seventeen general circulation models used in the ENSEMBLES project. A number of what we could call Tier-2 indicators were derived

¹⁰ <https://www.pik-potsdam.de/research/projects/activities/biosphere-water-modelling/lpjml>

from the exercise and reported for the global scale: Tree cover, Net Primary Production (NPP), heterotrophic respiration, evaporation, river runoff and incidence of fire.

With respect to extreme events, a number of impact models have been used in ENSEMBLES to define the nature of extreme events and their impacts. These impact assessments were carried out across a number of regions and topics, for example: potential changes in energy demand in the Mediterranean or changes of fire risk in *Fennoscandinavia*. To assess potential changes in energy demand a number of impact indicators were generated such as changes in cooling and heating degree days, mean change in cooling degree days and the standard deviation of change. To assess fire risk in *Fennoscandinavia* the Finnish Fire Index was used, using projections from a 100-year simulation with the SMHI-RCA (Swedish Meteorological and Hydrological Institute Rossby Centre regional atmospheric model) Regional Climate Model. In addition to the more global/regional efforts in providing climate change impact assessments, the ENSEMBLES project elaborated 11 more detailed case studies in Europe for which both climate change and impact indicators were generated. These are summarized in Table 2.

Table 2. Case study regions and downscaled indicators from the ENSEMBLES project

Study region	Indicator	Comment
Castilla León	Changes in phytoclimatic indices	Index represents the suitability of a certain species to live in a given region depending on its climate.
Spain	Mean and extreme precipitation.	Different ENSEMBLES RCMs used to reproduce the mean and extreme precipitation regimes in Spanish hydrological basins.
Andalucia	Changes in bioclimatic and drought indices	Percentage changes in four bioclimatic types (humid, semi-humid, dry and semi-arid) in Andalucía.
North sea	Decadal, monthly and daily means of 10 m wind components (u & v),	Approach consisted of a multi linear regression (MLR) model for spatial downscaling and a multi-variate auto regression (mvAr) model to generate highly temporal time series of wind components.
Rhine basin	Annual maxima of 10-day precipitation sums	Data to used in driving a hydrological model of the Rhine basin to study potential changes in the occurrence of extreme river discharges.
Alps	Changes in winter snow water equivalent	Ensemble mean, minimum and maximum based on six ENSEMBLES regional climate change scenarios assuming SRES A1B emissions
Northern Italy	Changes in temperature extremes	Statistical downscaling applied to several GCMs to construct probability density function (PDFs) of changes in temperature extremes over Northern Italy
Scandinavia	Frequency of second- and third-generations of bark beetles	Indicator resulted from impact modelling that used ENSEMBLES outputs of climate data as input.
Romania	Changes in extreme precipitation	Example: Mean frequency (number of days) of summer daily precipitation exceeding 15 mm/day at the Calarasi station (Romania)
Danube	Changes in river flow extremes	Changes in river flow extremes are associated to the atmospheric predictors of sea level pressure (SLP), geopotential, temperature, specific and relative humidity
Mediterranean	Changes in temperature and rainfall extremes	Example of indices determined: frequency of hot days ($T_{max} > 35^{\circ}\text{C}$), tropical nights ($T_{min} > 20^{\circ}\text{C}$) and length of maximum dry spell

2.5 ISIMIP: The Inter-Sectoral Impact Model Intercomparison Project

2.5.1 Objective, data, coverage and scenarios

ISIMIP¹¹ is a community-driven modelling effort with the goal of providing cross-sectoral global impact assessments, based on Representative Concentration Pathways (RCPs) and socio-economic Shared Socio-Economic Pathways (SSPs) scenarios. Its aim is similar to those of model intercomparison initiatives that are sector-specific, for example AgMIP¹² or waterMIP¹³ for the cases of agriculture and water respectively and which are included in the ISIMIP network. The first efforts of ISIMIP were devoted to the elaboration of a common climate dataset and bias correction to serve as input to the different impact models. This was achieved during the project fast track (until May 2013). During this phase a total of 5 GCMs has been used, as well as approximately 30 impact models covering the sectors of agriculture, biomes, water, health (restricted to malaria) and coastal infrastructure. In order to guarantee a minimum consistency of model outputs, a set of basic requirements was adopted by impact modellers during the fast track phase. All RCP concentration scenarios are to be run using data from one GCM. Four additional GCMs are only considered together with those RCPs producing the highest and lowest end-of-century forcing (RCP8.5 and RCP2.6 respectively). If applicable, only the middle-of-the-road socio-economic scenario (SSP2) is used in the minimal setting. Highly relevant sensitivities (e.g. to CO₂ fertilization) are also considered. Bias corrected climate data from the GCMs participating in Coupled Model Intercomparison Project Phase 5 (CMIP5) are provided. Data cover the time period from 1950 to 2099.

2.5.2 Climate change and impact indicators

The full output dataset of the ISIMIP fast track is available via an Earth System Grid Federation (ESGS) node¹⁴. Due to the large number of impact models and sectors assessed, the outputs of ISIMIP that could be considered to be climate impact indicators are substantial. Figure 6 illustrates the diversity of impacts indicators (understood in this case as model output variables) provided by the ISIMIP initiative as classified according to our Tier-framework. Output variables related to impact modelling in the water sector dominate the “indicator set”. These were found to be mostly dominated by Tier-2 indicators such as run off, soil moisture and irrigation demand. Tier-1 indicators include snowfall, rainfall, snow water equivalent and evapotranspiration. The biomes sector accounts only for Tier-2 indicators or model output variables (e.g. NPP, vegetation type or leaf area index). Output variables for the agricultural sector were divided into those emanating from biophysical modelling (Tier-2) and those resulting from agro-economic modelling (Tier-2 and 3). Regarding the latest, more than half were identified to be Tier-3 indicators, for example, average producer prices, total calorie consumption, water and land prices.

The two least represented sectors in terms of number of output variables are the sectors of health (in ISIMIP fast track restricted to the malaria issue) and coastal

¹¹ <https://www.isimip.org/>

¹² <http://www.agmip.org/>

¹³ <http://www.eu-watch.org/>

¹⁴ esg.pik-potsdam.de

infrastructure. For the latter, within the ISIMIP, only one model was used of which the output could be regarded as Tier-3 indicators: expected number of people flooded annually, expected sea-flood costs, adaptation costs of building, upgrading and maintaining dikes.




Figure 6. Number of Tier-1, -2 and -3 indicators for sector specific output variables

2.6 IMPACT 2C: Quantifying projected impacts under 2°C warming

2.6.1 Objective, data, coverage and scenarios

The project IMPACT2C (2011-2015) provided information and evidence on the impacts of 2 °C global warming for Europe and other key vulnerable global regions. The project aimed to consider the impacts from a cross-sectoral perspective, e.g. for particularly vulnerable areas that are subject to multiple impacts where cumulative effects may arise and in relation to cross-cutting themes. The work flows from climate information, its uncertainty processing, via the evaluation of impacts, vulnerabilities and risks, to cross-sectoral assessments and synthesis highlighting risks, trade-offs, synergies and costs at a pan-European level.

A global warming of 2 °C relative to pre-industrial climate has been proposed as a threshold which society should endeavour to remain below, in order to limit the dangerous effects of anthropogenic climate change. The IMPACT2C project started comparing the new RCP model runs to the A1B scenario, looking at the possible changes in regional climate under this target level of global warming.

2.6.2 Climate change and impact indicators

The possible changes have been investigated by analysing Tier-1 climate change impact indicators for Europe, i.e. robust changes in mean and extreme temperature, precipitation, winds and surface energy budgets. The project results (Vautard *et al.* 2014) indicate a large likelihood that most of Europe will experience a greater increase in heat extremes in Southern Europe, a robust increase in heavy precipitation and an increase in extreme winds in winter in Central Europe. The findings of the analysis of Tier-1 climate change indicators revealed also strong distributional patterns across Europe, which are important in the subsequent impact assessments. As a second step,

the project used a range of models to assess the 2°C global warming effects on water, energy, infrastructure, coasts, tourism, forestry, agriculture, ecosystems services, and health and air quality-climate interactions. The findings, dominated by Tier-2 and Tier-3 indicators, are presented as an interactive web-atlas.¹⁵

2.7 PESETA I and II: Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up Analysis

2.7.1 Objective, data, coverage and scenarios

The objective of the Joint Research Centre (JRC) PESETA II project (Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up Analysis) is to make a consistent multi-sectoral assessment of the impacts of climate change in Europe for the 2071-2100 time horizon. The project methodology has two distinctive features. Firstly, it is based on bottom-up biophysical impact models results. Bottom-up models take into account the relationship between climate change and biophysical impacts in a structural way, modelling all the relevant interactions and mechanisms. Secondly, the assessment is made in a consistent way, where all biophysical impact models use the same climate data.

For the JRC PESETA II study climate simulation runs were obtained from the ENSEMBLES project (see above). Runs were driven by the SRES (Special Report on Emission Scenarios) A1B emission scenario and the so called E1 emission scenario. The E1 scenario was developed within ENSEMBLES (Van der Linden and Mitchell 2009) as an attempt to match the European Union target of keeping global anthropogenic warming below 2°C above pre-industrial levels. Climate change runs were available for two resolutions, 25 and 50 km. A total of 7 RCMs and 6 GCMs are used to obtain climate change runs. As for the resolution of biophysical and economic impacts, these are largely variable. In case of agriculture (using the JRC owned BioMA (Biophysical Models Applications) framework)¹⁶ results are available for a 25 km grid cell, for the case of impacts on tourism the output resolution is the NUTS (Nomenclature of territorial units for statistics)-2 level. Economic impacts of climate change are only available for a highly aggregated level, such as major European regions (e.g., Northern Europe or Central Europe).

2.7.2 Climate change and impact indicators

The PESETA I and II projects have determined climate change and impact indicators across all the Tiers considered. Most of the Tier-1 indicators have been calculated on a daily basis, although some are also available on monthly (in regard to Forest species habitat suitability) and yearly basis (in particular in case of forest fire analysis). The indicators are basically several variations of temperature, precipitation, humidity and wind variables. Of particular interest are results from the PESETA project regarding the biophysical impacts (Tier-2) resulting from the projected changes in climate. These are summarized in Table 3 and serve as inputs to the determination of economic impacts (Tier-3) using the GEM-E3 (General Equilibrium Model for Economy – Energy – Environment) model¹⁷.

¹⁵ <https://www.atlas.impact2c.eu/en/>

¹⁶ <http://bioma.jrc.ec.europa.eu/>

¹⁷ <https://ec.europa.eu/jrc/en/gem-e3/model>

The outputs of biophysical models can be better linked to the Tier-2 indicators. The consistency of climate models and scenarios used, allows for inter-indicator comparability. The economic estimates of climate impacts (Tier-3) produced by PESETA are in principle very valuable since it is rather uncommon to find such a comprehensive and extensive sectoral coverage in economic impact assessments of climate change. The main disadvantage of the PESETA results is the highly aggregated spatial nature of Tier-3 indicators.

Table 3. Biophysical output used to run the GEM-E3 model by sector/theme investigated.

Sector/theme	Biophysical model output
Agriculture	Yield change
Energy	Heating and cooling demands
River floods	Residential building damages Production activity losses
Forest fires	Burnt area Reconstruction costs
Transport infrastructure	Changes in cost of road asphalt binder application and bridge scouring Net change in costs related to extreme flooding and winter conditions
Coastal areas	Migration costs Sea-flood costs
Tourism	Tourism expenditure
Human health	Hours lost due to morbidity and mortality Additional health expenditures Warmer temperature Mortality

2.8 Preliminary take-home messages

As noted in the introduction, there is no fully unambiguous definition of what a climate change impact indicator is. A dictionary definition that an indicator is “a sign that shows the condition or existence of something” or “a pointer or light that shows the state or condition of something” is generally fulfilled by all the variables and metrics that have been explored in the previous sections. An Essential Climate Variable (ECV) has been specified as a physical, chemical or biological variable or a group of linked variables that critically contributes to the characterization of the Earth’s climate (Bojinski *et al.* 2014). The EEA specifically includes the narrative that provides the wider context for the “something” in the indicator definition (EEA 2012a) whereas for example EMSEMBLES and ISIMIP primarily provide scientific data and outputs of models that can be used in exploring climate change impacts by others in a non-scientific context, e.g. policy context. For ESPON Climate, an impact indicator is the combination of an exposure (mostly climatic) and a sensitivity indicator (mostly of socio-economic character).

The review in Sections 2.1-2.7 has given an overview of the current state and availability of climate change and impact indicators (see Table 4 for study-specific indicator definitions). Indicators at Tier-1 are abundant and there are several parallel data sets that are good indicators of climate change as a phenomenon. The ECVs have been selected by virtue of their reliability and systematic quality controlled monitoring. From an impact perspective, one of their main roles is that they provide indication on the pressures that climate change exerts on natural and human systems.

Table 4. Understanding of climate change and impact indicators in the context of the evaluated studies.

Study	Indicators of climate change	Climate impact
EEA 2012a	Climate variables aggregated either in time or according to a given threshold	Combination of an exposure indicator (mostly a climate variable) and a sensitivity indicator (mostly a socio-economic variable).
ESPO Climate	Exposure indicator (mostly climate variables)	Combination of an exposure indicator (mostly a climate variable) and a sensitivity indicator (mostly a socio-economic variable).
ETC/ACC 2012-2012		Combination of an exposure and sensitivity indicators.
ENSEMBLES	Climate variables aggregated either in time or according to a given threshold	Output of biophysical modelling.
ISIMIP		Output of a biophysical or coupled biophysical and economic model.
IMPACT2C		Output of a biophysical or coupled biophysical and economic model.
PESETA I and II		Output of a biophysical or coupled biophysical and economic model.

Tier-2 indicators have been developed in numerous research projects that have aimed at linking observed changes in the climate with changes in biophysical systems. The review has shown that there is less fully standardized data that can be used to derive indicators that provide standardized information across wide geographical scales although there are numerous studies of particular regions and cases that link observed or projected changes in the climate to corresponding changes in the biophysical environment. One of the challenges lies in the geographical and temporal differences in the links between climate and the biophysical systems. Some systems may be sensitive to, for example, winter extremes whereas the duration of particular conditions may be more important for other regions.

There is a definitive lack of Tier-3 indicators. This is not primarily due to a lack of data on human systems in general. Statistics of societal changes are abundantly available, but the difficulties lie in identifying and verifying causal relationships between climate and societal changes. The data on human systems is systemic in the sense that it reflects changes in numerous different driving forces, only some of which are related to climate change. Even when there appears to be a fairly direct link between a particular set of observations and climatic conditions there are a number of confounding factors that may question the validity of the indicator as a sign of the impacts of climate change. For example monetary damage caused by floods and storms are clearly linked to extreme weather events. The monetary damage, as measured by the level of compensation paid by insurance companies, does not, however, reflect only impacts of climate change, but also the value of assets in affected areas, which may increase independently of climate change (Barredo 2010, Visser *et al.* 2012, Smith and Katz 2013). Therefore it appears that Tier-3 indicators are often more useful when they are framed in terms of vulnerability or adaptive capacity rather than in terms of actual impacts, which require detailed site specific analyses to deal with the question of attribution.

2.8.1 Challenges in specifying Tier-3 indicators and suggestions to improve the societal relevance of Tier-1 and Tier-2 indicators

The CLIPC project revealed that there were fewer Tier-3 indicators than Tier-2 and Tier-1 indicators. A similar distribution of indicators classified into Tiers 1, 2 and 3 was also seen in the review of existing indicator studies by the EEA and ISIMIP (see

sections 2.1 and 2.5). Yet it can be expected that Tier-3 indicators are the most appealing for those engaged in developing concrete policies and measures for adaptation (and mitigation).

One reason for the smaller number of indicators that describe socio-economic impacts of climate change is that these often require an analysis “further down the chain” combining information on purely bio-physical conditions with those describing the society and their effects on human activities. These entail more sources of uncertainty and tend to be restricted to smaller regions, although some Europe-wide analyses, for example on estimating the costs of climate change impacts in selected sectors, have been conducted (ClimateCost project, Watkiss (2011)). Possible Tier-3 indicators that attempt to project conditions to the future ideally require detailed scenarios of specific socio-economic variables.

An example that illustrates some of these challenges can be taken from the projections of flood risks and their effects on society (see Flörke et al. (2011) for a Europe-wide analysis). Models have been developed that simulate the risk of floods in a specific area for present-day and future climatic conditions (which could be based on a Tier-2 indicator). Future flood projections are dependent on assumptions about adaptation in the water management. Combining projections of flood areas with information on infrastructure, housing and population would allow defining a Tier-3 indicator; however, spatially detailed data on the latter is required to allow a spatial matching with projected flood zones. Projections of future floods would ideally be matched with scenarios of infrastructure, housing and population, again on a spatial scale that allows the matching with projected flood zones.

To overcome the possible underrepresentation of Tier-3 indicators, two approaches can be considered to increase the societal relevance of Tier-1 and Tier-2 indicators, without directly specifying new indicators that would fall in the Tier-3 category:

- The societal relevance of indicators (including Tier-1 and 2) can be described in a text section accompanying the indicator data, maps and graphs. This has been done for the EEA’s impact indicators (see section 2.1 – EEA 2012a), in which the text description for each indicator starts with a few paragraphs that outline the background and explains in general terms in which way an indicator is important. For example, for the Tier-1 indicator of snow cover, one can mention its relevance for transport and tourism/recreation.
- Spatially explicit Tier-1 and Tier-2 indicators can be overlaid with information about socio-economic conditions to identify regions where a high exposure to a bio-physical impact coincides with e.g. low regional financial resources or a large proportion of elderly. This has been done in indicator-based vulnerability assessments Europe-wide (e.g. ESPON-Climate, Greiving et al. (2011) – see section 2.2 above) and for European regions (e.g. for selected sectors in the Nordic region, Carter et al. (2016)). Developing a web-based mapping tool that would allow users themselves to do this overlaying is one of the directions that climate information portals could take. An example of such a web-based mapping tool has been presented by Carter et al. (2016)¹⁸ and by the CLIPC project¹⁹.

One of the main challenges in linking Tier-1 and Tier-2 indicators to socio-economic data is the attribution of changes to climate change. As long as the linking is exploratory and user driven, it can be seen as a search for possible connections

¹⁸ See <http://www.iav-mapping.net/U-C-IAV>

¹⁹ <http://www.clipc.eu/indicator-toolkit>

between climate change and socio-economic conditions. If one wishes to formalize a possible relationship and call it a Tier-3 indicator that expresses socio-economic consequences of climate change, the demands on rigorous evidence for the relationship increases considerably. The uncertainties need to be recognized and properly dealt with and this may not yet be possible. There is also a need to identify ownership and interest in regular updating of the indicator and the underlying data, meaning that any European socio-economic indicator should most likely rely on information that is produced by, for example, Eurostat.

2.8.2 Impact indicators and decision making

One of the justifications for developing indicators is that they should support decision making. The EEA (2012a), the ESPON Climate (Greiving *et al.* 2011) and the Urban vulnerability study (ETC/ACC 2010, ETC-CCA and ETC-SIA 2012) make this link explicit whereas research based projects such as ENSEMBLES, ISIMIP or IMPACT2C implicitly assume that the indicators that they produce or that can be derived from their output also serve decision making. The relationship between the decision making and indicators varies among the tiers (Table 5). Tier-1 indicators have been designed to provide information that is relevant for operational decisions or to give general background information for policy development. Tier-3 indicators can help focusing risk management strategies justifying policy initiatives, but are less useful for specific decisions. For example, indicators of overall average economic losses associated with climate change can support the planning and implementation of adaptation policies, but do not provide particularly useful guidance for specifying specific subsidies that increase adaptive capacity. Tier-2 indicators can justify action, but can also partially guide specific decisions. For example, the indicators of flood risk can be used to prioritize investment opportunities within river basins. The context determines the demand for indicators, but also how an indicator should be interpreted and displayed. Thus, there is a demand for very specific and spatially disaggregated Tier-1 indicators to support concrete decisions. General Tier-3 indicators are in greatest demand at a national or European aggregate level, where they can guide policy development. Tier-2 indicators are demanded both at a detailed level to guide design and at a general level to justify policies.

Table 5. The use of information from different Tiers in decision making.

Type of decision making	Use of indicator Tier-1	Tier-2	Tier-3
Implementation of specific measures	Use to determine standards or thresholds, for example resilience of buildings or other infrastructure to extreme weather events, sea level rise etc.	Focus, design and prioritization of measures	General contextual understanding and justification, little direct use at the level of individual measures
Implementation of programmes	Design and focus of programmes	Design, focus and evaluation of programmes	Design, focus and evaluation of programmes
General policy development	General contextual understanding and justification for climate policies	General contextual understanding and sector specific justification for focus and design of policies including targets or threshold values as well as policy evaluation	Justification for focus and design of policies, possibly targets or threshold values for policies and their evaluation

3 Criteria for examining climate impact indicators

3.1 General development of criteria

Indicators have been developed in many fields and a number of criteria have been presented to determine the usefulness of indicators. These include the “SMART” (Doran 1981) criteria list where an indicator should satisfy the following criteria:

- Specific,
- Measurable (and also reliable, comparable and contextually appropriate),
- Achievable (i.e. cost effective),
- Relevant and
- Time-bounded and sensitive.

More elaborate checklists have also been introduced (McDonald 2013). These include criteria such as (i) Clarity of Focus and Meaning, i.e. the degree to which a single indicator is unambiguous and reflects or represents what is to be examined accurately or (ii) Opportunity to Detect Unexpected or Unintended Findings, i.e. the degree to which an indicator (or set of indicators) allows for documentation of unexpected or unintended consequences.

Niemeijer and de Groot (2008) proposed a conceptual framework to select environmental indicators. They reviewed previous criteria commonly used to identify environmental indicators and found the criteria used in four or more cases to be: analytically soundness (strong scientific and conceptual basis); availability of historical record (existing historical record of comparative data); time-bound (sensitive to changes within policy time frames); measurability (measurable in qualitative or quantitative terms); resource demand (achievable in terms of the available resources) and relevance (relevance for the issue and target audience at hand).

Donnelly and colleagues (2007) focused on the development of environmental indicators and other methods for the provision of information as required under the EU Strategic Environmental Assessment (SEA). The list of criteria for indicator selection was found to be rather similar to the cases described above, comprising criteria such as: policy relevance, covering a range of environmental receptors, relevance to the plan, showing trends, understandable, well founded in scientific terms, enables to prioritize key issues, adaptable and able to identify conflicts.

For earth observation data, the principle “Data and derived products shall have associated with them an indicator of quality to enable users to assess their suitability for particular applications, i.e., their ‘fitness for purpose.’ ” (QA4EO 2010) is a key requirement that leads to many of the criteria listed. Concerning some specific data sets, the Quality Assurance Framework for Earth Observation developed for the Global Earth Observation System of Systems (GEOSS)²⁰ provides a basic set of criteria that all published earth observation data should fulfill. They have also been made operational for climate related data.

Previous work on criteria for the evaluation of climate change impact indicators includes: the Environment Protection Agency (EPA) report on Climate Change indicators in the USA (EPA 2012), the EEA report on climate change, impacts and vulnerability (EEA 2012a) whose criteria are described in Hildén and Marx (2013), the Streamlining European Biodiversity Indicators (SEBI) also published by the EEA

²⁰ <http://qa4eo.org>

Table 6. Collection of criteria used to evaluate climate indicators from existing reports. Criteria were divided in two groups describing (i) the *scientific adequacy and feasibility* (marked orange) and (ii) the *relevance, usability and scope of use* (marked in blue).

EEA(2012a) and Hildén and Marx (2013)¹⁾	SEBI and EEA (2012b)	IPCC (2013)	EPA (2012)
Relevance for climate change Causal link to climate change Sensitivity towards change	Cause-effect relationship Information on cause-effect relationships should be achievable and quantifiable in order to link pressures, state and response indicators.	Data quality Data are collected using scientifically valid data methods and can support sound conclusions	Connection to climate change The relationship to climate change is easily explained.
Methodological validity (including uncertainty) Transparency Valid model base Uncertainty	Methodological well founded Clear Well defined Relatively simple Cause-effect relationships should be achievable and quantifiable	Representativeness Indicator reflects the issue it is intended to characterize.	Transparent, reproducible, and objective The data and analysis are scientifically objective and methods are transparent.
Data availability Availability and regular updating Spatial coverage/resolution Time series length/temporal resolution	Routinely collected data Routinely collected Verifiable Scientifically acceptable	Sensitivity Indicator Can distinguish meaningful differences in conditions.	Broad geographic cover The spatial scale is adequately supported with data that are representative of the region/area.
Broad acceptability Ineligibility Participatory development	Spatial coverage Indicators should ideally be pan-European and include adjacent marine areas	Decision-support Indicator provides useful information for decision-making.	Peer-reviewed data The data are credible, reliable, and have been published and peer-reviewed.
Policy relevance For EU policies Policy targets	Time series Indicators should show temporal trends		Uncertainty Information on sources of uncertainty is available.
	Country comparison As far as possible, it should be possible to make valid comparisons between countries		Trends over time Long-term data are available to show trends over time.
	Timeliness Able to detect changes in systems in timeframes and on scales that are relevant to the decisions		Feasible to construct The indicator can be constructed or reproduced/allows for routine updates.
	Acceptance and intelligibility The power of an indicator depends on its broad acceptance.		Understandable to the public Depiction of observations and are understandable to the average reader.
	Policy relevance Relevance in terms of general concern Relevance in terms of specific decisions		Usefulness Informs issues of national importance and addresses issues important to human or natural systems.
	Biodiversity relevant Address key properties of biodiversity or related issues as pressures, state, impacts and responses.		Actual observations The data consist of actual measurements (observations) or derivations thereof.
	Progress towards target Indicators should show clear progress towards the 2010 target.		

1) Criteria identified for observed and projected data

(2012b), and the Indicators of Climate Change in California (ICCC) report (Kandir *et al.* 2013), (see Table 6).

The criteria used in the different frameworks presented in Table 6 are very similar, suggesting that it is possible to identify a set of basic criteria that climate change impact indicators should fulfil. For example, the criterion described as *policy relevance* is shared by the EEA and the SEBI frameworks and also in the ICCC (Kandir *et al.* 2013) under the term *decision support*. Issues of data quality and availability are also naturally part of all the frameworks. In an attempt to guarantee the quality of the indicators used EPA (2012) restricts the use of indicators to only those that have been through a peer-reviewed process. EEA (2012b) relaxes somehow this condition while imposing that indicators used have to be available for long time-series in order to capture temporal trends. Other types of criteria are rather unique to a particular framework, driven by the objectives and scope of the reports. For example, lists of criteria proposed by the Hildén and Marx (2013) highlight the need for climate indicators to be relevant for EU policies. The same can be observed in the EEA (2012b) and ICCC (Kandir *et al.* 2013) reports. Although this criterion appears to be important when selecting impact indicators, it is not expected to be easy to evaluate, unlike for example a quality criterion or the existence of long time-series of data.

Policy relevance can be determined by the extent to which an indicator serves the design and implementation of specific policies. All climate change indicators may be relevant for climate change policies at a general level (awareness, fostering public support for measures, see Table 5, section 2.8.2), but some indicators can also be used in guiding the implementation or judging the effectiveness of specific policies. For example, a Tier-2 indicator defined as the number or proportion of buildings subject to severe flood risk can be used to guide land use policies and to evaluate the success of such policies.

One way to ensure that indicators fulfil the *Policy relevance* criterion would be to make the selection of indicators a participatory process, where relevant audiences and users of indicators are invited to participate. It is worth mentioning the bias of the investigating frameworks towards the identification of indicators focused either on broad audiences (EPA 2012) or decisions makers and general public (EEA 2012b). For some data such as earth observations there have been efforts to develop criteria that seek to ensure the validity and ‘fitness-for-purpose’ of different sensors’ products for climate change (Bates and Privette 2012). The suggested approach focuses in particular on a ‘maturity matrix’ that offers a systematic means of assessing a Climate Data Record’s (CDR’s) ‘usability’ for long term monitoring of the climate and it stresses technical data quality, but also access and software readiness.

We cluster the criteria in Table 6 in two broad groups. The first is *scientific adequacy and feasibility*, and it refers to the methodological details. The second group we call *relevance, usability and scope of use*, and deals with the informative power of the indicator, its data availability, accessibility and the easiness to inform on particular uses. The first set of criteria can be understood to primarily examine the scientific base, while the second emphasizes different aspects of use. The separation of criteria into these broad dimensions is useful as it highlights that indicators should be evaluated from several perspectives that demand different approaches. For example, while judgments on the scientific criteria of an indicator can be made by checking to what extent the underlying methodology corresponds to accepted practice, the usefulness or the policy relevance of an indicator can only be judged using some measure of the user demand.

3.2 Scientific adequacy and feasibility

Criteria under *scientific adequacy and feasibility* are used to evaluate issues such as the methodological transparency and scientific soundness of the approach (see Table 7). The most straightforward way to ensure that these demands are satisfied is to use only methods and data that have gone through peer review processes (see EPA 2012). The drawback of such a strict criterion is the reduction of potential indicators. There are also border line cases, for example not all statistical data collections have necessarily gone through peer review. The criterion *methodological transparency* is used to evaluate the method underlying a particular climate change or impact indicator. One of the key issues is the scientific documentation of the relationship, i.e. is the relation between the indicator and the impact it aims to portray scientifically established and documented. The matrix intends also to clarify the potential sources of uncertainty of a given indicator. The uncertainty issue is transversal to the datasets used to calculate an indicator as well as the method employed. Other archetypical questions to be answered

Table 7. Scientific adequacy and feasibility criteria to document climate change and impact indicators.

Criterion	Specification
Methodological Description	This criterion is used to assess the methodology to produce a climate change and impact indicator. A publication or report in which the methodology is described should be available. Optimally the required methodology should be presented in peer review publications and be publicly available. This nevertheless is not always the case, and therefore some flexibility is required.
Conceptual framework of the indicator	Climate change and impact indicators are expected to differ considerably in terms of their complexity. Some will be obtained via relatively simple mathematical operations; others will be a product of complex process modelling. It is useful to document the complexity according to five pre-established categories: 1 - Transformation of a single climate variable 2 - Metric combining several climate variables 3 - Metric aggregating climate and non-climate data 4 - Metric derived from bio-physical data other than climate 5 - Output of biophysical or economic model Other - free text The simplicity/complexity information provides background (meta) information, but cannot as such be used to judge the merit or worth of an indicator. Generally simplicity is to be preferred, but some of the relevant processes may be so complex that a too simple indicator would provide spurious information.
Scientifically documented relationship	Usually an indicator of impact or change is a proxy measure that provides information on more complex phenomena. For example, global mean temperature is used as a widely accepted metric to assess the state of the climate system. This criterion is used to judge the strength of the assumed relationship between the indicator and the impact according to 4 pre-established categories. Scientific soundness (SDRI) can be categorized into 1 - A solid, agreed theoretic framework linking indicator and a statistical correlation between indicator and impact has been established. 2 - A solid, agreed theoretic framework linking indicator and impact is agreed, but the statistic relationship is poor. 3 - A statistical correlation between indicator and impact has been established but an agreed scientific explanation is yet missing. 4 - Circumstantial evidence about the relationship is known and accepted within the scientific community. Other – free text
Treatment of uncertainty (data and method of the indicator)	Since the evolution of the climate systems is uncertain, it is relevant to assess the extent to which an indicator is able to deal with uncertainty. There is uncertainty in both the input data and the methods. This criterion is used to evaluate the ability of the indicator to deal with uncertainty. We note that “uncertainty” is used in a broad manner, including statistical data and modelling uncertainties. The evaluation of the criterion is therefore based on qualitative considerations. In general indicators that are able to recognize and deal with uncertainty are preferred to those that neglect uncertainty altogether.

during the evaluation of the scientific adequacy and feasibility are, how complex²¹ are the indicators, and can the indicator be directly applied to different geographic regions and still keep its informative power.

3.3 Usability, relevance and scope of use

The second group of criteria is related to the usability, relevance and scope of use of an indicator. The criteria in Table 8 summarize the aspects to be investigated. The evaluation of the usability, relevance and scope will depend on the intended audience. For example, indicators that are good for raising public awareness may not be the same as indicators that assist implementation of adaptation action. An evaluation based on criteria such as data availability or updating frequency, provides information on the ease of producing the indicator. The criteria also allow judgments to be made on what the indicator illustrates and how intelligible it is, i.e. to what extent it is intuitively understandable without auxiliary descriptions.

In the framework of the CLIPC project a list of preliminary criteria were presented to a wider audience during a workshop hosted at the European Environmental Agency (EEA) in May 2015 (Annex 1). The list of criteria was refined based on the recommendations from EEA and external experts. The form to document climate change and impact indicators according to these criteria can be accessed online²².

²¹ For example are indicators derived from one simple formula applied to a standard climate variable, are they products of a combination of different climate variables and social-economic data, or are they a result of an impact modelling exercise

²² <https://docs.google.com/forms/d/1SSWYuMdYPK1WdY9uOANDZPqUJjwJYrcCKPN9qqjt1KE/viewform>

Table 8. Usability, relevance and scope of use criteria to document climate change and impact indicators.

Criterion	Specification
Input data, source, availability and type ISAT	<p>The criterion is based on details of the underlying data that is used to derive a climate change or impact indicator, the data sources required, the availability of the data and the type of data.</p> <p>Of particular interest is the availability status of the data. The following options have been identified:</p> <ul style="list-style-type: none"> 1 - Public domain 2 - Within consortium 3 - Through purchasing 4 - Restrictions <p>The CLIPC will have strong preference for indicators that are based on data in the public domain. For development and testing purposes consortium owned data can be valuable.</p> <p>Another important aspect is the type of data that the indicator is based on. The following options have been identified:</p> <ul style="list-style-type: none"> 1 - Observed in-situ bio-physical data 2 - Observed in-situ socio-economic data 3 - Model projection 4 - Model reanalysis 5 - Satellite data 6 - Other. <p>The type of data mainly serves to classify the indicator, there is no particular preference for the type of data, although ideally indicators based on different types of data can be used to develop broader indicator based descriptions of topics, for example by combining observations and model projections on the same topic.</p>
Updating frequency, time frame, resolution, ensemble details and spatial extent of underlying data UTRS	<p>These criteria are used to examine the updating frequency of the underlying data required for deriving an indicator, the time frame for which the data is available, its temporal and spatial resolution and spatial extent. The spatial extent of the data-set are classified into categories</p> <ul style="list-style-type: none"> 1 - Global 2 - Europe 3 - Sub-European 4 - Other <p>In general there is a preference for indicators that build on long data series with frequent updating, good spatial resolution and wide geographical coverage. Such ideal indicators are very rare, and the criteria can be used to determine acceptable trade-offs.</p> <p>Ensembles:</p> <ul style="list-style-type: none"> 1 - Multi-model-ensembles (one scenario and multiple models) 2 - Multi-scenario-ensembles (one model and multiple scenarios) 3 - Multi-parameterization-ensembles (one model and multiple parameterizations) 4 - Multi-member-ensembles (one model and one scenario and multiple realizations) 5 - Others
Fitness for purpose of time-series length of underlying data FPID	<p>This criterion is used to judge how well the available times series fits the purpose of the indicators. For example, a time series that only captures climatic variability over a decade is insufficient for indicating climate trends. The following categories have been identified:</p> <ul style="list-style-type: none"> 1 - Data allows to discern inter-annual variability only of climate or impact 2 - Data adequate to discern decadal variability of climate or impact 3 - Data allows for statements on the long-term evolution of climate or impact 4 - Data provides only a snapshot for a particular point in time or between particular time slices 5 - Other

Criterion	Specification
Indicator availability and storage IAS	In addition to the input data used for deriving the indicator, one can judge the databases in which the climate change or impact indicator is already available with respect to availability, updating frequency, time frame, resolution and spatial extent of the indicator. Some pre-calculated indicators may be based on aggregation of original data to a particular administrative or other geographic region.
Fitness for purpose of time-series length of indicator FPI	Similar to the criterion FPID, it is relevant to examine the fitness for purpose of already calculated indicators as available in databases. The categories for the criterion follow those of the FPID criterion.
Scope of use SU	An indicator, being a climate change or impact, can serve several purposes and users. The scope of use of the indicator is thereby important to consider. This criterion thus mainly serves the classifying of indicators so that their merits can be considered in their appropriate contexts. The sub-criteria recognize four basic uses and four groups of users. Uses: 1 - Research 2 - Planning 3 - Policy development 4 - Awareness raising 5 - Others, which Users: 1 - Climate scientists 2 - Impact researchers 3 - Intermediate organizations 4 - Societal end users In general indicators that are widely applicable and that correspond to the needs of many types of users are preferable to those serving a very narrow use and limited user groups. However, there may be special operational indicators serving for example concrete flood management that are highly important despite their very focused and restricted use.
Focus on adaptive/coping capacity dimensions CF	This criterion is used to assess to what extent the methodology used to derive the indicator includes factors that are relevant for adaptive or coping capacity. Key words indicating this in the methodological description of the indicator are: coping, adaptive, adaptation, capacity, and any mention of technological or physical 'defences' that mediate or prevent the occurrence of e.g. extreme events like flooding. The criterion can be used to group indicators rather than make a judgment of their merit although there is a great demand for indicators that are able to support conclusion about the evolution of adaptive/coping capacity. Judgments on the fulfilment of the criterion are based on qualitative considerations. Inclusion of adaptive capacity: Yes No Not sure Conceptual framework of the indicator: 1 - Methodology basically follows hazard/risk approach. (hazard+vulnerability = risk) 2 - Methodology basically follows IPCC climate change approach. (exposure+sensitivity = impact) 3 - Not clear which basic approach is underlying the methodology. 4 - Methodology follows another basic approach. (name it below in "Other") 5 - The previous options are not applicable to the indicator.

4 Strengths and weaknesses of documented indicators

A key objective of this report is to evaluate strengths and weaknesses of the documented indicators. This is of course partly a subjective endeavour in the sense that a judgment of a particular strength of an impact indicator generally refers to criteria, only some of which can be measured fully objectively.

4.1 Indicator database

Indicator compilation efforts returned a total of 81 climate change and impacts indicators consistently documented according to the criteria in section 3. Figure 7 shows the number of indicators documented according the Tier classification of (Fig. 1). The bulk of the indicators documented have been documented as Tier-1, that is, those informing mostly on changes of the climatic system. Next on the list are Tier-2 indicators, although already in substantially lower numbers. Tier-3 indicators constitute the least documented types of indicators. In general, the frequency of Tier-1 to 3 indicators resembles that of comparable efforts deriving impact metrics on societal and economic systems from indicators of change in the climate system. This reflects partly the availability of data sets in a form that allows calculation of indicators (easily accessible, standardized methods, long data series and wide spatial coverage). As noted in Section 2.8 there are also numerous socio-economic data bases that meet similar criteria. The main reason that they have not lead to a comparable number of climate change impact indicators is likely to be the difficulties in attributing changes in the data to climatic variables. Potential indicators thus often fail on the criterion “Scientifically documented relationship”.

As for the distribution of indicators across investigated themes, Figure 8 shows the number of indicators allocated on a thematic basis. Most of the documented indicators were identified for the water theme, followed by the urban and rural themes. About 43 indicators have been documented as exclusive for one theme (mostly water). 19 indicators have been documented as touching two themes, the same amount of indicators that has been recorded as useful for all the three themes. This shows




Figure 7. Number of indicators documented according to Tier classification.

that cross-thematic indicators emerged from the documentation without the need for having a specific theme dedicated to that purpose. Tier-1 indicators have been documented as belonging to two or more themes but also as unique to a particular theme. In terms of indicators allocated specifically to one theme, the most numerous were found for the water theme followed by the rural theme.

Figure 9 shows the following data categories: Model projections, Model reanalysis, Observed in-situ biophysical data, Satellite data and Observed in-situ socio economic data. Model projections dominate the data types, followed by observed in-situ biophysical data model reanalysis. Satellite observations, and to a greater extent, datasets from socio-economic in-situ observations, constitute a rather small fraction of data required for the indicators. The complete list of indicator documented can be found in Table 9.

Table 9. Catalogue of potential impact and climate change indicators for CLIPC. The colour coding reflects the indicator Tier. Blue for Tier-1; Green for Tier-2; Yellow for Tier-3.

Indicators	
Arctic and Baltic Sea ice extent	100 years flood return level
Cold days	Bathing water quality
Cold nights	Chilling Units
Cold spell duration index	Chlorophyll-a-concentration
Consecutive dry days	Climatic favourability of tree species
Consecutive wet days	Distribution of marine species
Diurnal temperature range	Freshwater biodiversity and water quality
Frost days	Growing Degree Days
Greenland ice sheet mass balance	Growing season for agriculture
Heavy precipitation days	Growing season length of vegetation
Ice days	Hazardous substances in marine organisms
Lake and river ice cover duration	Heating degree-days
Lake and river ice phenology	Intensity of urban heat island with city size
Lake Ice extension	Land-cover extension below projected sea-level
Mass balance of glaciers	Moth Phenology Index
Max 1 day precipitation	Reconnaissance Drought Index
Max 5 day precipitation	River flood occurrence
Maximum of daily maximum temperature	River flow
Maximum of daily minimum temperature	River flow change
Mean precipitation	River flow droughts
Minimum of daily maximum temperature	Sea level change ¹⁾
Minimum of daily minimum temperature	The length of thermal growing season
Number of wet days	Water scarcity
Nutrients in transitional, coastal and marine waters	Water-limited crop productivity
Ocean acidification	Water-limited crop yield
Ocean heat content	Annual average damage from river floods
Permafrost thickness	Annual olive-crop yield
Precipitation extremes	Average annual heat-related deaths per 100,000 habitats
Rainfall Deciles	Coastal flood damage and adaptation costs
Sea surface temperature	Irrigation water requirement
Simple daily precipitation intensity	Natural disasters
Snow cover extension	People affected by floods
Snow Water Equivalent	Percentage change in arrivals/departures due to global warming
Spring snow cover extension	Potential impact of river flooding on major roads
Standardized Snow Pack Index	Potential impact of river flooding on railways
Storm surges	Potential impact of river flooding on settlements
Summer days	
Total precipitation	
Tropical nights	
Very heavy precipitation days	
Very wet days	
Warm days	
Warm nights	
Warm spell duration index	
Warm spell duration index	
Water temperature	

¹⁾ The ocean and their level can be seen as a fundamental part of the climate system (Tier-1), but the sea level change can be seen as a consequence of the melting of the ice sheets and glaciers (Tier-2).




Figure 8. Number of indicators documented by theme.




Figure 9. Data types underlying the calculation of indicators.

4.2 Scientific and technical evaluation of indicators

The evaluation of indicators is done using the information gathered according to the two groups of criteria detailed in Section 3. The first is about basic strength and weaknesses related to the quality and characteristics of the indicator with particular reference to criteria of scientific adequacy and feasibility (Section 3.2) but also criteria such as availability and length of time series (Section 3.3). These can be considered to be scientific and technical strengths and weaknesses and the strengths should clearly outweigh the weaknesses for an indicator to be included in a portal. The second group of questions concerns strengths and weaknesses in terms of the applicability of the indicators for specific purposes that have been identified among user groups. The criteria that inform this assessment will depend on the specific uses being considered. These strengths and weaknesses can guide a user to choose the indicators that best suit their purposes.

The key criteria for judging the fundamental strengths and weaknesses of an indicator are the strength of the scientifically documented relationship between the indicator and what it is expected to indicate, the methodological transparency, the recognition of and ability to deal with uncertainty, the (public) availability of relevant data, the updating frequency, the length of the time series and the spatial resolution and coverage of the indicator.

4.2.1 Scientifically documented relationship

Arguably one of the most important criteria regarding impact indicators is the one of “Scientific documented relationship” (see also section 3.2 defining the indicator criteria to be documented). A *strong scientifically documented relationship* means that there is scientific evidence of a process that links the indicator to climate change as opposed to, for example, a weak correlation that upon closer examination may turn out to be spurious. In this particular respect most of the potential indicators examined in this report are based on a solid theoretical and statistical relationship between impact and indicator (Figure 10). For some of indicators only circumstantial evidence about the relationship between indicator and impact is documented. There are also indicators for which a solid theoretical relationship between indicator and impact has been established, although there is yet only poor evidence of a statistical relation between indicator and impact.

Examples of indicators for which the relationship with impact is both theoretic and statistically established included: water-limited crop productivity, length of thermal growing season, moth phenology index, distribution of marine species. Indicators for which only circumstantial evidence with the impact is reported are: water scarcity, land elevation below projected sea-level or standardized snow pack index. Information on this criterion is as yet not available for all indicators. This can mean that either the chosen classes do not capture the full extent of the potential relationships between indicator and impact, or, more likely, that the understanding of impact and indicator is very different across the persons providing the information.




Figure 10. Scientific documented relationship between impact indicator and impact.

4.2.2 Methodological transparency

A *sufficient methodological transparency* means that the way the indicator is being produced is traceable and published in such a way that all relevant aspects of the method can be scrutinized and reproduced independently. For the indicators examined it has been possible to trace the methodological base (see Figure 11) as well as relevant publications describing the calculation methods. The methodological basis to derive climate change and impact indicators shows great variability. Most indicators (about 40% of the total number of indicators) are based on the use or transformation of a single climatic variable. The second most common methodological basis of indicators is that of combining several climatic variables. Combining these two methodological bases would mean that about 60% of the documented indicators would only require climate data for their reproduction, which corresponds to the dominance of the Tier-1 indicators.

Indicators with a methodological basis (Figure 11) that require the combination of climate and non-climate data constitute only about 24% of the indicators. Indicators derived from data other than climate, for example, moth phenology observations or water quality; represent about 12% of the cases. Few (2%) of the examined indicators are based on bio-physical or economic models. The collected information also shows that there is a need to document this aspect in detail in metadata for indicators as the methodological base varies considerably and efforts to aggregate and or rank indicators must take into account the methodological and epistemological base of the indicators.




Figure 11. Methodological basis of the documented indicators

4.2.3 Recognition of and ability to deal with uncertainty

An *appropriate recognition of, and ability to deal with uncertainty* means that there is a description of relevant uncertainties that may affect the interpretation of the indicator. The description of uncertainty was separated in those indicators introduced by the method for calculating the indicator, and the uncertainty that is “inherited” by the data used for indicator calculation. Regarding the first, it was observed that for 30 of the indicators documented, the information is missing. Many of these indicators are simple climate indicators for which the uncertainty is mainly stemming from the input data source. Hence, the uncertainty was documented according to criterion “uncertainty from underlying data”. For the indicators with documentation on the uncertainty of the method, the descriptions provided help to inform the potential user on key uncertainties of the indicator method. A problem is that the description need to be further standardized in terms of language and aspects covered. We provide

in Box.1 an example of a description discerning on the uncertainty of the indicator “Land elevation below projected sea-level (observations)”.

Harmonizing the heterogeneous descriptions of uncertainty across documented indicators will better inform potential users and this is clearly an item that needs to be included in the metadata on indicators.

As for the issue of data uncertainty for indicator calculation (Figure 12), it was observed that for about 25% of the indicators data is available in the form of multi-model and multi-scenario ensembles (“Both 1 and 2” bar in Figure 12). For about 18 and 5% of the indicators, only multi-model and multi-scenario ensemble data is available, respectively. There are many indicators for which the information on ensemble data for their calculation is missing.

Box 1. Uncertainty for the indicator “Land elevation below projected sea-level (observations)”

There are largely two sources of uncertainty. The first is anchored in the detail of elevation model used. There are several products available and they can vary considerably in their vertical accuracy. The most common is the global digital elevation model available (STRM90). For example the vertical accuracy is about 7 m in regions of Thailand and 4 m in coastal regions of USA (Gorokhovich and Voustianiouk 2006).

The second source of uncertainty is found on the rules used to obtain the flood extent. For example, it is arguable to assume that a land cell is flooded if its neighbour cell is also flooded. In order to account for this drawback, two types of connectivity are usually considered. A so called 4-side rule assumes that a land cell will be flooded if at least 4 of its neighbouring cells are also flooded. A so called 8-side rule is similar to the one before but assuming that a land cell will be flooded if all the neighbouring cells are also flooded.




Figure 12. Number of indicators for which ensemble input data is available.

4.2.4 Public availability of relevant data

The *public availability of relevant data* is closely linked with the methodological transparency. Indicators which use data that is publicly available have a comparative advantage to those indicators that partly or fully depend on data that can only be accessed by a limited research team or obtained at great costs. Approximately 70% of the indicators can be calculated with data that is publicly accessible, while 17% of the indicators would require the purchase or negotiation of at least one dataset for their calculation (Figure 13). As for the remaining indicators, information gathered until now does not allow for a statement on the status of data availability. Non-public data can be used in exploratory phases and indicator testing.




Figure 13. Availability of data for indicator calculation in the public domain.

4.2.5 Updating frequency of relevant data, length of time series and spatial resolution

The *updating frequency* is relevant for ensuring that indicators are maintained and provide up to date information on the phenomena they indicate. A regular (yearly) update based on standardized monitoring is to be preferred over an indicator that is occasionally updated depending on, for example, availability of (irregular) funding for dedicated research projects. Of the examined indicators many suffer from less than regular updating. This is particularly the case when the indicators are developed and presented as the output of specific research projects and not maintained by organizations responsible for monitoring of statistical data.

The length of the time series is related to questions of attribution and uncertainty. A short time series is generally insufficient for making conclusive inference on the links between climate change and the indicator. A short time series can be accepted when the indicator is considered to be exploratory and a generator of hypotheses rather than an indicator supporting decision making. Of the examined indicators several Tier-1 indicators show time series that span many decades, even more than a century and projections are also commonly presented until 2100. Historical data for Tier-2 indicators are in many cases shorter and Tier-3 indicators even shorter. This suggests a need to develop, and in some cases reconstruct, data in order to be able to document long term changes in Tier-2 and 3 indicators as this affects possibilities to examine attribution to climate change.




Figure 14. Fitness for purpose of indicator input data.

Figure 14 gives an overview of the fitness for purpose of data time series required for indicator calculation. A large majority of the data is documented as being available in a length that is adequate to make statements on the long-term evolution of climate and climate impacts. It is nevertheless interesting to note that a considerable amount of input data for indicator calculation (about 20%) only allows for a snapshot of the impact between two particular time slices.

The *spatial resolution* of an indicator is also related to questions of attribution and uncertainty. The examined indicators display a wide range of variation with respect to spatial resolution, with some available at a fine scale (Table 10). Documented indicators presented a very heterogeneous picture in regard to the spatial resolution on which the underlying data is available. Of those indicators available in a grid format, average resolution ranges from 234 to 0.004 km. Some indicators result from point measurements or are derived by making use of station data information and do therefore not have a well-defined spatial resolution. A very small set of indicators had a resolution that matched a particular administrative region.

The indicators with highest spatial resolution are derived from satellite imagery (Table 10). The remaining documented indicators are defined at spatial resolution greater than 1 km.

Table 10. Indicators with the highest documented average resolution.

Indicator	Average resolution ¹⁾
Land elevation below projected sea-level (observations)	0.004 km
Lake Ice Extent (observations)	0.08 km
Intensity of urban heat island with city size (observations)	0.5 km
Chlorophyll-a concentration (observations)	0.7 km
Sea surface temperature (observations)	1 km

¹⁾ Average resolution means in this case the average spatial resolution of the data sources required for the calculation of the indicator.

4.2.6 Indicator-by-indicator evaluation

The criteria used in the previous section also allow for discerning the strengths and weaknesses of individual or group of indicators. In the following we make this kind of informed judgment based on the available information that we have gathered on the individual indicators with respect to the criteria. Table 11 suggests that all indicators have strengths and weaknesses, but their nature varies. Thus an overall view

of the impacts of climate change will benefit from a simultaneous use of indicators of different tiers in the same way as is done in, for example, the EEA indicator report (EEA 2012a). Such collections serve general awareness-raising in particular, and help to attract attention to specific topics. More specific uses are likely to require the use of several closely related indicators. Examples of indicator-by-indicator evaluation in the light of the fundamental aspects described before are shown in Table 11.

A detailed documentation of all relevant aspects of an indicator is time consuming and requires iteration between persons that are involved in calculating and maintaining the indicators in order to achieve coherence in the interpretation and judgment of the criteria. In particular, it needs to be ensured that the metadata on the indicators is systematically collected. In addition, it is meaningful to make aggregate judgments on the strengths and weakness of indicators in order to identify areas where there is a need of further development.

Table 11. Examples of judgment on main strengths and weaknesses of indicators

Indicator/tier	Strengths	Weaknesses	Comment
Global long-term anomalies of average temperature/ Tier-1	Scientifically sound and technically sophisticated procedure for providing information on progress of climate change.	Highly aggregated information that is difficult to relate to any specific impacts of climate change.	Spatial disaggregated information on temperatures are available for regions/countries.
Consecutive dry days/ Tier-1	Transparent calculation based on available information, detailed spatial resolution for observations gives overview of situation.	The effects of the length of a drought depend, among other things, on the region and season of its occurrence. Therefore it is difficult to compare across regions.	Relative changes within each region provide an indication of the direction of development.
Growing season for agriculture/ Tier-2	Well defined basic concept that has clear general relationship to agricultural production. Transparent calculation based on easily accessible data at different spatial resolutions.	Provides overview but does not alone indicate the actual development of agriculture as confounding factors, notably precipitation, influences it.	Can be combined with other indicators to identify the conditions for the development of agriculture.
Economic losses related to natural disasters/ Tier-3	Based on insurance payments, reflects actual costs incurred and therefore easy to link to costs of conceivable measures to reduce vulnerability.	Underlying data not fully publicly available, more detailed analyses of indicator and factors affecting it are therefore seriously hampered. Issues of attribution arise as economic conditions affect development of indicator.	More detailed information is available for individual countries/regions, but have so far not been standardized and made available for the whole of Europe.
Annual heat-related deaths/ Tier-3	Statistical relationship identified for many cities based on physiological considerations. Data increasingly available at different geographical scales.	Available data series are generally relatively short. Significant uncertainties exist and relationship appears to vary between regions/localities. Projections therefore highly uncertain.	Projections can be used for general indication of vulnerability rather than indications of actual deaths.

4.3 User needs

The scientific and technical aspects of an indicator can be complemented with considerations of the needs of specific users. A quick reading of Scope of Use criteria (see criterion in Table 8) returns that mostly climate impact indicators can be used for research and for raising societal awareness, see Figure 15. The remaining possible categories of uses rank substantially lower and account approximately for the same number of indicators. One should mention here that double counting of indicators for different uses was allowed.




Figure 15. Number of indicators documented by main use.

We further contrast the proposed uses of the indicators in Figure 15 with those extracted in an online survey questioning *climate scientists, impact scientists, intermediary organizations*²³ and *societal end users*. The survey was conducted in the CLIPC project and provides the first indications of particular user requirements for climate change and impact indicators across user groups (de Groot *et al.* 2014, de Groot *et al.* 2015). Despite several user groups being involved in answering the questionnaire, three main purposes for climate data and impact indicators were identified to be common for the user groups. These are:

1. To give advice on climate data and climate impact indicators to others.
2. To support the development of adaptation strategies and plans.
3. To perform risk and vulnerability assessments.

These purposes are general and do not suggest a direct preference for any particular kind of indicators, although purposes 2 and 3 above underline the importance of Tier-2 and Tier-3 indicators (see Figure 7). Some purposes appear to be more specific for particular user groups. For example intermediary organizations see raising awareness as a specific use of climate data and impact indicators, whereas impact and climate researchers wish to use climate data and indicators as input for research on climate change. Table 12 shows the matrix of the top three purposes of data and impact indicators according to user group. Some of the uses put quite distinct requirements on the available data and indicators. Indicators that are used to raise awareness should

²³ Intermediary organisations or boundary organisations assist stakeholders in decision making (see also de Groot A., Swart R., Hygen H.O., Bensted R., Cauchy A., Betgen C. & Dubois G. 2014. User requirements, part 1, Strategies for user consultation and engagement and user requirements: Synthesis from past efforts, CLIPC Deliverable (D2.1).)

preferably be easy to understand with general knowledge. Indicators that are used for (impact) research purposes often need to be available in a spatially disaggregated form to allow linking with other variables that include spatial information.

In addition, expert judgment of the CLIPC consortium partners places indicators documented as being mostly of potential use for “Impact researchers”. Almost every indicator gathered was perceived as potentially useful for impact research. “Societal end user” is the user category for which the smallest fraction of the collected indicators was considered useful, although about 30% of indicators are documented as potentially useful.

Table 12. Top purposes for data and indicators according to potential CLIPC users identified in a user consultation questionnaire.

User groups	Top three purposes for climate data and indicators		
	1st	2nd	3rd
Societal end users	Support the development of adaptation strategies and plans	Create awareness	Make risk and vulnerability assessments
Intermediary organizations	Give advice on data and climate impact indicators to others	Support the development of adaptation strategies and plans	Create awareness
Impact researchers	Make risk and vulnerability assessments	Input in research on climate change	Support the development of adaptation strategies and plans ¹⁾
Climate scientists	Give advice on data and climate impact indicators to others	Input in research on climate change	Mix of awareness raising, adaptation and risk assessment ²⁾

¹⁾ The second and third purposes in the user category “Impact Researchers” presented the same number of answers, meaning that the placement of the 2nd or 3rd purposes is in this case arbitrary.

²⁾ The same number of answers for purposes “Create awareness”, “Make risk and vulnerability assessments” and “Support the development of adaptation strategies and plans”.




Figure 16. Number of indicators documented by potential main user category identified in CLIPC.

Attending both to the results of the indicator documentation and the main uses of indicators, the current documentation of indicators already makes good progress in meeting the demand of indicators to be used for research purposes and awareness raising (shaded boxes in Table 12). Regarding other envisioned uses of indicators by potential users, these can at some extent be also addressed by the indicator sample in the current database. Nevertheless, due to the non-existence, at this point, of better details of what the use of an indicator for the development of adaptation plans entails, it is hard to make more concrete judgments of the potential miss-match between the indicators gathered and the envisioned uses of indicators by potential users.

A potential way-out from the lack of detailed information from users is to think about what criteria from Table 7 and Table 8 are relevant for an indicator to be used for the top-purposes in Table 12. For each use in Table 12 we have identified criteria that can be useful in judging indicators from a user perspective. Examples are provided in Table 13, together with a brief explanation on how the criteria relate to the rationale of the indicator being used for the top-purposes identified. Furthermore, a column in Table 13 is added with suggestions on how documented criteria could help discerning strengths and weaknesses of indicators in the context of a particular use.

For the use of indicators in *supporting the development of adaptation strategies and plans*, criteria such as the ability to display *adaptation/coping capacity* can be expected to be significant. Indicators that can be readily used in assessments, or easily adapted/ altered to suit the specificities of the assessment, have particular strengths. Figure 17 shows indicator shares in regard to the inclusion of the adaptive/coping capacity dimension. The potential impact indicators examined appear to be weak in accounting explicitly for adaptive capacity. Only about 15% of the documented indicators have been coded as incorporating adaptive capacity. Half of the indicators do not account directly for adaptive capacity. This is understandable since many of the indicators belong to Tier-1 indicators, which mostly (although not without exceptions) do not usually incorporate a measure of the capacity of society or ecosystems to sustain change.




Figure 17. Fraction of indicators that account for adaptive capacity.

The results indicate that there is a demand for indicators that provide ways to measure adaptive capacity. The availability of such indicators would strengthen adaptation planning.

In *elaborating risk and vulnerability assessments* the availability of long time series is considered to be a particular strength as it allows consideration of, for example, return times based on empirical evidence.

The use of data and indicators as *input in research to climate change* generally requires possibilities to manipulate the data or indicators further, at the very least in the form of customized graphical display. Thus a key strength is the availability of the data and indicator values in the public domain. Indicators that are not openly available can be referred to but not used as input. Research focusing on future impacts of climate change is dependent on projections. The availability of the data and the indicator is thus vital also here. In addition data and indicators that are available based on ensembles of models and scenarios are to be preferred over indicators that are based on single models and single scenarios. The ensemble data allow an exploration of variability and uncertainties that remain largely hidden in single data series.

The purpose of using indicators for *raising awareness* is in one sense the most general one, but at the same time very demanding, as these indicators have to resonate with

Table 13. Examples of criteria used to evaluate strengths and weaknesses of indicators with respect to a specific use.

Top-purpose of the indicator	Selected criteria from Table 7 and Table 8	Rationale	Strength (1) & Weakness (0) (evaluation pairs)
Supporting the development of adaptation strategies and plans	Presence/absence of adaptive/coping capacity	An ability of the indicator to include a dimension of adaptive/coping capacity makes it useful for this top-purpose is a comparative strength	(1) If indicator documented with "Yes" in criterion CF (Table 8) or (0) If indicator documented with "No" in criterion CF
Production of risk and vulnerability assessments	Length of time series	A long time-series of the indicator makes it useful for this top-purpose and gives information on its strength.	(1) If indicator documented with "3" in criterion FPI (Table 8) or (0) If indicator documented with "1, 2 or 4" in criterion FPI
Input in research on climate change	Data availability Ensembles details	The accessibility and availability of input data and indicators for further analysis and manipulation determines a basic usefulness and strength for use in research. For projected indicators the availability based on ensemble modelling and/or multiple scenarios is a strength.	(1) If indicator documented as public domain in criteria ISAT (Table 8) or (0) If indicator documented with other than public domain -- (1) If indicator documented with both with "1" and "2" in ensemble details in criteria UTRS (Table 8) or (0) If indicator documented other than "1" and "2"
Creating awareness	Main use of the indicator Indicator allocated to two or more themes	The degree to which the documented indicators have been used to raise awareness suggests makes it useful for this top-purpose and highlights a strength. The ability of an indicator to be relevant across several themes is assumed to reflect a strength.	(1) If indicator documented with "4" in criterion on uses in SU (Table 8) or (0) If indicator documented with "1,2 or 3" -- (1) If indicator documented in more than one theme or (0) If indicator documented as being exclusive to 1 theme

a diverse audience that does not necessarily have the expertise to judge the validity of indicator based claims or to understand the underlying processes. The first criterion is somewhat circular, but assumes that an evolutionary process has operated so that indicators whose *main use has been to raise awareness* have particular strengths in this regard. Since awareness raising should focus on key messages and not cause information overload, parsimony is a particular strength. Thus indicators that are relevant across multiple themes are likely to have strengths relative to more focused indicators that are relevant for a narrow theme.

Following the rationale exposed, it is now possible to proceed with a tentative match of a particular indicator and a very tentative evaluation of the relative strengths and weaknesses in regard to a particular use, and in parallel to what user category might be more interested in the indicator. Table 14 shows the summary of applying criteria from Table 13 to the indicator "Growing Season for Agriculture".

The identified criteria in Table 13 do not provide an exhaustive view of the strengths and weaknesses of indicators from a user perspective. A user perspective on the strength and weaknesses ultimately needs to be based on a holistic view of the indicator, weighting the different criteria in a suitable, partly subjective, way.

Table 14. Preliminary evaluation of the strengths and weaknesses based on the potential use for the “Growing season for agriculture” indicator.

Use of the indicator	Strength (1 or 2), Weakness (0), insufficient data (NA)
Supporting the development of adaptation strategies and plans	1
Production of risk and vulnerability assessments	1
Input in research on climate change	1/NA
Creating awareness	1/0

Nevertheless, we do find the exercise useful as a starting point to identify how to better match the user-groups perspectives on the *use* of climate impact indicators, and their adequacy for a specific use in the light of particular strengths and weaknesses. Accordingly, Table 14 highlights that the indicator “Growing Season for Agriculture” is adequate for the uses of “Supporting the development of adaptation strategies and plans” and the “Production of risk and vulnerability assessments”. Its use as input for research score 1/NA, meaning that while the input data sources are all in public domain, there is insufficient data in the documentation regarding the existence of ensemble-projections of the underlying databases for the calculation of the indicator. In regard to the use of “Creating awareness”, the indicator has been perceived as specific for one of the investigated themes (rural).

The complete table of examined indicators can be found in Annex 2. It is important to mention that the current classification with respect to use specific strengths is only indicative i) because the statements on potential uses of indicator by users are only general and ii) the existence of some lack of harmonization in the documentation, the mapping between evaluation criteria for a particular use cannot be used to make very specific statements on the strengths and weaknesses of individual indicators. Additional criteria and a stronger emphasis on context related uses such as the development of adaptation measures for specific locations would probably strengthen the differences.

Most of the indicators have been used in raising awareness of climate change. Noticeably, no substantial differences across Tiers are observed in this regard. We do, however, lack information on which indicators have been the most compelling ones although several international organisations and scientific institutions have tried to identify the characteristics of ‘good indicators’ (Hammond *et al.* 1995, Hart Environmental Data 1998, Meadows 1998, OECD 2003).²⁴ For example, the Lowell Center for Sustainable Development underlines relevance, understandability, and usability at the level of communities (Hart Environmental Data 1998). Such characteristics are particularly important for local awareness-raising. As such, many of the Tier-3 indicators are relevant and understandable and would also be useable, but are often hampered by the lack of community specific data, which may explain why they have not yet been used extensively in awareness raising, compared with, for example, many Tier-1 indicators.

The sample of indicators gathered at the time of writing have been observed to match the user needs for using indicators as input for climate research and for the purposes of awareness raising. The uses of supporting the elaboration of adaptation strategies and vulnerability studies can already be supported by the indicators gathered, although in these cases it is still preliminary to make definitive judgments in the light of drawbacks previously highlighted. Further interaction with potential users of indicators and subsequent updates of indicator documentation would help to clarify this.

²⁴ <http://www.rscproject.org/indicators/index.php?page=what-are-the-characteristics-of-a-good-indicators-or-indicator-sets> [October 13 2016]

5 Conclusions

The main conclusions and perspectives for further works can be summarized as follows:

The analysis of strengths and weaknesses of indicators has used a grouping into three tiers. In this categorization, Tier-1 indicators provide information on the past and future evolution of the climate system. Tier-2 indicators quantify the impacts of climate change in bio-physical systems. Tier-3 indicators primarily provide information on the socio-economic systems affected by climate change. Tier-3 indicators usually build on Tier-1 and Tier-2, and make the bridge from bio-physical change to social or economic loss/gain. This grouping served to highlight the relative scarcity of indicators that explicitly link climate change to socio-economic consequences.

The criteria that were used to identify strengths and weaknesses build on commonly used indicator criteria. Climate change indicators are special in that an important part of the indicators are also used to project future development. This implies a heavy reliance on modelling and puts special demands on, for example, the treatment of uncertainty.

The large collection of information on criteria for climate change and impact indicators across themes and tiers has identified general strengths and weaknesses of indicators. The key criteria for judging the fundamental strengths and weaknesses of indicator are the strength of the scientifically documented relationship between the indicator and what it is expected to indicate, the methodological transparency, the recognition of and ability to deal with uncertainty, the (public) availability of relevant data, the updating frequency, the length of the time series and the spatial resolution and coverage of the indicator.

A general strength is that there is an abundance of Tier-1 indicators that are based on publicly available data, well-established scientific theories, long time series, high spatial resolution and the availability of projections based on the results of ensembles of models calculated for several different scenarios. Good Tier-2 indicators are less well available, but several exist, and they are also based on high quality open source data that are regularly updated. It is a priority to develop new Tier-2 indicators that are based on regular monitoring and that can explore novel data sources such as those based on earth observations.

The poor availability of Tier-3 indicators is an obvious general weakness. One of the main reasons for the lack of Tier-3 indicators is that it is difficult to quantitatively attribute economic and wider societal development to climatic factors. This is natural as climate is only one of numerous factors affecting societal development. Therefore it appears that Tier-3 indicators are currently often more useful when they are framed in terms of vulnerability or adaptive capacity rather than in terms of actual impacts, which require detailed site specific analyses in order to deal with the question of attribution.

There is an obvious need to continue with exploratory work that can establish links between societal conditions and climatic factors. This can be achieved by examining socio-economic statistical information as dependent variables in the light of information from Tier-1 and Tier-2 climate indicators. Such studies demand sufficient spatial resolution in order to display the variation that is needed to explore possible relationships. Eventually the studies may lead to new Tier-3 indicators, but such studies will in any case improve the base for modelling and projections of impacts.

In the scanning work conducted for this report it was not possible to fully harmonise the use of the evaluation criteria across indicators. For specific indicators there was also missing or incomplete information on some criteria.

The sample of indicators gathered have been observed to match the user needs for using indicators as input for climate research and for the purposes of awareness raising. The uses of supporting the elaboration of adaptation strategies and vulnerability studies can already be supported by the indicators gathered, although in these cases it is still preliminary to make definitive judgments.

REFERENCES

- Barredo J.I. 2010. No upward trend in normalised windstorm losses in Europe: 1970–2008. *Nat. Hazards Earth Syst. Sci.* 10: 97–104.
- Bates J.J. & Privette J.L. 2012. A maturity model for assessing the completeness of climate data records. *Eos, Transactions American Geophysical Union* 93: 441–441.
- Bojinski S., Verstraete M., Peterson T.C., Richter C., Simmons A. & Zemp M. 2014. The Concept of Essential Climate Variables in Support of Climate Research, Applications, and Policy. *Bulletin of the American Meteorological Society* 95: 1431–1443.
- Carter T.R., Fronzek S., Inkinen A., Lahtinen I., Lahtinen M., Mela H., O'Brien K.L., Rosentrater L.D., Ruuhela R., Simonsson L. & Terama E. 2016. Characterising vulnerability of the elderly to climate change in the Nordic region. *Regional Environmental Change* 16: 43–58.
- Ciscar J.C., Feyen L., Soria A., Lavalle C., Raes F., Perry M., Nemry F., Demirel H., Rozsai M., Dosio A., Donatelli M., Srivastava A., Fumagalli D., Niemeyer S., Shrestha S., Ciaian P., Himics M., Van Doorslaer B., Barrios S., Ibáñez N., Forzieri G., Rojas R., Bianchi A., Dowling P., Camia A., Libertà G., San Miguel J., de Rigo D., Caudullo G., Barredo J.I., Paci D., Pycroft J., Saveyn B., Van Regemorter D., Revesz T., Vandyck T., Vrontisi Z., Baranzelli C., Vandecasteele I., Batista e Silva F. & Ibarreta D. 2014. *Climate Impacts in Europe. The JRC PESETA II Project*. JRC Scientific and Policy Reports, available at <http://ftp.jrc.es/EURdoc/JRC87011.pdf>.
- Costa L. & Kropp J.P. 2013. Linking components of vulnerability in theoretic frameworks and case studies. *Sustainability Science* 8: 1–9.
- Costa L. & Hildén M. 2015. *A review of climate impact indicators across themes: Description of strengths and weaknesses, technical requirements and mismatch from user expectations*. CLIPC deliverable, D.7.1, available at <http://www.clipc.eu/>.
- de Groot A., Swart R., Hygen H.O., Bensted R., Cauchy A., Betgen C. & Dubois G. 2014. User requirements, part 1, Strategies for user consultation and engagement and user requirements: Synthesis from past efforts, CLIPC Deliverable (D2.1), available at <http://www.clipc.eu/>.
- de Groot A., Betgen C., Dubois G., Roth E., Dhenain S., Swart R. & Mañez M. 2015. *User requirements, part 2, Climate (impact) data requirements of different user groups*, CLIPC Deliverable (D2.2). available at <http://www.clipc.eu/>.
- Donnelly A., Jones M., O'Mahony T. & Byrne G. 2007. Selecting environmental indicator for use in strategic environmental assessment. *Environmental Impact Assessment Review* 27: 161–175.
- Doran G.T. 1981. There's a S.M.A.R.T. way to write management's goals and objectives. *Management Review* 70: 35–36.
- EEA. 2012a. *Climate change, impacts and vulnerability in Europe 2012. An indicator-based report*. EEA Report, Copenhagen, Denmark, available at <http://www.eea.europa.eu/publications/climate-impacts-and-vulnerability-2012>.
- EEA. 2012b. *Streamlining European biodiversity indicators 2020: Building a future on lessons learnt from the SEBI 2010 process*. EEA Technical report, EEA, Copenhagen, Denmark.
- EPA. 2012. *Climate change indicators in the United States, Technical Documentation*. 2nd ed., available at <https://www3.epa.gov/climatechange/pdfs/CI-technical-documentation-2012.pdf>.
- ETC-CCA & ETC-SIA. 2012. *Urban Vulnerability Indicators. A joint report of ETC-CCA and ETC-SIA*. Technical Report, available at http://cca.eionet.europa.eu/docs/TP_3-2012.
- ETC/ACC. 2010. *Urban Regions: Vulnerabilities, Vulnerability Assessments by Indicators and Adaptation Options for Climate Change Impacts (Scoping Study)*. Technical Paper, available at http://acm.eionet.europa.eu/reports/docs/ETCACC_TP_2010_12_Urban_CC_Vuln_Adapt.pdf.
- Flörke M., Wimmer F., Laaser C., Vidaurre R., Tröltzsch J., Dworak T., Stein U., Marinova N., Jaspers F., Ludwig F., Swart R., Giupponi C., Bosello F. & Mysiak J. 2011. *Climate Adaptation—modelling water scenarios and sectoral impacts, Final Report*. Center for Environmental Systems Research, available at http://ec.europa.eu/environment/archives/water/adaptation/pdf/ClimWatAdapt_final_report.pdf.
- Füssel H.-M. & Klein R.J.T. 2006. Climate Change Vulnerability Assessments: An Evolution of Conceptual Thinking. *Climatic Change* 75: 301–329.
- Gorokhovich Y. & Voustianiouk A. 2006. Accuracy assessment of the processed SRTM-based elevation data by CGIAR using field data from USA and Thailand and its relation to the terrain characteristics. *Remote Sensing of Environment* 104: 409–415.
- Greiving S., Flex F., Lindner C., Lückenköter J., Schmidt-Thomé P., Klein J., Tarvainen T., Jarva J., Backman B., Luoma S., Langeland O., Langset B., Medby P., Davoudi S., Tranos E., Holsten A., Kropp J., Walter C., Lissner T., Roithmeier O.M.K., Juhola S.P.N., Peltonen L., Vehmas J., Sauri D., Serra A., Olcina J., March H., Martín-Vide J., Vera F., Padilla E., Serra-Llobet A., Csete M., Pálvölgyi T., Göncz A., Király D., Schneller K., Staub F., Peleanu I., Petrisor A.-I., Dzurdenik J., Tesliar J., Visy E., Bouwman A., Knoop J., Ligtoet W., van van Minnen J., Kruse S., Pütz M., Stiffler M. & Baumgartner D. 2011. *ESPO Climate: Climate Change and Territorial Effects on Regions and Local Economies*. Scientific Report, ESPON & IRPUD, TU Dortmund University, Germany.
- Hammond A., Adriaanse A., Rodenburg E., Bryant D. & Woodward R. 1995. *Environmental Indicators: A systematic approach to measuring and reporting on environmental policy performance in the context of sustainable development*. World Resources Institute, Washington, USA, available at http://pdf.wri.org/environmentalindicators_bw.pdf.

- Hart Environmental Data. 1998. *Sustainable community indicators trainer's workshop*, available at <http://www.sustainablemeasures.com/Training/pdf/HEDTrMan.pdf>.
- Hildén M. & Marx A. 2013. *Evaluation of climate change state, impact and vulnerability indicators*. ETC CCA Technical Paper.
- Hinkel J. 2011. "Indicators of vulnerability and adaptive capacity": Towards a clarification of the science-policy interface. *Global Environmental Change* 21: 198-208.
- IPCC. 2000. *IPCC special report emission scenarios - summary for policy makers*. A special report of working group II of the Intergovernmental panel on climate change, Geneva, Switzerland, available at <http://www.ipcc.ch/ipccreports/sres/emission/index.php?idp=0>.
- Kandir T., Mazur L., Milanec C. & Randles K. 2013. *Indicators of climate changes in Californien*. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment, available at <http://cnap.ucsd.edu/pdffiles/ClimateChangeIndicatorsReport2013.pdf>.
- McDonald G. 2013. *Criteria for Selection of High-Performing Indicators A Checklist to Inform Monitoring and Evaluation*.
- Meadows D. 1998. *Indicators and information systems for sustainable development*. A report to the Balaton group, The Sustainability Institute, Hartland Four Corners, USA, available at http://www.iisd.org/pdf/s_ind_2.pdf.
- Niemeijer D. & de Groot R.S. 2008. A conceptual framework for selecting environmental indicator sets. *Ecological Indicators* 8: 14-25.
- OECD. 2003. *OECD Environmental Indicators - Development, measurement and use*. Reference paper, Organisation for Economic Co-operation and Development, available at <http://www.oecd.org/environment/indicators-modelling-outlooks/24993546.pdf>.
- Palmer W.C. 1965. *Meteorological Drought*. U.S. Department of Commerce Weather Bureau, available at <http://www.ncdc.noaa.gov/temp-and-precip/drought/docs/palmer.pdf>.
- Prahl B.F., Rybski D., Burghoff O. & Kropp J.P. 2015. Comparison of storm damage functions and their performance. *Nat. Hazards Earth Syst. Sci.* 15: 769-788.
- Preston B.L., Smith T.F., Brooke C., Gorddard R., Measham T.G., Withycombe G., McInnes K., Abbs D., Beveridge B. & Morrison C. 2008. *Mapping Climate Change Vulnerability in the Sydney Coastal Councils Group*. Prepared for the Sydney Coastal Councils Group. available at <http://www.sydneycoastalcouncils.com.au/sites/default/files/systapproachphaseonereport.pdf>.
- QA4EO. 2010. *A quality assurance framework for earth observation: Principles, Version 4.0*. available at http://qa4eo.org/docs/QA4EO_Principles_v4.0.pdf.
- Schellnhuber H.J., Frieler K. & Kabat P. 2014. The elephant, the blind, and the intersectoral intercomparison of climate impacts. *Proceedings of the National Academy of Sciences* 111: 3225-3227.
- Smith A.B. & Katz R.W. 2013. US billion-dollar weather and climate disasters: data sources, trends, accuracy and biases. *Natural Hazards* 67: 387-410.
- Van der Linden P. & Mitchell J.F.B. 2009. *ENSEMBLES: Climate Change and its Impacts: Summary of research and results from the ENSEMBLES project*. Met Office Hadley Centre, Exeter, UK.
- Vautard R., Gobiet A., Sobolowski S., Kjellström E., Stegehuis A., Watkiss P., Mendlik T., Landgren O., Nikulin G., Teichmann C. & Jacob D. 2014. The European climate under a 2°C global warming. *Environmental Research Letters* 9: 034006.
- Visser H., Bouwman A., Peterson A. & Ligtvoet W. 2012. *A statistical study of weather-related disasters: Past, present and future*. PBL Netherlands Environmental Assessment Agency, The Hague, available at http://www.pbl.nl/sites/default/files/cms/publicaties/PBL_2012_Weather%20Disasters_555076001.pdf.
- Watkiss P. 2011. *The ClimateCost Project. Final Report1: Europe*. Stockholm Environment Institute.

ANNEX I

Expert workshop at the EEA

**Joint CLIPC - EEA meeting 13th to 14th May 2015 Copenhagen
Venue: EEA facilities at Copenhagen (Address:
Kongens Nytorv 6, 1050 Copenhagen K)**

Contents:

1. Background
2. The Objectives of the workshop
3. The EEA's expectations on the CLIPC indicator toolbox
4. Overview of the CLIPC project
5. The TGICA and the DDC: How to guarantee a consistent set of up-to-date scenarios for use in climate impacts assessments: Relevance for indicators of climate change?
6. JRC's strategy regarding climate change impact data and services: Prospects and developments
7. General indicator requirements and the experience of using criteria to evaluate indicators by the EEA with special attention to climate and climate impact indicators
8. A first set of criteria for CLIPC indicators: Example of how the CLIPC criteria could work in practice
9. General discussion on criteria
10. Priority users/user groups for CLIPC and what preliminary demands they may put on data and impact indicators to be provided
11. Brainstorming envisioned key features the CLIPC toolbox
12. General requirements
13. Technical requirements
14. Specific characteristics serving particular user groups
15. Climate scientists
16. Impact scientists
17. Intermediaries
18. Processes for user engagement
19. Next steps
20. List of participants

Background

The CLIPC project will provide access to climate information of direct relevance to a wide variety of users, from scientists to policy makers and private sector decision makers. Information will include data from satellite and in-situ observations, climate models and re-analyses, transformed data products and climate change impact indicators.

This particular workshop focused on criteria to be used for evaluating and screening climate and climate impact indicators to be included in the CLIPC toolbox. An agreement on criteria is a required outcome of report D7.1: *A review of climate impact indicators across specific themes and description of strengths, weaknesses and technical requirements*. In addition, the workshop discussed the envisioned functionalities of the CLIPC toolbox using as a starting point the identification of key users/user groups

undertaken in D2.1: *Synthesis of user requirements from past efforts and user involvement strategy on providing climate (impact) data*).

In developing requirements for CLIPC indicators the workshop reflected on ongoing and planned activities by the European Environmental Agency (EEA), the Joint Research Center (JRC), Copernicus Climate Services and the IPCC Task Group on data and scenario support for Impact and Climate Analysis (TGICA) and the IPCC Data Distribution Center (DDC) in order to make sure that both functionalities and data/indicator requirements can be harmonized with those developed elsewhere. The work will also reflect on related and relevant EU-projects projects such as CLIMSAVE, IMPRESSIONS and other pre-operational Copernicus projects addressing projections of climate change (and impacts). Experiences of indicator development and presentation will be fully used to avoid duplication of work.

This report is organized in the order in which the topics were dealt with at the workshop. The first half was devoted to the general expectations and ways of developing and using criteria for indicators of climate change and impacts of climate change. The second half of the workshop was devoted to a discussion on functionalities of the toolbox.

The Objectives of the workshop

The objectives of the workshop were to:

1. Discuss and agree on criteria to screen, evaluate and assess the strengths and weaknesses of climate and climate impact indicators and underlying data to be included in the CLIPC toolbox.
2. Discuss the functionalities of the CLIPC toolbox and how it will bring added value relative to other indicators and indicator tools.

To achieve its objectives the workshop was initiated with presentation from different perspectives on what CLIPC could achieve with respect to indicators of climate change.

The EEA's expectations on the CLIPC indicator toolbox
André Jol, EEA

The EEA's expectations were presented by André Jol who noted that the new multiannual work programme 2014-2018 guides EEA's activities and focal areas. Among these societal transitions have received particular attention and will be central in the work. In the State of the Environment Report of 2015 climate change will be one thematic issue. The portal Climate-ADAPT includes indicator information mainly from EEA (report on CC impacts, published in 2012) and maps (map viewer) and by 2016 climate indicators are expected to be updated closely linked with a new report due in 2016, building on the content and experiences with the 2012 report. This provides opportunities to link directly with the work in CLIPC. The Tier-1 indicators are fairly well placed and operational, but there is great interest in achieving progress in Tier-2 and Tier-3 indicators.

Climate-ADAPT is central for the EEA and it will link to work of JRC on CC impacts and adaptation and in the future also to the Copernicus climate services. On June 23 a expert meeting will be held on climate adaptation portals as part of EEA's work with member countries.

He summarised the main expectations from EEA towards the CLIPC project :

- Contribute to the EEA climate change impact indicators (on EEA web site) and 2016 indicator-based assessment report
- Contribute to Climate-ADAPT (e.g. map viewer)
- Help define the future linkages between the Copernicus climate change service and Climate-ADAPT
- Make effective use of linkages between CLIPC consortium and ETC CCA lead and partner organisations
- Participation by EEA in CLIPC advisory group and expert group meetings

Overview of the CLIPC project

Martin Juckes, CLIPC /STFC

Martin Juckes provided an overview of the CLIPC . He noted that CLIPC can be seen as a prototype for part of the future activities of the Copernicus Climate Change Services. The CLIPC will follow the Earth System Grid Federation (ESGF; <http://esgf.org/>) that develops, deploys and maintains software infrastructure for the management, dissemination, and analysis of model output and observational data. Another important connection is the IS-ENES climate4impact portal (<http://climate4impact.eu/impactportal/general/index.jsp>), which is oriented towards climate change impact modellers, impact and adaptation consultants, and other experts using climate change data. Specific goals are to provide

- harmonized access to data from many sources
- information on data value and limitations
- indices of climate change & impacts
- a knowledge base of authoritative information
- a toolkit to update indices and indicators

CLIPC makes assessment and data available for assessments but will not make its own assessment. A brief discussion noted the need to be clear about distinctions between indicators and indices.

The TGICA and the DDC: How to guarantee a consistent set of up-to-date scenarios for use in climate impacts assessments: Relevance for indicators of climate change?

Tim Carter, SYKE

Tim Carter described The Task Group on Scenarios for Climate and Impact Assessment (TGICIA) and the Data Distribution Centre (DDC) noting that TGICIA covers all WGs and that the information needs are catered for through the DDC, for which rigorous quality control has been set up. An important task is to provide technical guidelines, interpretation of data, with all guidance rigorously peer reviewed and transparent criteria for linking data sets. TGICIA has Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP) on its agenda, but DDC not yet covered impact model information.

Tim Carter noted that the interest in data is expanding as the expert community is widening; also utilities and other users are increasingly requesting information, but users may not find the data sufficiently detailed. The link to Copernicus Climate services will thus be important for a wider user community.

The possibilities to share insights on user needs and user surveys were discussed, along with possibilities for organizing a meeting partly joint TGICA – CLIPC meeting for southern Europe.

JRC's strategy regarding climate change impact data and services: Prospects and developments
Nadine Gobron, CLIPC /JRC

JRC's current work on climate change impact data were presented by Nadine Gobron who noted that JRC's key areas include:

- Development of the knowledge base Climate-ADAPT
- Estimating costs of future climate change
- Developing coherent integrated assessments

JRC is, in particular, the key map provider for Climate-ADAPT. New tools are also being developed including time series based on earth observation data. JRC has carried out the Peseta II project (<http://ftp.jrc.es/EURdoc/JRC87011.pdf>), which has led to additional work on climate change impacts to be carried out in 2014-2015 on topics such as:

- costs of droughts
- impacts in coastal areas
- impact on ecosystems & services
- global impacts with implications for Europe

On earth observations development work continues on:

- Mitigation and adaptation
- Quality control

For quality control the project: QA4ECV – quality control for ECV (<http://www.qa4ecv.eu/>) attempts to bridge the gap between end-users of satellite data and the satellite data products by developing an internationally acceptable Quality Assurance (QA) framework that provides understandable and traceable quality information for satellite data used in currently evolving climate and air quality services.

General indicator requirements and the experience of using criteria to evaluate indicators by the EEA with special attention to climate and climate impact indicators
Hans-Martin Füssel, EEA

Hans-Martin Füssel presented the EEA approach to indicators and the requirements that emerge from the chosen approach. In the context of climate change key interests are to:

- present climate change- providing the general context
- present climate related hazards
- assess the impacts of climate change on society, human health and ecosystems
- assess the effectiveness of climate risk management (with a focus on adaptation)

This leads to demands on spatial coverage and resolution. The spatial coverage should be as wide as possible (taking into account the EEA member countries), and the resolution sufficient to identify relevant changes. In addition indicators should be relevant for EU policies. Indicators should thus meet the following criteria.

- Thematic and policy relevance
- Full geographic coverage of relevant variables
- Appropriate geographical aggregation
- Long time series
- Reliable data series
- Clear methodology

As far as possible indicators should provide observations of historical development, projections for future development, and information on uncertainties.

In 2016-2017 the EU is likely to revise its adaptation strategy leading to specific demands in the accompanying 'impact assessment' of the strategy according to EU procedures for all new proposed policies. The planned 2016 EEA report on CC impacts will be one input to this 'impact assessment'.

It will be beneficial for CLIPC to further explore work that has been going on in projects such as Impact2c, PesetaII/III and to reflect on the question of attribution to climate change in considering indicators. He also stressed the importance of narratives that are an integral part of the EEA indicators.

For the development of indicators the EEA sees a need to link with many expert communities and to consider users involved in country level risk assessments. For future work it will be relevant to consider possibilities to expand the number of indicators and develop links to future Copernicus climate services. There is also ongoing development between JRC and EEA to ensure consistent approaches in their assessments and indicators, including easier access to data.

Development is also going on under Eurocordex regarding high resolution data with different bias correction methods but at the same time introducing some new uncertainties in bias correction.

Finally in the discussion it was noted that going through the past indicators and earlier data with a new framework can provide useful additional information.

A first set of criteria for CLIPC indicators: Example of how the CLIPC criteria could work in practice

Luis Costa, CLIPC /PIK

Luis Costa presented the application of the preliminary criteria for indicators. The main idea is to have a systematic framework that can be used to arrive at clear conclusions in D 7.1 on strengths and weaknesses of climate and climate impact indicators and underlying data. The aim is to provide a proof of concept of indicator criteria. A general starting point is the grouping of indicators into three tiers and the grouping of the criteria into two main groups: Scientific adequacy and feasibility and Usability, relevance and scope of use. In addition there is a consideration of impact functions which can be seen to relate indicators of different tiers to one another, or be used to develop new composite indicators.

General discussion on criteria

The discussion raised as a particular issue the link between the impact functions and the criteria and how to deal with that link. The need to consider some form of a numerical scale for the criteria was also raised.

In CLIPC there will be a need to consider possibilities to combine indicators thereby possibly producing new indicators. It was, however, noted, that these user driven combinations should not be considered as "indicators" in the sense of those that have been evaluated using the criteria.

For the input variables there is a need to achieve specificity with standard reference names ensuring traceability and transparency. The criteria to be stressed in particular are those that related to the quality of underlying data [thresholds, standard disclaimer, benchmarks and "references to authoritative sources"]. It was noted that verifiability should be emphasized for impact indicators and also the recognition of limits impact/indicator functions with respect to time interval and geographical region especially in the context of impact functions which have been developed for specific locations with specific data. The (limits of) transferability should be flagged through criteria. Based on criteria a distinction can be made between research/exploratory

work that may contribute to future indicators as opposed to “real indicators” that fulfill selection criteria.

Priority users/user groups for CLIPC and what preliminary demands they may put on data and impact indicators to be provided

Annemarie Groot, CLIPC /Alterra

Annemarie Groot presented the priority user groups and the user consultation strategies and user requirements that have been employed in other projects. She concluded that a pragmatic approach is needed in selecting priority user groups. Potential users can be placed in three circles dependent on the involvement in the CLIPC project and related projects. The inner circle consist of those already involved in projects of CLIPC partners, the second of users already involved in other similar European and national projects and finally the potential users of interest recognised by various partners but not necessary involved in any projects that has direct links with CLIPC . The user needs can be specified by identifying four categories, according to expected requirements and capabilities to handle climate change information:

1. Climate Scientists
2. Biophysical impact researchers
3. Boundary workers (or intermediary organizations) and socio-economic impact researchers
4. Societal end-users

The conclusion had been reached that the focus in identifying user need should be on the first three categories.

Brainstorming envisioned key features the CLIPC toolbox

Mikael Hildén (facilitator), CLIPC /SYKE

Using the priority user groups as guideline the workshop discussed what functionalities should be developed in CLIPC for the toolbox.

The discussion identified a number of general requirements and technical features that should be considered in developing the toolbox. In addition key features for the specific user groups were identified.

General requirements

User friendliness should be a basic starting point. Users could achieve guidance by registering according to the focus of their interest and the expressed interest would guide the user to relevant parts of the toolbox. There should also be opportunities for providing feed-back.

One way of guiding users is to take policy needs as a base for supporting the users' selection of topics in the toolbox; for example energy/bioenergy; climate data/impacts. The specific entry points should be supported by transparent meta-data explaining the base for the work.

The credibility of the contents of the toolbox needs to be ensured through:

- appropriate quality control and quality control procedures, including bench marking of quality with other related services and products
- verifiability of information and data provided
- disclaimers on data/indicators as appropriate

The toolbox should preferably include exploratory tools for analyzing the indicators that would allow comparison of indicators: across topics; across different time intervals and across different areas. It could also allow users to bring in their “own” data

to compare with what is available in the toolbox. This will require standards for data input and comparison but also disclaimers on the use of data for such comparisons. A distinction has to be made between 'indicators that have been approved by the project to be included in the portal based on QA/QC procedures and "User indicators & indexes", which are only exploratory products, not "approved indicators" even if they use information and data included in the toolbox.

In order to guide users there is a need to reflect on what limitations should be built into the toolbox that would stop users from creating combinations and analyses that are scientifically unjustified and potentially misleading. This is closely related to the question on what post processing opportunities CLIPC will provide. With extensive post processing opportunities there is a need for built in "warnings" on combinations of data or explanations for recommended combinations.

Different types of tools have different demands in this respect. Thus visualisations can be largely predefined giving users "controlled" ability to modify data through spatial and temporal aggregation. Opportunities for statistical analysis and overlay of, for example, uncertainties are more challenging in that they require the user to be experienced and aware of caveats.

The toolbox should provide free and open access to the available material and ensure its traceability and transparency. A review team is needed for checking all data and indicators that are proposed to be included in the toolbox.

Technical requirements

The amount of data and type of indicators should be taken into account in selecting server for the toolbox. The server must be able to cope with numerous simultaneous users requesting downloads of indicator information and data.

Registration of users according to needs could also lead to different user interfaces which are based on user profiles/areas of interest. There could also be a system for flexible data discovery (search function) but also (partly) predefined selection of products and indicators from the portal which the user can reach by specifying broad themes (see general requirements, user friendliness).

A help desk function should be included in the design of the portal. This could also include a general wish list for the management of tool box, and information on updates and new developments. Informing regular users can be considered. For example MyOcean regularly sends out information on new developments and products to registered users.

The toolbox should be able to automatically inform users of processes, in particular, it could provide information on processing time for "heavy requests" involving large amounts of data.

Specific characteristics serving particular user groups:

Climate scientists

Need for specific and detailed data; will wish to have maximum options to explore data further by analysing it using different user driven tools for treating the data, including scatter plots, free choice of timelines and other technical treatment. Flexibility with many choices in examining the data is a key to usefulness from the climate scientists point of view.

Climate scientists are also likely to wish opportunities that allow sharing of files, and the extraction of subsets of data for areas & issues

Impact scientists

Impact scientists are likely to benefit from partly predefined analyses of particular data and indicators, and to wish to have explanations and visualisations of climate

data (Tier-1) indicators in particular. They could also wish to see pointers to similar / related data starting from some topic. This can be achieved by clearly labelling specific information according to areas/topics of interest.

Impact scientists are dependent on good metadata when reporting analyses involving the combination of different indicators to get insights into Tier-2 and 3 of the indicators, and should also be required to contribute to the development of metadata.

Impact research will have a particular interest in considerations of links between impacts and adaptation action, and how to monitor measures improving adaptation or adaptive capacity. Therefore indicators or tools that allows the exploration of the available information in the light of, for example, the EU-adaptation strategy at Tier-2 and 3, will be of particular interest for impact scientists.

Intermediaries

Intermediaries are particularly likely to benefit from a toolbox that provides as many finalized products as possible. This means for example:

- Predefined maps/graphs of specific indicators with explanations and interpretations of plots provided.
- Predefined time slices (with possibilities for users to easily adjust them to their own preferences; or with time sliders to view changes over time)
- Possibilities to zoom different geographical levels: Regional (NUTS3), national, European wide aggregation
- Predefined aggregations of indicators developed by experts; possibly allowing users to define weights by users;
- Some (limited) possibilities for developing “indicators on the fly” to allow exploratory work with respect to relationships between indicators.
- Vivid examples based on/linked with the indicators, narratives and success stories and interesting cases

Intermediaries are also likely to benefit from information of (causal) links between indicators, but also from social/cognitive links (“those who viewed this also looked for...”) and indicators that can guide and inform steps towards adaptation.

Processes for user engagement

The workshop noted that there is a special need to develop processes for user engagement in the toolbox. An important function will be to include features that engage users, allowing them to make their feedback visible and to directing and guide user feedback with, for example FAQs.

CLIPC should link with activities such as CharmE that has focused on how to allow users to view or create annotations that describe how climate data has been used and what has been learned. For CLIPC the analogue is to describe the use impact data and indicators.

The point noted under general and technical requirements concerning category specific user registration can provide different entry points that take user need into account, and in so doing guiding users to key topics of her/his interest, and providing specific avenues for engagement.

In the discussion it was noted that the EIONET is a specific forum where the CLIPC can be marketed in particular to “intermediaries”, but it will require concrete examples of what the toolbox can provide.¹

¹ CLIPC was presented at the annual EIONET workshop on CC IVA held 24 June, EEA, Copenhagen. Interested countries were asked to contact CLIPC .

Next steps

It was noted that CLIPC can be seen as a prototype for services that Copernicus will develop further. Particular attention will have to be devoted to QA/QC procedures. There is thus a need to organize meetings between CLIPC and organisations with relevant tasks in Copernicus where the contents of climate services and the links to the development of CLIPC can be discussed further. Relevant discussion partners are also the other pre-operational climate change service projects², in particular those developing projections and/or predictions and that already have experience of user involvement activities. Lessons learnt should be assessed from these activities in order to be more focused and effective in CLIPC (and for the CCCS as a whole). There are also opportunities to identify and talk with EU-wide 'sectoral' organisations that maintain many relevant indicators including WHO, ECDC, ISDR but also those related to water and ecosystem issues. EEA can facilitate, through networks and systems managed by EEA colleagues, including WISE and BISE such discussions.

There is also a need to initiate the processes for integration /convergence between CLIPC and EEA activities. This will call for user meeting and smaller specific meetings on necessary steps in 2015 to track progress in CLIPC and to identify opportunities for establishing more formal links between CLIPC and the EEA Climate-ADAPT and indicator work.

There will also be a need to consider widely links to different activities that are potentially relevant for the production of indicators, for example the ISI-MIP (<http://www.isi-mip.org/>) which brings together impact models across sectors and scales to create consistent and comprehensive projections of the impacts of different levels of climate change. Also the outcomes of several finalized EU-projects such as CLIMSAVE and ongoing EU projects such as ToPDAd, IMPRESSIONS and BASE need to be considered.

List of participants

Surname	First name	Organisation
Barring	Lars	SMHI
Carter	Tim	SYKE
de Groot	Annemarie	Alterra
Fons-Esteve	Jaume	UAB
Fronzek	Stefan	SYKE
Füssel	Hans-Martin	EEA
Gobron	Nadine	JRC
Hildén	Mikael	SYKE
Jol	André	EEA
Juckes	Martin	STFC
Kurnick	Blaz	EEA
Luojus	Kari	FMI
Lüickenkötter	Johannes	TUDO
McCormick	Niall	JRC
Swart	Rob	Alterra
Teichmann	Claas	CSC
Thépaut	Jean-Noël	ECMWF

² <http://www.copernicus.eu/pages-principales/projects/other-fp7-projects/climate-change/>

ANNEX 2

Investigation of indicators according to the potential uses

Table 15. Investigation of indicator according to the potential uses identified in the CLIPC project. The number 1 and 0 (zero) are a preliminary indications of indicator strength (1) or weakness (0) to a given use. NAs highlight that information is incomplete.

Identified uses of indicators	Supporting the development of adaptation strategies and plans	Production of risk and vulnerability assessments	Input in research on climate change		Creating awareness	
			Ensemble details	Data availability	Main use of the indicator	Indicator allocated to two or more themes
Criteria used from Tables 7 and 8	Presence/absence of adaptive/coping capacity	Length of the time series				
Arctic and Baltic Sea ice	0	1	0	1	1	0
Bathing water quality	0	0	NA	1	0	0
Chlorophyll in transitional, coastal and marine waters	0	0	NA	1	0	0
Chlorophyll-a concentration (observations)	0	0	NA	1	1	0
Climatic favourability of tree species (projections)	1	0	1	NA	0	0
Coastal flood damage and adaptation costs (projections)	1	NA	NA	0	0	1
Cold days	NA	1	0	NA	0	1
Cold nights	NA	0	0	NA	0	1
Cold spell duration index	NA	1	1	NA	0	1
Consecutive dry days	NA	1	0	NA	1	1
Consecutive wet days	NA	1	1	NA	0	1
Distribution of marine species	1	1	NA	0	0	0
Diurnal temperature range	NA	NA	1	NA	0	1
Extremely wet days	NA	NA	1	NA	1	1
Floods and health	0	0	NA	0	0	0
Freshwater biodiversity and water quality	0	1	NA	1	1	0
Frost days	NA	NA	1	NA	1	1
Glaciers mass balance	0	1	0	1	1	0
Sea level change	0	NA	NA	1	1	1
Greenland ice sheet	0	1	NA	1	1	0
Grow season length	NA	NA	1	NA	1	0
Growing season for agriculture	1	1	NA	1	1	0
Hazardous substances in marine organisms	0	0	NA	1	1	0
heavy precipitation days	NA	NA	1	NA	1	1
Ice days	NA	NA	1	NA	1	1
Lake and river ice cover	0	0	NA	0	1	0
Lake and river ice phenology	0	1	NA	1	1	0
Lake Ice Extent	0	0	NA	1	0	0
Land elevation below projected sea-level	0	NA	NA	0	1	1
Max 1 day precipitation	NA	1	1	NA	0	1
Max 5 day precipitation	NA	NA	1	NA	0	1
Maximum of daily minimum temperature	NA	1	1	NA	0	1
Maximum of daily maximum temperature	NA	1	0	NA	0	1
Mean precipitation	0	0	0	1	1	1
Minimum of daily minimum temperature	NA	1	1	NA	0	1
Minimum of daily maximum temperature	NA	1	1	NA	0	1
Moth Phenology Index	0	NA	NA	NA	1	0
Number of wet days	NA	NA	1	NA	0	1
Nutrients in transitional, coastal and marine waters	0	NA	NA	1	0	0

Identified uses of indicators	Supporting the development of adaptation strategies and plans	Production of risk and vulnerability assessments	Input in research on climate change		Creating awareness	
			Ensemble details	Data availability	Main use of the indicator	Indicator allocated to two or more themes
Criteria used from Tables 7 and 8	Presence/absence of adaptive/coping capacity	Length of the time series				
Observed development of ocean acidification	1	1	NA	1	0	0
Ocean heat content	0	1	NA	1	1	0
Permafrost	0	1	0	1	1	0
Precipitation extremes	0	1	NA	1	1	1
Projected change in average annual and seasonal river flow	0	1	0	1	0	0
Projected change in river floods with a return period of 100 years	0	0	0	1	1	0
Projected changes in water-limited crop yield	1	1	0	1	0	0
Projection of ocean acidification	0	1	NA	1	0	0
River floods	0	0	NA	1	1	0
River flow	0	1	NA	1	1	0
River flow (projected)	0	0	NA	1	0	0
River flow drought	0	1	NA	1	0	1
Sea level change (observations)	0	1	NA	1	1	0
Sea level change (projections)	0	NA	NA	NA	0	0
Sea surface temperature (observations)	0	0	NA	1	1	0
Simple daily intensity	NA	NA	1	NA	1	1
Snow cover (observations and projections)	0	1	NA	1	1	0
Snow extent (observations)	0	0	NA	1	0	1
Standardized SnowPack Index	0	0	NA	1	1	0
Snow Water Equivalent	0	0	NA	1	1	0
Storm surges	0	NA	NA	NA	0	0
Summer days	NA	NA	1	NA	0	1
The length of thermal growing season	0	0	NA	1	0	0
Total wet-day precipitation	NA	1	1	NA	1	1
Tropical nights	NA	1	1	NA	1	1
Very heavy precipitation days	NA	1	1	NA	1	1
Very wet days	NA	1	1	NA	1	1
Warm days	NA	0	0	NA	0	1
Warm nights	NA	0	0	NA	0	1
Warm spell duration index	NA	1	1	NA	1	1
Water scarcity	0	NA	NA	NA	0	1
Water temperature (observations)	0	1	NA	0	1	0
Water temperature (projections)	0	0	NA	0	1	0
Water-limited crop productivity (projections)	1	NA	0	1	1	1
Irrigation water requirement	0	1	0	1	1	1
Ocean acidification	0	1	NA	1	1	0
Intensity of urban heat island with city size	0	NA	NA	0	0	0
Heating degree-days	1	0	NA	1	1	0
Rainfall Deciles (observations)	1	NA	NA	NA	0	1
Reconnaissance Drought Index	1	NA	NA	NA	1	1
Annual average damage from river floods	0	NA	NA	0	1	0
Average annual heat-related deaths per 100,000 habitats	1	0	1	0	1	0
Growing Degree Days	1	NA	NA	NA	1	0
Chilling Units (observations)	1	NA	NA	NA	0	0
Potential impact of river flooding on major roads	0	1	0	1	0	0

Identified uses of indicators	Supporting the development of adaptation strategies and plans	Production of risk and vulnerability assessments	Input in research on climate change		Creating awareness	
			Ensemble details	Data availability	Main use of the indicator	Indicator allocated to two or more themes
Criteria used from Tables 7 and 8	Presence/absence of adaptive/coping capacity	Length of the time series				
Potential impact of river flooding on railways	0	I	0	I	0	0
Potential impact of river flooding on settlements	0	I	0	I	0	0
Percentage change in arrivals/departures due to global warming	0	I	0	0	I	0
Annual olive-crop yield	I	NA	NA	NA	0	0
Natural disasters	0	0	NA	I	I	I

Schematic illustration of SRES scenarios




Figure 18. Main characteristics of the four SRES storylines and scenario family (as in SRES IPCC 2000, Figure SPM-1).

- The A1 storyline and scenario family describes a future world of very rapid economic growth, global population that peaks in mid-century and declines thereafter, and the rapid introduction of new and more efficient technologies. Major underlying themes are convergence among regions, capacity building, and increased cultural and social interactions, with a substantial reduction in regional differences in per capita income. The A1 scenario family develops into three groups that describe alternative directions of technological change in the energy system. The three A1 groups are distinguished by their technological emphasis: fossil intensive (A1FI), non-fossil energy sources (A1T), or a balance across all sources (A1B).
- The A2 storyline and scenario family describes a very heterogeneous world. The underlying theme is self-reliance and preservation of local identities. Fertility patterns across regions converge very slowly, which results in continuously increasing global population. Economic development is primarily regionally oriented and per capita economic growth and technological change are more fragmented and slower than in other storylines.
- The B1 storyline and scenario family describes a convergent world with the same global population that peaks in mid-century and declines thereafter, as in the A1 storyline, but with rapid changes in economic structures toward a service and information economy, with reductions in material intensity, and the introduction of clean and resource-efficient technologies. The emphasis is on global solutions to economic, social, and environmental sustainability, including improved equity, but without additional climate initiatives.

- The B2 storyline and scenario family describes a world in which the emphasis is on local solutions to economic, social, and environmental sustainability. It is a world with continuously increasing global population at a rate lower than A2, intermediate levels of economic development, and less rapid and more diverse technological change than in the B1 and A1 storylines. While the scenario is also oriented toward environmental protection and social equity, it focuses on local and regional levels.
(as in IPCC 2000,Box SPM-1)



ISBN 978-952-11-4650-3 (PDF)

ISSN 1796-1726 (online)