

and focusing on “melancholia,” which includes only the most severe cases and clearly respond to antidepressants and/or electroconvulsive therapy [3].

According to Bech, a psychiatrist and expert in psychometrics, the total score on the Hamilton depression scale should not be used in antidepressant RCTs; he recommends using the version with six items [4]. Finally, there remains much controversy as to whether antidepressants increase suicidal ideation, but I opine that this debate is rather pointless. Most people with suicidal ideation do not attempt suicide; those who do are statistical outliers [5]. Similarly, most suicide attempters do not complete suicide; those who do are “probably” statistical outliers, although we have very limited understanding of those who complete suicide since most of them have never seen a psychiatrist [5]. In addition, more than half of the patients in the author’s US hospital for patients with severe mental illnesses such as schizophrenia, mania, severe depression, and catatonia have a history of attempted suicide and will thus never be included in an RCT, which is quite problematic because data related to their treatment are very limited.

Jose de Leon

University of Kentucky Mental Health Research Center at Eastern  
State Hospital  
Room 3A15A, 1350 Bull Lea Road  
Lexington, KY 40511, USA  
Tel.: +1-859-246-8440; fax: +1-859-246-8446.  
E-mail address: [jdeleon@uky.edu](mailto:jdeleon@uky.edu)

## References

- [1] Ebrahim S, Bance S, Athale A, Malachowski C, Ioannidis JP. Meta-analyses with industry involvement are massively published and report no caveats for antidepressants. *J Clin Epidemiol* 2016;70:155–63.
- [2] Feinstein AR. What can clinicians contribute to mutual challenges in medical statistics? *J Clin Epidemiol* 1999;52:365–9.
- [3] de Leon J. Is Psychiatry only neurology? Or only abnormal psychology? Déjà vu after 100 years. *Acta Neuropsychiatr* 2015;27(2):69–81.
- [4] Bech P. Is the antidepressive effect of second-generation antidepressants a myth? *Psychol Med* 2010;40:181–6.
- [5] de Leon J, Baca-García E, Blasco-Fontecilla H. From the serotonin model of suicide to a mental pain model of suicide. *Psychother Psychosom* 2015;84:323–9.

<http://dx.doi.org/10.1016/j.jclinepi.2016.03.001>

## Many continuous variables such as the duration of the common cold should be analyzed using the relative scale



Alba et al. [1] compared the heterogeneity of meta-analyses on the basis of the type of outcome, and concluded that meta-analyses that evaluate continuous outcomes

showed substantially higher  $I^2$  values than meta-analyses of binary outcomes. However, in a proper comparison, like should be compared with like.

First, their measure of effect on binary outcomes was a relative effect (risk ratio, odds ratio), whereas their measure of effect on the continuous outcomes was an absolute effect or effect in standard deviation units (standardized mean difference). Second, there is great variability in the kind of phenomena that are measured by continuous outcomes. For example, quite diverse scales are used in psychiatry, and they are analyzed as continuous variables, yet those outcomes are quite different from, say, the occurrence of lung cancer (binary) or the duration of the common cold (continuous). The latter two directly measure biological phenomena and they might substantially differ from arbitrary scales. These two issues were not considered by Alba et al. [1].

Effects in binary outcomes are usually better captured by the relative scale than by an absolute scale, for example, the 10-fold increase in lung cancer risk by smoking. Similarly, the relative effect can better capture the differences between the compared groups in many biological phenomena that are measured as continuous outcomes.

Our Cochrane review on vitamin C and the common cold showed the shortest average duration of colds in a placebo group of children to be 2.6 days, whereas the longest was 14 days (Analysis 2.1.2 in [2]). Such a great variation in cold duration in the placebo groups is explained by variations in viruses and in operational outcome definitions between different trials, and so forth. An ideal treatment might shorten the long colds by 10 days, whereas such an effect on short colds is impossible. Thus, the absolute effect scale is not feasible for analyzing effects on the duration of colds. Therefore, we used the relative scale in our analysis. We calculated that the heterogeneity in the effect of vitamin C on cold duration over 14 trials on children was  $I^2 = 27%$  ( $P = 0.17$ ; Analysis 2.1.2 in [2]). The same set of studies gives  $I^2 = 46%$  ( $P = 0.03$ ) when the analysis is done on the absolute scale. The latter  $I^2$  is 19 percentage points more (1.7-fold) than the  $I^2$  of the relative scale (See Appendix). Thus, meta-analyses on different scales can lead to divergent  $I^2$  values. Different metrics may have biased the comparison of meta-analyses on binary and continuous outcomes [1].

Alba et al. cited a meta-analysis on zinc and the common cold [3] as a specific example of great heterogeneity in meta-analyses on continuous outcomes. However, much of the heterogeneity in that particular meta-analysis was explained by the apples and oranges problem [4]. Heterogeneity over 13 trials on zinc lozenges and the common cold ( $I^2 = 89%$ ) was largely explained by the dose of zinc and the type of zinc salt used in the lozenges [5]. Thus, separating apples from the oranges can explain a substantial part of heterogeneity in some meta-analyses. For the

previously mentioned reasons, the conclusions of Alba et al. [1] do not seem justified.

### Supplementary data

Supplementary data related to this article can be found online at <http://dx.doi.org/10.1016/j.jclinepi.2016.03.020>.

Harri Hemilä  
Department of Public Health  
University of Helsinki  
POB 20, Tukholmankatu 8 B  
FIN-00014, Helsinki, Finland  
Tel.: +358-41-532-9987.  
E-mail address: [harri.hemila@helsinki.fi](mailto:harri.hemila@helsinki.fi)

### References

- [1] Alba AC, Alexander PE, Chang J, MacIsaac J, DeFry S, Guyatt GH. High statistical heterogeneity is more frequent in meta-analysis of continuous than binary outcomes. *J Clin Epidemiol* 2016;70:129–35.
- [2] Hemilä H, Chalker E. Vitamin C for preventing and treating the common cold. *Cochrane Database Syst Rev* 2013;CD000980.
- [3] Science M, Johnstone J, Roth DE, Guyatt G, Loeb M. Zinc for the treatment of the common cold: a systematic review and meta-analysis of randomized controlled trials. *CMAJ* 2012;184:E551–61.
- [4] Hemilä H. Zinc acetate lozenges may shorten common cold duration by up to 40%. *CMAJ* 2012. eLetter. Available at: [http://www.cmaj.ca/content/184/10/E551.abstract/reply#cmaj\\_el\\_706238](http://www.cmaj.ca/content/184/10/E551.abstract/reply#cmaj_el_706238). Accessed April 22, 2016.
- [5] Hemilä H. Zinc lozenges may shorten the duration of colds: a systematic review. *Open Respir Med J* 2011;5:51–8. <http://dx.doi.org/10.2174/1874306401105010051>.

<http://dx.doi.org/10.1016/j.jclinepi.2016.03.020>