

## Accepted Manuscript

Formation of cyanuric acid from cyanate adsorbed at gold electrodes

William Cheuquepán, Antonio Rodes, José M. Orts, Juan M. Feliu

PII: S1388-2481(16)30269-7  
DOI: doi:[10.1016/j.elecom.2016.11.005](https://doi.org/10.1016/j.elecom.2016.11.005)  
Reference: ELECOM 5819

To appear in: *Electrochemistry Communications*

Received date: 31 October 2016  
Revised date: 8 November 2016  
Accepted date: 8 November 2016



Please cite this article as: William Cheuquepán, Antonio Rodes, José M. Orts, Juan M. Feliu, Formation of cyanuric acid from cyanate adsorbed at gold electrodes, *Electrochemistry Communications* (2016), doi:[10.1016/j.elecom.2016.11.005](https://doi.org/10.1016/j.elecom.2016.11.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Formation of cyanuric acid from cyanate adsorbed at gold electrodes****William Cheuquepán\*, Antonio Rodes\*#, José M. Orts<sup>&\*#</sup> and Juan M. Feliu\*#**

Departamento de Química Física# and Instituto Universitario de Electroquímica\*

Universidad de Alicante, Apartado 99, E-03080 Alicante, SPAIN

&amp;Corresponding author:

Email: jm.orts@ua.es

Phone/Fax: (+34) 965909814

**Abstract**

We report the formation of cyanuric acid species at Au(111) electrodes in cyanate-containing solutions, due to the electroless trimerization of isocyanic acid. Similar experimental bands in the range between 1300 and 1900  $\text{cm}^{-1}$  are observed in Surface Enhanced Infrared Reflection Spectroscopy experiments under Attenuated Total Reflection conditions (ATR-SEIRAS) with solutions containing either cyanate anions at high concentration or cyanuric acid. The experimental frequencies agree well with those obtained from Density Functional Theory (DFT) calculations for the adsorbed cyanurate anion, bonded to the metal in a tridentate configuration, with its molecular plane perpendicular to the metal surface.

**Keywords: ATR-SEIRAS; cyanate; cyanuric acid; cyanurate; DFT; Au(111)**

## 1. Introduction.

We have previously reported [1] a study of the adsorption of cyanate anions at Au(111) and Au(100) single crystal electrodes, combining external reflection spectroelectrochemical experiments and Density Functional Density (DFT) calculations. Taking into account the calculated optimized adsorbate geometries, adsorption energies and vibrational harmonic frequencies, we assigned the adsorbate bands found experimentally between 2100 and 2300  $\text{cm}^{-1}$ . In agreement with previous works [2-5], these bands correspond to the asymmetric OCN stretch of N-bonded, specifically adsorbed isocyanate anions, mainly occupying on-top sites, with their molecular axis preferentially oriented in the direction perpendicular to the metal surfaces.

In this paper we report new spectral features observed experimentally using in situ Surface Enhanced Infrared Reflection Spectroscopy under Attenuated Total Reflection conditions (ATR-SEIRAS) and Au(111)-25nm thin-layer electrodes. The ATR-SEIRAS technique [6-7] is especially advantageous for studying interfacial species, because of the strong exaltation of the infrared absorption by adsorbates and double-layer solvent molecules. An additional advantage is the removal of interferences due to the bulk solvent. Both advantages help in the observation of new absorption bands not detected in the external reflection experiments [1-5], that suggest the formation of new adsorbed chemical species. We show in this report that the observed bands are characteristic of species derived from cyanuric acid ( $\text{C}_3\text{N}_3\text{O}_3\text{H}_3$ ), a very stable cyclic trimer of isocyanic acid (HNCO). Cyanuric acid [8] can be synthesized by thermal condensation of three urea molecules, with simultaneous release of three ammonia molecules, or starting from metal cyanates in alkaline solutions. Cyanuric acid behaves as a weak acid. From its pKa values (6.88, 11.40, 13.5) [9] the only species that are expected to exist in significant amounts both in solution and at the electrified interface are the undissociated

acid and the cyanurate monoanion ( $C_3N_3O_3H_2^-$ ), that results from the first acid dissociation equilibrium. In this anion the negative charge on the N atom is stabilized by the two neighbouring carbonyl groups.

In order to support this interpretation, we have compared the voltammetric behaviour and the ATR-SEIRAS spectra obtained with solutions containing either cyanate salts or cyanuric acid. We have also carried out periodic DFT calculations of geometry and frequencies for the cyanurate anion adsorbed at the electrode surface. These calculations will provide the basis for the assignment of the experimental vibrational bands.

## 2. Experimental and computational details.

Working solutions were obtained by dissolving either sodium cyanate (96%, Sigma-Aldrich) or cyanuric acid (98%, Sigma-Aldrich), in a solution of sodium perchlorate (99.99%, Sigma-Aldrich) prepared in ultrapure water (18.2  $M\Omega\cdot cm$ , Elga-Vivendi). These solutions were deaerated with Ar (N50, Air Liquide) and blanketed with this gas during the experiments. Solutions in deuterated water were prepared with deuterium oxide (99.9 atom %D, Aldrich) which was used as received. In all the spectroelectrochemical experiments a Ag/AgCl/KCl (sat) electrode and a gold wire were used as the reference and the counter electrode, respectively. Details for the ATR-SEIRAS experiments regarding both the preparation of the gold thin film electrodes and the collection of the in situ infrared spectra can be found in previous works [10,11].

Projector-augmented-wave [12-13] DFT calculations of the cyanurate anion adsorbed on a model Au(111) surface were carried out using the VASP [14-17] code (version 4.6), with the PBE [18-19] functional.

The slab used for modeling a (3x3) cell of the Au(111) surface consisted of 4 metallic layers (with 9 Au atoms per layer). The structure of one adsorbed cyanurate species was optimized while the positions of the metal atoms were kept fixed at their bulk equilibrium distances (0.29520 nm) calculated with the same functional. A vacuum region (of more than 12 Angstrom) was used for separating the slabs in the Z direction.

Sampling of the Brillouin zone used a Monkhorst-Pack [20] (3x3x1) scheme and for smearing we used the Methfessel-Paxton [21] method (second order,  $\sigma=0.2$  eV). The convergence criteria used were:  $10^{-5}$  eV for electronic convergence, and 0.02 eV/Angstrom for the forces.


### 3. Experimental results.

Curve b in Figure 1 shows the cyclic voltammogram obtained with a gold thin film electrode with preferential (111) orientation (denoted Au(111)-25 nm in the following [7]) in contact with a 10 mM NaOCN+ 0.08 M NaClO<sub>4</sub> solution. As previously reported for Au(111) and Au(100) electrodes [1], a significant charge contribution associated with the presence of cyanate is superimposed on the voltammetric response of the cyanate-free solution (curve a) for potentials above -0.20 V. No significant faradic oxidation currents that could be related to the irreversible oxidation of cyanate anions can be detected in the explored potential region. A broad feature centered at 0.25 V and a peak at ca. 0.35 V are observed in the positive going sweep for the 10 mM solution.

Figure 2 shows a series of potential-dependent ATR-SEIRA spectra collected for the Au(111)-25 nm electrode in contact with 10 mM NaOCN + 0.08 M NaClO<sub>4</sub> solutions prepared in water (A) or deuterium oxide (B). Spectra are referred to that collected in

the same solution at -0.5 V. A potential-dependent feature is observed for adsorbed cyanate at 2150-2230  $\text{cm}^{-1}$ [1-5]. In addition, several positive-going features appear between 1800 and 1300  $\text{cm}^{-1}$ . First, a clearcut feature appears at ca. 1716  $\text{cm}^{-1}$  in the spectrum collected at -0.40 V, being blueshifted up to 1743  $\text{cm}^{-1}$  at 0.60 V. This feature is accompanied by a shoulder at ca. 1781  $\text{cm}^{-1}$ . Moreover, additional features are observed at 1654, 1496, 1388 and 1330  $\text{cm}^{-1}$ . The intensities of all these features increase with the electrode potential up to 0.60 V and decrease down to zero when the electrode potential is stepped back to -0.50 V (not shown). In order to avoid interferences from the bending modes of interfacial water, additional experiments were carried out in deuterium oxide. Potential-dependent absorption bands are also clearly observed. Positive-going features appear (panel B in Figure 2) at ca. 2213, 1789, 1720, 1650, 1581, 1480 and 1438  $\text{cm}^{-1}$  (all these frequency values measured in the spectrum collected at 0.60 V), some of them showing a band frequency strongly dependent on the electrode potential.

Except for the band at ca. 2213  $\text{cm}^{-1}$ , none of the positive-going features observed between 1850 and 1300  $\text{cm}^{-1}$  can be attributed to vibrational modes of adsorbed cyanate, pointing to the formation of other adsorbates. A relevant observation is that the most intense features appear in the spectral region for carbonyl groups [22]. This suggests the possibility of the trimerization of adsorbed cyanate, via its equilibrium with isocyanic acid, to form cyanuric acid or cyanurate as the final product.



Since this is a highly stable compound [8], spectroelectrochemical experiments in 0.08 M  $\text{NaClO}_4$  solutions containing various concentrations of cyanuric acid were carried out. In this way, curve c in Figure 1 corresponds to the voltammogram recorded in a 3.3 mM cyanuric acid-containing solution. Again, the voltammogram shows an excess of

charge density with respect to the blank solution, attributable to reversible specific anion adsorption-desorption processes. Remark that the main feature in the cyanuric acid-containing solution appears at ca. 0.25 V and fits with the shoulder observed in the cyanate-containing solution.

ATR-SEIRAS experiments have been carried out in cyanuric acid-containing solutions with the Au(111)-25 nm electrode. In this work we limit our discussion to the spectra collected during a potentiostatic dosing experiment at 0.20 V. This electrode potential was chosen as typical for the region where charge excess associated to the presence of cyanuric acid is observed in the corresponding cyclic voltammogram (curve c in Figure 1). The time-dependent spectra obtained under these conditions for a 10 mM solution are reported in Figure 3. Note that the reference spectrum in this experiment was collected at 0.20 V in the blank solution just before adding cyanuric acid. A few minutes after dosing, positive-going bands develop in the absorbance spectra whose intensities steadily increase with time. These bands appear at ca. 1743  $\text{cm}^{-1}$  (with a shoulder at 1781  $\text{cm}^{-1}$ ), 1648, 1580, 1446 and 1396  $\text{cm}^{-1}$ , nearly at the same frequencies as the features for the 10 mM cyanate solution. No absorbance band is observed at ca. 2200  $\text{cm}^{-1}$  that could be ascribed to adsorbed cyanate. In addition, none of the positive-going bands reported in Figure 3 are observed when the dosing experiment was carried out at -0.50 V. This indicates that at this potential, either no adsorption of cyanurate takes place, or the adsorbates have their molecular planes essentially parallel to the metal surface. The observation of changes in the spectra of interfacial water suggests that the latter possibility could be the case.

#### 4. Computational results and discussion.

According to the pKa values for cyanuric acid, the cyanurate monoanion is expected to be the predominant anionic species derived from cyanuric acid in solution. The most stable adsorbed geometry found for this species involves bonding to the Au(111) surface through two oxygen atoms and the deprotonated N atom, all three being located in positions close to top sites, and with the molecular plane aligned with the metal surface dense rows. This orientation is compatible with the observation of adsorbate absorption bands in the ATR-SEIRA spectra (because of the surface selection rule [23]). Table 1 summarizes the experimental and calculated frequencies for the adsorbed cyanurate anion, together with their assignments, based on the visualization of the normal modes using Jmol [24]. It can be seen that most of the main experimental features can be attributed to vibrations of adsorbed cyanurate. Within the uncertainty typical of the experimental measurements (a resolution of  $8\text{ cm}^{-1}$ ) and the errors intrinsic of the calculated frequencies (around 2% of the frequency values), a very good agreement exists between our calculated and experimental values. On this basis we can conclude that on these Au(111) electrode surfaces, the trimerization of isocyanic acid/cyanate takes place in an electroless process, producing adsorbed cyanurate. Some experimental features still remain to be explained. It must be taken into account that the calculated frequencies correspond to a surface coverage of 1 cyanurate per 9 surface metal atoms. Because of the ability of this compound to participate in hydrogen bonds (with cyanurate, cyanuric acid, cyanate and water, either coadsorbed or in the vicinity of the electrode surface), some effects on the vibrational frequencies are to be expected with increasing surface coverage, including also collective vibrational modes. These points will be explored elsewhere.



## 5. Conclusions.

ATR-SEIRAS experiments evidence the formation of adsorbed cyanurate anions at Au(111)-25 nm electrodes in contact with cyanate-containing solutions. This latter species coexists with N-bonded cyanate anions. The main experimental vibrational bands between 1800 and 1300  $\text{cm}^{-1}$  can be explained on the basis of the optimized geometry and harmonic vibrational frequencies obtained from DFT calculations for adsorbed cyanurate bonded to the metal in a tridentate fashion, with the molecular plane perpendicular to the electrode surface.

## ACKNOWLEDGEMENTS

The authors acknowledge funding by Ministerio de Economía y Competitividad (project CTQ2013-44083-P) and University of Alicante. William Cheuquepán is grateful for the award of a F.P.I. grant associated to project CTQ2009-13142.

## REFERENCES

1. W. Cheuquepán, J.M. Orts, A. Rodes, J.M. Feliu, "DFT and spectroelectrochemical study of cyanate adsorption on gold single crystal electrodes in neutral medium", *J. Electroanal. Chem.*, in press. Electronic version: DOI:10.1016/j.jelechem.2016.10.011 (available on-line 8-OCT-2016).
2. O. Hofmann, K. Doblhofer, and H. Gerischer, "Infrared reflection-absorption measurements on emersed gold electrodes", *J. Electroanal. Chem. Interfacial Electrochem.*, 161 (1984) 337-344.
3. D. S. Corrigan and M. J. Weaver, "Adsorption and oxidation of benzoic acid, benzoate, and cyanate at gold and platinum electrodes as probed by potential-difference infrared spectroscopy", *Langmuir*, 4 (1988) 599-606.
4. D. S. Corrigan and M. J. Weaver, "The interpretation of solution electrolyte vibrational bands in potential-difference infrared spectroscopy", *J. Electroanal. Chem.*, 239 (1988) 55-66.
5. M. Bron and R. Holze, "Cyanate and thiocyanate adsorption at copper and gold electrodes as probed by in situ infrared and surface-enhanced Raman spectroscopy", *J. Electroanal. Chem.*, 385 (1995) 105-113.
6. M. Osawa, "Dynamic processes in electrochemical reactions studied by surface-enhanced infrared absorption spectroscopy (SEIRAS)", *Bull. Chem. Soc. Jpn.*, 70 (1997) 2861-2880.
7. T. Wandlowski, K. Ataka, S. Pronkin, D. Diesing, "Surface enhanced infrared spectroscopy-Au(111-20 nm)/sulphuric acid - new aspects and challenges", *Electrochim. Acta*, 49 (2004) 1233-1247.
8. G.B. Seifert, "Cyanuric acid and cyanurates", *Russ. J. Coord. Chem.* 28 (2002) 301-324.

9. CRC Handbook of Chemistry and Physics, 89th ed, David R. Lide, editor, CRC Press, Boca Raton, 2008.
10. J. M. Delgado, J. M. Orts, J. M. Pérez, A. Rodes, "Sputtered thin-film gold electrodes for in situ ATR-SEIRAS and SERS studies", *J. Electroanal. Chem.*, 617 (2008) 130-140.
11. W. Cheuquepan, J. M. Perez, J. M. Orts, and A. Rodes, "Spectroelectrochemical and DFT Study of Thiourea Adsorption on Gold Electrodes in Acid Media", *J. Phys. Chem. C*, 118 (2014) 19070-19084.
12. P.E.Bloechl, "Projector augmented-wave method", *Phys. Rev. B* 50 (1994) 17953-17979.
13. G. Kresse, D. Joubert, "From ultrasoft pseudopotentials to the projector augmented-wave method", *Phys. Rev. B*. 59 (1999) 1758-1775.
14. G. Kresse, J. Hafner. "Ab initio molecular dynamics for liquid metals", *Phys.Rev. B* 47 (1993) 558-561.
15. G. Kresse, J. Hafner, "Ab initio molecular dynamics simulation of the liquid-metal-amorphous-semiconductor transition in germanium", *Phys. Rev. B* 49 (1994) 14251-14269.
16. G. Kresse, J. Furthmueller, "Efficient iterative schemes for ab-initio total-energy calculations using a plane-wave basis set", *Phys.Rev. B* 54 (1996) 11169-11186.
17. G. Kresse, J. Furthmueller, "Efficiency of ab-initio total energy calculations using a plane-wave basis set", *Comput.Mater.Sci.* 6 (1996) 15-50
18. J.P. Perdew, K. Burke, M. Ernzerhof, "Generalized gradient approximation made simple", *Phys.Rev.Lett.* 77 (1996) 3865-3868.
19. J.P. Perdew, K. Burke, M. Ernzerhof, "Generalized gradient approximation made simple. Erratum". *Phys.Rev.Lett.* 78 (1997) 1396.

20. H.J. Monkhorst, J.D. Pack, "Special points for Brillouin-zone integrations", *Phys.Rev. B* 13 (1976) 5188-5192.
21. M. Methfessel, A.T. Paxton, "High-precision sampling for Brillouin-zone integration in metals", *Phys.Rev. B* 40 (1989) 3616-3621.
22. G. Socrates, *Infrared and Raman Characteristic Group Frequencies: Tables and Charts*, 3rd Edition, Wiley, Chichester, 2004.
23. M. Osawa, K. Ataka, K. Yoshii, and Y. Nishikawa, "Surface-enhanced Infrared Spectroscopy : the origin of the absorption enhancement and band selection rule in the infrared spectra of molecules adsorbed on fine metal particles", *Appl. Spectrosc.*, 47 (1993) 1497-1502.
24. "Jmol: an open-source Java viewer for chemical structures in 3D". <http://www.jmol.org>, (2015). (Version 14.2.12\_2015.02.11).

Table 1. Experimental and calculated wavenumbers (in  $\text{cm}^{-1}$ ) corresponding to adsorbed cyanurate on Au(111)-25nm electrode surfaces.

Experimental			Calculated	Assignment
A	B	C		
1781	1789	1781	1769	Str CO + bend CNH
1743	1720	1743	1769	Str CO + bend CNH
1654	1650	1646	1627	Sym str OCNCO + bend CNH
	1581	1577	1553	Asym str OCNCO + bend CNH
1496	1488			
1446	1438	1446	1416	Asym str NCN + bend CNH
1388		1396	1397	Sym str NCN + bend CNH

Calculated values correspond to non-deuterated cyanurate.

A) cyanate in water solution

B) cyanate in  $\text{D}_2\text{O}$  solution

C) cyanuric acid solution

## FIGURE CAPTIONS.

Figure 1. Stationary cyclic voltammograms of aAu(111)-25nm thin-layer electrode in contact with 0.08 M NaClO<sub>4</sub> solutions. a) Blank electrolyte; b) with added 10mM NaOCN, and c) with added 0.1mM cyanuric acid.

Figure 2. Series of ATR-SEIRA spectra collected with aAu(111)-25nm thin-layer electrode in contact with 0.08 M NaClO<sub>4</sub> + 10 mM NaOCN solutions in: A)H<sub>2</sub>O; and B) D<sub>2</sub>O. 100 interferograms were co-added with a resolution of 8 cm<sup>-1</sup> at the indicated potentials and referred to the single beam spectrum collected at -0.50 V.

Figure 3. Time-dependent ATR-SEIRA spectra obtained with a Au(111)-25nm thin-layer electrode in contact with a 0.08 M NaClO<sub>4</sub> solution, after dosing cyanuric acid in amount to reach a concentration of 0.1 mM. Dosage potential: 0.20 V. 100 interferograms were coadded with a resolution of 8 cm<sup>-1</sup> at the indicated times and referred to the single beam spectrum collected at the same potential just before the addition of the cyanuric acid.




Fig. 1




Fig. 2






Fig. 3

**Highlights:**

- cyanate anions trimerize in an electroless process yielding cyanuric acid on Au(111) electrodes

- ATR-SEIRAS and DFT calculations agree with specifically adsorbed cyanurate on Au(111) electrodes formed either from cyanuric acid or from cyanate in solution.