



ELSEVIER

Contents lists available at ScienceDirect

Deep-Sea Research II

journal homepage: www.elsevier.com/locate/dsr2

Short-term metabolic and growth responses of the cold-water coral *Lophelia pertusa* to ocean acidification



S.J. Hennige^a, L.C. Wicks^a, N.A. Kamenos^b, D.C.E. Bakker^c, H.S. Findlay^d,
C. Dumousseaud^e, J.M. Roberts^{a,f,g,*}

^a Centre for Marine Biodiversity and Biotechnology, Heriot-Watt University, Edinburgh EH14 4AS, UK

^b School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ, UK

^c School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

^d Plymouth Marine Laboratory, The Hoe, Plymouth PL13DH, UK

^e National Oceanographic Centre Southampton, University of Southampton Waterfront Campus, European Way, Southampton SO143ZH, UK

^f Center for Marine Science, University of North Carolina Wilmington, 601 S. College Road, Wilmington, NC 28403-5928, USA

^g Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll PA37 1QA, UK

ARTICLE INFO

Available online 22 July 2013

Keywords:

Climate change

Lophelia pertusa

Deep-sea coral

Respiration

Growth

Calcification

Mingulay Reef Complex

ABSTRACT

Cold-water corals are associated with high local biodiversity, but despite their importance as ecosystem engineers, little is known about how these organisms will respond to projected ocean acidification. Since preindustrial times, average ocean pH has decreased from 8.2 to ~8.1, and predicted CO₂ emissions will decrease by up to another 0.3 pH units by the end of the century. This decrease in pH may have a wide range of impacts upon marine life, and in particular upon calcifiers such as cold-water corals. *Lophelia pertusa* is the most widespread cold-water coral (CWC) species, frequently found in the North Atlantic. Here, we present the first short-term (21 days) data on the effects of increased CO₂ (750 ppm) upon the metabolism of freshly collected *L. pertusa* from Mingulay Reef Complex, Scotland, for comparison with net calcification. Over 21 days, corals exposed to increased CO₂ conditions had significantly lower respiration rates (11.4 ± 1.39 SE, $\mu\text{mol O}_2 \text{ g}^{-1}$ tissue dry weight h^{-1}) than corals in control conditions (28.6 ± 7.30 SE $\mu\text{mol O}_2 \text{ g}^{-1}$ tissue dry weight h^{-1}). There was no corresponding change in calcification rates between treatments, measured using the alkalinity anomaly technique and ¹⁴C uptake. The decrease in respiration rate and maintenance of calcification rate indicates an energetic imbalance, likely facilitated by utilisation of lipid reserves. These data from freshly collected *L. pertusa* from the Mingulay Reef Complex will help define the impact of ocean acidification upon the growth, physiology and structural integrity of this key reef framework forming species.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Cold-water corals are among the most three-dimensionally complex deep-sea habitats known and are associated with high local biodiversity (Roberts et al., 2006, 2009). However, their remoteness and the relatively short history of ecological research in these habitats mean that to date, we have little information on how these ecosystems will fare in the face of predicted future climate change. Similar to their tropical counterparts, cold-water corals (referred to as CWC herein) are under potential threat from increasing sea temperatures and ocean acidification.

Ocean acidification, also referred to as the ‘other CO₂ problem’ or the ‘evil twin of global warming’, is caused by CO₂ dissolving into the oceans. As atmospheric CO₂ levels increase, more CO₂ dissolves into the oceans and forms carbonic acid, which dissociates to form hydrogen and bicarbonate ions. This process has led to a decrease in pH by 0.1 units since the industrial revolution, and increasing amounts of atmospheric CO₂ are projected to further decrease ocean pH by another 0.3–0.4 pH units by the end of the century in addition to altering seawater carbonate chemistry (Caldeira and Wickett, 2003; Guinotte et al., 2006; Kleypas et al., 1999; Orr et al., 2005). The shift in carbonate chemistry associated with ocean acidification also reduces the saturation state of aragonite, which is a naturally occurring polymorph of calcium carbonate from which most framework-building corals build their skeletons. The aragonite saturation depth or ‘horizon’ (ASH) (Guinotte et al., 2006) is predicted to become shallower (shoal), making it more difficult for calcifying organisms near this depth to

* Corresponding author at: Centre for Marine Biodiversity and Biotechnology, Heriot-Watt University, Edinburgh EH14 4AS, UK. Tel.: +44 131 451 3463.

E-mail addresses: s.hennige@hw.ac.uk (S.J. Hennige),
J.M.Roberts@hw.ac.uk (J.M. Roberts).

maintain their calcified structures, thus affecting net reef growth. CWC are under particular threat as they inhabit a much larger bathymetric range than tropical corals, and as such are closer to the ASH (Fautin et al., 2009; Form and Riebesell, 2012; Guinotte et al., 2006; Kleypas, 2006; McCulloch et al., 2012a; Roberts et al., 2006; Thresher et al., 2011; Turley et al., 2007). Only 5% of CWC are found below the ASH at present (Form and Riebesell, 2012), as net calcification below the ASH would require considerable energetic input at the detriment of other energetic processes. Shoaling of the horizon could thus have potentially severe consequences for CWC.

To date, only a handful of studies have been conducted on the responses of *Lophelia pertusa* to ocean acidification. These focus on growth rate, ranging from very short term 24 hour experiments (Maier et al., 2009) on freshly collected coral, to longer-term 6 month (Form and Riebesell, 2012) experiments on laboratory-kept specimens. Ocean acidification has very varied effects on the growth of calcifying organisms, with different phyla, and even species within phyla showing highly variable responses in experimental studies to date (Ries et al., 2009; Wicks and Roberts, 2012). For scleractinian corals (i.e. calcareous skeleton forming corals), many studies demonstrate a reduction in growth (net calcification rates) in response to ocean acidification (Gattuso et al., 1998; Kleypas and Langdon, 2006; Krief et al., 2010; Langdon and Atkinson, 2005; Marubini et al., 2003). However, corals can actively increase the pH in the organic matrix where calcification occurs. Thus even in situations where the surrounding water is undersaturated with respect to aragonite, calcification can still occur (McCulloch et al., 2012a, 2012b). The process of increasing pH at calcification sites within corals is driven by Ca^{2+} ATPase, which pumps Ca^{2+} ions into the sub-calicoblastic space in exchange for H^{+} ions (Allemand et al., 2004; Cohen and McConnaughey, 2003; Mass et al., 2012). However, this process is energy intensive and thus may require increased food intake (Edmunds, 2011). The low abundance of CWC below the ASH suggests that increased energetic demands cannot usually be met, and that dissolution of exposed skeletal framework may be greater than net calcification by tissue-covered skeleton.

To fully understand the impact of increased CO_2 on live CWC, it is important to combine growth rates with measures of metabolism, which has not been done previously. The research presented here addressed the question of whether ocean acidification will impact upon the metabolism and net calcification rate of the CWC *Lophelia pertusa* in a short-term experiment. To this end, respiration and net

calcification rates were calculated in corals exposed to increased CO_2 conditions every 7 days for a total period of 21 days.

2. Methods

2.1. Sample collection

Colonies of *Lophelia pertusa* were collected from Area 1 within the Mingulay Reef Complex (Roberts et al., 2005, 2009), $56^{\circ} 49.38' \text{N}$, $7^{\circ} 22.56' \text{W}$ (Figs. 1 and 2), during RRS *Discovery* cruise 366/7 in July 2011 (Achterberg and Richier, 2012). The Mingulay complex is a relatively shallow inshore seascape of *Lophelia* reefs that developed through the Holocene with oldest currently dated coral from within the reef mounds at 7.68 ka (Douarin et al., in press). They are the only known inshore *Lophelia* reef in UK waters.

Colonies were collected using a modified video assisted van-Veen grab (Dodds et al., 2007). All colonies were collected from 141–167 m. Upon return to the surface, corals were placed in a holding tank at ambient seabed temperature for 2 days, to recover from collection. Corals were then carefully fragmented into smaller pieces for experiments. These fragments had 5–20 polyps, and were taken from the top of sampled colonies to ensure that relatively young polyps were used consistently, as polyp age can determine physiological response with younger polyps known to be the fastest-growing (Maier et al., 2009). Fragments were attached to pre-labelled bases made of PVC pipe with Grotech Korafix epoxy.

2.2. Treatments

Two tanks (each ~ 350 L seawater within 430 L volume ($0.9 \times 0.6 \times 0.8$ m)) were established for experiments; one at ambient reef conditions of 9.5°C , 380 ppm and the other at 9.5°C , 750 ppm in accordance with the IPCC IS92a CO_2 emission scenario. Elevated CO_2 750 ppm gas was purchased pre-mixed (BOC) and generously bubbled directly into a corner of the experimental tank near a powerhead to ensure gas and water mixing and dispersal throughout the tank. To check that bubbling was sufficient, tank pH was checked with a Mettler-Toledo SevenGo SG2 pH meter. Ambient air was pumped into the control tank and checked for consistency with a Li-820 gas analyser (Licor). Both tanks were equipped with chillers, filtration units and powerheads to provide adequate temperature control, filtration and




Fig. 1. Location of Mingulay Reef Complex with three-dimensional shaded colour multibeam bathymetry. Sample area has been circled on the multibeam map. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Digital still of live *Lophelia pertusa* at the Mingulay Reef Complex taken during the RRS *James Cook* 073 cruise (May 2012).

circulation. Water circulation in this closed system from the tank to the filtration unit and chiller was 300 l h^{-1} . Thirty per cent water changes were conducted on each tank every 2 days. Tank temperatures and salinity were checked throughout the experimental period with a YSI (30) salinity and temperature meter. Salinity and temperature for the control and experimental tanks were 35.2 ± 0.04 and 35.3 ± 0.03 , and 9.72 ± 0.13 °C and 9.86 ± 0.07 °C respectively. Corals ($n=32$ in each treatment) were fed a mixture of live *Artemia* and *Skeletonema marinoi* every two days. Time points were Time Zero, +7 days, +14 days and +21 days. Following measurements at time zero, 750 ppm gas bubbling was initiated in the treatment tank. Fragments from individual colonies were split evenly between treatments and time points to avoid colony pseudo replication.

2.3. Site and experimental carbonate chemistry

Mingulay carbonate chemistry was assessed on two cruises; on D366/7 (July–August 2011) and on RRS *James Cook* 073 (Roberts et al., 2013) (May–June 2012) in surface waters (< 20 m) and near the reef crest (~ 120 m). Borosilicate glass bottles with ground glass stoppers were used to collect seawater from Niskin bottles on the CTD rosette, and sample bottles were rinsed and filled according to standard procedures detailed in Dickson et al. (2007). Samples were poisoned with mercuric chloride, and duplicate samples were taken from the same Niskin bottle. Samples were bought to room temperature (approx. 23 °C) and analysed for total inorganic carbon and total alkalinity within 24 h of collection. Total Alkalinity (A_T) and Dissolved Inorganic Carbon (C_T) were calculated according to depth-specific salinity, and also normalised to a salinity of 35 for comparative purposes across depths where salinity changed. A_T was corrected for the addition of mercuric chloride. Carbonate parameters were calculated using CO2sys (Pierrot et al., 2006) with dissociation constants from Mehrbach et al. (1973), refit by Dickson and Millero (1987) and KSO₄ using Dickson (1990). A_T and C_T samples were collected (as described above) from the middle of experimental tanks at each time point. Although there was only one experimental tank per treatment, the turnover of seawater between time points (through regular maintenance water changes) ensured that carbonate chemistry was not pseudo-replicated across time points. For full details of instruments used to assess A_T and C_T for experimental purposes, and for site carbonate chemistry, please refer to Supplementary Material.

2.4. Physiology

Rates of oxygen consumption were assessed in coral fragments placed within 220 mL incubation chambers fitted with oxygen optodes connected to a temperature-compensated oxygen analyser (Oxy-4 Mini with Temp-4, Presens & Loligo systems). Magnetic stirrers ensured homogeneity of oxygen around the coral fragments. Chambers were filled with tank seawater and corals were allowed to acclimate to the conditions for 2 h. Ten chambers were used per treatment at each time point; 8 for respiration, and 2 as seawater ‘blanks’ in order to measure (and subsequently subtract) background microbial respiration. Prior to respiration measures, corals were not fed for 48 h. Once chamber lids were attached ensuring no headspace, oxygen consumption was recorded for a 40-min period for each fragment, during which oxygen saturation did not fall below 80%. Following incubations at $T+21$ days, fragments were removed and preserved at -20 °C for subsequent weight determination.

The ash-free dry mass (AFDM) of each sample was determined by adding homogenised material to a pre-weighed porcelain crucible and noting the weight of the crucible and the sample. The crucibles were covered and placed in a muffle furnace (Nabertherm Controller B170). The temperature of the furnace increased to 450 °C over a 30-min period and then remained at that temperature for a further 4 h. After this time the crucibles were re-weighed and the difference in the weight of the sample minus the ashed weight gave the amount of organic matter or AFDM.

2.5. Net calcification

Two methods were used to assess *L. pertusa* growth through net calcification; the alkalinity anomaly technique (Smith and Key, 1975) and uptake of labelled carbon (Marshall and Wright, 1998). To determine if there was any increase or decrease in tissue mass, the AFDM was compared to dry coral weight. For additional comparison to recent CWC studies (Form and Riebesell, 2012; Maier et al., 2012, 2009), coral polyp numbers were also counted to determine calcification per polyp. Different fragments were used for both techniques (Table 1).

2.5.1. Alkalinity anomaly

Following techniques from Smith and Key (1975), and Ohde and Hossain (2004) calcification rates were calculated in *L. pertusa* by measuring the change in seawater alkalinity in respiration chambers housing coral fragments for 4 h. Samples of incubation water were taken at the beginning and end of the experimental period, and total alkalinity determined using an automatic titrator

Table 1

Mean (\pm S.E.) polyp number and dry skeletal weights for fragments used in respiration/alkalinity anomaly technique (AA_T) measurements ($N=8$) and ¹⁴C experiments ($N=5$). Respiration/AA_T fragments were the same used at all time points.

	Respiration/AA _T		¹⁴ C
	Polyp no.	Dry skeletal weight (g)	Dry skeletal weight (g)
Time 0			
380 ppm	N/A	N/A	3.80 (0.42)
750 ppm	N/A	N/A	3.15 (1.02)
Time +21 days			
380 ppm	8.63 (1.22)	9.73 (3.00)	2.90 (0.39)
750 ppm	12.2 (2.37)	11.7 (2.31)	2.11 (0.62)

(Metrohm 702 SM Titrino). Calcification ($\mu\text{mol CaCO}_3 \text{ g}^{-1} \text{ h}^{-1}$) was estimated following Eq. (1),

$$\text{Calcification} = 0.5(\Delta\text{TA}) \cdot V / \Delta T / \text{AFDM} \quad (1)$$

where ΔTA is the change of total alkalinity ($\mu\text{mol/l}$), V is the volume of experimental seawater (L) and ΔT is the experimental period (h). Calcification rates were calculated per hour (and extrapolated to per day) through a linear function. However, this may potentially underestimate growth rates if all newly accreted material contributes to calcification. Further study is needed to calculate whether *L. pertusa* growth over time is better represented by linear or exponential growth functions. With regard to nutrients, corrections applied to total alkalinity to account for the release of nutrients during incubations are considered negligible, especially for tropical corals (Riebesell et al., 2010; Smith, 1995). Considering that the $\sim 10\%$ underestimation of net calcification rate due to nutrient omission is small compared to natural variation in CWC calcification (Maier et al., 2012), nutrients were considered negligible for experiments here. Changes in aragonite, C_T and pH during incubation were not quantified. For comparison to other CWC studies, calcification rates were also expressed as a percentage compared to initial skeletal weight of the corals to give growth as $\% \text{ d}^{-1}$.

2.5.2. Radioisotope

Calcification rates were measured by incorporation of ^{14}C (from sodium bicarbonate, CHNaO_3), into new coral skeleton. Fragments of live corals (different from those used for respiration/alkalinity anomaly) from both treatments were placed in 30 ml filtered seawater (FSW) maintained at tank temperature (9.5°C) in 50 ml falcon tubes (20 ml headspace). After 1 h acclimation, 120 μl of ^{14}C stock solution was added to each tube to a final activity of 3 kBq/ml (0.08101 uCi/ml) (Al-Horani et al., 2005). Two controls were included; one of a dead coral fragment (previously placed in a 4% formalin solution), and one of Filtered Sea Water (FSW). Tubes were kept in floating tube racks, so ship movement and a pump maintained gentle movement of the tubes to ensure isotope mixing. Immediately following addition of the radioisotope, 100 μl of sample water was removed and added to 4 ml of Optisafe scintillation cocktail with 200 μl β -phenylethylamine to assess total activity. Following 6 h incubation, another 100 μl was taken to assess the decrease in total activity in incubation water. Coral fragments were then placed in FSW for 1 h to remove unbound tracer. This step was repeated before corals were preserved at -20°C for analysis following the expedition. No samples of water carbonate chemistry were taken from the experiment tubes before or after incubation.

To determine the amount of ^{14}C incorporated in skeleton, frozen samples were dried overnight at 60°C . Samples were weighed and tissue was then hydrolysed with a known volume of 2 N NaOH at 90°C . Hydrolysate (100 μl) was sampled for uptake of ^{14}C in coral tissue. The skeleton was rinsed with distilled water and dried overnight again for re-weighing. Tissue dry weight was calculated as the difference between skeleton and tissue weight, and skeleton weight. To calculate uptake of ^{14}C , skeletons were placed in flasks fitted with an outlet to a swinnex filter holder which contained a GF/C filter impregnated with β -phenylethylamine. Excess acid (HCl) was added to flasks to dissolve the skeleton and evolve CO_2 . Evolved ^{14}C was trapped in the GF/C filter, and following complete skeletal dissolution, the filter was placed in 4 ml of Optisafe scintillation cocktail for counting on a Packard 1900CA Tri-Carb Liquid Scintillation Analyser.

3. Results

3.1. Mingulay seawater parameters

From CTD casts in June 2012, salinity increased with depth at Mingulay whereas temperature slightly decreased from the surface to the reef crest (ca. 120 m) (Table 2). C_T increased with depth, both when normalised to the changing salinity and when normalised to 35 for comparison with surface values (Table 2). A_T increased with depth when using depth-specific salinity values, but when normalised to 35, a decrease in A_T was observed with depth. *In situ* pH_T decreased with depth as pCO_2 increased (Table 2). Deeper CTD casts taken in July 2011 at the side of the Mingulay Reef mounds correlate with Table 1; such that A_T and C_T increased with depth (A_T 2332.7 ± 1.18 ; DIC 2149.0 ± 2.08 at 172 m).

Table 2

Mean (\pm S.E.) environmental conditions in the surface water (0–20 m, $N=45$) and in the deep water (100–120 m, $N=34$) immediately above the Mingulay reef complex in May 2012. Measured values include salinity, temperature, total alkalinity (TA) and dissolved inorganic carbon (C_T). Also shown are A_T and C_T normalised to a salinity of 35 (nA_T and nC_T , respectively); and calculated values for *in situ* pH on the total scale (pH_T), pCO_2 and saturation states of calcite (Ω_{Calcite}) and aragonite ($\Omega_{\text{Aragonite}}$).

	Surface (0–20 m)	Reef top (100–120 m)
Salinity	34.99 (± 0.003)	35.27 (± 0.005)
Temperature ($^\circ\text{C}$)	9.556 (± 0.020)	9.304 (± 0.003)
A_T ($\mu\text{mol kg}^{-1}$)	2303.4 (± 0.8)	2312.5 (± 0.8)
C_T ($\mu\text{mol kg}^{-1}$)	2079.0 (± 1.0)	2117.4 (± 1.6)
nA_T ($S=35$) ($\mu\text{mol kg}^{-1}$)	2303.8 (± 0.8)	2294.6 (± 0.7)
nC_T ($S=35$) ($\mu\text{mol kg}^{-1}$)	2079.4 (± 1.0)	2101.1 (± 1.4)
pH_T (<i>in situ</i>)	8.140 (± 0.0027)	8.074 (± 0.0036)
pCO_2 (μatm)	309.0 (± 2.2)	365.5 (± 3.4)
Ω_{Calcite}	3.78 (± 0.02)	3.29 (± 0.02)
$\Omega_{\text{Aragonite}}$	2.40 (± 0.01)	2.09 (± 0.01)

Table 3

Carbonate chemistry of experimental tanks at 380 ppm, 9.5°C and 750 ppm, 9.5°C ($N=5$ at each time point from same tank location) \pm SD. Dissolved inorganic carbon (C_T) was measured as described above. Total Alkalinity (A_T), pH on the total scale (pH_T) and aragonite saturation ($\Omega_{\text{Aragonite}}$) were calculated using C_T and CO_2 . 750 ppm gas bubbling was initiated after time zero measurements.

	380 ppm 9.5°C	750 ppm 9.5°C
Time 0		
A_T ($\mu\text{mol kg}^{-1}$)	2307.8 (1.60)	2317.8 (1.81)
C_T ($\mu\text{mol kg}^{-1}$)	2124.7 (1.3)	2131 (35.2)
$\Omega_{\text{Aragonite}}$	2.13 (0.19)	2.06 (0.01)
pH_T (<i>in situ</i>)	8.07 (0.02)	8.07 (0.01)
Time +7 days		
A_T ($\mu\text{mol kg}^{-1}$)	2327.9 (2.59)	2150.6 (2.88)
C_T ($\mu\text{mol kg}^{-1}$)	2140 (2.2)	2078 (2.7)
$\Omega_{\text{Aragonite}}$	2.07 (0.01)	1.05 (0.01)
pH_T (<i>in situ</i>)	8.06 (0.01)	7.77 (0.01)
Time +14 days		
A_T ($\mu\text{mol kg}^{-1}$)	2330.1 (0.91)	2170.7 (2.08)
C_T ($\mu\text{mol kg}^{-1}$)	2144 (1.4)	2098 (1.9)
$\Omega_{\text{Aragonite}}$	2.07 (0.01)	1.06 (0.01)
pH_T (<i>in situ</i>)	8.07 (0.01)	7.77 (0.01)
Time +21 days		
A_T ($\mu\text{mol kg}^{-1}$)	2295.6 (0.26)	2141.6 (1.72)
C_T ($\mu\text{mol kg}^{-1}$)	2112 (0.8)	2071 (1.7)
$\Omega_{\text{Aragonite}}$	2.02 (0.01)	1.03 (0.01)
pH_T (<i>in situ</i>)	8.06 (0.01)	7.77 (0.01)

3.2. Physiology

The respiration rates for coral fragments at Time Zero before fragments were exposed to experimental conditions was $23 \mu\text{mol O}_2 \text{ g}^{-1} \text{ h}^{-1}$. For fragments in the control treatment at 380 ppm, 9.5°C , pH 8.07 (Table 3), there was no significant change in respiration rates from Time Zero to 21 days. For fragments exposed to increased CO_2 at 750 ppm, 9.5°C , pH 7.77 (Table 3), respiration rates were significantly lower than control fragments at 14 days with $12.1 \pm 1.30 \text{ SE}$ at 750 ppm versus $23.7 \pm 5.13 \text{ SE}$ at 380 ppm, and at 21 days with $11.441.39 \text{ SE}$ at 750 ppm versus $28.6 \pm 7.30 \text{ SE}$ at 380 ppm $\mu\text{mol O}_2 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$ (two sample t-test; $t=2.19$, $p < 0.05$; $t=2.31$, $p < 0.05$ respectively), see Fig. 3. Microbial respiration typically ranged from 0.005 to $1.3 \mu\text{mol O}_2 \text{ h}^{-1}$, but on average was $\sim 0.5 \mu\text{mol O}_2 \text{ h}^{-1}$ in the experimental chambers.

There was no difference in the relationship between AFDM and whole coral (skeleton+tissue) dry weight between 380 ppm and 750 ppm treatments after 21 days, so tissue mass did not change




Fig. 3. Respiration rate ($\mu\text{mol O}_2 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$) \pm SE of *L. pertusa* fragments ($N=8$) exposed to control conditions (380 ppm, 9.5°C) and elevated CO_2 conditions (750 ppm, 9.5°C) over 21 days. CO_2 conditions were 380 ppm for both treatments at T0.




Fig. 4. Relationships between coral tissue dry weight and (A) combined skeletal and tissue dry weight and (B) fragment polyp number for fragments from control and experimental treatments. Relationships include combined 380 and 750 ppm corals. Relationships between *L. pertusa* respiration and (C) tissue dry weight and (D) polyp number are from 380 ppm and 750 ppm treatments independently. Additional fragments not used for respiration are used to derive weight relationships in panels (A) and (B).

in coral fragments under acidified conditions compared to control individuals. Fragments from 380 ppm and 750 ppm were thus combined for correlation analysis between AFDM and whole coral dry weight. The linear regression was strong ($r^2=0.85$, Fig. 4A, at $T+21$ days; $y=0.0428x+0.08473$) even with the presence of one anomalous point. If this point was removed from the regression, $y=0.0415x+0.0849$, $r^2=0.97$, $p < 0.0001$, with tissue dry weight approximately 5% of dry coral weight.

Respiration rates for each treatment at $T+21$ days were compared to AFDM, with positive relationships for both 380 ppm ($r^2=0.62$, $p=0.04$) and 750 ppm ($r^2=0.78$, $p=0.008$) fragments (Fig. 4C). However, these regressions were partially driven by one larger coral in each treatment. Comparisons of AFDM and respiration rates to fragment polyp numbers elicited poor relationships; 380 ppm ($r^2=0.14$, $p=0.40$), 750 ppm ($r^2=0.08$, $p=0.59$) (Fig. 4D).

3.3. Net calcification

Net calcification rates at time zero, measured with the alkalinity anomaly technique, were ca. $1.5 \mu\text{mol CaCO}_3 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$. This did not change significantly for either treatment over the 21 days, or between treatments at any time point (Fig. 5A). Changes in A_T (blank corrected) across both treatments were on average $34.7 (\pm 3.34) \mu\text{mol kg}^{-1}$ and ranged from 9 to $59 \mu\text{mol kg}^{-1}$. Growth rates measured as $\% \text{ d}^{-1}$ did not change significantly over the 21 days either, and averaged $\sim 0.028\% \text{ d}^{-1}$ (Fig. 5B). Calcification rate as determined by ^{14}C uptake did not differ between treatment and control fragments at either the experiment start (380 ppm; $19.3 \times 10^{-2} (\pm \text{SE } 2.76)$, 750 ppm; $21.8 \times 10^{-2} (\pm \text{SE } 7.36) \mu\text{mol CaCO}_3 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$) or end (380 ppm; $1.68 \times 10^{-2} (\pm \text{SE } 0.60)$, 750 ppm; $4.38 \times 10^{-2} (\pm \text{SE } 1.6) \mu\text{mol CaCO}_3 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$). However, there was a significant decrease in calcification rates measured by ^{14}C uptake in 380 ppm fragments over time (380 ppm, two sample t-test; $t=7.28$, $p=0.001$), which was not significant in 750 ppm fragments due to very high variability.




Fig. 5. Net calcification rate \pm SE of *L. pertusa* fragments exposed to control conditions (380 ppm, 9 °C) and elevated CO₂ conditions (750 ppm, 9 °C) over 21 days calculated from the alkalinity anomaly technique expressed as (A) $\mu\text{mol CaCO}_3 \text{ g}^{-1} \text{ tissue dry weight h}^{-1}$, and (B) as % d⁻¹.

This decrease was not observed through the alkalinity anomaly technique. ¹⁴C uptake into *L. pertusa* tissue was approximately the same for all treatments and time points (ca. $60 \times 10^{-2} \mu\text{mol } ^{14}\text{C g}^{-1} \text{ tissue dry weight h}^{-1}$) and did not significantly change between treatments or over time.

4. Discussion

4.1. Physiology, calcification and ocean acidification

Lophelia pertusa respiration rate was significantly lower in fragments exposed to increased CO₂ than in control fragments after 2 weeks. Metabolic respiration is needed to produce ATP, which in turn is used to support energy-requiring processes (Al-Horani et al., 2003). This includes calcification, tissue production, mucus-production (Wild et al., 2009) (which accounts for a large proportion of tropical corals' carbon budget (Muscatine et al., 1984)) and driving the transport protein Ca²⁺-ATP-ase. This transport protein is essential for calcification to occur, as it actively pumps Ca²⁺ ions into the sub-calicoblastic space in exchange for H⁺ ions, thereby increasing Ca²⁺ concentration and making conditions favourable for calcification to occur by increasing pH (McCulloch et al., 2012a). The decreased respiration found in the present study under acidified conditions may indicate a decreased requirement of ATP production by *L. pertusa* under increased levels of CO₂ after a few weeks. This may be due to a change in energetic requirements, a reallocation of resources, or changing metabolic pathways (Findlay et al., 2011). Form and Riebesell (2012) noted discrepancies in *L. pertusa* physiology between very short (24 h) and longer-term (months) experiments; whereby responses observed during the first 24 h were no longer exhibited after some months. This indicates that *L. pertusa* may have both 'shock' and acclimation responses to changes in CO₂ conditions. The two-week period observed here before any noticeable differences were observed between experimental and control fragments may indicate a switching of metabolic pathways cued by extended exposure to elevated CO₂.

However, due to coral respiration in closed-system experimental chambers, it is likely that there will have been a significant rise in C_T and corresponding decrease in pH during the respiration incubation periods (Form and Riebesell, 2012). Although unquantified in this experiment, reductions in pH may have matched those reported by Maier et al. (2009) where decreases over 24 h (in smaller vessels) ranged from 0.1–0.5 pH units. For low pH treatments (such as the 750 ppm), reductions could be potentially larger due to the reduced buffering capacity. Although this is important to consider, the fact that respiration was not significantly lower than T₀ in the experimental treatment at T+7 days, but only after 14 days, indicates that it was likely to be a result of experimental conditions rather than changes during incubations. This is supported by observations that there are no significant differences in respiration rates in corals in the first 30 min of respiration incubation compared to after T+4 h in the same incubation water (Hennige, S., personal observation).

Respiration rates for *L. pertusa* documented in this study are higher than those reported by Dodds et al. (2007). However, this experiment was conducted on freshly collected *L. pertusa* fragments, and not on colonies kept in aquaria conditions for several months. Indeed, respiration rates by Dodds et al. (2007) are very comparable to rates observed in *L. pertusa* fragments that have been maintained in aquaria at Heriot-Watt University for more than a year (Hennige, S., personal observation). This is likely due to the feeding limitations of aquaria in comparison to the *in situ* food supply. Freshly collected fragments will be acclimated to the conditions they were removed from *in situ*, but may exhibit an unquantified and unknown element of stress due to collection and pressure differences. However, while long-term aquarium fragments may not have any unknown 'stress' and can be kept at very specific conditions, they may be acclimated to some unrepresentative aquarium conditions. Both approaches are equally valid but it has to be considered that freshly collected corals may not always exhibit similar physiological responses to long-term aquarium corals. Results here were normalised to AFDM in preference to polyp number, as respiration rates of fragments from both treatments significantly correlated with tissue mass, and not polyp number. Although coral polyps are the centres of heterotrophic feeding, dry tissue weight, which includes both polyp tissue and the surrounding coenosarc (which also respire), is therefore a potentially more useful normalisation parameter than polyp number. The relationship between AFDM and dry coral weight means that dry skeletal weight can provide a convenient proxy of tissue weight, without the need to kill the coral.

Net calcification rates as measured by the alkalinity anomaly technique and ¹⁴C uptake, did not differ between control and experimental fragments, even though experimental fragments were at a much lower aragonite saturation level (still > 1). This complements a 24-h study by Maier et al. (2012) and a longer-term experiment by Form and Riebesell (2012), where growth rates of *L. pertusa* did not significantly change under different CO₂ conditions.

The stable net calcification rates reported in this study are in contrast to the reduced respiration in experimental fragments. This suggests an energetic imbalance between the production of ATP, and the use of ATP to actively provide Ca²⁺ needed for calcification. In a recent study, Kaniewska et al. (2012) found that under high CO₂ conditions, metabolic activity in the tropical coral *Acropora millepora* was suppressed after a period of weeks, and changes were observed in genes regulating membrane cytoskeletal interactions and cytoskeletal remodelling. Results by Kaniewska et al. (2012) also indicated a possible breakdown of lipid reserves, which could provide the energy needed to maintain net calcification rates even with suppressed metabolism as observed. This highlights the importance of considering energetic budgets and wider cellular processes in studies such as this

(Findlay et al., 2009; Kaniewska et al. 2012). The consistent AFDM: coral dry weight between fragments exposed here to ambient and 750 ppm conditions did not indicate any major change in tissue mass in coral fragments from either treatment. However, it is probable that an increase in lipid breakdown would not be apparent through changes in tissue mass over this relatively short time period.

Net calcification rates reported here from the alkalinity anomaly technique are comparable to previously published rates by Maier et al. (2012, 2009), and Form and Riebesell (2012). Net calcification rates reported here for *L. pertusa* were slightly lower than those reported by Maier et al. (2009) where ^{45}Ca was used to determine net calcification, within variability of alkalinity anomaly derived data in Maier et al. (2012), and slightly higher than those reported in Form and Riebesell (2012). Net calcification rates in *L. pertusa* are often highly variable, even within studies where fragments are collected from the same locale and time of year. It is perhaps expected then, that differences exist between *L. pertusa* from different reefs, depths, and age. Further differences may be elicited through irregular growth episodes or ‘spurts’ (Form and Riebesell, 2012; Mortensen et al., 2001), which is typical in scleractinian corals.

The calcification rates measured by ^{14}C uptake were significantly lower than rates calculated through the alkalinity anomaly technique. This is not surprising, as labelled bicarbonate is not guaranteed to be taken up solely into the coral skeleton as calcium carbonate, and its uptake can be significantly less than ^{45}Ca (Marshall and Wright, 1998). The reduction in ^{14}C uptake from time zero to the end of the experiment in both control and experimental fragments indicates that some physiological process may have altered during this time. This may indicate a degree of acclimation to tank conditions (changing food source/change in circulation) that may have impacted the control of internal coral chemistry and cellular processes, so less ^{14}C bicarbonate was incorporated into the skeleton. Non-quantified changes in carbonate chemistry during radioisotope experimental incubations may also have impacted upon these calcification rates. Considering the similarity of ^{14}C incubations to the ^{45}Ca incubations by Maier et al. (2009), it is possible that pH may have changed during incubations by up to 0.5 units, even though incubations here were four times shorter. However, considering that calcification rates (as measured through the alkalinity anomaly technique) did not change over the course of the 21 days, it is likely the reduction in ^{14}C uptake from T0 represents experimentally induced and not incubation induced changes.

In general, the impact of ocean acidification upon scleractinian corals; both tropical and cold, seems to be inconsistent, with different species exhibiting negative (Ohde and Hossain, 2004), no measurable response (Reynaud et al., 2003), or variable responses (Gattuso et al., 1998) to a change in conditions (Wicks and Roberts, 2012). This is further complicated by suggestions that corals may be more or less susceptible to ocean acidification depending upon their ontogenetic stage. Albright et al. (2010) demonstrated that the tropical coral *Acropora palmata* is negatively impacted by increasing CO_2 with respect to fertilisation, settlement of larvae, and growth of juveniles. Impacts on these aspects in *L. pertusa*, and indeed any CWC remain unknown to date. Scleractinian responses to ocean acidification may also not be observable in physiology, and may be enacted through changes in biomineralisation (Cohen and Holcomb, 2009; Holcomb et al., 2010). However, when comparing existing ocean acidification studies, care has to be taken, as methodologies often differ with respect to length of exposure to increased CO_2 levels, the speed at which organisms are subjected to change (instantly versus increasing intermediate levels) and the way in which pH is reduced, i.e. acid addition versus CO_2 bubbling (Wicks and Roberts, 2012). A particularly interesting

area for future focus may be whether the internal pH upregulation noted in certain coral species (Anagnostou et al., 2012; McCulloch et al., 2012a; McCulloch et al., 2012b; Venn et al., 2009) is consistent across species, and across different simulated future conditions.

4.2. Environmental conditions at Mingulay

The increases observed in C_T and A_T from the surface to the reef crest were primarily driven by increasing salinity with depth. When normalised to a salinity of 35, A_T decreased from the surface to the reef. Proximity to the reef, where active calcification is occurring, would explain this decrease in alkalinity through calcification (Kleypas and Langdon, 2006), as for every mole of CaCO_3 produced by the coral, total alkalinity of the water decreases by two moles.

Differences observed between normalised C_T indicate that there is either an addition of C_T at the reef, or a drawdown of C_T at the surface. This may be a combination of phytoplankton photosynthesis at the surface (Riebesell, 2004), and respiration from the coral reef, which would act to increase C_T . It has been noted in tropical corals that as $\Omega_{\text{Aragonite}}$ decreases from 3 or 4 (Kleypas and Langdon, 2006) to ca. 2, significant reduction in calcification rates can occur (Wicks and Roberts, 2012). The $\Omega_{\text{Aragonite}}$ at Mingulay (Table 1), which is considered relatively shallow for a *Lophelia pertusa* reef (Roberts et al., 2006), is about half that of many tropical reefs (Kleypas and Langdon, 2006). However, $\Omega_{\text{Aragonite}}$ is still > 1 , so dissolution should not occur.

At Mingulay Reef, corals experience a periodic downwelling of surface water (Davies et al., 2009). Recently characterised in terms of changing seawater carbonate chemistry (Findlay et al., 2013), this surface water will bring with it an influx of food from the surface, as well as periods of warmer water. Thus, corals at Mingulay are exposed to regularly fluctuating water conditions, both in terms of temperature and also water chemistry. This raises the possibility that corals at Mingulay may have higher tolerance for changing seawater conditions than those in more stable, bathyal environments. These and related questions form the basis of on-going investigations.

4.3. Conclusions

From this experiment, we present the first short-term data on the effects of increased CO_2 , (750 ppm) upon the metabolism of freshly collected *L. pertusa* from Mingulay Reef Complex, Scotland, and its comparison with net calcification rates. The sustained net calcification rates of *L. pertusa* under elevated CO_2 conditions corresponds with other studies, but the observed decrease in respiration rate highlights an energetic imbalance, whereby *L. pertusa* may be forced to use energetic reserves to maintain calcification rates. However, the observed decrease in respiration in response to ocean acidification is potentially detrimental in the longer term, as expending energetic reserves is a finite strategy. Thus, it is crucial to perform longer-term experiments on *Lophelia pertusa* metabolism and growth to assess the acclimation potential, and ultimately the success, of this deep-sea ecosystem engineer to predicted increases of CO_2 and warming. Finally it is important to note that the rapid rise in atmospheric CO_2 is not only causing ocean acidification but warming. Further studies examining the combined effects of warming and acidification alongside other predicted stressors are urgently needed if we are to truly appreciate the significance of global climatic change on cold-water corals and other vulnerable marine ecosystems.

Acknowledgments

This paper is a contribution to the UK Ocean Acidification Research Programme (NE/H017305/1 to JMR; funded by the Natural Environment Research Council, the Department for Energy and Climate Change, and the Department for Environment, Food and Rural Affairs) and the European Commission's Seventh Framework Programme projects EPOCA (Grant agreement no. 211384) and HERMIONE (Grant agreement no. 226354). SJH, LCW and JMR acknowledge support from Heriot-Watt University's Climate Change Theme. We thank Juan Moreno-Navas for GIS assistance and multibeam maps, the Scottish Association for Marine Science for the loan of coral sampling equipment, and the captains, crew and scientific participants of RRS *Discovery* cruise 366/7 and RRS *James Cook* cruise 073 for assistance at sea.

Appendix A. Supplementary Information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dsr2.2013.07.005>.

References

- Achterberg, E., Richier, S., 2012. D366/7 Cruise Report, UK Ocean Acidification Research Program. (www.bodc.ac.uk).
- Albright, R., Mason, B., Miller, M., Langdon, C., 2010. Ocean acidification compromises recruitment success of the threatened Caribbean coral *Acropora palmata*. *Proc. Natl. Acad. Sci. USA* 107, 20400–20404.
- Al-Horani, F.A., Al-Moghrabi, S.M., de Beer, D., 2003. The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*. *Mar. Biol.* 142, 419–426.
- Al-Horani, F.A., Ferdelman, T., Al-Moghrabi, S.M., De Beer, D., 2005. Spatial distribution of calcification and photosynthesis in the scleractinian coral *Galaxea fascicularis*. *Coral Reefs* 142, 419–426.
- Allemand, D., Ferrier-Pages, C., Furla, P., Houlbrèque, F., Puverel, S., Reynaud, S., Tambutté, E., Tambutté, S., Zoccola, D., 2004. Biomineralisation in reef-building corals: from molecular mechanisms to environmental control. *C. R. Palevol* 3, 453–467.
- Anagnostou, E., Huang, K.F., You, C.F., Sikes, E.L., Sherrell, R.M., 2012. Evaluation of boron isotope ratio as a pH proxy in the deep sea coral *Desmophyllum dianthus*: evidence of physiological pH adjustment. *Earth Planet. Sc. Lett.* 349–350, 251–260.
- Caldeira, K., Wickett, M.E., 2003. Anthropogenic carbon and ocean pH. *Nature* 425, 365.
- Cohen, A.L., Holcomb, M., 2009. Why corals care about ocean acidification: uncovering the mechanism. *Oceanography* 22, 118–127.
- Cohen, A.L., McConnaughey, T.A., 2003. Geochemical perspective on coral mineralization. In: Dove, P.M., de Yoreo, J.J., Weiner, S. (Eds.), *Biomaterialization, Reviews in Mineralogy and Geochemistry*. Mineralogical Society of America, pp. 151–187.
- Davies, A.J., Duineveld, G., Lavaleye, M., Bergman, M., van Haren, H., Roberts, J.M., 2009. Downwelling and deep-water bottom currents as food supply mechanisms to the cold-water coral *Lophelia pertusa* (Scleractinia) at the Mingulay Reef Complex. *Limnol. Oceanogr.* 54, 620–629.
- Dickson, A.G., Millero, F.J., 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res.* 1 34, 1733–1743.
- Dickson, A.G., 1990. Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K. *Deep Sea Res.* 37, 755–766.
- Dickson, A.G., Sabine, C.L., Christian, J.R., 2007. Guide to Best Practices for Ocean CO₂ Measurements. PICES Special Publication 3, IOCCP Report No. 8, 191 pp.
- Dodds, L.A., Roberts, J.M., Taylor, A.C., Marubini, F., 2007. Metabolic tolerance of the cold-water coral *Lophelia pertusa* (Scleractinia) to temperature and dissolved oxygen change. *J. Exp. Mar. Biol. Ecol.* 349, 205–214.
- Douarin, M., Elliot, M., Noble, S.R., Sinclair, D., Henry, L.-A., Long, D., Moreton, S.G., Roberts, J.M., 2013. Growth of north-east Atlantic cold-water coral reefs and mounds during the Holocene: a high resolution U series and ¹⁴C chronology. *Earth Planet. Sci. Lett.* <http://dx.doi.org/10.1016/j.epsl.2013.05.023>, in press.
- Edmunds, P.J., 2011. Zooplanktivory ameliorates the effects of ocean acidification on the reef coral *Porites* spp. *Limnol. Oceanogr.* 56, 2402–2410.
- Fautin, D.G., Guinotte, J.M., Orr, J.C., 2009. Comparative depth distribution of corallimorpharians and scleractinians (Cnidaria: Anthozoa). *Mar. Ecol. Prog. Ser.* 397, 63–70.
- Findlay, H.S., Kendall, M.A., Spicer, J.L., Widdicombe, S., 2009. Future high CO₂ in the intertidal may compromise adult barnacle *Semibalanus balanoides* survival and embryonic development rate. *Mar. Ecol. Prog. Ser.* 389, 193–202.
- Findlay, H.S., Wood, H.L., Kendall, M.A., Spicer, J.L., Twitchett, R.J., Widdicombe, S., 2011. Comparing the impact of high CO₂ on calcium carbonate structures in different marine organisms. *Mar. Biol. Res.* 7, 565–575.
- Findlay, H.S., Wicks, L., Moreno-Navas, J., Hennige, S.J., Huvenne, V., Woodward, E.M.S., Roberts, J.M., 2013. Tidal down-welling and implications of future ocean acidification and warming on cold-water coral reefs. *Global Change Biol.* 19, 2708–2719. <http://dx.doi.org/10.1111/gcb.12256>.
- Form, A.U., Riebesell, U., 2012. Acclimation to ocean acidification during long-term CO₂ exposure in the cold-water coral *Lophelia pertusa*. *Global Change Biol.* 18, 843–853.
- Gattuso, J.P., Frankignoulle, M., Bourge, I., Romaine, S., Buddemeier, R.W., 1998. Effect of calcium carbonate saturation of seawater on coral calcification. *Global Planet. Change* 18, 37–46.
- Guinotte, J.M., Orr, J.C., Cairns, S., Freiwald, A., 2006. Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Front. Ecol.* 4, 141–146.
- Holcomb, M., McCorkle, D.C., Cohen, A.L., 2010. Long-term effects of nutrient and CO₂ enrichment on the temperate coral *Astrangia poculata* (Ellis and Solander, 1786). *J. Exp. Mar. Biol. Ecol.* 386, 27–33.
- Kaniewska, P., Campbell, P.R., Kline, D.I., Rodriguez-Lanetty, M., Miller, D.J., Dove, S., Hoegh-Guldberg, O., 2012. Major cellular and physiological impacts of ocean acidification on a reef building coral. *PLoS One* 7 (4), e34659. <http://dx.doi.org/10.1371/journal.pone.0034659>.
- Kleypas, J.A., Buddemeier, R.W., Archer, D., Gattuso, J.P., Langdon, C., Opdyke, B.N., 1999. Geochemical consequences of increased atmospheric CO₂ on coral reefs. *Science* 284, 118–120.
- Kleypas, J.A., 2006. Constraints on predicting coral reef response to climate change. In: Aronson, R. (Ed.), *Geological Approaches to Coral Reef Ecology*. Springer-Verlag, New York, pp. 386–424.
- Kleypas, J.A., Langdon, C., 2006. Coral reefs and changing seawater chemistry. In: Phinney, J.T., Hoegh-Guldberg, O., Kleypas, J., Skirving, W., Strong, A. (Eds.), *Coral Reefs and Climate Change: Science and Management*. AGU Monograph Series, Coastal and Estuarine Studies. American Geophysical Union, Washington DC, pp. 73–110.
- Krief, S., Hendy, E.J., Fine, M., Yam, R., Meibom, A., Foster, G.L., Shemesh, A., 2010. Physiological and isotopic responses of scleractinian corals to ocean acidification. *Geochim. Cosmochim. Acta* 74, 4988–5001.
- Langdon, C., Atkinson, M.J., 2005. Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *J. Geophys. Res. Oceans* 110, 1–54.
- Maier, C., Hegeman, J., Weinbauer, M.G., Gattuso, J.P., 2009. Calcification of the cold-water coral *Lophelia pertusa*, under ambient and reduced pH. *Biogeosciences* 6, 1671–1680.
- Maier, C., Watremez, P., Taviani, M., Weinbauer, M.G., Gattuso, J.P., 2012. Calcification rates and the effect of ocean acidification on Mediterranean cold-water corals. *Proc. R. Soc. B* 279, 1716–1723.
- Marubini, F., Ferrier-Pagès, C., Cuif, J.P., 2003. Suppression of skeletal growth in scleractinian corals by decreasing ambient carbonate-ion concentration: a cross-family comparison. *Proc. R. Soc. B* 270, 179–184.
- Marshall, A.T., Wright, A., 1998. Coral calcification: autoradiography of a scleractinian coral *Galaxea fascicularis* after incubation in Ca-45 and C-14. *Coral Reefs* 17, 37–47.
- Mass, T., Drake, J.L., Haramaty, L., Rosenthal, Y., Schofield, O.M.E., Sherrell, R.M., Falkowski, P.G., 2012. Aragonite precipitation by “proto-polyyps” in coral cell cultures. *PLoS One* 7, e35049. <http://dx.doi.org/10.1371/journal.pone.0035049>.
- McCulloch, M., Trotter, J., Montagna, P., Falter, J., Dunbar, R., Freiwald, A., Forsterra, G., Lopez Correa, M., Maier, C., Ruggeberg, A., Taviani, M., 2012a. Resilience of cold-water scleractinian corals to ocean acidification: boron isotopic systematics of pH and saturation state up-regulation. *Geochim. Cosmochim. Acta* 87, 21–34.
- McCulloch, M., Falter, J., Trotter, J., Montagna, P., 2012b. Coral resilience to ocean acidification and global warming through pH up-regulation. *Nat. Clim. Change* 2, 623–627.
- Mehrbach, C., Culbertson, C.H., Hawley, J.E., Pytkowicz, R.M., 1973. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol. Oceanogr.* 18, 897–907.
- Mortensen, P., Hovland, M., Fosså, J., Furevik, D., 2001. Distribution, abundance and size of *Lophelia pertusa* coral reefs in mid-Norway in relation to seabed characteristics. *J. Mar. Biol. Assoc. UK* 81, 581–597.
- Muscantine, L., Falkowski, P.G., Porter, J.W., Dubinsky, Z., 1984. Fate of photosynthetic fixed carbon in light- and shade-adapted colonies of the symbiotic coral *Stylophora pistillata*. *Proc. R. Soc. London B* 222, 181–202.
- Ohde, S., Hossain, M., 2004. Effect of CaCO₃ (aragonite) saturation state of seawater on calcification of *Porites* coral. *Geochim. J.* 38, 613–621.
- Orr, J.C., Fabry, V., Aumont, O., Bopp, L., Doney, S., 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437, 681–686.
- Pierrot, D., Lewis, E., Wallace, D.W.R., 2006. MS Excel Program Developed for CO₂ System Calculations: ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee.
- Reynaud, S., Leclercq, N., Romaine, S., 2003. Interacting effects of CO₂ partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Global Change Biol.* 9, 1660–1668.
- Riebesell, U., 2004. Effects of CO₂ enrichment on marine phytoplankton. *J. Oceanogr.* 60, 719–729.
- Riebesell, U., Fabry, V.J., Hansson, L., Gattuso, J.P., 2010. Guide to Best Practices for Ocean Acidification Research and Data Reporting. Publications Office of the European Union, Luxembourg.

- Ries, J.B., Cohen, A.L., McCorkle, D.C., 2009. Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* 37, 1131–1134.
- Roberts, J.M., Shipboard Party, 2013. RRS James Cook 073 Cruise Report. <www.bodc.ac.uk>.
- Roberts, J.M., Brown, C.J., Long, D., Bates, C.R., 2005. Acoustic mapping using a multibeam echosounder reveals cold-water coral reefs and surrounding habitats. *Coral Reefs* 24, 654–669.
- Roberts, J.M., Wheeler, A.J., Freiwald, A., 2006. Reefs of the deep: the biology and geology of cold-water coral ecosystems. *Science* 312, 543–547.
- Roberts, J.M., Wheeler, A.J., Freiwald, A., Cairns, S.D., 2009. *Cold-water Corals: The Biology and Geology of Deep-sea Coral Habitats*. Cambridge University Press, Cambridge, p. 334.
- Roberts, J.M., Davies, A.J., Henry, L.-A., Duineveld, G.C.A., Lavaleye, M.S.S., Dodds, L.A., Maier, C., van Soest, R.W.M., Bergman, M.I.N., Hühnerbach, V., Huvenne, V., Sinclair, D., Watmough, T., Long, D., Green, S., van Haren, H., 2009. Mingulay reef complex: an interdisciplinary study of cold-water coral habitat, hydrography and biodiversity. *Mar. Ecol. Prog. Ser.* 397, 139–151.
- Smith, S.V., Key, G.S., 1975. Carbon dioxide and metabolism in marine environments. *Limnol. Oceanogr.* 20, 493–495.
- Smith, S.V., 1995. Reflections on the Measurement and Significance of Carbon Metabolism on Coral Reefs. Kansas Geological Survey Open-File Report Series 95, pp. 1–18.
- Thresher, R.E., Tilbrook, B., Fallon, S., Wilson, N.C., Adkins, J., 2011. Effects of chronic low carbonate saturation levels on the distribution, growth and skeletal chemistry of deep-sea corals and other seamount megabenthos. *Mar. Ecol. Prog. Ser.* 442, 87–99.
- Turley, C.M., Roberts, J.M., Guinotte, J.M., 2007. Perspective. Corals in deep-water: will the unseen hand of ocean acidification destroy cold-water ecosystems? *Coral Reefs* 26, 445–448.
- Venn, A., Tambutte, E., Lotto, S., Zoccola, D., Allemand, D., Tambutte, S., 2009. Imaging intracellular pH in a reef coral and symbiotic anemone. *Proc. Natl. Acad. Sci. USA* 106, 16574–16579.
- Wicks, L.C., Roberts, J.M., 2012. Benthic invertebrates in a high CO₂ world. *Oceanography & Marine Biology: An Annual Review*, vol. 50. CRC Press, Boca Raton, Florida, USA, pp. 127–188.
- Wild, C., Wehrmann, L.M., Mayr, C., Schöttner, S.I., Allers, E., Lundälv, T., 2009. Microbial degradation of cold-water coral-derived organic matter: potential implication for organic C cycling in the water column above Tisler Reef. *Aquat. Biol.* 7, 71–80.