



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Biogeochemistry

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa17140>

Paper:

Nejstgaard, J., Tang, K., Steinke, M., Dutz, J., Koski, M., Antajan, E. & Long, J. (2007). Zooplankton grazing on Phaeocystis: a quantitative review and future challenges. *Biogeochemistry*, 83(1-3), 147-172.

<http://dx.doi.org/10.1007/s10533-007-9098-y>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Zooplankton grazing on *Phaeocystis*: a quantitative review and future challenges

Jens C. Nejstgaard · Kam W. Tang ·
Michael Steinke · Jörg Dutz · Marja Koski ·
Elvire Antajan · Jeremy D. Long

Received: 23 May 2006 / Accepted: 16 November 2006 / Published online: 20 April 2007
© Springer Science+Business Media B.V. 2007

Abstract The worldwide colony-forming haptophyte phytoplankton *Phaeocystis* spp. are key organisms in trophic and biogeochemical processes in the ocean. Many organisms from protists to fish ingest cells and/or colonies of *Phaeocystis*. Reports on specific mortality of *Phaeocystis* in natural plankton or mixed prey due to grazing by zooplankton, especially protozooplankton, are still limited. Reported feeding rates vary widely for both crustaceans and protists feeding on even the same *Phaeocystis* types and sizes. Quantitative analysis of available data showed that: (1) laboratory-derived crustacean grazing rates on monocultures of *Phaeocystis* may have been overestimated compared to feeding in natural plankton communities, and should be treated with caution;

(2) formation of colonies by *P. globosa* appeared to reduce predation by small copepods (e.g., *Acartia*, *Pseudocalanus*, *Temora* and *Centropages*), whereas large copepods (e.g., *Calanus* spp.) were able to feed on colonies of *Phaeocystis pouchetii*; (3) physiological differences between different growth states, species, strains, cell types, and laboratory culture versus natural assemblages may explain most of the variations in reported feeding rates; (4) chemical signaling between predator and prey may be a major factor controlling grazing on *Phaeocystis*; (5) it is unclear to what extent different zooplankton, especially protozooplankton, feed on the different life forms of *Phaeocystis* in situ. To better understand the mechanisms controlling zooplankton grazing in situ, future studies

J. C. Nejstgaard (✉)
UNIFOB AS, Department of Biology, University of
Bergen, Bergen High Technology Centre, 5020 Bergen,
Norway
e-mail: jens.nejstgaard@bio.uib.no

K. W. Tang
Virginia Institute of Marine Science, 1208 Grete Road,
Gloucester Point, VA, 23062, USA

M. Steinke
Department of Biological Sciences, University of Essex,
Wivenhoe Park, Colchester, CO4 3SQ, UK

J. Dutz
Institute for Baltic Sea Research, Seestraße 15,
18119 Rostock Warnemunde, Germany

M. Koski
Danish Institute for Fisheries Research, Kavalergården 6,
2920 Charlottenlund, Denmark

E. Antajan
Laboratoire d'Océanographie de Villefranche, Université
Pierre et Marie Curie-Paris6, 06230 Villefranche-sur-Mer,
France

E. Antajan
Laboratoire d'Océanographie de Villefranche, CNRS,
06230 Villefranche-sur-Mer, France

J. D. Long
Northeastern University Marine Science Center,
430 Nahant Road, Nahant, MA, 01908, USA

should aim at quantifying specific feeding rates on different *Phaeocystis* species, strains, cell types, prey sizes and growth states, and account for chemical signaling between the predator and prey. Recently developed molecular tools are promising approaches to achieve this goal in the future.

Keywords Colony formation · DMS · Gut pigment · Molecular methods · Microzooplankton · *Phaeocystis* · *antarctica* · Predator defense

Introduction

The haptophyte *Phaeocystis* is a dominant phytoplankton genus in tropical to polar seas (Baumann et al. 1994). They are key species in marine food webs and biogeochemical cycles, e.g., as major producers of carbon and climatically important sulfide compounds (Liss et al. 1994; Alderkamp et al. this volume; Stefels et al. this volume). Three major species: *P. globosa* Scherffel, *P. pouchetii* (Hariot) Lagerheim and *P. antarctica* Karsten (Medlin and Zingone this volume) exist in two main morphotypes: small single cells and mucilaginous colonies (see Rousseau et al. this volume for details on the different cell and morphotypes).

The success of *Phaeocystis* has been ascribed to a number of factors, including escaping from grazing by its dramatic ability to shift morphotype between solitary flagellates of a few micrometers to large mucous colonies up to several cm in diameter (Weisse et al. 1994; Chen et al. 2002; Schoemann et al. 2005; Veldhuis and Wassmann 2005; Rousseau et al. this volume). For a phytoplankton bloom to form, the sum of growth and accumulation must be larger than the sum of loss due to horizontal and vertical advection, sinking, lysis and predation (e.g., Smayda 1997; Banse 1994). Advection is beyond the control of all phytoplankton, and thus not likely to be a strong selective force for *Phaeocystis* (but see Seuront et al. this volume). Neither does the growth rate of *Phaeocystis* appear to be exceptionally high relative to other bloom-forming phytoplankton, such as diatoms (Hegarty and Villareal 1988). Indeed, in many field and mesocosm studies *Phaeocystis* blooms co-occurred with, or followed the demise of, diatom blooms (Peperzak et al. 1998; Goffart et al. 2000; Rousseau et al. 2002; Tungaraza et al. 2003; Larsen

et al. 2004). Sinking loss of *Phaeocystis* also seems to be small for the large colonies (Reigstad and Wassmann this volume). This could be partly due to the balloon-like characteristics of the colonies: colonial cells are embedded in a thin mucous skin whereas the interior of the colonies is hollow (van Rijssel et al. 1997). Although single cells are susceptible to viral lysis, an intact mucous skin may protect colonial *Phaeocystis* cells from viral lysis and other infections (reviewed by Brussaard et al. this volume). A remaining possible explanation for the success of *Phaeocystis* despite its moderate growth rate is its ability to reduce grazing mortality, which is the focus of this review article.

Grazing on *Phaeocystis* (mainly *P. globosa* and *P. pouchetii*) has been studied since the beginning of the last century (e.g., Lebour 1920, 1922), and a wide range of organisms have been reported to be able to ingest *Phaeocystis* (Table 1). Unfortunately, most of the early studies were based on light microscopy of gut contents of the grazers, and neither feeding rate nor detailed description of the prey species was available (see Notes in Table 1, and further discussion below). It was not until the 1980s that direct quantitative studies on zooplankton feeding on *Phaeocystis* were reported (reviewed in Peperzak 2002; Rousseau et al. 2000; Schoemann et al. 2005; Weisse et al. 1994). Published grazing rates span wide ranges even for the same predators feeding on the same *Phaeocystis* species, morphotype and food concentration. This indicates that the widely accepted view of grazing vulnerability as a function of predator-to-prey size ratio and prey abundance (Frost 1972; Hansen et al. 1994a) for a given predator type (Hansen et al. 1997) may be compounded by other factors controlling grazing on *Phaeocystis*.

Although copepods and other organisms may ingest *Phaeocystis* spp. (Table 1), at least when they are in palatable condition (Estep et al. 1990; Long and Hay 2006), there are also reports of reduced zooplankton feeding and abundance during blooms of *Phaeocystis* (Table 2) and low reproductive output in copepods, even when feeding rates are relatively high (Verity and Smayda 1989; Turner et al. 2002; Klein Breteler and Koski 2003; Long and Hay 2006). Several authors have previously reviewed various negative effects of *Phaeocystis* on different organisms (e.g., Weisse et al. 1994; Turner et al. 2002; Schoemann et al. 2005); and we provide an updated summary in Table 2. However,

Table 1 Organisms recorded to ingest *Phaeocystis* spp.

Grazer	Developmental stage	<i>Phaeocystis</i> species	type/stage	Observation method	References
<i>Dinoflagellates</i>					
<i>Amphidinium</i> sp.	–	sp.	“remains of spores”	in situ gut microscopy	Lebour (1922)
<i>Gymnodinium rhomboides</i> Schütt	–	sp.	–	in situ gut microscopy	Lebour (1922)
<i>Gymnodinium</i> sp.	–	<i>globosa</i>	single cells	incubation experiments	Tang et al. (2001)
<i>Gymnodinium triangularis</i> Lebour	–	sp.	–	in situ gut microscopy	Lebour (1922)
<i>Gyrodinium dominans</i> Hulbert	–	<i>globosa</i>	single cells	incubation experiments	Tang et al. (2001)
<i>Gyrodinium</i> cf. <i>spirale</i>	–	<i>globosa</i>	small colony	photomicrograph	Stelfox-Widdicombe et al. (2004)
<i>Noctiluca scintillans</i> (Macartney)	–	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995), Jakobsen and Tang (2002), Weisse et al. (1994)
Kofoed & Swezy					
<i>Oxyrrhis marina</i> Dujardin	–	cf. <i>globosa</i>	single cells	incubation experiments	Hansen et al. (1993)
<i>Ciliates</i>					
<i>Euplotes</i> sp.	–	<i>antarctica</i>	single cells and colonies	fluorescently labeled algae	Shields and Smith (2005)
<i>Euplotes</i> sp.	–	<i>globosa</i>	single cells and colonies	incubation experiments	Long (2004)
<i>Helicostomella subulata</i> (Ehrenberg)	–	<i>globosa</i>	single cells	live microscopy	Admiraal and Venekamp (1986)
<i>Lohmanniella oviformis</i> Leegaard	–	<i>globosa</i>	single cells	incubation experiments	Tang et al. (2001)
<i>Rinosrombidium conicum</i> (Kahl)	–	<i>globosa</i>	single cells	incubation experiments	Tang et al. (2001)
Petz & Foissner					
<i>Strombidinopsis acuminatum</i>	–	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1995, 1993)
Fauré-Fremiet					
<i>Strombidium</i> sp.	–	<i>globosa</i>	single cells	incubation experiments	Verity (2000)
<i>Strombidium vestitum</i> Leegaard	–	<i>globosa</i>	single cells	incubation experiments	Tang et al. (2001)
<i>Tintinnopsis beroidea</i> Stein	–	<i>globosa</i>	single cells	live microscopy	Admiraal and Venekamp (1986)
<i>Rotifers</i>					
<i>Synchaeta vorax</i> Rousselet	–	<i>globosa</i>	eating at colonies	live microscopy	Hollowday (1949)
<i>Polychaets</i>					
<i>Polydora pulchra</i> Carazzi	nectochaeta larvae	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Tomopteris</i> sp.	Larvae	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Molluscs</i>					
<i>Macoma balthica</i> (L.)	Mussels??	sp.	ingest single cells	laboratory incubation	Kamerians (1994)
<i>Mytilus edulis</i> L.	Mussels 33–58 mm	cf. <i>globosa</i>	cultured, ingested single cells	flow chamber, cell counts	Smaal and Twisk (1997)

Table 1 continued

Grazer	Developmental stage	<i>Phaeocystis</i> species	type/stage	Observation method	References
<i>Mytilus edulis</i> L.	Mussels > 3 cm	<i>globosa</i>	able to filter colonies	?	Petri and Vareschi (1997)
<i>Nassarius reticulatus</i> (L)	veliger larvae	<i>pouchetii</i>	cultured, probably single cells	incubation, visual inspection	Fretter and Montgomery (1968)
<i>Rissoa inconspicua</i> Alder	veliger larvae	<i>pouchetii</i>	cultured, probably single cells	incubation, visual inspection	Fretter and Montgomery (1968)
<i>Cladocera</i>					
<i>Evadne nordmanni</i> Lovén	–	sp.	cells and “spores”	in situ gut microscopy	Lebour (1922)
<i>Copepods, calanoids</i>					
<i>Acartia clausi</i> (Gunn.)	copepodites ?	sp.	“green remains, probably <i>Phaeocystis</i> in many”	in situ gut microscopy	Lebour (1922)
<i>Acartia hudsonica</i> Pinhey	CVI female	<i>pouchetii</i>	single cells and colonies	incubation experiments	Verity and Smayda (1989)
<i>Acartia longiremis</i> (Lilljeborg)	CVI female	<i>pouchetii</i>	ingested only colonies < 100 µm	incubation experiments	Hansen et al. (1994b)
<i>Acartia tonsa</i> Dana	CVI female	<i>globosa, pouchetii</i>	single cells and colonies	incubation experiments	Tang et al. (2001), Tang and Simó (2003), Verity (2000), Verity and Smayda (1989)
<i>Anomalocera pattersoni</i> Tempelton	copepodites ?	sp.	“ <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Calanoides</i> sp.	Copepodites	<i>Antarctica</i>	fluorescently labeled single cells	gut microscopy	K. Tang unpublished
<i>Calanus finmarchicus</i> (Gunn.)	copepodites ?	sp.	cells and “remains”	in situ gut microscopy	Lebour (1922)
<i>Calanus finmarchicus</i> (Gunn.)	CI-V	<i>pouchetii</i>	single cells and colonies	incubation experiments	Hansen et al. (1990), Tande and Båmstedt (1987)
<i>Calanus glacialis</i> Jaschnov	CIII-CV	<i>pouchetii</i>	colonies	incubation experiments	Estep et al. (1990)
<i>Calanus helgolandicus</i> (Claus)	copepodites > 280 µm	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995), Turner et al. (2002)
<i>Calanus hyperboreus</i> (Krøyer)	CIIV-CVI female	<i>pouchetii</i>	single cells and colonies	incubation experiments	Estep et al. (1990), Huntley et al. (1987), Tande and Båmstedt (1987)
<i>Centropages hamatus</i> Lilljeborg	copepodites > 280 µm	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1995)
<i>Centropages hamatus</i> Lilljeborg	copepodites ?	cf. <i>globosa</i>	colonies? “ <i>Phaeocystis</i> bladders”	in situ gut microscopy	Jones and Haq (1963)
<i>Centropages typicus</i> Krøyer	copepodites ?	sp.	“spores”	in situ gut microscopy	Lebour (1922)
<i>Eucalanus pileatus</i> Giesbrecht	CVI female	<i>globosa</i>	single cells and colonies	incubation experiments	Long and Hay (2006)
<i>Metridia gerlachi</i> Giesbrecht	copepodites ?	<i>pouchetii</i> (in Antarctica)	colonies 0.05–1.5 mm	incubation experiments	Schmack (1983)

Table 1 continued

Grazer	Developmental stage	<i>Phaeocystis</i> species	type/stage	Observation method	References
<i>Microcalanus</i> sp.	Copepodites	<i>antarctica</i>	fluorescently labeled single cells	gut microscopy	K. Tang unpublished
<i>Pareuchaeta antarctica</i> (Giesbrecht)	copepodites ?	<i>pouchetii</i> (in Antarctica)	colonies 0.05–1.5 mm	incubation experiments	Schmack (1983)
<i>Pseudocalanus elongatus</i> (Boeck)	copepodites > 280 μ m	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Pseudodiaptomus pelagicus</i> Herrick	CVI female	<i>globosa</i>	single cells and colonies	incubation experiments	Long and Hay (2006)
<i>Temora longicornis</i> Müller	nauplii to adults	<i>globosa</i>	single cells and colonies	incubation experiments	Dutz and Koski (2006), Koski et al. (2005), Hansen (1992, 1995)
<i>Temora longicornis</i> Müller	copepodites ?	cf. <i>globosa</i>	colonies? “ <i>Phaeocystis</i> bladders”	in situ gut microscopy	Jones and Haq (1963)
<i>Temora longicornis</i> Müller	copepodites ?	sp.	“green cells, probably <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Temora stylifera</i> Dana	adult female	appeared similar to <i>globosa</i>	cultured single cells and colonies	incubation experiments	Turner et al. (2002)
<i>Copepods, cyclopoids</i>					
<i>Corycaeus anglicus</i> Lubbock	copepodites ?	sp.	“ <i>Phaeocystis</i> in several”	in situ gut microscopy	Lebour (1922)
<i>Oithona nana</i> Giesbrecht	copepodites ?	cf. <i>globosa</i>	colonies? “ <i>Phaeocystis</i> bladders”	in situ gut microscopy	Jones and Haq (1963)
<i>Oithona</i> sp.	copepodites	<i>antarctica</i>	fluorescently labeled single cells	gut microscopy	K. Tang unpublished
<i>Oncaea curvata</i> (Gunn.)	adult female	cf. <i>pouchetii</i> (in Antarctica)	colonies	incubation experiments	Metz (1998)
<i>Copepods, harpacticoids</i>					
<i>Euterpina acutifrons</i> (Dana)	copepodites ?	sp.	“flagellates...especially <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Cirripedes</i>					
<i>Balanus crenatus</i> Bruguière	nauplii	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
“Cirripedes”	nauplii	sp.	“probably <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Longipeda minor</i> T. & A. Scott	adult female	sp.	colonies	incubation, visual inspection	Nicholls (1935)
<i>Longipeda scotti</i> G.O.Sars	adult female	sp.	colonies	incubation, visual inspection	Nicholls (1935)

Table 1 continued

Grazer	Developmental stage	<i>Phaeocystis</i> species	type/stage	Observation method	References
<i>Amphipods</i>					
<i>Pontogeneia antarctica</i> Chevreux	hatchlings (2 mm)- adults	<i>antarctica</i>	“green plastic debris... probably indicate ingestion of <i>Phaeocystis antarctica</i> ”	in situ gut microscopy	Richardson and Whitaker (1979)
<i>Euphausiids</i>					
<i>Euphausia superba</i> Dana	35–45 mm	sp.	single cells and colonies < 500 µm	in situ gut microscopy	Sieburth (1960)
<i>Euphausia superba</i> Dana	carapace ca 5 mm	<i>pouchetii</i>	ingested only colonies > 20 µm	incubation experiments	Haberman et al. (2003)
<i>Thysanoessa inermis</i> (Krøyer)	carapace ca 5 mm	<i>pouchetii</i>	ingested only colonies > 20 µm	incubation experiments	Hansen et al. (1994b)
<i>Thysanoessa raschii</i> (M. Sars)	carapace ca 5 mm	<i>pouchetii</i>	ingested only colonies > 20 µm	incubation experiments	Hansen et al. (1994b)
<i>Decapods</i>					
Crab (unidentified)	zoëa	sp.	“ <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Galathea intermedia</i> Lilljeborg	megalopa	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Porcellana</i> sp.	“larva”	sp.	“green spores, probably <i>Phaeocystis</i> ”	in situ gut microscopy	Lebour (1922)
<i>Portunus depurator</i> (Linnaeus)	zoëa III-V	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Portunus holsatus</i> (Fabricius)	megalopa	cf. <i>globosa</i>	single cells and colonies	incubation experiments	Hansen (1992, 1995)
<i>Appendicularians</i>					
<i>Megalocercus huxleyi</i> (Ritter)	-	sp.	small colony	in situ faecal pellet microscopy	Gorsky et al. (1999)
<i>Fish</i>					
<i>Gadus morhua</i> L.	larvae < 0.2 mg/ 4 mm	<i>pouchetii</i>	colonies	in situ gut microscopy	Krogstad (1989), Løken (1990)
<i>Pleuronectes flesus</i> L.	larvae < 11 mm	sp.	colonies	incubation, visual inspection and in situ gut microscopy	Lebour (1920)
<i>Scomber scombrus</i> L.	large from fish catches	<i>globosa</i>	“ <i>phytoplankton chiefly P. globosa</i> ” associated with jelly probably colonies	in situ gut microscopy	Bullen (1908)

(–) denotes no information/not applicable. (?) denotes likely but unclear. Note that species composition within the genus *Phaeocystis* has been debated (Baumann et al. 1994; Sournia 1988). Because the solitary cells of *P. globosa*, *P. pouchetii* and others may not be distinguishable by light microscopy, reports before 1980 could be misleading [e.g., the species in Antarctica reported by Schmack (1983) and Metz (1998) may likely be *P. antarctica*, rather than *P. pouchetii*]. Where multiple references are available (mostly for calanoid copepods) only selected representative references are shown here. See text for more references.

some of these data were derived from ecologically unrealistic organisms or assays and should be interpreted with caution (see discussion below). Further, positive correlations between *Phaeocystis* abundance and mesozooplankton in the field have also been reported, and some of the negative effects have been debated and ascribed to other organisms (see Notes in Table 2). Furthermore, results from fatty acid analysis indicate that dominating crustacean zooplankton may derive a major part of their diet from *P. pouchetii* in northern latitudes (Sargent et al. 1985) and from *P. globosa* in lower latitudes (Hamm and Rousseau 2003), suggesting that *Phaeocystis* do enter the food web.

A number of factors have been shown to control predation on this enigmatic genus: predator-to-prey size ratio, predator species and stage, prey species, morphotype, growth state and abundance (reviewed in Weisse et al. 1994; Rousseau et al. 2000; Peperzak 2002; Schoemann et al. 2005). However, to our knowledge, a quantitative analysis of the interactive effect of these factors has not been attempted. Using available literature, we attempt to quantify the relative importance and interactions among some of these factors in controlling zooplankton grazing on *Phaeocystis* spp. We also review recent findings on various unique aspects of the life history and physiology of *Phaeocystis* spp., and how they affect zooplankton grazing. Last, we discuss the need for new methodological approaches to address some of the remaining questions concerning zooplankton grazing on *Phaeocystis*.

Grazing on *Phaeocystis*: quantitative patterns in published data

Reports on copepods and other crustaceans dominate available zooplankton grazing data on *Phaeocystis* spp. These data show a wide range of rates, even for similar predator-prey combinations. We therefore attempt to summarize available quantitative data in relation to some of the mechanisms proposed to control the feeding on *Phaeocystis*. The shortage of quantitative feeding studies on microzooplankton and *Phaeocystis* was pointed out already a decade ago (Weisse et al. 1994), and the number of such studies is still limited, especially for protozoan microzooplankton, whether using laboratory cultures (Table 3)

or natural plankton assemblages (Table 4) as food. Due to the shortage of protozooplankton data a general statistical analysis of the factors controlling their grazing on *Phaeocystis* cannot yet be performed, but the available data will be discussed further below.

Crustacean zooplankton grazing on *Phaeocystis*: is there a general pattern?

It is generally assumed that a shift from single cell to colony may be part of *Phaeocystis* defense against grazing by microzooplankton (Weisse et al. 1994; Rousseau et al. 2000; Peperzak 2002; Schoemann et al. 2005). Because of the ability of *Phaeocystis* to vary its functional prey size by forming colonies, it is often considered as dual species (sensu Turner et al. 2002): colonies can be referred to as macroplankton and are assumed to be suitable food for zooplankton and nekton, while the solitary nanoplankton are assumed to be vigorously grazed by microzooplankton (Lebour 1922; Hollowday 1949; Admiraal and Venekamp 1986; Weisse and Scheffel-Möser 1990). On the other hand an increase in size may result in a more suitable prey size range for larger predators such as crustacean zooplankton. For palatable prey, ingestion rates generally increase from small prey to the optimum prey size before the prey becomes too large to handle or ingest (Frost 1972; Hansen et al. 1994a). Thus, size may be a major factor controlling predation on *Phaeocystis*.

Many studies showed that the size of *Phaeocystis* strongly influences the feeding rates of differently sized metazoan predators. Hansen et al. (1990) reported that *Calanus finmarchicus* copepodites (CIV–V) showed lower feeding rate on *P. pouchetii* colonies of >100 µm than similarly sized diatoms, while in the 30–100 µm size range feeding rates on diatoms and colonies were similar. Tande and Båmstedt (1987) showed that copepodite stage V of *C. finmarchicus* fed equally well on *P. pouchetii* single cells and the diatom *Chaetoceros furcellatus*, whereas Huntley et al. (1987), using the larger copepod *C. hyperboreus* CIV–V, reported much higher grazing rates on colonies 25–200 µm and >200 µm than on single cells of *P. pouchetii* or the diatom *Chaetoceros socialis*. In a study with grazers of different sizes feeding on *P. pouchetii*, Hansen et al. (1994b) observed an optimal predator-to-prey size ratio for feeding and an upper size ratio above which no ingestion

Table 2 Negative effects on various organisms ascribed to *Phaeocystis* spp., with notes on contradictory reports

Effect	Note	References
<i>Protoists and other microbes</i>		
Defence against viral attacks by intact colony membranes		Reviewed by Brussaard et al. (this volume)
Antibiotic to bacteria in the acid environment of penguin guts.	A	Sieburth (1960, 1961)
Haemolytic activity in laboratory, or correlated with abundance of <i>P. pouchetii</i> during blooms of natural planktonic mesocosms		Stabell et al. (1999), van Rijssel et al. (this volume)
Suggested allelopathy, e.g., possible negative interactions by <i>P. pouchetii</i> on <i>T. nordenskiöldii</i> and <i>Skeletonema costatum</i> in vitro. Rapid decrease in abundance of <i>Thalassiosira nordenskiöldii</i> and <i>Ebria</i> sp. during a bloom of <i>P. pouchetii</i> .		Barnard et al. (1984), Smayda (1973)
Allelopathy, filtrates from mesocosm blooms of <i>P. pouchetii</i> and laboratory cultures of <i>P. globosa</i> resulting in reduced growth and lysis of the cryptophyte phytoplankton <i>Rhodomonas baltica</i> .		Long (Unpublished data)
Lack of feeding of the ciliates <i>Mesodinium pulex</i> and <i>Strombidium elegans</i> in laboratory experiments despite high concentrations of <i>P. globosa</i> and no alternative food.		Hansen et al. (1993), Tang et al. (2001)
Low biomass of bacteria and protozooplankton during blooms of colonial <i>Phaeocystis</i>	B	van Boekel et al. (1992)
<i>Metazooplankton</i>		
Many copepods show low abundances, development, gut content and/or much lower feeding activity compared to alternative food such as diatoms, during blooms of <i>Phaeocystis</i>	C	Bautista et al. (1992), Breton et al. (1999), Daro (1985), Davies et al. (1992), Frangoulis et al. (2001), Gasparini et al. (2000), Hansen and van Boekel (1991), Turner (1994), Weisse et al. (1994)
Low feeding by <i>Calanus</i> spp. and <i>Metridia longa</i> on actively growing colonies of <i>P. pouchetii</i>	D	Estep et al. (1990)
Low feeding by nauplii on some cell types of <i>P. globosa</i>		Dutz and Koski (2006)
Low reproductive output from <i>Acartia</i> spp., <i>Temora longicornis</i> , <i>Pseudodiaptomus pelagicus</i> and <i>Eucalanus pileatus</i> feeding on <i>Phaeocystis</i> , despite sometimes high ingestion rates.	E	Klein Breteler and Koski (2003), Long and Hay (2006), Tang et al. (2001), Turner et al. (2002), Verity and Smayda (1989)
Inhibiting copepod feeding by transparent exopolymer particles (TEP) derived from a <i>Phaeocystis globosa</i> .		Dutz et al. (2005)
Suggested reduction of zooplankton feeding due to mucus secretion from colonies of <i>P. globosa</i> .		Seuront et al. (this volume)
Not a good food organism for the barnacle <i>Balanus balanoides</i>		Cook and Gabbott (1972)
<i>Other organisms</i>		
Oxygen deficiency due to sedimenting <i>Phaeocystis</i> killing less mobile fauna in the sediments.		Rogers and Lockwood (1990)
Colonies clogging gills and thus reducing feeding in several mussels, especially smaller, e.g., <i>P. globosa</i> colonies reduced feeding in <i>Mytilus edulis</i> <3 cm, while larger where able to filter colonies.		Blauw (this volume), Kopp (1978), Meixner (1981), Petri et al. (1999), Pieters et al. (1980), Smaal and Twisk (1997)
Inhibited growth in the bryozoan <i>Electra pilosa</i> possibly due to <i>Phaeocystis</i> .		Jebram (1980)
Toxic to sea urchin larvae		Hansen et al. (2003, 2004)
Toxic to blowflies (<i>Calliphora omissa</i>)		Stabell et al. (1999)

Table 2 continued

Effect	Note	References
Migrating herring hindered by a “barrier” of <i>Phaeocystis</i> blooms	F	Savage (1930, 1932)
Toxic to cod larvae		Aanesen et al. (1998), Eilertsen and Raa (1995), Hansen et al. (2004)
Skin and eye irritation of humans		Dunne et al. (1984)

Notes:

A: It has also been reported that the high concentrations of acrylate in *Phaeocystis* colonies do not inhibit surrounding bacteria and may be absorbed to the mucus matrix (Noordkamp et al. 2000)

B: But when solitary cells are abundant during blooms (Admiraal and Venekamp 1986; Weisse and Scheffel-Möser 1990), or during the breakdown (van Boekel et al. 1992), microzooplankton such as ciliates may be numerous

C: There are also some reports on apparently increased abundances of mesozooplankton during periods of blooms of *Phaeocystis* (Fransz and Gieskes 1984; Fransz et al. 1992; Weisse et al. 1986)

D: But in a later bloom stage the copepods switch to selectively feed upon the senescent *P. pouchetii*, rather than the diatoms (Estep et al. 1990)

Table 3 Quantitative laboratory studies on protozooplankton grazing on *Phaeocystis*

Grazer	Alternative prey	Feeding on <i>P. globosa</i>	References
<i>Oxyrrhis marina</i>	No	Yes	Hansen et al. (1993)
<i>Strombidinopsis acuminatum</i>	No	Yes	Hansen et al. (1993)
<i>Strombidium elegans</i>	No	No	Hansen et al. (1993)
<i>Strombidium</i> sp.	No	Yes	Verity (2000)
<i>Noctiluca scintillans</i>	No	Yes	Jakobsen and Tang (2002)
<i>Gyrodinium dominans</i>	No	Yes	Tang and Simó (2003)
<i>Lohmanniella oviformis</i>	<i>Isochrysis galbana</i>	Lower	Tang et al. (2001)
<i>Rimostrombidium conicum</i>	<i>Isochrysis galbana</i>	Higher	Tang et al. (2001)
<i>Strombidium vestitum</i>	<i>Isochrysis galbana</i>	Higher	Tang et al. (2001)
<i>Gyrodinium dominans</i>	<i>Isochrysis galbana</i>	Higher	Tang et al. (2001)
<i>Gymnodinium</i> sp.	<i>Isochrysis galbana</i>	Lower	Tang et al. (2001)
<i>Mesodinium pulex</i>	<i>Gymnodinium</i> sp.	No	Tang et al. (2001)

All experiments are based on particle disappearance and cell counts of cultured *P. globosa*, or cf. *globosa* in Hansen et al. (1993). Notes are given on whether the grazer feed (yes) or not at all (no) on *Phaeocystis*, or showed higher or lower feeding rates on *Phaeocystis*, compared to alternative prey offered in parallel single prey species incubations

occurs. These results were also supported by Levinson et al. (2000), who reported ingestion of single *P. pouchetii* by the smaller females of *C. finmarchus* but not by the larger *C. glacialis* and *C. hyperboreus*. Also, the krill *Euphausia superba* (35–40 mm) showed similar feeding on 50–100 µm *P. antarctica* colonies compared to the diatom *Thalassiosira antarctica*, whereas both single cells and larger colonies of *P. antarctica* were grazed at lower rates compared to the diatom (Haberman et al. 2003). Thus, data such as these support the notion that substantial shifts in size between *Phaeocystis* life forms affect grazing by

differently sized predators. However, there is still a considerable variation in feeding rates on various forms of *Phaeocystis*, ranging from zero to expected maximum rates (sensu Hansen et al. 1997) for a number of predators (reviewed in Weisse et al. 1994; Schoemann et al. 2005).

In order to evaluate the relative importance of factors other than size and how they may help explain the observed variations in grazing rates, we compiled the available quantitative data, converted them into carbon units, and performed a series of correspondence and cluster analyses described below.

Table 4 Literature inferring quantitative evidence of microzooplankton community grazing (g) and grazing impact (percentage of average standing stock (SS) removed d^{-1}) on *Phaeocystis* solitary cells in natural plankton

<i>Phaeocystis</i> spp.	Area	Time	Analysis	μ (d^{-1})	g (d^{-1})	Impact (% SS d^{-1})	Note	References
<i>MESOCOSM (natural plankton indoor tank)</i>								
<i>globosa</i>	South North Sea	Sep	Cell Counts	? - 0.3 - ?	0.3-0.7	26-48		Brussard et al. (2005)
<i>FIELD</i>								
<i>cf. globosa?</i>	South North Sea	Apr-May	Cell Counts	0.8-2.3	0.9-4.2	59-98	A	Weisse and Scheffel-Möser (1990)
<i>cf. globosa?</i>	South North Sea	Feb-Jul	Total Chl <i>a</i>	-0.3-0.4	0.0-0.6	0-46	B	Brussard et al. (1995)
<i>globosa</i>	South North Sea	April	Total Chl <i>a</i>	0.1-0.7	0.3-1.1	24-68		Stelfox-Widdicombe et al. (2004)
<i>pouchetti</i>	North Norway fjord	April	Total Chl <i>a</i>	0.3-0.5	0.21-0.36	19-30	C	Archer et al. (2000)
<i>pouchetti</i>	Labrador Sea	May-Jun	Total Chl <i>a</i>	0.2-0.9	0.0-0.3	0-27	D	Wolfe et al. (2000)
<i>antarctica</i>	Ross Sea	Oct-Apr	Chl <i>a</i> HPLC FC	?	0-0.23	0-21	E	Caron et al. (2000)
<i>antarctica</i>	Antarctica	Nov-Mar	Chl <i>a</i> HPLC FC	0.1-0.6	0.1-0.3	14-26	E	Landry et al. (2002)
<i>antarctica</i>	Antarctica	Dec	Chl <i>a</i> HPLC FC	0.2-0.9	0.1-1.0	10-63	E	Selph et al. (2001)

Specific phytoplankton growth rate (μ) is also shown. All rates are quantified by the dilution technique (Landry and Hassett 1982). Prey concentrations were analyzed by cell counts, total chlorophyll *a* fractions (Total Chl *a*), high-performance liquid chromatography (HPLC), or flow cytometry (FC)

A: Incubation time only 5-14 h during the day

B: Instantaneous grazing (g) max $\leq 0.1 d^{-1}$ when the phytoplankton was dominated by *Phaeocystis*

C: Additional experiment with FLA (*Chlorella*). Only 5 of 13 tested ciliates ingested the labeled phytoplankton

D: Also measured dimethyl sulphide (DMS) dynamics. During *P. pouchetti* bloom g was 0.0 and 0.3

E: No notes on specific *Phaeocystis* rates

Treatment of literature data

Many reports of crustacean grazing on *Phaeocystis* spp. are based on non-prey-specific gut fluorescence during blooms of *Phaeocystis*. An unknown fraction of such fluorescence may stem from prey other than *Phaeocystis*. Indeed, the often co-occurring diatoms may even be a preferred prey (Gasparini et al. 2000). In addition, more than half the total biomass (and chlorophyll) of the *Phaeocystis* colony may be due to diatoms and other organisms living in and on the colonies during late bloom stages (Sazhin et al. this volume). Therefore, we exclude data based on non-specific gut pigments of grazers feeding on mixed or natural food suspensions. We converted all grazing rate measurements into common currency of carbon units, using either (in the preferred order): (1) directly reported data, (2) data reported in accompanying papers, (3) personal communications with the authors, or (4) common conversion factors (such as those used in Hansen et al. 1997). For quantitative and statistical analyses we included only grazing data from studies that reported experiment location (field or laboratory study), grazer species, stage or size, and *Phaeocystis* average size (or size range). To be included in the statistical analysis the sources also must have reported *Phaeocystis* species, abundance, morphotype and growth state (exponential or stationary—either given in the text or assumed from the state of the bloom). See Table 5 for further details on the variables. The sources used are listed in Table 6.

Statistical methods

To investigate relationships between crustacean grazing rates on *Phaeocystis* and experimental conditions, a multiple correspondence analysis (MCA) followed by a hierarchical cluster analysis (HCA) was performed using SPAD 3.5 software (Lebart et al. 1988). The combination of MCA and cluster analysis is a common way to explore relationships among a large number of variables and to facilitate interpretation of the correspondence analysis results (Lebart et al. 2000). MCA uses a contingency table as data, which provides a simultaneous representation of the observations (rows) and variables (column) in a factorial space. This form of multivariate analysis describes the total inertia (or variability) of a multidimensional

Table 5 Active structural variables used in the multiple correspondence analysis (MCA). *n* denotes the number of observations from each source

Description of variables: modalities	Label	<i>n</i>
<i>Experiment location</i>		
Field	Field	91
Laboratory	Laboratory	205
<i>Crustacean predators</i>		
<i>Acartia clausi</i>	Acl	13
<i>Acartia hudsonica</i>	Ahu	5
<i>Acartia tonsa</i>	Ato	12
<i>Pseudocalanus elongatus</i>	Pel	13
<i>Temora longicornis</i>	Tlo	35
<i>Temora stylifera</i>	Tst	15
<i>Metridia longa</i>	Mlo	1
<i>Centropages hamatus</i>	Cha	14
<i>Calanus finmarchicus</i>	Cfi	105
<i>Calanus glacialis</i>	Cgl	8
<i>Calanus hyperboreus</i>	Chy	75
<i>Phaeocystis species</i>		
<i>globosa</i>	<i>P. globosa</i>	107
<i>pouchetii</i>	<i>P. pouchetii</i>	189
<i>Phaeocystis growth</i>		
Exponential	Exponential	277
Stationary	Stationary	19
<i>Phaeocystis abundance</i> (Ab, $\mu\text{g C l}^{-1}$)		
0.1–125	Ab1	81
125–250	Ab2	91
250–500	Ab3	37
500–1000	Ab4	71
1000–1810	Ab5	16
<i>C-specific ingestion</i> (CI, d^{-1})		
0–0.01	CI1	83
0.01–0.1	CI2	151
0.1–1.5	CI3	62
<i>Phaeocystis selection</i>		
Positive selection	For	14
Negative selection	Against	76
Null selection	Null	5
Single prey	Single	201
<i>Predator-to-prey size ratio</i> (<i>P:p</i>)		
0.8–4	P:p1	57
4–16	P:p2	90
16–64	P:p3	49
64–256	P:p4	25
256–1120	P:p5	75

Table 6 Sources of data used for the multiple correspondence analysis (MCA) and hierarchical cluster analysis (HCA) presented in Fig. 1, and quantitative summary in Fig. 2, respectively

References	Location	<i>Phaeocystis</i>	Fig. 1 <i>n</i>	Fig. 2 <i>n</i>
Antajan unpublished	Field	<i>globosa</i>	6	6
Dutz and Koski (2006)	Lab	<i>globosa</i>	7	7
Estep et al. (1990)	Field	<i>pouchetii</i>	25	25
Gasparini et al. (2000)	Field	<i>globosa</i>	55	55
Haberman et al. (2003)	Lab	<i>antarctica</i>		3
Hansen et al. (1990)	Lab	<i>pouchetii</i>	62	62
Hansen et al. (1993)	Lab	<i>globosa</i>	5	5
Hansen et al. (1994b)	Lab	<i>pouchetii</i>		20
Huntley et al. (1987)	Lab	<i>pouchetii</i>	4	4
Klein Breteler and Koski (2003)	Field	<i>globosa</i>		4
Koski et al. (2005)	Field	<i>globosa</i>	2	6
Koski unpublished	Field	<i>pouchetii</i>	3	3
Tande and Båmstedt (1987)	Lab	<i>pouchetii</i>	95	95
Tang et al. (2001)	Lab	<i>globosa</i>		4
Turner et al. (2002)	Lab	<i>globosa</i>	15	15
Verity and Smayda (1989)	Lab	<i>globosa</i>	9	9
Verity (2000)	Lab	<i>globosa</i>	8	12
Weisse (1983)	Lab	<i>globosa</i>		18

n denotes the number of observations from each source

set of data, in a sample of fewer dimensions (or axes) that is the best summary of the information contained in the data (Greenacre 1984; Everitt and Dunn 2001). Our data set contains both categorical and continuous variables that were used in the MCA as active (Table 5) and illustrative variables (PL, ESD, CI and Ab defined below), respectively. The factorial axes of the MCA were computed using active (categorical) variables whereas the continuous illustrative (or supplementary) variables were simply projected into this factorial plane without participating in its computation. The location of these continuous illustrative variables, is shown as arrows along the factorial axes in Fig. 1A, and expresses their linkage to the pattern of the categorical active variables displayed by the factorial axes (Lebart et al. 2000). Observations with missing data for carbon-specific ingestion and experimental characteristics were excluded from the MCA. Only four observations with *Phaeocystis antarctica* were available, and were therefore also excluded. A data matrix of the remaining 296 observations with eight active and four illustrative variables were used for the analyses (Table 5). The size of predators (prosome length; PL) and prey (equivalent spherical diameter; ESD) were treated as illustrative variables.

These continuous variables were used to calculate a predator-to-prey size ratio (P:p) with five categories (or modalities) used in the statistical treatment. Similarly, C-specific ingestion (CI) and *Phaeocystis* abundance (Ab) were transformed into categorical variables of three and five categories, respectively, to be tested both as illustrative and active variables in the MCA. The metric used in the MCA is based on the χ^2 . This was also the metric used in the following cluster analysis. The type of HCA used here is an agglomerative clustering, i.e., a procedure that successively groups the closest objects into clusters, which then are grouped into larger clusters of higher rank (Legendre and Legendre 1998). The programme identifies: (i) the cluster (group of observations) which has the smallest within-group variance and the greater variance between groups, and (ii) the descriptors (variables) that are highly representative of each cluster. The descriptors of the observations used in the clustering were their factorial coordinates on the first six axes obtained in the MCA. These first six axes explained about 58% of the total variability (inertia) in the data, and the additional amount of variability explained by the axes decreased markedly after the first six axes.




Fig. 1 First factorial plane of MCA of data on crustacean grazing experiment on *Phaeocystis*. (A) Projections of continuous illustrative variables in the correlation circle (radius 1) and ordination of active variables: *Phaeocystis* species (Δ), growth (\triangleleft) and abundance (\circ), crustacean species (∇), predator-to-prey

size ratio (\triangleright), experiment location (\square), selection (\diamond). (B) Ordination of data (\circ) labelled according to their C-specific ingestion characteristic and delimitation of the four groups designed by the hierarchical clustering (χ^2 distance). See Table 5 for label identification

Results from statistical analysis

The result of the MCA indicated that the first and second axes accumulate about 19% and 12% of the total variability, respectively (Fig. 1A). Two variables basically contribute to the first axis: the selection

(“Against” and “Single”, cumulated 18% of the contribution to axis 1) and the experiment location (“Lab” and “Field”, cumulated 17.5% of the contribution). Thus, this axis differentiates laboratory experiments where *Phaeocystis* was the only prey (“Single”), as opposed to field experiments where

Table 7 Characterization of the most typical properties of the four groups designed by the hierarchical cluster analysis defined in Fig. 1 panel B

Group <i>n</i>	1			2			3			4		
	82			15			27			172		
	Modality	gr/mod	mod/gr	Modality	gr/mod	mod/gr	Modality	gr/mod	mod/gr	Modality	gr/mod	mod/gr
Group	Against	95	88	For	100	93	P:p4	96	89	Single	84	98
characteristics	Field	80	89	Stationary	79	100	Tst	100	56	<i>P. pouchetii</i>	87	95
	P:p1	96	67	Ab1	19	100	<i>P. globosa</i>	25	100	Lab	82	98
	<i>P. globosa</i>	66	87	Field	16	100	Tlo	34	44	P:p5	100	44
	CI1	55	56	P:p2	14	87	Lab	13	100	Chy	96	42
	Cha	100	17	Cgl	38	20	CI3	19	44	Cfi	88	53
	Acl	100	16	<i>P. pouchetii</i>	7	93				Ab4	86	35
	Pe1	100	16							Exponential	62	100
	Ab1	48	48							P:p3	88	25
	Tlo	63	27							Ab5	100	9
	Ab2	44	49							CI3	77	28
										CI2	66	58

Gr/mod corresponds to the percentage of the modality belonging to the group, and mod/gr corresponds to the percentage of the group belonging to the modality. Modality label as in Table 5. *n* denotes the number of observations

alternative prey were available and *Phaeocystis* was selected against (“Against”). The predator-to-prey size ratio modality “P:p1” (size ratio < 4) also has an important contribution to the first axis (10%), showing that small copepods (such as *Centropages hamatus*, *Acartia clausi* and *Pseudocalanus elongatus*) tend to reject *Phaeocystis* in situ. On axis 2, the two modalities “Stationary” and “For” have the greatest contribution (18% and 20%, respectively). This axis separates grazing experiments where *Phaeocystis* in a stationary growth phase was selected positively by predators in situ. In the first factorial plane we also observed that *Phaeocystis* concentration and predator-to-prey size ratio in lab experiments were higher than what was generally observed in situ (Fig. 1A) and that higher carbon-specific ingestions were mostly related to laboratory experiments (Fig. 1B).

The combined analysis of the MCA and HCA clearly distinguished four groups of observations (Fig. 1B and Table 7). Groups 1 and 2 bring together 97% of the grazing experiments performed in the field and are separated from groups 3 and 4, which comprise 96% of the lab experiments with *Phaeocystis* provided as single prey. Group 1 consists of field experiments with small-size copepods selecting against *P. globosa*. Group 2 includes all field experiments with positive selection for *P. pouchetii* (93% of

the group), and is characterized by a slightly higher predator-to-prey size ratio (87% of them had a P:p-ratio of 4–16, “P:p2”). In these experiments *Phaeocystis* was in the stationary growth phase and in relatively low abundance (“Ab1” = abundance between 0.1 and 125 $\mu\text{g C l}^{-1}$). The group 3 is characterized by lab grazing experiments of *Temora* spp. on *P. globosa*. The predator-to-prey size ratio in these laboratory experiments was higher (89% of them had a P:p ratio of 64–256, “P:p4”) than what was commonly observed in grazing experiments on natural plankton. Similarly we observed higher carbon-specific ingestion than what was observed in situ for this copepod genus (group 1). Group 4 brings together the lab grazing experiments on *P. pouchetii* by larger copepods (*Calanus* spp.). This group includes the highest *Phaeocystis* abundance and predator-to-prey size ratio tested in lab experiments and grouped the highest carbon-specific ingestion estimates.

In conclusion, the MCA and the HCA analyses suggested that: (1) the lowest grazing rates from the field (often zero) were recorded for small copepods (*Acartia*, *Pseudocalanus*, *Temora* and *Centropages*) in blooms of *Phaeocystis globosa* colonies, (2) large copepods (e.g., *Calanus*) feeding on *Phaeocystis pouchetii* had higher grazing rates, especially in lab studies, when no alternative food was present and (3)

P. pouchetii in a stationary growth phase was positively selected by large copepods.

It is important to note that grazing data for large copepods (*Calanus* spp. and *Metridia longa*) were only available for *P. pouchetii*, whereas grazing studies with small copepods (*Acartia*, *Centropages*, *Pseudocalanus* and *Temora*) were limited to *P. globosa*. Thus, the presently available data do not allow us to determine whether the different results for *P. pouchetii* and *P. globosa* are due to differences between the two species, or differences between the predators. In order to test for species-specific differences, we need to compare grazing on both species simultaneously using similar methodologies in future studies.

Quantitative results from data on crustacean grazing

Because the statistical analysis suggested a large difference between feeding rates in laboratory and field investigations, we compared the average specific ingestion rates in all available field and laboratory experiments (Table 6), and for five groups of predator-to-prey size ratios separately (Fig. 2). These comparisons revealed that: (1) overall feeding (carbon specific ingestion) on *Phaeocystis* spp. was significantly lower [$p < 10^{-8}$, analysis of variation (ANOVA)] in all field (average 2.5% d^{-1}) studies with natural plankton, compared to laboratory studies (average 11% d^{-1}), (2) the highest average carbon-specific ingestion rate (23% d^{-1}) on *Phaeocystis* spp. was found when the predator-to-prey size ratio (P:p) was 4–16 in lab studies, whereas there was no clear size ratio trend in field studies, and (3) ingestion was low ($\leq 2\%$ d^{-1}) when P:p was < 4 in both lab and field, indicating that the upper effective size limit for prey equivalent spherical diameter was $\frac{1}{4}$ of the predator prosome length.

Thus, crustacean grazers show a much lower grazing on *Phaeocystis* in the field than in the laboratory. One reason for this discrepancy could be that fewer than 5% of the laboratory studies offered the copepods alternative prey to *Phaeocystis* whereas alternative prey were available in the field. Alternatively, laboratory cultures might not display the same chemical grazing cues (and possible grazing deterrents) as *Phaeocystis* growing in natural plankton. Haptophytes may lose their inhibitory effects in vitro, and toxicity may be species-, strain- or growth-condition-




Fig. 2 Crustaceans (all stages of copepods and krill) grazing on *Phaeocystis pouchetii*, *P. globosa* or *P. antarctica*, in laboratory and field experiments. Average daily carbon-specific ingestion rates (percent $\mu\text{g C } \mu\text{g C}^{-1} \text{d}^{-1}$) are presented for five different predator-to-prey size categories [derived from predator prosome or carapace length and average equivalent spherical diameter (ESD) of the *Phaeocystis* prey, solitary cells or colonies]. Error bars denote the standard error (SE) of the average; the number of observations (n) for each category are given above each column. Note the high daily ration for P:p size ratio 64–256 in the field is due to a single (30%) value (with that value excluded, the rate is $3.0 \pm 1.0\%$)

specific (Edvardsen and Paasche 1998). Support for this hypothesis comes from laboratory studies that failed to recreate toxicity in the lab that was observed in the field, even with haptophytes isolated from highly toxic blooms, such as the *Chrysochromulina polylepis* bloom in 1988 (Nielsen et al. 1990). Likewise, the antipredation effect revealed by Estep et al. (1990) in actively growing field-collected *Phaeocystis pouchetii* apparently disappeared during the first 12 h in experimental containers. It may be that laboratory or other in vitro studies underestimate the negative effects of potentially toxic haptophytes in situ, but this requires explicit evaluation.

Ideally, future investigations on the ecology of zooplankton feeding should preferably be conducted in situ, although laboratory experiments are needed to investigate the effects of potentially important signaling substances between *Phaeocystis* and different predators in presence of realistic alternative prey.

Grazing by protozooplankton and other microzooplankton

General trends in grazing by protozoan microzooplankton on *Phaeocystis* are difficult to assess since the number of quantitative investigations is limited (Tables 3 and 4). Very few studies have investigated

Table 8 Microzooplankton grazing on *Phaeocystis globosa* solitary cells during blooms in the English Channel off northern France May 2003 and April 2004 (JC Nejtgaard, AF Sazhin and LF Artigas unpubl.)

Date	Prey type (μm)	r^2	μ (d^{-1})	G (d^{-1})	Grazing impact (% SS d^{-1})
Apr 2003	Chl <i>a</i> (>0.45)	0.58	$0.24 \pm 0.07^{**}$	$0.61 \pm 0.14^{***}$	46
	Diatoms (5-99)	0.45	$1.34 \pm 0.18^{***}$	$1.24 \pm 0.40^{**}$	71
	<i>Phaeocystis</i> (2-8)	0.04	0.08 ± 0.16 ns	0.21 ± 0.30 ns	19
May 2004	Chl <i>a</i> (>0.45)	0.13	$-1.17 \pm 0.12^{***}$	0.18 ± 0.18 ns	17
	Diatoms (5-99)	0.18	0.16 ± 0.09 ns	0.19 ± 0.13 ns	18
	<i>Phaeocystis</i> (2-8)	0.04	$-1.14 \pm 0.20^{***}$	-0.23 ± 0.34 ns	0

Dilution experiments were performed and analyzed as described in Nejtgaard et al. (2001). Specific phytoplankton growth rate (μ), microzooplankton grazing coefficient (g), grazing impact (percentage of average standing stock removed d^{-1}). \pm SE for the mean. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ for μ or $g = 0$. Prey types are given with equivalent spherical diameter size ranges (μm)

microzooplankton as potential grazers on colonial *Phaeocystis* spp. The colonies are protected by a tough skin (Hamm et al. 1999) and microzooplanktoners are generally assumed to be too small to prey actively on colonial forms of *Phaeocystis* (Hansen et al. 1994a; Weisse et al. 1994; Tang 2003). In accordance, Irigoien et al. (2003) reported that nauplii of *Calanus finmarchicus* did not ingest colonies of *Phaeocystis* in the Irminger sea. However, the protists *Noctiluca scintillans* (Weisse et al. 1994; Jakobsen and Tang 2002) and apparently *Gyrodinium cf. spirale* (Stelfox-Widdicombe et al. 2004) are able to ingest small colonies, while some tintinnids and rotifers “attack” and “hover around colonies”, possibly ingesting released cells (Hollowday 1949; Admiraal and Venekamp 1986). The abundance of microzooplankton has also been reported to decline or remain low during natural blooms of colonial *Phaeocystis*, but increases rapidly during the breakdown of the colonies at the end of the bloom (e.g., Peperzak et al. 1998; van Boekel et al. 1992). This suggests that actively growing colonies are inferior prey for microzooplankton, while single cells released from decaying colonies may be a suitable food source. However, an alternative explanation for the low microzooplankton abundance during *Phaeocystis* blooms is the strong selective predation pressure from larger zooplankton such as copepods (Hansen et al. 1993; Gasparini et al. 2000).

It is clear from the literature that at least some microzooplankton may readily ingest single-celled *Phaeocystis* (Tables 3 and 4), and chemical cues from microzooplankton grazing have been shown to induce colony formation (Long and Hay 2006) and enlargement in *P. globosa* (Tang 2003), supporting the

assumption that microzooplankton graze on the single cells but not the colonies. However, some of the reported grazing rates in laboratory studies were low, with or without alternative prey present (Table 3). Also, in several incubation experiments with natural plankton the microzooplankton community grazing rates were very low ($\leq 0.1 \text{ d}^{-1}$) during dominance of *Phaeocystis* (Table 4). It is well known that some microzooplankton feed selectively (e.g., Verity 1988; González et al. 1990; Strom and Loukos 1998; Archer et al. 2000), and it was recently shown that copepod nauplii ingest some clones and cell morphotypes of *P. globosa* at low rates (Dutz and Koski 2006). Results showing selection against single cells by microzooplankton in the sea was obtained in recent dilution experiments in the English Channel (Table 8). These experiments yielded zero or close to zero microzooplankton grazing rates on single-celled *P. globosa*, but high microzooplankton grazing rates on diatoms based on cell counts, while bulk measurements of chlorophyll *a* (as used in most of the experiments in Table 4) indicated intermediate grazing values. This suggests that microzooplankton feeding on *Phaeocystis* may be highly variable in the sea, and reliable grazing rate measurements will therefore require taxon-specific quantitative methods, rather than bulk measurements (see further discussion under Future challenges).

Cell-type and life-stage-specific interactions with grazers

In the species *P. globosa*, three solitary, flagellated cell types (micro-, meso-, and macroflagellates)

differing in size and morphology have been described (Kornmann 1955; Peperzak et al. 2000). The ecological roles of these different cell types are largely unknown, partly due to difficulties in the identification and quantification of the flagellated cell type in field samples (Rousseau et al. 1994; Peperzak et al. 2000; Schoemann et al. 2005; Rousseau et al. this volume). A recent study showed that the grazing mortality of single cells may depend on the cell type, with moderate to high ingestion by *Temora longicornis* nauplii on non-flagellated cell types, but rejection of a flagellate type (Dutz and Koski 2006). This rejection was ascribed to the production of chitinous threads and the cohesion of threads into pentagonal star-like structures which are a typical feature of mesoflagellates (Chretiennot-Dinet et al. 1997; Peperzak et al. 2000).

The variable cell-type-specific grazing mortality of *P. globosa* offers important insight into the natural history of this phytoplankton species. Although flagellated and non-flagellated cell types co-occur throughout the year, blooms of *Phaeocystis* in temperate waters are generally dominated by the colonial cell type and by colonies (Peperzak et al. 2000; Rousseau et al. this volume). At the end of such blooms, single cells are released from the colonies presumably due to nutrient deficiency or irradiance limitation. The liberated cells appear to suffer a high mortality due to microzooplankton grazing, cell lysis and/or perhaps sedimentation (van Boekel et al. 1992; Brussaard et al. 1995; Riebesell et al. 1995; Brussaard et al. this volume; Reigstad & Wassmann this volume). Concurrent to the liberation and disappearance of single cells, an increasing abundance of micro/mesoflagellates has been observed in mesocosms and in the field (Veldhuis et al. 1986; Peperzak et al. 2000; Escaravage et al. 1995; Nejstgaard et al. 2006). The formation of intracolony flagellated cells may bridge the disappearance of colonial cells and the appearance of flagellated cells (Peperzak et al. 2000). These haploid micro/mesoflagellates probably function as gametes or spores to survive unfavourable conditions during the warm summer months (Veldhuis et al. 1986; Peperzak et al. 2000). Reduced vulnerability of these cells to microzooplankton grazing could explain the accumulation of single flagellated cells despite a high grazing pressure at the end of *Phaeocystis* blooms, and be part of these phytoplankters' strategy to increase survival and foster bloom formation when

favourable conditions return. Future studies should investigate the relevance of cell type- and life-stage-specific selection on wider range of zooplankton organisms.

Colony formation and its potential role in morphological defense

The presence of the two distinctly different morphotypes in *Phaeocystis* has long intrigued scientists, and much remains unknown about their respective biological roles and regulation of transition between morphotypes (Rousseau et al. 1994; Lancelot et al. 1998; Rousseau et al. this volume). The prevalence of both morphotypes in natural *Phaeocystis* blooms in contrasting water types prompts the idea that *Phaeocystis* colony development is regulated by common factors (Lancelot and Rousseau 1994) and contributes to the success of the genus in marine systems (Rousseau et al. 1994; Lancelot et al. 1998; Rousseau et al. this volume). The production of the mucilaginous structure represents a substantial energy investment by the cells; the mucilaginous matrix may account for >50% of the total organic carbon of a *Phaeocystis* population (Rousseau et al. 1990; Mathot et al. 2000). The large colony size may also pose a problem of diffusion limitation to nutrient and oxygen uptake, although a modeling study shows that the colonies are unlikely to be limited by diffusion (Ploug et al. 1999). Thus, compared to solitary *Phaeocystis* cells, it is likely that the formation of colonies involves some additional costs, but that the drastic increase in size may be an effective defense mechanism against (relatively) small grazers, not unlike the highly successful freshwater colonial chlorophyte *Scenedesmus* (Lüring 2003; e.g., Lüring and Van Donk 1997), cyanobacteria *Microcystis aeruginosa* and *Sphaerocystis* sp. (e.g., De Bernardi and Giussani 1990), and for different copepods feeding on the large colony-forming diatom *Thalassiosira partheneia* (Schnack 1983).

Some studies suggest that microzooplankton prefer and perform better on single cells than on colonies. When incubated with a mixture of *Phaeocystis globosa* single cells and colonies, microzooplankton reduced the single cells to very low abundance, and the grazer population subsequently declined even when colonies were abundant, presumably due to starvation (Jakobsen and Tang 2002). When restricted

to a single diet of either single cells or colonies, the ciliate *Euplotes* grew three times faster on single cells (Long 2004). In contrast, the copepod *Acartia tonsa* and mixed mesozooplankton had higher feeding rates on colonies compared to single cells (Long and Hay 2006). The differential susceptibility of morphologies to grazers could allow *Phaeocystis* to use transformations as an adaptive, inducible response towards grazers. Two recent studies tested the role of chemical cues for inducing transformations using either incubators separated by a permeable membrane (Tang 2003) or filtrate experiments (Long 2004). In one study chemical cues from grazers triggered colony enlargement (Tang 2003). In the other study microzooplankton cues enhanced colony formation while cues from macrozooplankton suppressed colony formation (Long 2004). These observations point to the involvement of a grazing-related chemical signal in morphological defense, but the chemical characteristics of this signal remain unknown (Long and Hay 2006; Tang 2003). Colony formation by *Phaeocystis* could affect grazing even after the demise of a bloom. For example, the degrading colony matrix, making the surrounding water gelatinous, may continue to prevent grazing losses of single cells by microzooplankton (Seuront et al. this volume), and similarly, transparent exopolymer particles formed by coagulation of colony-derived carbohydrates could also inhibit copepod grazing (Dutz et al. 2005).

Chemical defense

Macroalgae commonly use chemical defense against grazers (Hay and Fenical 1988; Pohnert 2004), and similar examples may exist in phytoplankton (see review by Pohnert 2004; and further examples in Yoshida et al. 2004; Pohnert 2005). A few observations suggest that chemical defense could be important for survival of *Phaeocystis* during exponential growth (Weisse et al. 1994; Estep et al. 1990; Long and Hay 2006). Recently, a number of toxins have been isolated from *Phaeocystis* (reviewed by van Rijssel et al. this volume), and *Phaeocystis* is reportedly toxic to some aquatic organisms, including phytoplankton, sea urchin and cod larvae (Table 2). However, specific chemical zooplankton feeding deterrents from *Phaeocystis* have not yet been isolated and characterized so their existence is still

hypothetical. Several of the reported toxic effects were obtained using organisms (or even extracts) that do not interact in nature, and the ecological relevance of such results is unknown.

Does DMS affect grazing on *Phaeocystis*?

Phaeocystis is a prominent producer of the secondary metabolite dimethylsulphoniopropionate (DMSP) that can be enzymatically cleaved to acrylate and the volatile trace gas dimethyl sulphide (DMS; Keller et al. 1989; Schoemann et al. 2005; Stefels et al. this volume). DMSP is a multifunctional compound that probably assists several physiological processes related to salinity-, temperature- and light-stress in algal cells (Stefels 2000). This compound and its cleavage products have also been suggested to function as grazing deterrents (Strom et al. 2003; Wolfe et al. 1997) and it is possible that they provide a competitive advantage to *Phaeocystis* when non-DMSP-producing alternative prey are available to potential predators. Microzooplankton grazing greatly increases the production of DMS in *Emiliania huxleyi* (Wolfe and Steinke 1996) and microzooplankton are suspected to have caused the conversion of *Phaeocystis*-DMSP to DMS in the southern North Sea (Archer et al. 2003). The grazing-induced production of DMS may be analogous to the production of volatile infochemicals during herbivore grazing on terrestrial plants that utilize such trace gases to attract carnivores (Dicke and Sabelis 1988; Steinke et al. 2002; Hay et al. 2004). Such indirect defense mechanisms involve interactions on three trophic levels that include plants, herbivores and carnivores. Seabirds are attracted by gradients of DMS in the atmosphere (Nevitt et al. 1995; Nevitt and Bonadonna 2005), and copepods are chemosensitive to DMS and react with search behavior when encountering plumes of this compound (Steinke et al. 2006). If DMS produced during grazing makes the microzooplankton grazers more susceptible to predation by copepods, then investment into such an indirect defense mechanism could be beneficial for the phytoplankton.

However, the ecological consequences of the production of DMS and associated compounds are controversial and explicit tests of possible defensive functions for *Phaeocystis* are required. For example, Tang and Simó (2003) did not observe negative

effects when the heterotrophic dinoflagellate *Gyrodinium dominans* grazed on *P. globosa* single cells, and the grazers retained >40% of the ingested DMSP. Furthermore, the grazing-induced production of DMS by unicellular phytoplankton is a result of grazing on individual cells and, as a consequence, these ingested cells will not directly benefit from the potential signaling to carnivorous enemies of the microzooplankton. It is possible that such indirect defense mechanisms are beneficial for *Phaeocystis* populations but they would also be of benefit to other members of the phytoplankton community that may not invest in chemical defenses (the cheater problem: Lewis 1986). One may argue that the cheater problem does not exist within a bloom of genetically closely related cells. However, molecular studies have shown that the genetic variation in phytoplankton populations is high, even within a monospecific diatom bloom (Rynearson and Armbrust 2005). It is impossible to predict functional diversity from such molecular data and there is very little information on the effect of genetic variation on the ecophysiological fitness in *Phaeocystis*.

Evidence is accumulating that chemical communication influences the composition and dynamics of pelagic communities (Pohnert 2004). The production of DMSP in *Phaeocystis* is probably only one example where a secondary metabolite benefits the physiology of individual cells and its catabolic products could affect the structure and function of *Phaeocystis*-dominated food webs. This is a complex research area that needs to be addressed with more detailed studies in the future.

Does nutritional value affect grazing on *Phaeocystis*?

Copepods may feed selectively based on nutritional quality of the prey (e.g., Houde and Roman 1987). However, it is not clear how food quality of the different forms of *Phaeocystis* affects zooplankton grazing. It has been suggested that *Phaeocystis* spp. are of low nutritional value due to their low content of polyunsaturated fatty acids (PUFA) (Al-Hasan et al. 1990; Claustre et al. 1990; Rogers and Lockwood 1990; Nichols et al. 1991; Cotonnec et al. 2001; Tang et al. 2001; Turner et al. 2002). This contrasts with the high survival and development rates observed in other

studies (Verity and Smayda 1989), at least for some strains of *P. globosa* (Dutz and Koski 2006). High amounts of C₁₈-PUFA were detected in a *P. globosa* bloom in the North Sea (Hamm and Rousseau 2003) and in a *P. pouchetii* bloom in Balsfjorden in northern Norway (Hamm et al. 2001). The mucus produced by colonies is generally considered to be of low nutritional value, refractory (Thingstad and Billen 1994), and of low carbohydrate content (van Rijssel et al. 1997). However, the composition of the mucus material may vary during a bloom cycle (Alderkamp et al. 2006), and colonization of this material by pennate diatoms and other organisms on and inside older colonies (Hamm and Rousseau 2003; Sazhin et al. this volume) may increase its nutritional value. Besides PUFA and carbohydrates, the effects of other important nutritional components of *Phaeocystis* such as amino acids, sterols and vitamins, on copepod feeding and reproduction are still poorly known.

Survival of gut passage?

Some phytoplankton can survive gut passage, a trait that is often overlooked in zooplankton studies. For instance, large fractions of the chlorophytes *Chlamydomonas reinhardtii* and *Selenastrum capricornutum* survive passing the guts of *Daphnia*. This is considered a defense mechanism by the algae to reduce grazing pressure when their growth rates are low (van Donk et al. 1997). Further, the colonial chlorophyte *Sphaerocystis Schroeteri* survives gut passage in *Daphnia* and benefits from grazer-released nutrients. This interaction is hypothesized to affect bloom formation in this species (Porter 1976). If cells of *Phaeocystis* may survive gut passage of zooplankton, this would have significant implications for the understanding of the grazing interactions. Because, net removal rates of the prey from the population will decrease, and if cells may produce defense chemicals upon predation and surviving gut passage it would affect the chemical antipredatory mechanisms suggested above. But, to our knowledge there are no such studies for *Phaeocystis* or any other haptophyte phytoplankton. However, small intact colonies of *P. pouchetii* have been observed fluorescing inside copepod pellets in the Barents Sea (P. Verity and P. Wassmann, personal communication) suggesting that surviving gut passage might be possible for *Phaeocystis*.

Conclusions and future challenges

Although over 100 publications have examined zooplankton grazing on *Phaeocystis* spp., more than 90% report on copepods or other crustacean grazers, so quantitative data for other grazers are limited. Grazing on *Phaeocystis* was often determined by feeding on only single types of cultured phytoplankton in the laboratory. Field reports often do not provide enough information to evaluate even some of the most basic factors known to affect zooplankton grazing simultaneously, such as predator-to-prey size ratio, food concentration and feeding rates on alternative prey. Further, most reports on crustacean grazing in the field are based on non-prey-specific grazing estimations such as bulk gut pigments, that cannot be used to determine specific feeding on *Phaeocystis* in the mixed natural plankton assemblages.

Nevertheless, our analysis of the present literature suggests that: (1) laboratory-derived crustacean grazing rates on monocultures of *Phaeocystis* may have been overestimated compared to feeding in natural plankton communities, and should be treated with caution; (2) formation of colonies by *P. globosa* appeared to reduce predation by small copepods (e.g., *Acartia*, *Pseudocalanus*, *Temora* and *Centropages*), whereas large copepods (e.g., *Calanus* spp.) were able to feed on colonies of *Phaeocystis pouchetii*; (3) physiological differences between different growth states, species, strains, cell types, and laboratory culture versus natural assemblages may explain most of the variations in reported feeding rates; (4) chemical signaling between predator and prey may be a major factor controlling grazing on *Phaeocystis*; (5) it is unclear to what extent different zooplankton, especially protozooplankton, feed on the different life forms of *Phaeocystis* in situ.

In the present literature there is a dichotomy between data showing high feeding rates on both single cells and colonies of *Phaeocystis pouchetii* by larger copepods, especially in the laboratory studies, and avoidance of colonies of *Phaeocystis globosa* by other smaller copepods in the field. This imbalance in basic quantitative data could initially be remedied by applying classical bottle incubation methods with a range of predators, preferably using natural plankton in situ, analyzed with methods that resolve the feeding rates on the different life forms (and prey sizes) of different species of *Phaeocystis* from alternative prey

types. Such analyses could be done by microscopy (Verity and Paffenhöfer 1996; Båmstedt et al. 2000) corrected for food web cascades (Nejstgaard et al. 2001), and/or possibly analyzed by a combination of more recent methods such as flow cytometry (Collier and Campbell 1999; Jonker et al. 2000), computer/video-aided plankton counting devices (See et al. 2005), and specific molecular probing (Caron 2005) to increase the taxonomic precision. However, the ability of chemical cues to rapidly alter colony formation indicates that grazing estimates based on cell counts could grossly misrepresent actual grazing rates.

For example, zooplankton grazing may induce or suppress colony formation and enlargement (Tang 2003; Long 2004; Long and Hay 2006), and accumulating data suggests that chemical signaling between different levels of predators of prey can play an important role influencing the dynamics in pelagic communities (e.g., reviewed by Pohnert 2004; Yoshida et al. 2004). This suggests that chemical signaling in artificially increased grazer densities over long incubations could confound data from bottle experiments by changing the relative abundance of different life forms in the grazing bottle due to factors other than direct ingestion of the cells. Further, recent results suggesting that feeding on *Phaeocystis* may even be cell type- or life-stage-specific (Dutz and Koski 2006), and that new methods accounting for such differences are necessary.

Thus, the ideal way to assess feeding on *Phaeocystis* would be a direct quantification of feeding rates on specific prey by assessing individual predators that have been feeding undisturbed in situ. Molecular quantitative analysis of the prey in the guts and faecal pellets of the predators could potentially achieve this. Such approaches are already common in studies of e.g., terrestrial arthropods (Sheppard and Harwood 2005; Symondson 2002), and it has been shown that prey DNA from haptophyte phytoplankton can be detected in copepod guts and fecal pellets by standard polymerase chain reaction (PCR) targeting 18S ribosomal DNA (Nejstgaard et al. 2003). More recently, it was also shown that carnivorous diet may be inferred from the content of copepod fecal pellets using standard PCR (Vestheim et al. 2005), and that different phytoplankton DNA, including *Phaeocystis*, may be quantified in guts and fecal pellets from both appendicularians

and copepods using quantitative real-time PCR (Nejstgaard et al. 2005; Troedsson et al. 2007).

Thus the rapidly developing molecular approaches may help us obtain more-realistic ingestion rates of *Phaeocystis* and other live prey in situ. However, to further reveal and quantify mechanisms controlling the complex trophic interactions between *Phaeocystis* and its predators, new approaches need to be developed. In concert with the recent review by Pohnert (2004) we conclude that such development will probably be as complex as the interactions it tries to reveal. Developing successful innovative approaches and better quantitative analytical tools will depend largely upon multidisciplinary efforts between chemists, molecular biologists and plankton ecologists.

Acknowledgements We thank S. Gasparini for his valuable suggestions concerning MCA and HCA analyses. This work was supported by the Norwegian Research Council project 152714/120 30 to JCN, the US-NSF OPP award ANT-0440478 to KWT, the UK Natural Environment Research Council grants NER/I/S/2000/00897 and NE/B500282/1 to MS, the German Federal Ministry for Education and Research grants within the GLOBEC Germany project 03F0418C to JD, the Carlsberg Foundation to MK, the Belgian Research Program for sustainable management of the North Sea within the AMORE project MN/DD/20/21/22 to EA, and the EPA Star Fellowship U-91599501-0 and a National Parks Ecological Research Fellowship to JDL.

References

- Aanesen RT, Eilertsen HC, Stabell OB (1998) Light-induced toxic properties of the marine alga *Phaeocystis pouchetii* towards cod larvae. *Aquat Toxicol* 40:109–121
- Admiraal W, Venekamp LAH (1986) Significance of tintinnid grazing during blooms of *Phaeocystis pouchetii* (haptophyceae) in Dutch coastal waters. *Neth J Sea Res* 20:61–66
- Al-Hasan RH, Ali AM, Radwan SS (1990) Lipids, and their constituent fatty-acids, of *Phaeocystis* sp. from the Arabian Gulf. *Mar Biol* 105:9–14
- Alderkamp A-C, Buma AGJ and van Rijssel M (this volume) The carbohydrates of *Phaeocystis* and their degradation in the microbial food web. *Biogeochemistry*. doi:10.1007/s10533-007-9078-2
- Alderkamp A-C, Nejstgaard JC, Verity PG, Zirbel MJ, Sazhin AF, van Rijssel M (2006) Dynamics in carbohydrate composition of *Phaeocystis pouchetii* colonies during spring blooms in mesocosms. *J Sea Res* 55:169–181
- Archer SD, Stelfox-Widdicombe CE, Malin G, Burkill PH (2003) Is dimethyl sulphide production related to microzooplankton herbivory in the southern North Sea? *J. Plankton Res* 25:235–242
- Archer SD, Verity PG, Stefels J (2000) Impact of microzooplankton on the progression and fate of the spring bloom in fjords of northern Norway. *Aquat Microb Ecol* 22:27–41
- Båmstedt U, Gifford DJ, Irigoien X, Atkinson A, Roman M (2000) Feeding. In: Harris R, Wiebe P, Lenz J, Skjoldal HR, Huntley M (eds) ICES Zooplankton Methodology Manual. Academic Press, London, pp 297–399
- Banse K (1994) Grazing and zooplankton production as key controls of phytoplankton production in the open ocean. *Oceanography* 7:13–20
- Barnard WR, Andreae MO, Iverson RL (1984) Dimethylsulfide and *Phaeocystis pouchetii* in the Southeastern Bering Sea. *Cont Shelf Res* 3:103–113
- Baumann MEM, Lancelot C, Brandini FP, Sakshaug E, John DM (1994) The taxonomic identity of the cosmopolitan prymnesiophyte *Phaeocystis*: a morphological and eco-physiological approach. *J Mar Syst* 5:5
- Bautista B, Harris RP, Tranter PRG, Harbour D (1992) In situ copepod feeding and grazing rates during a spring bloom dominated by *Phaeocystis* sp. in the English Channel. *J Plankton Res* 14:691–703
- Blauw AN (this volume) *Phaeocystis* blooms forecasting in Dutch coastal waters, the impact of suspended matter. *Biogeochemistry*
- Breton E, Sautour B, Brylinski J-M (1999) No feeding on *Phaeocystis* sp. as solitary cells (post bloom period) by the copepod *Temora longicornis* in the coastal waters of the English Channel. *Hydrobiologia* 414:13–23
- Brussaard CPD, Bratbak G, Baudoux A-C and Ruardij P (this volume) *Phaeocystis* and its interaction with viruses. *Biogeochemistry*. doi:10.1007/s10533-007-9096-0
- Brussaard CPD, R. R, Noordeloos AAM, Cadée GC, Witte H, Kop AJ, Nieuwland G, van Duyl FC and Bak RPM (1995) Effects of grazing, sedimentation and phytoplankton cell lysis on the structure of a coastal pelagic food web. *Mar Ecol Prog Ser* 123
- Bullen GE (1908) Plankton studies in relation to the western mackerel fishery. *J Mar Biol Ass UK* 8:269–302
- Caron DA (2005) Introductory remarks: advances in the molecular ecology of protists. *J Eukaryot Microbiol* 52:81–82
- Caron DA, Dennett MR, Lonsdale DJ, Moran DM, Shalapyonok L (2000) Microzooplankton herbivory in the Ross Sea, Antarctica. *Deep Sea Res Part II Top Stud Oceanogr* 47:3249–3272
- Chen Y-Q, Wang N, Zhang P, Zhou H, Qu L-H (2002) Molecular evidence identifies bloom-forming *Phaeocystis* (Prymnesiophyta) from coastal waters of southeast China as *Phaeocystis globosa*. *Biochem Syst Ecol* 30:15–22
- Chretiennot-Dinet MJ, Giraud-Guille MM, Vaulot D, Putaux JL, Saito Y, Chanzy H (1997) The chitinous nature of filaments ejected by *Phaeocystis* (*Prymnesiophyceae*). *J Phycol* 33:666–672
- Claustre H, Poulet SA, Williams R, Marty JC, Coombs S, Benmlih F, Hapette AM, Martinjezequel V (1990) A biochemical investigation of a *Phaeocystis* sp. bloom in the Irish Sea. *J Mar Biol Ass U K* 70:197–207
- Collier JL, Campbell L (1999) Flow cytometry in molecular aquatic ecology. *Hydrobiologia* 401:33–53
- Cook PA, Gabbott PA (1972) Seasonal changes in the biochemical composition of adult barnacle, *Balanus balanoides*, and the possible relationships between biochemical

- composition and cold-tolerance. *J Mar Biol Ass U K* 52:805–825
- Cotonnec G, Brunet C, Sautour B, Thoumelin G (2001) Nutritive value and selection of food particles by copepods during a spring bloom of *Phaeocystis* sp. in the English Channel, as determined by pigment and fatty acid analyses. *J Plankton Res* 23:693–703
- Daro M-H (1985) Field study of the diel, selective and efficiency feeding of the marine copepod *Temora longicornis* in the Southern Bight of the North Sea. In: Van Grieken R, Wollast R (eds) *Proc progress in Belgian oceanographic research*, Brussels, pp 250–263
- Davies AG, Demadariaga I, Bautista B, Fernandez F, Harbour DS, Serret P, Tranter PRG (1992) The ecology of a coastal *Phaeocystis* bloom in the North-Western English-Channel in 1990. *J Mar Biol Ass UK* 72:691–708
- De Bernardi R, Giussani G (1990) Are blue-green algae suitable food for zooplankton? A review. *Hydrobiologia* 200/201:29–41
- Dicke M, Sabelis MW (1988) How plants obtain predatory mites as bodyguards. *Neth J Zool* 38:148–165
- Dunne T, Doyle J, Parker M (1984) A review of the incidence and effects of unusual marine blooms in Irish coastal waters. *ICES Spec Meet ICES CM 1984/D* 9:4
- Dutz J, Klein Breteler WCM, Kramer G (2005) Inhibition of copepod feeding by exudates and transparent exopolymer particles (TEP) derived from a *Phaeocystis globosa* dominated phytoplankton community. *Harmful Algae* 4:929
- Dutz J, Koski M (2006) Trophic significance of solitary cells of the prymnesiophyte *Phaeocystis globosa* depends on cell type. *Limnol Oceanogr* 51:1230–1238
- Edwardsen B, Paasche E (1998) Bloom dynamics and physiology of *Prymnesium* and *Chrysochromulina*. In: Anderson DM, Cembella AD, Hallegraeff GM (eds) *The physiological ecology of harmful algal blooms*. Springer Verlag, Heidelberg, pp 193–208
- Eilertsen HC, Raa J (1995) Phytoplankton toxins in sea-water. *J Mar Biotechnol* 3:115–119
- Escaravage V, Peperzak L, Prins TC, Peeters JCH, Joordens JCA (1995) The development of a *Phaeocystis* bloom in a mesocosm experiment in relation to nutrients, irradiance and coexisting algae. *Ophelia* 42:55–74
- Estep KW, Nejstgaard JC, Skjoldal HR, Rey F (1990) Predation by copepods upon natural populations of *Phaeocystis pouchetii* as a function of the physiological state of the prey. *Mar Ecol Prog Ser* 67:235–249
- Everitt BS, Dunn G (2001) *Applied multivariate data analysis*, 2nd edn. Oxford University Press, New York
- Frangoulis C, Belkhiria S, Goffart A, Hecq J-H (2001) Dynamics of copepod faecal pellets in relation to a *Phaeocystis* dominated phytoplankton bloom: characteristics, production and flux. *J Plankton Res* 23:75–88
- Fransz HG, Gieskes WWC (1984) The unbalance of phytoplankton and copepods in the North Sea. *J Cons Int Explor Mer* 183:218–225
- Fransz HG, Gonzalez SR, Cadée GC, Hansen FC (1992) Long-term change of *Temora longicornis* (copepoda, Calanoida) abundance in a Dutch tidal inlet (Marsdiep) in relation to eutrophication. *Neth J Sea Res* 30:23–32
- Fretter V, Montgomery MC (1968) The treatment of food by prosobranch veligers. *J Mar Biol Ass UK* 48:275–544
- Frost BW (1972) Effect of size and concentration of food particles on the feeding behaviour of the marine planktonic copepod *Calanus pacificus*. *Limnol Oceanogr* 17:805–815
- Gasparini S, Daro MH, Antajan E, Tackx M, Rousseau V, Parent J-Y, Lancelot C (2000) Mesozooplankton grazing during the *Phaeocystis globosa* bloom in the southern bight of the North Sea. *J Sea Res* 43:345–356
- Goffart A, Catalano G, Hecq JH (2000) Factors controlling the distribution of diatoms and *Phaeocystis* in the Ross Sea. *J Mar Syst* 27:161
- González JM, Sherr EB, Sherr BF (1990) Size-selective grazing on bacteria by natural assemblages of estuarine flagellates and ciliates. *Appl Environ Microbiol* 56:583–589
- Gorsky G, Chretiennot-Dinet M-J, Blanchot J, Palazzoli I (1999) Picoplankton and nanoplankton aggregation by appendicularians: fecal pellet contents of *Megalocercus huxleyi* in the equatorial Pacific. *J Geophys Res C* 104:3381–3390
- Greenacre MJ (1984) *Theory and applications of correspondence analysis*. Academic Press, London
- Haberman KL, Quetin LB, Ross RM (2003) Diet of the Antarctic krill (*Euphausia superba* Dana): I. Comparisons of grazing on *Phaeocystis antarctica* (Karsten) and *Thalassiosira antarctica* (Comber). *J Exp Mar Biol Ecol* 283:79–95
- Hamm CE, Reigstad M, Wexels Riser C, Mühlebac A, Wassmann P (2001) On the trophic fate of *Phaeocystis pouchetii*. VII. Sterols and fatty acids reveal sedimentation of *P. pouchetii*-derived organic matter via krill fecal strings. *Mar Ecol Prog Ser* 209:55–69
- Hamm CE, Rousseau V (2003) Composition, assimilation and degradation of *Phaeocystis globosa*-derived fatty acids in the North Sea. *J Sea Res* 50:271
- Hamm CE, Simson DA, Merkel R, Smetacek V (1999) Colonies of *Phaeocystis globosa* are protected by a thin but tough skin. *Mar Ecol Prog Ser* 187:101–111
- Hansen B, Bjørnsen PK, Hansen PJ (1994a) The size ratio between planktonic predators and their prey. *Limnol Oceanogr* 39:395–403
- Hansen B, Tande KS, Berggreen UC (1990) On the trophic fate of *Phaeocystis pouchetii* (Hariot). III. Functional responses in grazing demonstrated on juvenile stages of *Calanus finmarchicus* (Copepoda) fed diatoms and *Phaeocystis*. *J Plankton Res* 12:1173–1187
- Hansen B, Verity P, Falkenhaus T, Tande KS, Norrbin F (1994b) On the trophic fate of *Phaeocystis pouchetii* (Hariot). V. Trophic relationships between *Phaeocystis* and zooplankton: an assessment of methods and size dependence. *J Plankton Res*. 16:487–511
- Hansen E, Eilertsen HC, Ernstsens A, Genevieve A-M (2003) Anti-mitotic activity towards sea urchin embryos in extracts from the marine haptophycean *Phaeocystis pouchetii* (Hariot) Lagerheim collected along the coast of northern Norway. *Toxicol* 41:803–812
- Hansen E, Ernstsens A, Eilertsen HC (2004) Isolation and characterisation of a cytotoxic polyunsaturated aldehyde from the marine phytoplankton *Phaeocystis pouchetii* (Hariot) Lagerheim. *Toxicology* 199:207–217
- Hansen FC (1992) Zooplankton-Grazing an *Phaeocystis* mit besonderer Berücksichtigung der Calanoiden Copepoden.

- Institut für Meereskunde Kiel, Christian-Albrechts-Universität, Kiel, Kiel, p 137
- Hansen FC (1995) Trophic interactions between zooplankton and *Phaeocystis* cf. *globosa*. Helgol Meeresunters 49: 283–293
- Hansen FC, Reckermann M, Klein Breteler WCM, Riegman R (1993) *Phaeocystis* blooming enhanced by copepod predation on protozoa: evidence from incubation experiments. Mar Ecol Prog Ser 102:51–57
- Hansen FC, van Boekel WHM (1991) Grazing pressure of the calanoid copepod *Temora longicornis* on a *Phaeocystis* dominated spring bloom in a Dutch tidal inlet. Mar Ecol Prog Ser 78:123–129
- Hansen PJ, Bjørnsen PK, Hansen BW (1997) Zooplankton grazing and growth: scaling within the 2–2,000- μ m body size range. Limnol Oceanogr 42:687–704
- Hay ME, Fenical W (1988) Marine plant-herbivore interactions: the ecology of chemical defense. Annu Rev Ecol Syst 19:111–145
- Hay ME, Parker JD, Burkepile DE, Caudill CC, Wilson AE, Hallinan ZP, Chequer AD (2004) Mutualisms and aquatic community structure: the enemy of my enemy is my friend. Annu Rev Ecol Syst 35:175–197
- Hegarty SG, Villareal TA (1988) Effects of light level and N:P supply ratio on the competition between *Phaeocystis* cf. *pouchetii* (Hariot) Lagerheim (Prymnesiophyceae) and five diatom species. J Exp Mar Biol Ecol 226:241–258
- Hollowday ED (1949) A preliminary report on the Plymouth marine and brackish-water rotifera. J Mar Biol Ass UK 28:239–253
- Houde SEL, Roman MR (1987) Effects of food quality on the functional ingestion response of the copepod *Acartia tonsa*. Mar Ecol Prog Ser 40:69–77
- Huntley M, Tande K, Eilertsen HC (1987) On the trophic fate of *Phaeocystis pouchetii* (Hariot). II. Grazing rates of *Calanus hyperboreus* (Krøyer) on diatoms and different size categories of *Phaeocystis pouchetii*. J Exp Mar Biol Ecol 110:197–212
- Irigoin X, Titelman J, Harris RP, Harbour D, Castellani C (2003) Feeding of *Calanus finmarchicus* nauplii in the Irminger Sea. Mar Ecol Prog Ser 262:193–200
- Jakobsen HH, Tang KW (2002) Effects of protozoan grazing on colony formation in *Phaeocystis globosa* (Prymnesiophyceae) and the potential costs and benefits. Aquat Microb Ecol 27:261–227
- Jebram D (1980) Prospection for a sufficient nutrition for the cosmopolitan marine bryozoan *Electra pilosa* (Linnaeus). Zool Jahrb Syst 107:368–390
- Jones PGV, Haq SM (1963) The distribution of *Phaeocystis* in the Eastern Irish Sea. J Cons Int Explor Mer 28:8–20
- Jonker R, Groben R, Tarran G, Medlin L, Wilkins M, Garcia L, Zabala L, Boddy L (2000) Automated identification and characterisation of microbial populations using flow cytometry: the AIMS project. Sci Mar 64:225–234
- Kamermans P (1994) Nutritional value of solitary cells and colonies of *Phaeocystis* sp. for the bivalve *Macoma balthica* (L.). Ophelia 39:35–44
- Keller MD, Bellows WK, Guillard RRL (1989) Dimethyl sulphide production in marine phytoplankton. Acs Symp Ser 393:167–182
- Klein Breteler WCM, Koski M (2003) Development and grazing of *Temora longicornis* (Copepoda, Calanoida) nauplii during nutrient limited *Phaeocystis globosa* blooms in mesocosms. Hydrobiologia 491:185–192
- Kopp J (1978) Les phénomènes d’eaux colorées ayant affecté le littoral Bas-Normand au cours du mois de juin 1978. Inst Sci Tech Pêches Mar, Centre de Quistreham: 30
- Kornmann VP (1955) Beobachtungen an *Phaeocystis*-Kulturen. Helgoländer wiss Meeresunters 5:218–233
- Koski M, Dutz J, Klein Breteler WCM (2005) Selective grazing of *Temora longicornis* in different stages of a *Phaeocystis globosa* bloom: a mesocosm study. Harmful Algae 4:915
- Krogstad PK (1989) Produksjon av planteplankton til bruk som fôr i intensivt oppdrett av torskelarver (*Gadhus morhua* L.) med referanse i ekstensivt oppdrettssystem (poll). Institute of Fisheries, University of Tromsø, Tromsø, p 147
- Lancelot C, Keller MD, Rousseau V, Smith WOJ, Mathot S (1998) Autecology of the marine haptophyte *Phaeocystis* sp. In: Anderson DM, Cembella AD, Hallegraeff GM (eds) The physiological ecology of harmful algal blooms. Springer Verlag, Berlin, pp 209–224
- Lancelot C, Rousseau V (1994) Ecology of *Phaeocystis*: the key role of colony forms. In: Green JC, Leadbeater BSC (eds) The haptophyte algae, Clarendon Press, Oxford, pp 229–245
- Landry MR, Hassett RP (1982) Estimating the grazing impact of marine micro-zooplankton. Mar Biol 67:283–288
- Landry MR, Selph KE, Brown SL, et al (2002) Seasonal dynamics of phytoplankton in the Antarctic Polar Front region at 170° W Deep Sea Res Part II Top Stud Oceanogr 49:1843–1865
- Larsen A, Fonnes Flaten GA, Sandaa R-A, Castberg T, Thyrraug R, Erga SR, Jacquet S, Bratbak G (2004) Spring phytoplankton bloom dynamics in Norwegian coastal waters: Microbial community succession and diversity. Limnol Oceanogr 49:180–190
- Lebart L, Morineau A, Lambert T (1988) SPAD.N Système portable pour l’analyse des données. Manuel de référence. (SPAD.N a software for Correspondence Analysis User’s guide). CISIA, Saint-Mandé, France
- Lebart L, Morineau A, Piron M (2000) Statistique exploratoire multidimensionnelle. Dunod, Paris
- Lebour MV (1920) The food of young fish. No. III (1919). J Mar Biol Ass UK 12:261–324
- Lebour MV (1922) The food of planktonic organisms. J Mar Biol Ass UK 12:644–677
- Legendre P, Legendre L (1998) Numerical Ecology. Elsevier, Amsterdam
- Levinsen H, Turner JT, Nielsen TG, Hansen BW (2000) On the trophic coupling between protists and copepods in arctic marine ecosystems. Mar Ecol Prog Ser 204:65–77
- Lewis WMJ (1986) Evolutionary interpretations of allelochemical interactions in phytoplankton algae. Am Nat 127:184–194
- Liss PS, Malin G, Turner SM, Holligan PM (1994) Dimethyl sulphide and *Phaeocystis*: a review. J Mar Syst 5:41
- Long JD (2004) Plasticity of consumer-prey interactions in the sea: chemical signaling, learned aversions, and ecological consequences. Biology. Georgia Institute of Technology, Atlanta, p 111

- Long JD, Hay ME (2006) When intraspecific exceeds interspecific variance: effects of phytoplankton morphology and growth phase on copepod feeding and fitness. *Limnol Oceanogr* 51:988–996
- Løken S (1990) Næringsøkologi og vekst hos torskelarver (*Gadus morhua* L.) i poll og intensive system. Institute of Fisheries. University of Tromsø, Tromsø, p 86
- Lucas CE, Henderson GTD (1936) On the association of jellyfish and other organisms with catches of herring. *J Mar Biol Ass U K* 21:293–304
- Lürling M (2003) Phenotypic plasticity in the green algae *Desmodesmus* and *Scenedesmus* with special reference to the induction of defensive morphology. *Ann Limnol* 39:85–101
- Lürling M, Van Donk E (1997) Morphological changes in *Scenedesmus* induced by infochemicals released in situ from zooplankton grazers. *Limnol Oceanogr* 42:783–788
- Mathot S, Smith WO, Carlson CA, Garrison DL, Gowing MM, Vickers CL (2000) Carbon partitioning within *Phaeocystis antarctica* (Prymnesiophyceae) colonies in the Ross Sea, Antarctica. *J Phycol* 36:1049–1056
- Medlin L and Zingone A (this volume) A review: the genus *Phaeocystis* and its species. *Biogeochemistry*
- Meixner R (1981) Herzmuschel-Suche mit FA-Niobe im ostfriesischen Wattenmeer. *Inf Fischwirtsch* 28:133–134
- Metz C (1998) Feeding of *Oncaea curvata* (Poecilostomatoida, Copepoda). *Mar Ecol Prog Ser* 169:229–235
- Nejstgaard JC, Frischer ME, Raule CL, Gruebel R, Kohlberg KE, Verity PG (2003) Molecular detection of algal prey in copepod guts and faecal pellets. *Limnol Oceanogr Methods* 1:29–38
- Nejstgaard JC, Frischer ME, Simonelli P, Brakel M, Sazhin A, Artigas LF (2005) Comparison of molecular and traditional methods to quantify selective copepod feeding in situ ASLO, Summer Meeting. Santiago de Compostela, Spain
- Nejstgaard JC, Frischer ME, Verity PG, Anderson JT, Jacobsen A, Zirbel MJ, Larsen A, Martínez-Martínez J, Sazhin AF, Walters T, Bronk DA, Whipple SJ, Borett SR, Patten BC, Long JD (2006) Temporal patterns in planktonic food web development in mesocosms with added nutrients and *Phaeocystis*. *Mar Ecol Prog Ser* 321:99–121
- Nejstgaard JC, Naustvoll L-J, Sazhin A (2001) Correcting for underestimation of microzooplankton grazing in bottle incubation experiments with mesozooplankton. *Mar Ecol Prog Ser* 221:59–75
- Nevitt GA, Bonadonna F (2005) Sensitivity to dimethyl sulphide suggests a mechanism for olfactory navigation by seabirds. *Biol Lett* 1:303–305
- Nevitt GA, Veit RR, Kareiva P (1995) Dimethyl sulfide as a foraging cue for Antarctic Procellariiform seabirds. *Nature* 376:680–682
- Nicholls AG (1935) The larval stages of *Longipedia coronata* Claus, *L. scotti* G. O. Sars, and *L. minor* T. and A. Scott, with a description of the male of *L. scotti*. *J Mar Biol Ass UK* 20:29–45
- Nichols PD, Skerratt JH, Davidson A, Burton H, McMeekin TA (1991) Lipids of cultured *Phaeocystis pouchetii*: signatures for food-web, biogeochemical and environmental studies in Antarctica and the Southern-Ocean. *Phytochemistry* 30:3209–3214
- Nielsen TG, Kjørboe T, Bjørnsen PK (1990) Effects of a *Chrysochromulina polyylepis* subsurface bloom on the planktonic community. *Mar Ecol Prog Ser* 62:1–2
- Noordkamp DJB, Gieskes WWC, Gottschal JC, Forney LJ, van Rijssel M (2000) Acrylate in *Phaeocystis* colonies does not affect the surrounding bacteria. *J Sea Res* 43:287–296
- Peperzak L (2002) The wax and wane of *Phaeocystis globosa* blooms. *Wiskunde en Natuurwetenschappen. Rijksuniversiteit Groningen, Groningen*, p 254
- Peperzak L, Colijn F, Vrieling EG, Gieskes WWC, Peeters JCH (2000) Observations of flagellates in colonies of *Phaeocystis globosa* (Prymnesiophyceae); a hypothesis for their position in the life cycle. *J Plankton Res* 22:2181–2203
- Peperzak LM, Colijn F, Gieskes WWC, Peeters JCH (1998) Development of the diatom-*Phaeocystis* spring bloom in the Dutch coastal zone of the North Sea: the silicon depletion versus the daily irradiance threshold hypothesis. *J Plankton Res* 20:517
- Petri G, Donat FH, Vareschi E (1999) Investigations on uptake and utilization of *Phaeocystis globosa* colonies by the blue mussel *Mytilus edulis*. *Senckenb Marit* 29:117–118
- Petri G, Vareschi E (1997) Utilization of *Phaeocystis globosa* colonies by young *Mytilus edulis*. *Arch Fish Mar Res* 45:77–91
- Pieters H, Kluytmans JH, Zandee DI, Cadée GC (1980) Tissue composition and reproduction of *Mytilus edulis* in relation to food availability. *Neth J Sea Res* 14:349–361
- Ploug H, Stolte W, Jørgensen BB (1999) Diffusive boundary layers of the colony-forming plankton alga *Phaeocystis* sp—implications for nutrient uptake and cellular growth. *Limnol Oceanogr* 44:1959–1967
- Pohnert G (2004) Chemical defense strategies of marine organisms. In: Schulz SE (ed) *Topics in current chemistry*. Springer-Verlag GmbH, Berlin, pp 179–219
- Pohnert G (2005) Diatom/copepod interactions in plankton: the indirect chemical defense of unicellular algae. *ChemBioChem* 6:946–959
- Porter KG (1976) Enhancement of algal growth and productivity by grazing zooplankton. *Science* 192:1332–1334
- Reigstad M and Wassmann P (this volume) Does *Phaeocystis* spp. contribute significantly to vertical export of biogenic matter? *Biogeochemistry*
- Richardson MG, Whitaker TM (1979) An Antarctic fast-ice food chain: observations on the interaction of the amphipod *Pontogeneia antarctica* Chevreux with ice-associated micro-algae. *Br Antarct Surv Bull* 47:107–115
- Riebesell U, Reigstad M, Wassmann P, Noji T, Passow U (1995) On the trophic fate of *Phaeocystis pouchetii* (Hariot): 6. Significance of *Phaeocystis*-derived mucus for vertical flux. *Neth. J Sea Res* 33:193–203
- Rogers SI, Lockwood SJ (1990) Observations on coastal fish fauna during a spring bloom of *Phaeocystis pouchetii* in the eastern Irish Sea. *J Mar Biol Ass UK* 70:249–253
- Rousseau V, Becquevort S, Parent J-Y, Gasparini S, Daro M-H, Tackx M, Lancelot C (2000) Trophic efficiency of the planktonic food web in a coastal ecosystem dominated by *Phaeocystis* colonies. *J Sea Res* 43:357–372
- Rousseau V, Chrétiennot-Dinet M-J, Jacobsen A, Verity PG, Whipple SJ (this volume) The life cycle of *Phaeocystis*: state of knowledge and presumptive role in ecology. *Biogeochemistry*. doi:10.1007/s10533-007-9085-3

- Rousseau V, Leynaert A, Daoud N, Lancelot C (2002) Diatom succession, silicification and silicic acid availability in Belgian coastal waters (Southern North Sea). *Mar Ecol Prog Ser* 236:61–73
- Rousseau V, Mathot S, Lancelot C (1990) Calculating carbon biomass of *Phaeocystis* sp. from microscopic observations. *Mar Biol* 107:305–314
- Rousseau V, Vault D, Casotti R, Cariou V, Lenz J, Gunkel J, Baumann M (1994) The life cycle of *Phaeocystis* (Prymnesiophyceae): evidence and hypotheses. *J Mar Syst* 5:23
- Rynearson TA, Armbrust EV (2005) Maintenance of clonal diversity during a spring bloom of the centric diatom *Ditylum brightwellii*. *Mol Ecol* 14:1631–1640
- Sargent JR, Eilertsen HC, Falk-Petersen S, Taasen JP (1985) Carbon assimilation and lipid production in phytoplankton in northern Norwegian fjords. *Mar Biol* 85:109–116
- Savage RE (1930) The influence of *Phaeocystis* on the migrations of the herring. *Fish Invest II* 12:1–14
- Savage RE (1932) *Phaeocystis* and herring shoals. *J Ecol* 20:326–340
- Sazhin AF, Artigas LF, Nejstgaard JC, Frischer ME (this volume) The colonization of two *Phaeocystis* species (Prymnesiophyceae) by pennate diatoms and other protists: a significant contribution to colony biomass. *Biogeochemistry*
- Schnack SB (1983) On the feeding of copepods on *Thalassiosira partheneia* from the Northwest African upwelling area. *Mar Ecol Prog Ser* 11:49–53
- Schoemann V, Becquevort S, Stefels J, Rousseau V, Lancelot C (2005) *Phaeocystis* blooms in the global ocean and their controlling mechanisms: a review. *J Sea Res* 53:43
- See JH, Campbell L, Richardson TL, Pinckney JL, Shen R, Guinasso NL (2005) Combining new technologies for determination of phytoplankton community structure in the northern Gulf of Mexico. *J Phycol* 41:305–310
- Selph KE, Landry MR, Allen CB, et al (2001) Microbial community composition and growth dynamics in the Antarctic Polar Front and seasonal ice zone during late spring 1997. *Deep Sea Res Part II Top Stud Oceanogr* 48:4059–4080
- Seuront L, Lacheze C, Doubell MJ, Seymour JR, Mitchell JG (this volume) The influence of *Phaeocystis globosa* bloom dynamics on microscale spatial patterns of phytoplankton biomass and bulk-phase seawater viscosity. *Biogeochemistry*
- Sheppard SK, Harwood JD (2005) Advances in molecular predator–prey ecology. *Funct Ecol* 19:751–762
- Shields AR, Smith WJO (2005) A novel technique to examine the role of colonial *Phaeocystis antarctica* in the microbial loop of the Ross Sea, Antarctica. SCOR working group #120. Final meeting: *Phaeocystis*, major link in the biogeochemical cycling of climate-relevant elements. University of Groningen, Haren, the Netherlands, p 14
- Sieburth JM (1960) Acrylic acid, an “antibiotic” principle in *Phaeocystis* blooms in antarctic waters. *Science* 132:676–677
- Sieburth JM (1961) Antibiotic properties of acrylic acid, a factor in the gastrointestinal antibiosis of polar marine animals. *J Bacteriol* 82:72–79
- Smaal AC, Twisk F (1997) Filtration and absorption of *Phaeocystis* cf. *globosa* by the mussel *Mytilus edulis* L. *J Exp Mar Biol Ecol* 209:33–46
- Smayda TJ (1973) The growth of *Skeletonema costatum* during a winter-spring bloom in Narragansett Bay, Rhode Island. *Norw J Bot* 20:219–247
- Smayda TJ (1997) What is a bloom? A commentary. *Limnol Oceanogr* 42:1132–1136
- Sournia A (1988) *Phaeocystis* (Prymnesiophyceae): how many species? *Nova Hedwigia* 47:211–217
- Stabell OB, Aanesen RT, Eilertsen HC (1999) Toxic peculiarities of the marine alga *Phaeocystis pouchetii* detected by in vivo and in vitro bioassay methods. *Aquat Toxicol* 44:279–288
- Stefels J (2000) Physiological aspects of the production and conversion of DMSP in marine algae and higher plants. *J Sea Res* 43:183–197
- Stefels J, Steinke M, Turner S, Gill M and Belviso S (this volume) Environmental constraints on the production of the climatically active gas dimethylsulphide (DMS) and implications for ecosystem modelling. *Biogeochemistry*. doi:10.1007/s10533-007-9091-5
- Steinke M, Malin G, Liss PS (2002) Trophic interactions in the sea: an ecological role for climate relevant volatiles? *J Phycol* 38:630–638
- Steinke M, Stefels J, Stamhuis E (2006) Dimethyl sulfide triggers search behavior in copepods. *Limnol Oceanogr* 51:1925–1930
- Stelfox-Widdicombe CE, Archer SD, Burkill PH, Stefels J (2004) Microzooplankton grazing in *Phaeocystis* and diatom-dominated waters in the southern North Sea in spring. *J Sea Res* 51:37
- Strom S, Wolfe G, Slajer A, Lambert S, Clough J (2003) Chemical defense in the microplankton II: Inhibition of protist feeding by β -dimethylsulfoniopropionate (DMSP). *Limnol Oceanogr* 48:230–237
- Strom SL, Loukos H (1998) Selective feeding by protozoa: model and experimental behaviors and their consequences for population stability. *J Plankton Res* 20:831–846
- Symondson WOC (2002) Molecular identification of prey in predator diets. *Mol Ecol* 11:627–641
- Tande KS, Båmstedt U (1987) On the trophic fate of *Phaeocystis pouchetii* I. Copepod feeding rates on solitary cells and colonies of *P. pouchetii*. *Sarsia* 72:3–4
- Tang KW (2003) Grazing and colony size development in *Phaeocystis globosa* (Prymnesiophyceae): the role of a chemical signal. *J Plankton Res* 25:831–842
- Tang KW, Jakobsen HH, Visser AW (2001) *Phaeocystis globosa* (Prymnesiophyceae) and the planktonic food web: feeding, growth, and trophic interactions among grazers. *Limnol Oceanogr* 46:1860–1870
- Tang KW, Simó R (2003) Trophic uptake and transfer of DMSP in simple planktonic food chains. *Aquat Microb Ecol* 31:193–202
- Thingstad F, Billen G (1994) Microbial degradation of *Phaeocystis* material in the water column. *J Mar Syst* 5:55
- Troedsson CG, Frischer ME, Nejstgaard JC, Thompson EM (2007) Molecular quantification of differential ingestion and particle trapping rates by the appendicularian *Oikopleura dioica* as a function of prey size and shape. *Limnol Oceanogr* 52:416–427
- Tungaraza C, Rousseau V, Brion N, Lancelot C, Gichuki J, Baeyens W, Goeyens L (2003) Contrasting nitrogen uptake by diatom and *Phaeocystis*-dominated phytoplankton

- assemblages in the North Sea. *J Exp Mar Biol Ecol* 292:19–41
- Turner JT (1994) Planktonic Copepods of Boston Harbor, Massachusetts Bay and Cape Cod Bay, 1992. *Hydrobiologia* 293:405–413
- Turner JT, Ianora A, Esposito F, Carotenuto Y, Miralto A (2002) Zooplankton feeding ecology: does a diet of *Phaeocystis* support good copepod grazing, survival, egg production and egg hatching success? *J Plankton Res* 24:1185–1195
- van Boekel WHM, Hansen FC, Riegman R, Bak RPM (1992) Lysis-induced decline of a *Phaeocystis* spring bloom and coupling with the microbial foodweb. *Mar Ecol Prog Ser* 81:269–276
- van Donk E, Luerling M, Hessen DO, Lokhorst GM (1997) Altered cell wall morphology in nutrient-deficient phytoplankton and its impact on grazers. *Limnol Oceanogr* 42:357–364
- van Rijssel M, Alderkamp A-C, Nejtgaard JC, Sazhin AF, Verity PG (this volume) Haemolytic activity of living *Phaeocystis pouchetii* during mesocosm blooms. *Biogeochemistry*. doi:10.1007/s10533-007-9095-1
- van Rijssel M, Hamm CE, Gieskes WWC (1997) *Phaeocystis globosa* (Prymnesiophyceae) colonies: hollow structures built with small amounts of polysaccharides. *Eur J Phycol* 32:185–192
- Veldhuis MJW, Colijn F, Venekamp LAH (1986) The spring bloom of *Phaeocystis pouchetii* (Haptophyceae) in Dutch coastal waters. *Neth J Sea Res* 20:37–48
- Veldhuis MJW, Wassmann P (2005) Bloom dynamics and biological control of a high biomass HAB species in European coastal waters: a *Phaeocystis* case study. *Harm Algae* 4:805
- Verity PG (1988) Chemosensory behavior in marine planktonic ciliates. *Bull Marine Sci* 43:772–782
- Verity PG (2000) Grazing experiments and model simulations of the role of zooplankton in *Phaeocystis* food webs. *J Sea Res* 43:317–343
- Verity PG, Paffenhöfer G-A (1996) On assessment of prey ingestion by copepods. *J Plankton Res* 18:1767–1779
- Verity PG, Smayda TJ (1989) Nutritional value of *Phaeocystis pouchetii* (Prymnesiophyceae) and other phytoplankton for *Acartia* spp. (Copepoda): Ingestion, egg production, and growth of nauplii. *Mar Biol* 100:161–171
- Vestheim H, Edvardsen B, Kaartvedt S (2005) Assessing feeding of a carnivorous copepod using species specific PCR. *Mar Biol* 147:381–385
- Weisse T, Grimm N, Hickel W, Martens P (1986) Dynamics of *Phaeocystis pouchetii* blooms in the Wadden Sea of Sylt (German Bight, North Sea). *Est Coast Shelf Sci* 23:171
- Weisse T, Scheffel-Möser U (1990) Growth and grazing loss rates in single-celled *Phaeocystis* sp. (Prymnesiophyceae). *Mar Biol* 106:153–158
- Weisse T, Tande K, Verity P, Hansen F, Gieskes W (1994) The trophic significance of *Phaeocystis* blooms. *J Mar Syst* 5:67–79
- Wolfe GV, Levasseur M, Cantin G, Michaud S (2000) DMSP and DMS dynamics and microzooplankton grazing in the Labrador Sea: application of the dilution technique. *Deep-Sea Res Part I Oceanogr Res Pap* 47:2243–2264
- Wolfe GV, Steinke M (1996) Grazing-activated production of dimethyl sulfide (DMS) by two clones of *Emiliania huxleyi*. *Limnol Oceanogr* 41:1151–1160
- Wolfe GV, Steinke M, Kirst GO (1997) Grazing-activated chemical defence in a unicellular marine alga. *Nature* 387:894–897
- Wulff A (1934) Über Hydrographie und Oberflächenplankton nebst Verbreitung von *Phaeocystis* in der Deutschen Bucht im Mai 1933. *Ber Deutsch Wiss Komm Meeresforsch NF* 7:343–350
- Yoshida T, Hairston NG, Ellner SP (2004) Evolutionary trade-off between defence against grazing and competitive ability in a simple unicellular alga, *Chlorella vulgaris*. *Proc Roy Soc Lond Ser B Biol Sci* 271:1947–1953