



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Injury

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa26356>

Paper:

O'Reilly, G., Gabbe, B., Moore, L. & Cameron, P. (2016). Classifying, measuring and improving the quality of data in trauma registries: A review of the literature. *Injury*

<http://dx.doi.org/10.1016/j.injury.2016.01.007>


This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Accepted Manuscript

Title: Classifying, measuring and improving the quality of data in trauma registries: a review of the literature

Author: Gerard M O'Reilly Belinda Gabbe Lynne Moore
Peter A Cameron



PII: S0020-1383(16)00009-7
DOI: <http://dx.doi.org/doi:10.1016/j.injury.2016.01.007>
Reference: JINJ 6566

To appear in: *Injury, Int. J. Care Injured*

Received date: 24-7-2015
Revised date: 12-12-2015
Accepted date: 9-1-2016

Please cite this article as: O'Reilly GM, Gabbe B, Moore L, Cameron PA, Classifying, measuring and improving the quality of data in trauma registries: a review of the literature, *Injury* (2016), <http://dx.doi.org/10.1016/j.injury.2016.01.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Classifying, measuring and improving the quality of data in trauma registries: a review of the literature

(ii) Authors**1. Dr Gerard M O'Reilly (1,2)**

MBBS FACEM MPH MBiostat AStat

1. Adjunct Senior Research Fellow
2. Emergency Staff Specialist and International Operations Manager

2. Professor Belinda Gabbe (1,3)

BPhysio(Hons), M App Sc, M Biostat, PhD

1. Head of Pre-hospital, Emergency and Trauma Research
2. Professorial Chair, College of Medicine, Swansea University

3. Professor Lynne Moore (4)

PhD

1. Associate Professor

4. Professor Peter A Cameron (1,2,5)

MBBS MD FACEM

1. and 2. Professor Emergency Medicine, The Alfred Hospital, Monash University
5. Visiting Professor Emergency Medicine, Hamad Medical Corporation

(iii) Affiliations

1. Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Commercial Rd, Melbourne, Australia, 3004 (1, 2 and 4)
2. Emergency and Trauma Centre, Alfred Health, Commercial Rd, Melbourne, Victoria, Australia, 3004 (1 and 4)
3. Swansea University, United Kingdom (2)
4. University of Laval, Quebec, Canada (3)
5. Emergency Medicine, Hamad Medical Corporation, Doha, Qatar (4)

(iv) Source of Work

Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Faculty of Medicine, Monash University (Alfred Campus), Commercial Rd, Melbourne, Australia, 3004

(iv) Correspondence

Dr Gerard O'Reilly

**Emergency and Trauma Centre, The Alfred, Commercial Rd, Melbourne, Victoria, 3004,
Australia**

gerard.oreilly@monash.edu

Phone: +61 (0) 438 338 414

Title

Classifying, measuring and improving the quality of data in trauma registries: a review of the literature

Key Words

trauma registry, trauma database, trauma databank, injury registry, injury database, injury databank, data quality, accuracy, completeness, capture.

Structured Abstract**Introduction**

Globally, injury is a major cause of death and disability. Improvements in trauma care have been driven by trauma registries. The capacity of a trauma registry to inform improvements in the quality of trauma care is dependent upon the quality of data. The literature on data quality in disease registries is inconsistent and ambiguous; methods used for classifying, measuring, and improving data quality are not standardised. The aim of this study was to review the literature to determine the methods used to classify, measure and improve data quality in trauma registries.

Methods

A scoping review of the literature was performed. Databases were searched using the term “trauma registry” and its synonyms, combined with multiple terms denoting data quality. There was no restriction on year. Full-length manuscripts were included if the classification, measurement or

improvement of data quality in one or more trauma registries was a study objective. Data were abstracted regarding registry demographics, study design, data quality classification, and the reported methods used to measure and improve the pre-defined data quality dimensions of accuracy, completeness and capture.

Results

Sixty-nine publications met the inclusion criteria. Four publications classified data quality. The most frequently described methods for measuring data accuracy (n=47) were checks against other datasets (n=18) and checks of injury coding (n=17). The most frequently described methods for measuring data completeness (n=47) were the percentage of included cases, for a given variable or list of variables, for which there was an observation in the registry (n=29). The most frequently described methods for measuring data capture (n=37) were the percentage of cases in a linked reference dataset that were also captured in the primary dataset being evaluated (n=24). Most publications dealing with the measurement of a dimension of data quality did not specify the methods used; most publications dealing with the improvement of data quality did not specify the dimension being targeted.

Conclusion

The classification, measurement and improvement of data quality in trauma registries is inconsistent. To maintain confidence in the usefulness of trauma registries, the metrics and reporting of data quality need to be standardised.

Manuscript

Introduction

Injury is a major cause of disability and death. The burden of injury is of particular concern amongst young and working-age adults and in developing countries¹. In developed countries, trauma care systems have reduced morbidity and mortality from injury²⁻⁸. Improvements in the delivery of trauma care have been driven by the information contained in trauma registries, which are dedicated to storing data on the outcomes and processes of trauma care⁹. Trauma registries have played a very important role in developed trauma systems for more than thirty years¹⁰. Conversely, the use of trauma registries in developing countries, where the burden of injury is most prominent, remains sporadic^{11,12}.

The capacity of a trauma registry to inform improvements in the quality of trauma care depends upon the quality of its data. Incomplete and erroneous data are a threat to the use of trauma registries for comparing and benchmarking systems of trauma care^{10,13}. If the quality of data in a trauma registry is unknown, questionable or poor, it will be rendered less valuable as a tool for improving the quality of trauma care. Systems of trauma care, at the single-hospital or multi-hospital level, need methods to measure and improve the quality of new or established trauma registries; users of observational research based on registry data need to be confident of the quality of the source data.¹⁴

For trauma registries, there is no single definition or classification of data quality. Amongst datasets in general, the taxonomy of data quality varies considerably according to the type and intended use of the dataset being considered¹⁵⁻²¹. For example, one publication defined data quality as “fitness for purpose” and listed its six dimensions as: relevance, accuracy and reliability, timeliness, accessibility,

interpretability and coherence¹⁵. Another publication listed seven dimensions of data quality: valid, complete, consistent, unique, timely, accurate and precise¹⁶. Wang and Strong, in their iterative approach to developing a framework of data quality grouped a total of 15 dimensions under four categories of data quality: intrinsic, contextual, representational and accessibility¹⁷.

The literature dealing with data quality amongst *health*-related datasets is similarly inconsistent¹⁸⁻¹⁹. Several publications have provided a relatively simple set of definitions and classification for data quality^{20,21}. Sorensen et al included accuracy and completeness of variables, and completeness of registration, amongst their list of dimensions²⁰. Similarly, Arts et al noted that the two most cited data quality attributes are accuracy (the extent to which registered data are in conformity to the truth) and completeness (the extent to which all necessary data that could have been registered have actually been registered)²¹.

The aim of this study was to review the accessible literature to determine the methods used to classify, measure and improve data quality in trauma registries.

Methods

A scoping review of the literature was performed. Abstracts were sourced by searching MEDLINE, Embase and CINAHL databases on 12 February 2015. There were no year restrictions placed on the search. Key words used for the search were: “trauma registry”, “trauma registries”, “trauma database*”, “trauma databank*”, “injury registry”, “injury registries”, “injury database*”, and “injury databank*”. An additional subject heading search was undertaken in: MEDLINE using “Wounds and Injuries” AND “Registries”; Embase using “Registry” AND “Injury” OR “Trauma”; and CINAHL using “Registries, Trauma”. Finally, in each database, the

results of the search strategy described above were combined (using “AND”) with each of the keywords denoting data quality and its dimensions in previous literature¹⁵⁻²¹ (see Table 1).

From the results of the search, the authors included those full-length manuscripts which met the following criteria:

1. The trauma registry was a dedicated database, prospectively collecting data regarding injured patients at one or more health facilities, and
2. The classification, measurement or improvement of data quality in one or more trauma registries was a study objective.

Data on the following were abstracted from each full-text manuscript:

1. Characteristics of the trauma registry (source country, level of country development, jurisdiction, stage of registry existence)²²
2. Characteristics of the publication (year published, study design)
3. Did the publication provide a classification of data quality?
4. Did the study of trauma registry data quality deal with measuring data quality *or* improving data quality *or* both?
5. What dimensions of trauma registry data quality were named?
6. For those publications dealing with measuring data quality, what dimensions of data quality were measured, how were the measurements defined and what specific metrics were used?
7. For those publications dealing with the improvement of data quality, what dimensions of data quality were targeted and what domains of trauma registry function were described?

For the final question of the trauma registry domains described in publications looking at *improving* data quality (7), a modified version of the Trauma Registry Assessment Tool was used^{11,12,23}. The details of this tool and its validation have been published elsewhere¹². Under the broad headings of

physical resources, human resources and processes it comprises a checklist of the key components or domains which are generic to any trauma registry. The tool has previously been used to capture information on these domains of trauma registries, regardless of their context^{11,12,23}.

For the final two questions listed above (6 and 7), three key dimensions of data quality were chosen on the basis of being commonly cited in the disease registry literature and most clearly defined^{20,21}. For the purposes of this review, they were defined by the authors by merging the approaches of Sorensen et al and Arts et al^{20,21}.

1. Accuracy: the extent to which registered data are in conformity with the truth
(i.e. How true is the value of an observation or the content of a cell in a spreadsheet of data, compared to a gold standard?)
2. Completeness: the extent to which all necessary data have been registered on registered cases
(i.e. Is there an observation for a variable on a registered case?)
3. Capture: the extent to which all necessary cases that could have been registered have actually been registered
(i.e. How complete is the case ascertainment, given the registry inclusion criteria?)

Dimensions of data quality which did not fit under these headings were also recorded and grouped under the heading "Other".

Results

The initial search yielded 147 abstracts; 69 full text manuscripts matched the pre-defined inclusion criteria²⁴⁻⁹². Of the 78 excluded publications, 69 did not have the classification, measurement or improvement of data quality of a trauma registry as a study objective. For 8 of the excluded publications,

no full text journal article existed (mostly published conference abstracts). One potentially eligible abstract was excluded because the full text version was not available in English⁹³.

Although still few in number, the frequency of publications focusing on data quality in trauma registries has increased gradually over time, especially since 1998 (Figure 1). Almost half of the publications were from the US (n=34). Six of the seven publications from a trauma registry based in a developing country (level of development: 2, 3 or 4) were published after 2011. Most publications (n=44) were sourced from multi-hospital registries. More than one quarter of publications (n=18) dealt with data quality in the context of trauma registry establishment or feasibility. The aforementioned results are detailed for each publication in Supplementary File 1.

There were two perspectives of data quality in trauma registries and one literature review which conducted an audit of study methods and approaches to reporting and dealing with missing data in a multi-centre trauma registry^{38,48,53}. Four of the 69 publications provided a general definition or classification of at least one data quality dimension^{49,56,62,85}. Gomez et al defined “case ascertainment” as “systematic differences in the types of patients captured in registries”⁴⁹. Alexandrescu et al referenced Arts et al for the afore-mentioned definitions of accuracy and completeness^{21,56}. Data quality was defined as “the totality of features and characteristics of a data set that bear on its ability to satisfy the needs that result from the intended use of the data”^{21,56}. Hlaing et al classified “errors” into seven groups: “Out-of-Range Time Values”, “False Positive and False Negative Coding Errors”, “Errors of Commission or Omission”, “Duplication of Data”, “Errors in Demographic Data”, “Errors Because of Inconsistency in Coding” and “Errors Because of Incongruence in Coding”⁶². Meeuwisse et al referenced the USA Centers for Disease Control in its approach to classifying and defining the attributes of a surveillance system: simplicity (in structure and ease of use), flexibility (to adapt to changes), acceptability (the willingness of individuals to participate), sensitivity (the proportion of cases detected), predictive value positive

(the proportion detected who actually have the condition under surveillance), representativeness (occurrence over time and distribution) and timeliness (speed or delay between steps)^{85,94}.

All of the publications studied either the measurement (n=67) or improvement (n=62) of data quality. Almost all (n=68) of the publications covered at least one of the key dimensions of accuracy (n=49), completeness (n=48), or capture (n=38); most dealt with at least one other dimension of data quality (n=44). The aforementioned results are detailed for each publication in Supplementary File 2.

There were at least 120 different names for dimensions of data quality used in the 69 publications (Table 2); 83 of these described dimensions other than accuracy, completeness and capture.

Table 3 and 4 provide in detail the approaches used or described in measuring (Table 3) and improving (Table 4) data quality in one or more trauma registries.

Accuracy

In most of the publications (n=47) one or more different methods used to evaluate trauma registry accuracy were described. Simpler methods, which could be applied without additional data, included checks of internal consistency (whether an observation on one variable had a logical relationship with an observation on another related variable) (n=7) and domain checks (whether an observation on one variable was plausible) (n=3). More extensive checks included the evaluation of the inter-rater reliability among data collectors or coders (n=10). Additional approaches to evaluating accuracy included linking with other data sources of the same population (n=18) and verification against the source data (n=6).

Where a “gold standard” was defined, measures of sensitivity and specificity were used; otherwise measures of agreement (e.g. kappa statistic for categorical data; Bland-Altman plot for numerical data) were used. In more than a quarter of the publications considering the evaluation of accuracy, no method was specified (n=13) (Table 3).

Some publications were particularly noteworthy in that they clearly demonstrated the measurement of data accuracy using one or more of the methods described in the previous paragraph^{39,59,72}. Beretta et al performed checks of internal consistency, using the example of “ICP (intracranial pressure) lower than 20 mmHg in a patient without ICP monitoring” and domain, using the example of “hemoglobin oxygen saturation more than 100%”⁵⁹. Read-Allsop et al evaluated inter-rater reliability for AIS (Abbreviated Injury Scale) coding⁷². Following linkage between datasets, Newgard et al first performed checks of internal consistency and domain and then reported accuracy by measuring against an alternative database (sensitivity, specificity, exact agreement, median difference), followed by measures of agreement (kappa, modified kappa, Bland-Altman plot and Intraclass Correlation Coefficient)³⁹.

Few publications dealt with methods which specifically targeted the improvement of data accuracy. Comprehensive approaches to the training and feedback of staff involved in data collection, coding and input were provided by Auerbach et al and Read-Allsop et al^{72,74}. Protetch et al and Beretta et al identified the importance of integrating validation tools (checks of internal consistency and domain) into registry computer software^{52,59}.

Completeness

The majority of publications evaluating data completeness (n=47) measured this as the percentage of included cases, for a given variable or list of variables, for which there was an observation in the registry (n=32). There were several publications, demonstrating the measurement of data completeness in a trauma registry, which were notable for their high level of detail^{24,39,41}. Ringdal et al provided a benchmark level of completeness of 80%, but stressed that this was an arbitrarily chosen threshold and that “no justifications or guidelines for the acceptability of missing data in registry studies exist”⁴¹. Nottidge et al provided a detailed evaluation of data completeness in a feasibility study of a registry in Nigeria²⁴.

A number of the more recent publications examined approaches to handling missing data at the analysis stage, including multiple imputation (MI). MI uses the relationship between variables in collected data to estimate the distribution of the missing observations. Amongst the six publications focused upon improving data completeness for analysis, MI was promoted as the preferred method, but not as a remedy for the important step of rigorous data collection^{36,38,46,48,55,65}.

Capture

There were fewer publications dealing with evaluating case capture when compared to accuracy or completeness (n=37). The most commonly used approach was to link and compare different datasets (n=27). Capture was generally measured as the percentage of cases in the reference dataset that were also captured in the primary dataset being evaluated. Notable examples demonstrating the measurement of data capture using percentage of a reference dataset, were Newgard et al, Schootman et al and Goldberg et al^{43,80,91}.

Few publications were dedicated to methods of improving data capture at the data collection stage. One such publication was Wainiqolo et al³⁴, which provided in detail the implementation of supervision, cross-checking with hospital administrative records, and data collector feedback to improve data capture. Specific statistical methods to improve data capture in trauma registries operated concurrently with the methods used to measure data capture. Linkage methods were either “deterministic”⁸⁷, where cases were merged if there was an exact match (unlikely), or “probabilistic”^{39,43,83,87} (preferred), where various weights were applied according to the level of matching. Working in tandem with dataset linkage, “capture-recapture” methods were used to estimate the eligible population and the proportion of these eligible cases actually included in the registry^{32,43,80,83,86,87}. One publication urged caution on the capture-recapture approach, on the basis that the necessary assumption of dataset independence was flawed⁷⁸.

Linkage methods primarily used to improve data capture were also used to improve data accuracy and completeness³².

Other data quality domains

Five of the 42 publications dealing with the measurement of a data quality dimension other than accuracy, completeness or capture, actually specified the methods used. Ringdal et al measured “feasibility”, “attainability” and “difficulty” of data collection using a written questionnaire⁴¹. Scheetz et al measured “accessibility” and “usefulness” by recording the response rate to written requests for data⁵⁴.

Meeuwisse 1998 measured “usefulness” with a written questionnaire⁸⁵. Laing et al measured “user satisfaction” and the “ease” and “timeliness” of data entry²⁷. Wargo et al also measured “timeliness”, but of registry staff activities, including case completion²⁸.

Very few publications were specific regarding methods used to improve “other” dimensions of data quality. Wargo et al provided a detailed account of approaches to improve productivity and timeliness, whilst maintaining data accuracy²⁸.

Discussion

This study examined the peer-reviewed literature to determine approaches to the definition, classification, standardisation, measurement, improvement and reporting of data quality in a trauma registry. Most publications focusing on data quality were based on trauma registries in developed countries and many were large multi-hospital registries.

There were just four studies which provided a general definition or classification of data quality in trauma registries, amongst which the taxonomy was inconsistent. Most of the publications covered at least one of the key data quality dimensions: accuracy, completeness and capture. Most of the publications also

considered at least one other data quality dimension. A closer examination of Table 2 generated the following potential candidates for other dimensions: “usefulness” (utility, usability), “timeliness” (efficiency), “comparability” (standardisation) and “cost-effectiveness”.

Approximately one quarter of publications which considered the measurement of accuracy, completeness or capture, did not specify the method or indicator used. The vast majority of publications dealing with the measurement of at least one “other” dimension did not specify the method or indicator used. While most publications focused on improving data quality, few provided detail regarding which methods were linked to which dimensions of data quality.

Described approaches to *measuring* data accuracy varied considerably, from the relatively simple approaches of consistency and domain checks, through monitoring of coding reliability and agreement with other databases, to source data verification. For source data verification, the patient medical record remains the best source, or “gold standard”, although completeness cannot be assured. Specific methods targeting the *improvement* of data accuracy were rarely provided in detail. Approaches to *measuring* data completeness, namely measuring the percentage of cases with complete observations for each variable, were more uniform and more widely described, implemented and reported. Specifically described methods to *improve* data completeness were mostly statistical approaches, namely MI. The published approaches to *measuring* and *improving* data capture largely involved the linkage of alternative datasets, but is dependent upon the existence and accessibility of such alternative datasets. The best approaches to *improving* data capture are the practical interventions intended to also *improve* accuracy and completeness such as developing efficient feedback mechanisms with data collection staff.

Rigorous attention to data quality in trauma registries is critically important to the valid benchmarking of trauma systems. How data quality is managed can have a very large impact on trauma system evaluation and health policy^{36,38,46,53,60,95}. This recognition of the paramount importance of data quality is not new^{90,91}

and it is not confined to trauma registries⁹⁶. But to date, there has been no single accessible document in the published peer-reviewed medical literature dealing with the definitions, dimensions, standards, evaluation, improvement or reporting of data quality in a trauma registry. Furthermore, even where interventions for improving specific dimensions of data quality have been thoroughly tested and published, the evidence of widespread adoption as best practice remains sparse^{46,65,96}.

While the non-trauma literature has provided a resource for describing the range of data quality dimensions, it is inconsistent with regard to the definition and classification of data quality. It may be that trauma registries need a specific framework to classify data quality. Unless there is clarity on the matter of classification, the progress towards developing a simple approach and metric to evaluating data quality in trauma registries will be slow. In turn, achieving the capacity to universally *measure* data quality in trauma registries will allow the focus to shift to the best interventions to *improve* data quality. Finally, to allow confidence in the results commonly generated from the growing number of trauma registries, there needs to be a uniform and universal standard for reporting trauma registry data quality. In the absence of standards for trauma registry data quality, the adoption of registry-derived initiatives to improve the quality of *trauma care* will be slowed.

Although the term “feasibility” or “pilot” was not included in the search strategy, more than one quarter of the publications dealing with data quality in trauma registries, including six of the seven publications from developing countries, were feasibility studies, mostly by self-identification. The definition of feasibility is not entirely clear, but implies, in the event of establishing a trauma registry, the very basic entry point of the data quality spectrum, where there has been the “successful” collection, and reporting, of *some* data on *some* variables for *some* patients. For those looking to start a successful and high quality trauma registry, there is no universally accessible standardised checklist.

Proper attention to ensure data quality to inform improvements in trauma care is critically important on a global scale. One of the five pillars of The United Nations' "Global Plan" for the current "Decade of Action for Road Safety 2011-2020" is "Post crash response". Activities encouraged include Activity 2: "Develop hospital trauma care systems and evaluate the quality of care through the implementation of good practices on trauma care systems and quality assurance" and Activity 7: "Encourage research and development into improving post crash response"⁹⁷. To support these activities, there needs to be a global standardisation of trauma registry functions, including a global minimum dataset and dictionary, uniform approaches to risk adjustment and the standardisation of data quality monitoring and reporting. Furthermore, this standardisation must account for the resource challenges of the vast majority of countries for which the burden of injury is greatest.⁹⁸⁻¹⁰⁰

This study has some limitations. The impetus for the study question and review included the ambiguity regarding the definition and classification of data quality in trauma registries. This same ambiguity necessarily provided challenges in sampling, and analysing, the literature methodically. Some informative publications may have been missed, including documents dealing with the broader remit of trauma centre verification¹⁰¹⁻¹⁰³. The intention was to focus upon the content of peer-reviewed publications which were widely accessible in the medical literature and contained a specific focus upon data quality. Given the study purpose and the challenges of defining and classifying data quality, the use of a scoping review methodology was considered ideal.¹⁰⁴⁻¹⁰⁵

That these limitations exist heightens the need to develop a comprehensive literature on data quality in trauma registries. This study addressed the question of "how" data quality is defined, classified, measured and improved. A logical next step, sampling all trauma registry-based publications over a recent period, would be to examine "if" data quality is reported and "what" is the result of data quality measurement. Such an audit of data quality would require a checklist or scorecard, potentially informed by this scoping review. Only through the development of accessible published documents, informed by both consensus

and evidence, will there be the necessary standardised guidance for the evaluation and improvement of data quality in trauma registries, and the concomitant improvement in the quality of trauma care.

Conclusion

The definition and classification of trauma registry data quality is ambiguous. Publications dealing with the evaluation and improvement of data quality in trauma registries are limited in number, consistency of approach and detail. To maintain the confidence of clinicians and administrators in the usefulness of trauma registries for monitoring the quality of trauma care, the metrics and reporting of data quality need to be standardised. There is a clear need for additional research and a globally accessible document standardising the definition, classification, measurement, improvement and reporting of data quality in trauma registries. Publications of trauma-registry-based studies must, as a minimum, report on data accuracy, completeness and capture.

References

1. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2012; 380 (9859): 2095 – 2128.
2. Cameron PA, Gabbe B, Cooper DJ. A statewide system of trauma care in Victoria: effect on patient survival. *Med J Aust*. 2008; 189: 546-50.
3. Gabbe BJ, Simpson PM, Sutherland AM, Wolfe R, Fitzgerald MC, Judson R, Cameron PA. Improved functional outcomes for major trauma patients in a regionalized, inclusive trauma system. *Ann Surg*. 2012; 255(6): 1009-15.
4. Gabbe BJ, Lyons RA, Fitzgerald MC, Judson R, Richardson J, Cameron PA. Reduced Population Burden of Road Transport-Related Major Trauma After Introduction of an Inclusive Trauma System. *Ann Surg* 2015;261:565–572
5. Fuller G, Bouamra O, Woodford M, Jenks T, Patel H, Coats TH, Oakley P, Mendelow AD, Pigott T, Hutchinson PJ, Lecky F. Temporal trends in head injury outcomes from 2003 to 2009 in England and Wales. *British Journal of Neurosurgery*. 2011; 25(3): 414-421
6. Celso B, Tepas J, Langland-Orban, Pracht E, Papa L, Lottenberg L, Flint L. A Systematic Review and Meta-Analysis Comparing Outcome of Severely Injured Patients Treated in Trauma Centers Following the Establishment of Trauma Systems *J Trauma*. 2006;60:371–378.
7. Lansink KW, Leenen LP. Do designated trauma systems improve outcomes? *Curr Opin Crit Care*. 2007 Dec;13(6):686-90.

8. Mock CN, Adzotor KE, Conklin E, Denno DM, Jurkovich GJ. Trauma outcomes in the rural developing world: comparison with an urban level I trauma center. *J Trauma*. 1993; 35: 518-23.
9. Mock C, Juillard C, Brundage S, Goosen J, Joshipura M. Guidelines for trauma quality improvement programmes. Geneva, World Health Organization, 2009. http://whqlibdoc.who.int/publications/2009/9789241597746_eng.pdf
10. Zehtabchi S, Nishijima DK, McKay MP, Mann, NC. Trauma Registries: History, Logistics, Limitations, and Contributions to Emergency Medicine Research. *Academic Emergency Medicine*. 2011; 18 (6): 637-43.
11. O'Reilly GM, Cameron PA, Joshipura M. Global trauma registry mapping: A scoping review. *Injury* 2012; 43: 1148-53.
12. O'Reilly GM, Joshipura M, Cameron PA, Gruen R. Trauma registries in developing countries: A review of the published experience. *Injury*. 2013; 44 (6): 713-21.
13. Schluter PJ, Nathens A, Neal ML, Goble S, Cameron CM, Davey TM, McClure RJ. Trauma and Injury Severity Score (TRISS) coefficients 2009 revision. *Journal of Trauma - Injury, Infection and Critical Care*. 2010; 68(4): 761-770.
14. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP: The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007, 370:1453-1457.
15. Australian Bureau of Statistics (ABS). Quality dimensions of the Australian National Accounts. 52160.55.002. Published 2007. Accessed 17 January 2015 at www.abs.gov.au
16. Nousak R, Phelps R. A scorecard approach to improving data quality. 200x. PWC Consulting, Chapel Hill, NC, USA.
17. Wang RL, Strong DM. Beyond accuracy: what data quality means to data consumers. *Journal of Management Information Systems*. 1996; 12(4): 5-33.
18. The Canadian Institute of Health Information (CIHI). The CIHI data quality framework 2009. CIHI. Accessed 17 January 2015 at http://www.cihi.ca/CIHI-ext-portal/pdf/internet/DATA_QUALITY_FRAMEWORK_2009_EN
19. Australian Commission on Safety and Quality in Healthcare. Framework for Australian clinical quality registries. Sydney. ACSQHC. March 2014. Accessed 17 January 2015 at <http://www.safetyandquality.gov.au/wp-content/uploads/2014/09/Framework-for-Australian-Clinical-Quality-Registries.pdf>
20. Sorensen HT, Sabroe S, Olsew J. A framework for evaluation of secondary data sources for epidemiological research. *International Journal of Epidemiology*. 1996; 25: 435-442.
21. Arts DGT, De Keizer NF, Scheffer GJ. Defining and improving data quality in medical registries: a literature review, case study and generic framework. *J Am Med Inform Assoc*. 2002;9:600-611.
22. United Nations Development Program. Human Development Report 2013. The rise of the south: human progress in a diverse world. Accessed 1 June 2014 at http://hdr.undp.org/sites/default/files/reports/14/hdr2013_en_complete.pdf
23. Mock C, Lormand JD, Goosen J, Joshipura M, Peden M. Guidelines for essential trauma care. Geneva, World Health Organization, 2004. <http://whqlibdoc.who.int/publications/2004/9241546409.pdf>
24. Nottidge TE, Dim M, Udoinyang CI, Udoh IA. The Uyo Trauma Registry-developed for sustainable audit of trauma care and cause in Nigeria. *Tropical Doctor*. 2014; 44(1):14-8.
25. Lehmann W, Hoffmann M, Fensky F, Nuchtern J, Grossterlinden L, Aghayev E, Lehmann H, Stuby F, Rueger J.M. What Is the Frequency of Nerve Injuries Associated With Acetabular Fractures? *Clinical Orthopaedics and Related Research*. 2014.
26. Hu P, Galvagno SM, Sen A, Dutton R, Jordan S, Floccare D, Handley C, Shackelford S, Pasley J, MacKenzie C. Identification of dynamic prehospital changes with continuous vital signs acquisition. *Air Medical Journal*. 2014; 33(1): 27-33.
27. Laing GL, Bruce JL, Aldous C, Clarke DL. The design, construction and implementation of a computerised trauma registry in a developing South African metropolitan trauma service. *Injury* 2014; 45(1): 3-8.

28. Wargo C, Bolig N, Hixson H, McWilliams N, Rummerfield H, Stratton E, Woodruff T. Trauma Registry Reengineered. *J TRAUMA NURS*. 2014; 21(6): 287-90.
29. Olsen CS, Thomas AM, Cook LJ. Hospital charges associated with motorcycle crash factors: a quantile regression analysis. *INJ PREV*. 2014; 20(4): 276-80.
30. Stevens KA, Paruk F, Bachani AM, Wesson HH, Wekesa JM, Mburu J, Mwangi JM, Saidi H; Hyder AA. Establishing hospital-based trauma registry systems: lessons from Kenya. *Injury*. 2013; 44 Suppl 4:S70-4.
31. Sonshine DB, Shantz J, Kumah-Ametepey R, Coughlin RR, Gosselin RA. The implementation of a pilot femur fracture registry at Komfo Anokye Teaching Hospital: an analysis of data quality and barriers to collaborative capacity-building. *World Journal of Surgery*. 2013; 37(7):1506-12.
32. Burkhardt M, Nienaber U, Holstein JH, Culemann, U, Bouillon B, Aghayev E, Paffrath T; Maegele M, Pohlemann T, Lefering R. Trauma registry record linkage: methodological approach to benefit from complementary data using the example of the German Pelvic Injury Register and the TraumaRegister DGU(R); TraumaRegister DGU(R), and Pelvic Injury Register of the German Trauma Society. *BMC Medical Research Methodology*, 2013; 13: 30.
33. Taylor SL, Lee D, Nagler T, Lawless MB, Curri T, Palmieri TL. A validity review of the National Burn Repository. *Journal of Burn Care & Research*. 2013; 34(2):274-80.
34. Wainiqolo I; Kafoa B; McCaig E; Kool B; McIntyre R; Ameratunga S. Development and piloting of the Fiji Injury Surveillance in Hospitals System (TRIP Project-1). *Injury*. 2013; 44(1):126-31.
35. Zelle BA, Bhandari M, Sanchez AI, Probst C, Pape HC. Loss of follow-up in orthopaedic trauma: Is 80% follow-up still acceptable? *Journal of Orthopaedic Trauma*. 2013; 27(3): 177-181.
36. O'Reilly GM; Cameron PA; Jolley DJ. Which patients have missing data? An analysis of missingness in a trauma registry. *Injury*. 2012; 43(11):1917-23.
37. Ordonez CA, Pino LF, Tejada JW, Badiel M, Loaiza JH, Mata LV, Aboutanos MB. Experience of two first level hospitals in the southwest region of Colombia on the implementation of the Panamerican Trauma Society International Trauma Registry. *Revista do Colegio Brasileiro de Cirurgioes*. 2012; 39(4):255-62.
38. Haider AH, Saleem T, Leow JJ, Villegas CV, Kisat M, Schneider EB, Haut ER, Stevens KA, Cornwell EE 3rd, MacKenzie EJ, Efron DT. Influence of the National Trauma Data Bank on the study of trauma outcomes: is it time to set research best practices to further enhance its impact? *Journal of the American College of Surgeons*. 2012; 214(5):756-68.
39. Newgard CD, Zive D, Jui J, Weathers C, Daya M. Electronic versus manual data processing: evaluating the use of electronic health records in out-of-hospital clinical research. *Academic Emergency Medicine*. 2012; 19(2):217-27.
40. Haas B, Xiong W, Brennan-Barnes M, Gomez D, Nathens AB. Overcoming barriers to population-based injury research: development and validation of an ICD10-to-AIS algorithm. *Canadian Journal of Surgery*, 2012; 55 (1): 21-6.
41. Ringdal KG, Lossius HM, Jones JM, Lauritsen JM, Coats TJ, Palmer CS, Lefering R, Di Bartolomeo S, Dries DJ, Soreide K. Utstein Trauma Data Collaborators. Collecting core data in severely injured patients using a consensus trauma template: an international multicentre study. *Critical Care*. 2011; 15(5):R237.
42. Sears JM, Bowman SM, Adams D, Silverstein BA. Occupational injury surveillance using the Washington State Trauma Registry. *Journal of Occupational & Environmental Medicine*. 2011; 53(11):1243-50.
43. Newgard CD, Zive D, Malveau S, Leopold R, Worrall W, Sahni R. Developing a statewide emergency medical services database linked to hospital outcomes: a feasibility study. *Prehospital Emergency Care*. 2011;15(3):303-19.
44. Furlan JC, Fehlings MG. The National Trauma Registry as a Canadian spine trauma database: a validation study using an institutional clinical database. *Neuroepidemiology*. 2011; 37(2):96-101.

45. Bowman, Stephen M.; Aitken, Mary E. Assessing External Cause of Injury Coding Accuracy for Transport Injury Hospitalizations. *Perspectives in Health Information Management*. 2011; 8 (4): 1-7.
46. O'Reilly GM, Jolley DJ, Cameron PA, Gabbe B. Missing in action: a case study of the application of methods for dealing with missing data to trauma system benchmarking. *Academic Emergency Medicine*. 2010; 17(10):1122-9.
47. Leukhardt WH, Golob JF, McCoy AM, Fadlalla AM, Malangoni MA, Claridge JA. Follow-up disparities after trauma: a real problem for outcomes research. *American Journal of Surgery*. 2010; 199(3):348-52.
48. Newgard CD, Haukoos JS. Measuring Quality With Missing Data: The Invisible Threat to National Quality Initiatives.; *Academic Emergency Medicine*. 2010; 17 (10): 1130-3.
49. Gomez D, Xiong W, Haas B, Goble S, Ahmed N, Nathens AB. The missing dead: the problem of case ascertainment in the assessment of trauma center performance. *Journal of Trauma-Injury Infection & Critical Care*. 2009; 66(4):1218-24.
50. Finlay-Morreale HE, Tsuei BJ, Fisher BS, Davis K, Johannigman JA. Close is dead: determinants of firearm injury lethality in women. *J TRAUMA*. 2009; 66 (4): 1207-11.
51. Mackenzie CF; Hu P; Sen A; Dutton R; Seebode S; Floccare D; Scalea T. Automatic pre-hospital vital signs waveform and trend data capture fills quality management, triage and outcome prediction gaps. *AMIA. Annual Symposium Proceedings*. 2008; 318-22.
52. Protetch J, Chappel D. Trauma registry data validation: building objectivity. *Journal of Trauma Nursing*. 2008;15(2):67-71.
53. Moore L, Clark DE. The value of trauma registries. *Injury*. 2008; 39(6):686-95.
54. Scheetz LJ, Zhang J, Kolassa JE, Allen P, Allen M. Evaluation of injury databases as a preliminary step to developing a triage decision rule. *Journal of Nursing Scholarship*. 2008; 40(2):144-50.
55. Hayes JR, Groner JI. Using multiple imputation and propensity scores to test the effect of car seats and seat belt usage on injury severity from trauma registry data. *Journal of Pediatric Surgery*. 2008; 43(5):924-7.
56. Alexandrescu R, O'Brien SJ, Lyons RA, Lecky FE. A proposed approach in defining population-based rates of major injury from a trauma registry dataset: delineation of hospital catchment areas. *BMC Health Services Research*. 2008; 8:80.
57. Kardooni S, Haut ER, Chang DC, Pierce CA, Efron DT, Haider AH, Pronovost PJ, Cornwell EE. Hazards of benchmarking complications with the national trauma data bank: Numerators in search of denominators. *Journal of Trauma - Injury, Infection and Critical Care*. 2008; 64(2): 273-277.
58. Zhaohui S, Shuxia Z, Xinghua F, Shujun L, Yanpu L, Bin B, Zhongzhi S. The design and implementation of chinese maxillofacial trauma registry, analysis and injury severity score system. *J TRAUMA*. 2008; 64 (4): 1024-33.
59. Beretta L, Aldrovandi V, Grandi E, Citerio G, Stocchetti N. Improving the quality of data entry in a low-budget head injury database. *Acta Neurochirurgica*. 2007;149(9):903-9.
60. Datta I, Findlay C, Kortbeek JB, Hameed SM. Evaluation of a regional trauma registry. *Canadian Journal of Surgery*. 2007; 50(3):210-3.
61. Guice KS, Cassidy LD, Mann NC. State trauma registries: survey and update-2004. *Journal of Trauma-Injury Infection & Critical Care*. 2007; 62(2):424-35.
62. Hlaing T, Hollister L, Aaland M. Trauma registry data validation: Essential for quality trauma care. *Journal of Trauma-Injury Infection & Critical Care*. 2006; 61(6):1400-7.
63. Eastridge BJ, Jenkins D, Flaherty S, Schiller H, Holcomb JB. Trauma system development in a theater of war: Experiences from Operation Iraqi Freedom and Operation Enduring Freedom. *Journal of Trauma-Injury Infection & Critical Care*. 2006; 61(6):1366-72.
64. Burd RS, Jang TS, Nair SS. Predicting hospital mortality among injured children using a national trauma database. *Journal of Trauma-Injury Infection & Critical Care*. 2006; 60(4):792-801.

65. Newgard CD. The validity of using multiple imputation for missing out-of-hospital data in a state trauma registry. *Academic Emergency Medicine*. 2006;13(3):314-24.
66. Holcomb JB, Stansbury LG, Champion HR, Wade C, Bellamy RF. Understanding combat casualty care statistics. *Journal of Trauma-Injury Infection & Critical Care*. 2006; 60(2):397-401.
67. Bergeron E, Lavoie A, Moore L, Bamvita JM, Ratté S, Clas D. Paying the price of excluding patients from a trauma registry. *Journal of Trauma-Injury Infection & Critical Care*. 2006; 60(2):300-4.
68. McKenzie K, Walker S, Besenyei A, Aitken LM, Allison B. Assessing the concordance of trauma registry data and hospital records. *Health Information Management Journal*. 2005; 34(1):3-7.
69. Rossi PG, Farchi S, Chini F, Camilloni L, Borgia P, Guasticchi G. Road traffic injuries in Lazio, Italy: a descriptive analysis from an emergency department-based surveillance system. *ANN EMERG MED*. 2005; 46 (2): 152-7.
70. Petridou E, Dessypris N, Frangakis CE, Belechri M, Mavrou A, Trichopoulos D. Estimating the population burden of injuries: a comparison of household surveys and emergency department surveillance. *Epidemiology*. 2004; 15(4):428-32.
71. Joseph L; Belisle P; Tamim H; Sampalis JS. Selection bias found in interpreting analyses with missing data for the prehospital index for trauma. *Journal of Clinical Epidemiology*. 2004; 57(2):147-53.
72. Read-Allsopp C. Trauma registry. Establishing inter-rater reliability scoring in a state trauma system. *Journal of Trauma Nursing*. 2004; 11(1): 35-9.
73. Mikhail JN; Harris YD; Sorensen VJ. Injury severity scoring: influence of trauma surgeon involvement on accuracy. *Journal of Trauma Nursing*. 2003; 10(2):43-7.
74. Auerbach S; FitzPatrick MK; Garuffe A; Williams B; McMaster J; Stum M; Reilly P. Enhancing data quality through the formation of a trauma registry workgroup. *Journal of Trauma Nursing*. 2002; 9(2):46-50.
75. Lucas CE, Buechter KJ, Coscia RL, Hurst JM, Lane V, Meredith JW, Middleton JD, Mitchell FL Jr, Rinker CF, Tuggle D, Vlahos A, Wilberger J, Yu P. The effect of trauma program registry on reported mortality rates. *Journal of Trauma-Injury Infection & Critical Care*. 2001; 51(6):1122-6.
76. Al Naami MY, Sadik AA, Adam MA. Evaluation of trauma registry data in Asir region. *Saudi Medical Journal*. 2001; 22(5):438-43.
77. Wynn A, Wise M, Wright MJ, Razaat A, Wang Y, Steeb G, McSwain N, Buechter KJ, Hunt JP. Accuracy of administrative and trauma registry databases. *J TRAUMA*. 2001; 51 (3): 464-8.
78. Jarvis SN, Lowe PJ, Avery A, Levene S, Cormack RM. Children are not goldfish--mark/recapture techniques and their application to injury data. *Injury Prevention*. 6(1):46-50, 2000 Mar.
79. Sanidas EE, Valassiadou KE, Kafetzakis AG, Yannopoulos AT, Vlazakis SS, Markogiannakis HE, Karniadakis S, Alamanos G, Melissas JA, Tsiftsis DD. Organisation of a trauma registry in a regional Greek university hospital: the first two years experience. *European Journal of Surgery*. 2000; 166(1):13-7.
80. Schootman M, Harlan M, Fuortes L. Use of the capture-recapture method to estimate severe traumatic brain injury rates. *Journal of Trauma-Injury Infection & Critical Care*. 2000; 48(1):70-5.
81. Hunt JP, Cherr GS, Hunter C, Wright MJ, Wang Y, Steeb G, Buechter KJ, Meyer AA, Baker CC. Accuracy of administrative data in trauma: splenic injuries as an example. *J TRAUMA*; 2000; 49 (4): 679-88
82. FitzPatrick MK, Frankel HL, McMaster J, Heliger LA, Auerbach S, Reilly P, Schwab CW. Integration of concurrent trauma registry and performance improvement programs. *J TRAUMA NURS*. 2000; 7 (4): 92-7.
83. Clark DE, Hahn DR. Hospital trauma registries linked with population-based data. *Journal of Trauma-Injury Infection & Critical Care*. 1999; 47(3):448-54.
84. Owen JL, Bolenbaucher RM, Moore ML. Trauma registry databases: a comparison of data abstraction, interpretation, and entry at two level I trauma centers. *Journal of Trauma-Injury Infection & Critical Care*. 1999; 46(6):1100-4.

85. Meeuwisse WH, Love EJ. Development, implementation, and validation of the Canadian Intercollegiate Sport Injury Registry. *Clinical Journal of Sport Medicine*. 1998; 8(3):164-77.
86. Surkin J, Smith M, Penman A, Currier M, Harkey HL; Chang YF. Spinal cord injury incidence in Mississippi: a capture-recapture approach. *Journal of Trauma-Injury Infection & Critical Care*. 1998; 45(3):502-4.
87. Rosman DL. The feasibility of linking hospital and police road crash casualty records without names. *Accident Analysis & Prevention*. 1996; 28(2):271-4.
88. Vestrup JA, Phang PT, Vertesi L, Wing PC, Hamilton NE. The utility of a multicenter regional trauma registry. *Journal of Trauma-Injury Infection & Critical Care*. 1994; 37(3):375-8.
89. Tepas JJ, Ramenofsky ML, Barlow B, Gans BM, Harris BH, DiScala C, Butler K. National Pediatric Trauma Registry. *Journal of Pediatric Surgery*. 1989; 24(2):156-8.
90. Rossignol AM, Locke JA. An assessment of the completeness of the Massachusetts burn registry. *Public Health Reports*. 1983; 98(5): 492-496.
91. Goldberg J, Gelfand HM, Levy PS, Mullner R. An evaluation of the Illinois trauma registry: the completeness of case reporting. *Medical Care*. 18(5):520-31, 1980 May.
92. Charters AC, Bailey JA. Experience with a simplified trauma registry: profile of trauma at a university hospital. *Journal of Trauma-Injury Infection & Critical Care*. 1979; 19(1):13-7.
93. Gravseth HM, Wergeland E, Lund J. Underreporting of occupational injuries to the Labour Inspection. *Tidsskrift for Den Norske Laegeforening*. 2003; 123(15):2057-9.
94. Centers for Disease Control. Guidelines for evaluating surveillance systems. *MMWR* 1988;37(suppl S-5):1-18.
95. Rodenberg H. The Florida trauma system: assessment of a statewide database. *Injury* 1996;27:205-8.
96. Little RJ, D'Agostino R, Cohen ML, Dickersin K, Emerson SS, Farrar JT, Frangakis C, Hogan JW, Molenberghs G, Murphy SA, Neaton JD, Rotnitsky A, Scharfstein D, Shih WJ, Siegel JP, Stern H. The prevention and treatment of missing data in clinical trials. *N Engl J Med*. 2012; 367(14): 1355-1360.
97. World Health Organization. Global Plan for the Decade of Action for Road Safety 2011-2020. Geneva, World Health Organization, 2011. http://www.roadsafetyfund.org/UnDecadeOfAction/Documents/global_plan_en.pdf
98. World Health Organization. Violence and injury prevention: trauma registry and data. Accessed 1 June 2014 at http://www.who.int/violence_injury_prevention/services/gaci/activities/en/
99. O'Reilly GM, Gabbe BJ, Cameron PA. Trauma registry methodology: A survey of trauma registry custodians to determine current approaches. *Injury*. 2014 (in press). DOI: <http://dx.doi.org/10.1016/j.injury.2014.09.010>
100. O'Reilly GM, Gabbe B, Graaf S, Cameron PA. An interview of trauma registry custodians to determine lessons learnt. *Injury*. 2015 (in press).
101. Rotondo MF, Cribari C, Smith RS (eds). Resources for the Optimal Care of the Injured Patient (2014). American College of Surgeons, Committee on Trauma. Accessed 2 February 2015. <https://www.facs.org>
102. Royal Australasian College of Surgeons. Model Resource Criteria (2014). Accessed 2 February 2015. http://www.surgeons.org/media/309212/doc_2014-04-29_mrc_v2.0_for_web_-_full_document.pdf
103. The Trauma Audit and Research Network (TARN). Procedures Manual. May 2015. Salford, UK. Accessed 21 July 2015 at <https://www.tarn.ac.uk/content/downloads/53/Complete%20Procedures%20Manual%20-%20May%202015.pdf>
104. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int. J. Social Research Methodology*. 2005; 8(1): 19-32.
105. Armstrong R, Hall BJ, Doyle J, Waters E. 'Scoping the scope' of a Cochrane review. *J. Pub Health*. 2011; 33(1): 147-150.

Table 1: Dimensions of quality used in search terms

Figure 1: Number of publications by year

Table 2: Dimensions of data quality named in publications

Table 3: Approaches to measuring data quality by dimension

Table 4: Approaches to improving data quality (accuracy, completeness and/or capture)

Supplementary File 1: Included publications, by year of publication, source country, level of country development, registry jurisdiction and level of registry development.

Supplementary File 2: Description of publications by study design, study question and data quality dimension

Table 1: Dimensions of quality used in search terms

quality	Correctness	comparability
accuracy	consistency	uniqueness
completeness	usability	availability
capture	relevance	representativeness
coverage	timeliness	case ascertainment
missing	accessibility	error
reliability	interpretability	concordance
validity	coherence	precision
	verification	


Figure 1: Number of publications by year

Table 2: Dimensions of data quality named in publications

Accuracy		
Consistency	Variances	Misclassification
Errors	Reproducibility	Replicable
Validity	Precision	Noise
Differences	Agreement	Bias
Discrepancies	Verification	Logical
Variability	Sensitivity	Matching
Reliability	Concordance	Sloppy
Congruence	Misreporting	Commission
Homogeneity	Miscoding	Mistakes
Correctness	Out of range	Contradictory
Flawed	Confidence	Certainty
Completeness		
Dropouts	Missingness	Collectability
Omissions	Blank	Empty
Capture		
Completeness	Representativeness	Coverage
Case ascertainment	Over-reported	Under-reported
Generalisability	Omissions	Missing
	Duplication	
Other		
Importance	Accessibility	Timeliness
Relevance	Flexibility	Labour-intensive
Ease	Manageability	Credibility
Sustainability	Time-consuming	Deficiencies
Detail	Descriptive	Shortcomings
Usefulness	Promptness	Misleading
Efficiency	Acceptability	Real-time
Functionality	Value	Feasibility
Utility	Meaningful	Versatility
Standardisation	Simplicity	Optimal
Uniformity	Availability	Continuity
Integrity	User-friendly	Effectiveness
Comparability	Usability	Rigour
Lacking	Comprehensive	Indispensable
Applicability	Flow	Interpretability
Limitations	Granularity	Inexpensive
Scope	Cleanliness	Advantageous
Range	Attainability	Desirability
Nonsensible	Difficulty	Compliance
Cost-effectiveness	Value-add	Robustness
Misinformation	Contribute	Productivity
Maintenance	Performance	Satisfactory
Obscurity	Current	Speed
Informative	Adequacy	Ideal
Professional	Veracity	Compatibility
Stability	Security	Convenience
Excellence	Failure	Suitability
	Appropriateness	

Table 3: Approaches to measuring data quality by dimension

Dimension	Method ^a (of measurement used)	Number of publications	Indicator ^a (used to measure dimension)	Number of publications	References
Accuracy (n=47)	Source data verification	6	Sensitivity / specificity / accuracy (%)	4	44,52,60,85
			Unspecified	2	34,56
	Validation – other datasets	18	Sensitivity / specificity / accuracy (%)	11	32,39,42,45,62,68,70,77,81,85,86
			Agreement (e.g. kappa / ICC / Bland Altman plot)	4	26,39,45,65
			Unspecified	5	43,56,83,84,87
	Coding verification	17	Sensitivity / specificity / accuracy (%)	9	41,42,45,52,62,72,73,77,81
			Agreement (e.g. kappa)	2	40,72
			Unspecified	7	31,34,53,56,75,81,88
	Internal consistency check	7	Proportion of cases for a variable (%)	4	25,33,59,62
			Unspecified	3	39,54,56
	Domain check	3	Proportion of observations outside plausible range (%)	1	62
			Unspecified	2	39,59
	Inter-rater reliability	10	Proportion of observations in agreement (%)	3	25,72,85
Kappa			1	26	
Unspecified			6	52,53,54,73,84,88	
Unspecified	13	-	13	27,28,30,38,51,57,58,61,72,76,79,89,92	
		-	13	27,28,30,38,51,57,58,61,72,76,79,89,92	
Completeness (n=47)	Completeness of variables	32	% of cases with complete observations for each variable	29	24,27,29,31,32,33-36,39,41,43,46,47,48,54-57,59,60,62,64,65,71,76,77,81,85
			Unspecified	3	30,44,53
	Completeness of cases	10	% of cases with complete observations for all variables	9	25,36,42,46,60,63,64,71,76
			Unspecified	1	68
Unspecified	11	-	11	26,38,51,52,58,61,67,83,87,89,92	
Capture (n=37)	Linkages – other datasets	27	% of eligible population included in registry	24	24,27,32,34,39,42,43,45,56,63,67-70,77,78,80,81,83,85-87,90,91
			Unspecified	3	30,60,62
	Comparison of distributions	3	Difference in means / other	3	49,67,81
			Unspecified	0	-
	Case duplication	6	Number (%) or cases with more than one record	4	32,33,34,62
Unspecified			2	39,42	
Unspecified	9	-	9	29,31,38,51,53,61,66,79,92	
Other (n=42)	Specified ^b	5	-	5	27,28,41,54,85
			Unspecified	37	24,25,26,30,31,33,34,37,40,43,45,47,52,53,56,58-61,64,66,67,70,73-77,79,82,83,84,87,88,90-92

^aNot mutually exclusive (except “unspecified”). i.e. There may be more than one measurement method or indicator in a single publication

^bDescribed in the results section

Table 4: Approaches to improving data quality (accuracy, completeness and/or capture)

Domain	Items	Primary dimension(s)			Number of papers (n=69)	References
		Accuracy	Complete	Capture		
Physical Resources						
Equipment	Computer hardware	x	x	x	9	24,27,30,37,61,74,79,88,92
	Computer software	x	x	x	26	25,27,29-31,33,34,37,39,43,51-53,58,59,61,62,74-76,79,82,83,87-89
Human resources						
Staff – numbers		x	x	x	7	28,30,37,59,63,88,91
Staff – type / mix		x	x	x	19	27,30,33,34,37,61-63,73-76,79,82,85,88,89,91,92
Staff – training		x	x	x	21	27,30,33,34,37,45,52,53,59,62,63,68,72,73-75,84,85,88,91,92
Processes						
Administration	Ethics / consent	-	-	x	11	30,52-54,61,62,74,79,82,90,91
	Supervision	x	x	x	6	30,34,52,53,59,62
	Feedback	x	x	x	16	25,28,30,34,37,52,53,59,62,68,73,74,79,82,85,88
	Financial incentives / penalties	x	x	x	4	31,52,74,90
Organisation						
Inclusion criteria		-	-	x	22	27,31,32,34,39,45,53,56,61,66-68,70,76,79,80,83,84,86,88,91,92
Variables	Selection (minimum dataset)	x	x	-	33	24,26,27,30-32,34,37,39,43,45,48,51-54,56,58,59,61-65,76,79,82,84,85,87-89,91
	Definition / coding	x	x	x	19	25,26,30,34,42,51,53,54,57,59,61,66,67,74,80,84,88,90,92
Data capture	Form	x	x	-	12	24,30,34,59,62,63,66,79,85,88,89,92
	Collector type	x	x	-	15	24-27,30,31,34,37,62,70,73,79,85,88,92
	Collection mechanism	x	x	-	21	24,26-28,30,31,34,37,39,48,51,57,59,62,70,73,74,79,84,85,88
	Coding (e.g. ICD / AIS)	x	-	x	23	25,31,34,40,42,45,52-54,58,62,67,68,72,73,75,77,79,80,83,84,88,92
	Input	x	x	x	17	24,30,32,33,37,39,43,51,53,57-59,61,74,75,82,84
	Linkages	x	x	x	24	29-32,39,43,45,50,51,53,54,61,67-70,77,78,80,82,83,86,87,91
	Missing data ^a	-	x	-	15	26,29,35,36,38,39,43,46,48,53-55,64,65,71

^aMissing data methods at analysis stage: includes complete case analysis, single imputation, multiple imputation

Regarding the authors of the manuscript titled:

Classifying, measuring and improving the quality of data in trauma registries: a review of the literature

There are no known or perceived conflicts of interest.

Accepted Manuscript

Acknowledgements

There were no sources of funding for this study. Belinda Gabbe was supported by a Career Development Fellowship (GNT1048731) during the preparation of this manuscript.

Accepted Manuscript