



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa26445>

Paper:

Monnier, L., Colette, C., Dejager, S. & Owens, D. (2016). Near normal HbA1c with stable glucose homeostasis: The ultimate target/aim of diabetes therapy.

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Reviews in Endocrine and Metabolic Disorders

Near normal HbA1c with stable glucose homeostasis: The ultimate target/aim of diabetes therapy --Manuscript Draft--

Manuscript Number:	
Full Title:	Near normal HbA1c with stable glucose homeostasis: The ultimate target/aim of diabetes therapy
Article Type:	Review Article (invited)
Corresponding Author:	Louis Monnier FRANCE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Louis Monnier
First Author Secondary Information:	
Order of Authors:	Louis Monnier Claude Colette, PhD Sylvie Dejager, MD,PhD David Owens, MD
Order of Authors Secondary Information:	
Funding Information:	

Dear Editor,

Please find the review article entitled “New normal HbA1c with stable glucose homeostasis: The ultimate target/aim of diabetes therapy”, which has been solicited consecutively to my contribution as speaker at the Hanefeld symposium to be held in Dresden.

Hoping that this review article is in agreement with your expectation.

Looking forward to hearing from you,

With best regards,

Louis Monnier

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Near normal HbA1c with stable glucose homeostasis:

The ultimate target/aim of diabetes therapy

L Monnier¹, C Colette¹, S Dejager², DR Owens³

Address of authors

1 Institute of Clinical Research, University of Montpellier, France

2 Department of Endocrinology, Hospital Pitié Salpêtrière, Paris, France

3 Diabetes Research Group, Swansea University, United Kingdom

Short title: Stable glucose homeostasis and diabetes therapy

Total number of words in manuscript: 4582

Total number of words in abstract: 184

Total number of figures: 7

Corresponding author:

Professor Louis Monnier
Institute of Clinical Research
641 Avenue Doyen Giraud
34093 Montpellier Cedex 5 France
Tél: +33 411 759 891
e-mail: louis.monnier@inserm.fr

Abstract,

1
2 Achieving near normal glucose homeostasis implies that all components of dysglycemia that are present in
3 diabetes states be eliminated. Reducing ambient/overall hyperglycaemia is a pre-requisite to eliminate the risk of
4 development and progression of diabetes complications. More controversially however, are the relative and
5 related contributions of postprandial glucose excursions, glucose variability, hypoglycaemia and the dawn
6 phenomenon across the spectrum of dysglycemia. For instance, it is likely that the dawn phenomenon
7 contributes to ambient hyperglycaemia and that postprandial glucose excursions are at the cross road of ambient
8 hyperglycaemia and glucose variability with glucose fluctuations as causative risk factors for hypoglycaemia.
9 Proof-of-concept trials such as the ongoing FLAT-SUGAR study are necessary for gaining further insight into
10 the possible harmful effects of some of these features such as excessive glycaemic variability and glucose
11 excursions, still considered to be of minor relevance by several diabetologists. Whether their role will be more
12 thoroughly proven through further intervention trials with “hard” endpoints, remains to be seen. In the meantime
13 more consideration should be given to medications aimed at concomitantly reducing ambient/overall
14 hyperglycaemia and those additional abnormal glycaemic features of dysglycemia.
15
16
17
18
19
20
21
22

23 Keywords: glucose homeostasis; quality of diabetes therapy.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Introduction

1
2 The role of any antidiabetic treatment should be to achieve a near to normal glycaemic control based on the fact
3 that most observational, epidemiologic and several interventional studies [1,2,3,4] have shown that the incidence
4 of cardiovascular events, premature death and microvascular diabetic complications are associated with the
5 overall glycaemic exposure over time. The degree of sustained chronic (ambient) hyperglycaemia is quantified
6 by the determination of circulating HbA1c levels representing glycaemic control over a 2-to 3-month period of
7 time [3,4]. It has also been established that even in persons with a HbA1c level of 6.5% (diagnostic threshold for
8 diabetes), the risk of coronary heart disease and ischaemic stroke is 2- to -3 fold higher compared to those with a
9 value of 5.5% [7]. There is incontrovertible evidence that ambient hyperglycaemia is a key player in the
10 pathogenesis of diabetic complications at least of microvascular diseases, both in type 1 and type 2 diabetes.
11 Such an evidence has been provided by two landmark interventional trials: The Diabetes Control and
12 Complications Trial (DCCT) in type 1 diabetes [3] and the United Kingdom Prospective Diabetes Study
13 (UKPDS) in type 2 diabetes [4]. Both these studies have demonstrated a consistent benefit of intensive therapies
14 on the incidence of microvascular complications (DCCT) and of all related-diabetes complications (UKPDS).
15 The evidence is less clear for macrovascular complications because the ACCORD (Action to Control
16 Cardiovascular Risk in Diabetes) [8], the ADVANCE (Action in Diabetes and Vascular Disease: preterax and
17 diamicon modified release Controlled Evaluation) [9] and the VADT (Veterans Affairs Diabetes Trial) [10], i.e.
18 the three major randomised control interventional trials to compare intensive treatment with standard strategies,
19 have shown a modest benefit in terms of macrovascular outcomes or have even failed to demonstrate any
20 significant improvement in the short term [11]. In the ACCORD study [8] the risk of death, and especially as a
21 result of cardiovascular disease, was found to be greater on intensive than standard therapy bringing the study to
22 an early termination. The disappointing cardiovascular outcomes with intensive intervention led several authors
23 to challenge the then current therapeutic strategies employed in an attempt to achieve near normal glucose
24 control [12, 13]. However, these authors failed to acknowledge that the studies included in their meta-regression
25 analysis were of a relatively short duration up from 1 to 5.6 years. Since, it has been recognised that lowering-
26 glucose treatment requires a much longer period to demonstrate the cardiovascular benefit of improvement in
27 glucose control [14,15,16,17]. The pivotal role of hyperglycaemia as a causative factor of macro-vascular
28 complications was confirmed at least by five studies: the Diabetes Control and Complications
29 Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study in type 1 diabetes [14,15]
30 and the analyses of the extended follow-up of such studies as the UKPDS [16], ACCORD [17], VADT [18] and
31 the Steno-2 study [19] in type 2 diabetes. Figure 1 represents the time course of the benefit of lowering-glucose
32 strategies (intensive vs standard) on the relative risk of major cardiovascular outcomes in the different
33 interventional trials mentioned above. The observations indicate that a significant reduction in the relative risk of
34 cardiovascular disease becomes evident only when the duration of follow-up was beyond 10 years as seen in the
35 extended follow-up of the UKPDS [16] and the VADT [18] studies. Recently however, surprising results were
36 observed by the EMPA-REG OUTCOME Investigators [20] after a median observation time of 3.1 years. It is
37 highly likely that the improvements on cardiovascular outcomes in empagliflozin- treated subjects with type 2
38 diabetes are not simply due to the glucose-lowering effect of the drug but possibly due to improvements in blood
39 pressure, cardiac function and body weight.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Reverting to the results of the UKPDS [4], it has been demonstrated that the incidence of clinical complications
2 was positively and significantly associated with HbA1c and that there is no definitive threshold of HbA1c for
3 any type of diabetes complication. When setting the baseline risk of complications at 1 when the HbA1c level is
4 5.5%, the analysis of the UKPDS data indicates that this risk increases 1.5 and 2 fold higher when the mean
5 HbA1c level reaches 7 and 8%, respectively. There is also a remnant risk for adverse cardiovascular events even
6 when HbA1c levels range between 5.5 and 7%, i.e. below the current ADA [21] and IDF [22] recommended
7 target of <7% and even below 6.5% which is currently used for defining the presence of diabetes [23].
8
9

10 **2. Features of dysglycaemia involved in the evolution of type 2 diabetes**

11 The features of dysglycaemia in persons with type 2 diabetes can be roughly divided into the following
12 components (figure 2):
13

14 - Normal glucose exposure, which corresponds to the area between 0 and 100 mg/dL (0-5.6 mmol/L) in the
15 fasting state increasing to a postprandial peak <140 mg/dL (<7.8 mmol/L) before return to baseline values within
16 2-3 hours
17

18 - Additional excessive glucose exposure, which is only observed in patients with diabetes, in other words in
19 those with HbA1c levels $\geq 6.5\%$ according to American Diabetes Association (ADA) standards [23]. This
20 additional glucose exposure observed in persons with type 2 diabetes can be separated into three further
21 subcomponents: (a) the dawn phenomenon [24-26], (b) post-prandial hyperglycaemia [27-29] and (c) basal
22 hyperglycaemia. These aforementioned dysglycaemic states do not necessary occur simultaneously in the
23 evolution of type 2 diabetes and can vary in their contribution throughout the natural history of the disease
24 (figure 3) [30].
25
26
27
28
29
30
31
32
33

34 **2.1 The dawn phenomenon**

35 This phenomenon corresponds to a rise in plasma glucose and/or insulin requirement towards the end of the
36 nocturnal period, in the absence of any dietary (carbohydrate) intake. The dawn phenomenon is mainly due to
37 the circadian variation in hepatic glucose production, which starts to increase in the evening and reaching a peak
38 towards the end of an overnight fast and then declining during daytime until its late afternoon nadir [31]. The
39 two main consequences of this rise of circulating blood glucose overnight includes elevation of the early
40 morning fasting blood glucose and secondly abnormally high and delayed post-breakfast glucose excursions
41 referred to as the “extended dawn phenomenon” [27]. This latter phenomenon is postulated to be due to the
42 combined influence of an overproduction of glucose by the liver complemented by the intestinal hydrolysis of
43 breakfast carbohydrate. Both phenomena, and more specifically the dawn phenomenon is not observed in non-
44 diabetic subjects [32] since the hepatic glucose output in the early morning is counteracted by an increase in the
45 endogenous insulin secretion [33].
46
47
48
49
50
51
52
53

54 We have recently demonstrated [34], in a group of 50 well-controlled persons with type 2 diabetes with a HbA1c
55 ranging from 5.7 to 6.5% and predominantly treated with dietary measures alone (34 out of 50), that the “dawn
56 phenomenon” is evident whilst the mean postprandial glucose (131 mg/dL, 7.3 mmol/L) and overall 24-hour
57 glucose levels (115 mg/dl, 6.4 mmol/L) remained within the normal range in most of them (figure 4). This
58
59
60
61
62
63
64
65

1 suggests that the dawn phenomenon is likely to be the earliest expression of dysglycaemia in the natural history
2 of type 2 diabetes (figure 3). Furthermore, we have demonstrated that its contribution to the overall glucose
3 exposure cannot be neglected since 0.4% (expressed as percentage point) of the HbA1c in type 2 diabetes can be
4 explained by the dawn phenomenon [26].
5

6 **2.2 Postprandial hyperglycaemia**

7
8
9 This glycaemic state is defined as the increment (the AUCs of the glycaemic profiles) above the horizontal lines
10 set for each meal at pre-meal glucose values (figure 2).
11

12 In 2003 [28], we demonstrated that postprandial hyperglycaemia made the greatest contribution to the overall
13 hyperglycaemia (70%) in persons with type 2 diabetes treated with oral anti-diabetic agents. In 2007 this finding
14 was confirmed by analysing the 24-hour continuous glucose profiles of non-insulin-treated type 2 diabetic
15 subjects presenting at different levels of HbA1c [27]. As soon as HbA1c levels exceeded 6.5% we observed an
16 abnormal elevation of post-meal glucose levels. This phase of dysglycaemia has been recently confirmed (figure
17 4) in a study of 100 persons with type 2 diabetes (HbA1c < 7%), treated either with dietary measures alone or in
18 combination with oral anti-diabetic agents [34]. The difference between those exhibiting mild dysglycaemia
19 (HbA1c 6.5-6.9%, n = 50) compared with those with even better glycaemic control (HbA1c < 6.5%, n = 50) was
20 due to the greater post-meal excursions in the former group. In those with a HbA1c < 6.5%, the proportion of
21 individuals with an average two-hour post-meal glucose (mean of post-breakfast, post-lunch and post-dinner
22 values) above the upper limit of normal of 140 mg/dL (7.8 mmol/L) was less than a quarter. In contrast, more
23 than one half of those with an HbA1c level between 6.5% and 6.9% exceeded this threshold value [34].
24

25 Therefore, persons with a HbA1c \geq 6.5% but < 7%, i.e. “residual dysglycaemia” differ pathophysiologically
26 from those with a HbA1c < 6.5% by virtue of the greater frequency and magnitude of the post-meal glucose
27 excursions. In both groups, the “dawn phenomenon” was present whilst basal hyperglycaemia was absent.
28

29 Therefore, one can conclude that excess postprandial hyperglycaemia is the second abnormality in the
30 dysglycaemic continuum/spectrum of type 2 diabetes provided that the HbA1c levels remain below 7% (figure
31 3). The calculated absolute contribution of postprandial glucose excursions to the HbA1c level is usually of 0.6%
32 when the HbA1c is below 6.8%, which increases to a constant 1% beyond this level and remaining stable despite
33 ever increasing HbA1c [29] (figure 5).
34

35 According to these observations, one can hypothesize that eradication of both the dawn phenomenon and
36 postprandial hyperglycaemia would permit to decrease the HbA1c level by a total of 1%, i.e. 0.4% for the dawn
37 phenomenon [26] and 0.6% for the post-meal glucose increments. Consequently, such a reduction in those
38 individuals with HbA1c level below 6.8% should result in a near normal level of HbA1c (<5.8%). It should be
39 noted that the reduction can be even more pronounced in certain individuals with a fall of 1.4% (0.4% for the
40 dawn phenomenon and 1% for the post-meal glucose increments). Consequently, those with an HbA1c as high as
41 7.5% can achieve a near normal HbA1c level provided that both the dawn phenomenon and abnormal
42 postprandial glucose excursions are eradicated. Consequently the risk for the development and progression of
43 micro or macro-vascular complications could be either eradicated or at least strongly delayed according to the
44 data from the United Kingdom Prospective Study (UKPDS) [16] and as has been suggested by Zoungas et al
45 [35].
46
47
48
49
50
51
52
53
54
55
56
57
58

59 **2.3 Basal hyperglycaemia**

1 When the HbA1c is between 7.5 and 8%, the relative contribution of the postprandial and basal hyperglycaemia
2 to the overall hyperglycaemia becomes equivalent, whereas beyond 8% basal hyperglycaemia becomes
3 increasingly the predominant defect [28, 36]. This observation is simply due to the fact that the absolute
4 contribution of the postprandial hyperglycaemia across the increasing HbA1c spectrum beyond 7% remains
5 stable at approximately of 1% of HbA1c as mentioned above [29]. Therefore, any increase in glucose exposure
6 beyond this HbA1c threshold of 7% is due to a linear increase in absolute basal hyperglycaemia [28] evident
7 beyond 8% of HbA1c [28,29]. This therefore represents the final stage in the worsening of glycaemic control in
8 persons with type 2 diabetes, corresponding predominantly to a progressive deterioration of basal
9 hyperglycaemia during both the diurnal and nocturnal periods.

10 In summary, this stepwise deterioration of glycaemic control is illustrated schematically in figure 3 representing
11 the initial stage of an isolated dawn phenomenon (HbA1c below 6.5%), followed by the combination of a dawn
12 phenomenon with postprandial hyperglycaemia (HbA1c between 6.5 and 6.9%), i.e. when subjects are in the so-
13 called of “intermediate” stage of dysglycaemia. The final phase is reached with the addition of progressive basal
14 hyperglycaemia, when the HbA1c is $\geq 7\%$, being the last member of the “triumvirate” of abnormalities seen in
15 type 2 diabetes.

23 **3. Relationship between postprandial glucose excursions and glycaemic variability**

24
25
26 Glycaemic variability refers to glucose fluctuations from peaks to nadirs [37-40]. Peaks, especially in type 2
27 diabetes, usually correspond to postmeal glucose excursions while nadirs can indicate risk of hypoglycaemia
28 during the inter-prandial and nocturnal periods. The glycaemic variability has been extensively described and
29 investigated during the last decade mainly by using the newly developed technology of ambulatory continuous
30 glucose monitoring (CGM). At present, several methods are employed to quantifying the magnitude of glucose
31 variability [39-42] including: the standard deviation (SD) around the mean glucose value, MAGE (Mean
32 Amplitude of Glycaemic Excursions) and the MODD (Mean Of Daily Differences), the CONGA (Continuous
33 Overlapping Net Glycaemic Action), the M-index of Schlichtkrull [43] and the Lability Index [44]. These
34 parameters investigate either the within-day or between-day glucose variability for which CGM is required
35 available in a limited number of medical units. Therefore, one of the new challenges in the near future is to
36 develop simplified methods for the assessment of glycaemic variability. Such methods should be able to be
37 based on the measurement of capillary glucose concentrations by using the structured self-blood glucose
38 monitoring at accurately selected time-points. One of the approaches for addressing this issue is to consider that,
39 at least in type 2 diabetes, glycaemic variability and postprandial glucose excursions are inter-correlated. By
40 using CGM in a population of 63 patients with type 2 diabetes, Suh et al [45] recently demonstrated that
41 postprandial glucose excursions were strongly correlated with glycaemic variability, at least in a subset of
42 subjects who were under reasonable glycaemic control (HbA1c < 7.5%). These findings are in general
43 agreement with our own observations in a post hoc analysis of the results of a multicenter prospective
44 randomized trial [46] that was conducted in a small number (n = 30) of persons with type 2 diabetes treated with
45 metformin and secondarily allocated after randomization to one of two DPP-4 inhibitor preparations either
46 vildagliptin (n= 14) or sitagliptin (n = 16) as add-on therapy for a period of 8 weeks. In a post hoc analysis of the
47 results of this study, we observed that the changes in postprandial glucose increments from baseline to end point
48 were strongly and positively correlated with those of glycaemic variability (MAGE) [30]. In addition, a few
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 years ago, we noted that in most persons with type 2 diabetes, the highest postprandial glucose peaks recorded
2 were mid-morning or 1 or 2 hours after having breakfast [27]. These abnormally high glucose excursions after
3 breakfast may be referred as an "extended dawn phenomenon".

4 We then went on to investigate in people with non-insulin-treated type 2 diabetes (n = 271) the relationship
5 between the increment in blood glucose from the pre-breakfast value after an overnight fast to the 90-min post
6 breakfast value (x value) and the glycaemic variability assessed from the SD around the 24-hour mean glucose
7 value (y value). A highly significant positive relationship was observed between the two parameters: $r = 0.74$, p
8 < 0.0001 (figure 6). These results not only confirm that glycaemic variability and postprandial glucose
9 excursions are strongly linked but also provide a simpler method to CGM for quantifying the glycaemic
10 variability based on only two determinations of capillary glucose at pre and post breakfast time points sufficient
11 to allow a reliable assessment of within-day glucose fluctuations.
12
13
14
15
16

17 **4. Glucose variability: is this glycaemic disorder an important matter?**

18
19
20 At present, the role of glycaemic variability on the development and progression of cardiovascular diseases is a
21 subject of debate. In a recent issue of Diabetes Care, Hirsch and Bergenstal had a Point-Counterpoint debate
22 [47,48]. Hirsch provided arguments that glucose fluctuations are deleterious and that control of glycaemic
23 variability should be a primary treatment target [47]. Bergenstal instead argued that we should give preferential
24 consideration to other markers than glycaemic variability [48].
25
26
27

28 **4.1 Glycaemic variability: worthy of consideration?**

29
30
31 Ten years ago, Ceriello et al [49] epitomised postprandial excursions as "dangerous waves" for vascular
32 endothelial cells of vascular walls through the activation of oxidative stress, one of the key pathophysiological
33 mechanisms for the development of diabetic vascular complications [50,51]. Subsequently this hypothesis was
34 confirmed by us when we demonstrated that in non-insulin-using type 2 diabetic patients, the 24-hour urinary
35 excretion rate of 8-iso PGF₂α a marker of the activation of oxidative stress, was strongly correlated with the
36 Mean Amplitude of Glycaemic Excursions (MAGE) [52] which was later re-affirmed by Ceriello [53]. Using the
37 glucose clamp technique in persons with type 2 diabetes, upward and downward swings of glucose
38 concentrations are paralleled by oscillating plasma levels of nitrotyrosine, another marker of oxidative stress
39 [53]. More recently, we reported that the 24-hour urinary excretion rate of 8-isoPGF₂α depends equally on
40 the ambient hyperglycaemia and the glycaemic variability, as estimated by HbA1c levels and MAGE
41 respectively, in non-insulin-treated type 2 diabetes [54]. According to these and other observations, the
42 pathogenesis of diabetic complications appear to be a consequence of both chronic hyperglycaemia and short-
43 term excessive glucose variability resulting in enhanced glycation and activation of oxidative stress.
44
45
46
47
48
49
50
51

52 **4.2 Glycaemic variability: significance in diabetic complications?**

53
54
55 Key observations opposing the view that glycaemic variability is important in the development of diabetic
56 complications were mainly provided by two retrospective analyses of the Diabetic Control and Complications
57 Trial (DCCT) [55,56] and the HEART2D Study [57]. Analysis of the DCCT data set [55,56] concluded that
58 glucose variability has only a minor contribution to vascular complications. However, it should be noted that the
59
60
61
62
63
64
65

conclusions were limited to microvascular complications in persons with type 1 diabetes on insulin treatment. The HEART2D study was initially designed to answer whether control of basal hyperglycaemia or postprandial hyperglycaemia is best for reducing cardiovascular outcomes in patients with poorly controlled type 2 diabetes who had a history of myocardial infarction [57]. Participants were further assigned to either a basal insulin strategy targeting fasting and inter-prandial glycaemia or an insulin regimen with three daily injections of a rapid-acting insulin analogue at premeal times in order to attenuate postprandial glucose excursions. A similar lowering effect of ambient hyperglycaemia represented by HbA1c was observed with the two insulin regimens. No difference in the incidence of cardiovascular events was detected between the two regimens despite the lower postprandial glycaemia with the prandial insulins compared with the basal group at interim analysis, when the study was halted after a mean follow-up of 2.7 years. It was therefore concluded that better control of postprandial excursions and probably reduced glucose variability does not provide any benefit in terms of macrovascular outcomes.

4.3 Pros and cons of glycaemic variability as a risk factor of macrovascular diseases

From the aforementioned studies, there arises the question as to why glycaemic variability as an activator of oxidative stress in non-insulin-treated type 2 diabetes, did not appear to exert any significant influence on the cardiovascular outcome in those treated with insulin. We derived an answer to this question with a cross-sectional study that compared three groups of subjects including type 1 diabetes and type 2 diabetes treated either with oral hypoglycaemic agents or in combination with insulin [54]. The 24-hour urinary excretion rate of 8-isoPGF2 alpha was only elevated in those subjects treated with oral hypoglycaemic agents (OHAs) remaining within the normal range in the other two groups treated with insulin. These results were observed despite the fact that glycaemic variability (MAGE) was significantly higher in those on insulin versus those on oral therapy alone. In addition, in a subgroup of those with type 2 diabetes, the 24-hour urinary excretion rate of 8-isoPGF2alpha was evaluated at baseline whilst on treatment with OHAs and several months or years after initiation of insulin treatment. A drastic reduction in the 24-hour urinary excretion rates of 8-isoPGF2alpha was seen from a highly elevated level before initiation of insulin returning to within the normal range following the introduction of insulin treatment. These results strongly suggest that insulin *per se* exerts an inhibitory effect on the activation of oxidative stress [58].

From a clinical care point of view, these observations support the concept that treatments with insulin should be implemented as early as possible in the time course of type 2 diabetes [59], bearing in mind the clinical characteristics of the patient [13]. Therefore, randomized interventional control trials comparing 2 groups of subjects maintained at similar levels of HbA1c but submitted to therapeutic strategies aimed at reducing glycaemic variability in one group but not in the comparator, is needed to provide a clear answer as to whether glycaemic variability contributes to adverse cardiovascular outcomes in type 2 diabetes. The protocol of the ongoing FLAT-SUGAR Study has been designed to address this issue [60]. In this proof-of-concept study, participants will be randomized and enrolled into two groups according to whether they will be treated with a basal-bolus insulin regimen or a combination of basal insulin and a prandial GLP-1 receptor agonist. The primary end point will be the changes in glycaemic variability that should normally be lower in the latter group than in the former. The secondary end point will be changes in several biological markers of inflammation,

1 activation of the oxidative stress, renal and cardiac functions. Should the results of this preliminary study be
2 consistent with the hypothesis, this would lead to a long-term study with “hard” cardiovascular outcomes in
3 order to validate the potential role of glycaemic variability.
4

5 **5. Relationship between glycaemic variability and hypoglycaemia**

6
7
8 As hypoglycaemic episodes can contribute to the occurrence of adverse cardiovascular events [8, 61-66], it is
9 important to know whether glycaemic variability contributes to hypoglycaemia. It is well accepted that
10 hypoglycaemia is more frequent when mean glucose values are low, with recent data [67, 68] indicating that
11 increased glucose fluctuations around the mean glucose value can play an additional role in precipitating
12 hypoglycaemia. These results based on the DOVES trial [67] and the DCCT data set [68] have been confirmed
13 by our own data [69]. Similar results were observed in an unpublished analysis by ourselves of 828 day-patient
14 glycaemic profiles that were obtained from ambulatory CGM carried out in subjects with type 1 diabetes (331),
15 type 2 diabetes treated with insulin (216) and non-insulin-treated type 2 diabetes (222). In each group, the
16 frequency of hypoglycaemic episodes (defined as all interstitial glucose values < 56 mg/dL, 3.3 mmol/L) was
17 calculated after the 3 groups had been divided into 3 further subgroups according to whether the 24-hour mean
18 glucose value was < 150 mg/dL, (8.3 mmol/L), between 150 and 180 mg/dL (8.3-10.0 mmol/L) or > 180 mg/dL
19 (10 mmol/L). Finally, in each subset, the frequency of hypoglycaemic episodes was compared according to
20 whether the glycaemic variability (SD around the mean glucose value) was above or below the mean SD in each
21 selected subgroup i.e. 60 mg/dL in type 1 diabetes, 50 mg/dL in insulin-treated type 2 diabetes and 30 mg/dL in
22 non-insulin-treated subjects, values were rounded to the nearest ten. The results are illustrated in figure 7a for
23 type 1 diabetes, 7b insulin-treated type 2 diabetes, and 7c non-insulin-treated type 2 diabetes. The frequency of
24 hypoglycaemic episodes ranked according to decreasing ordinal scale were Type 1 diabetes > insulin-treated
25 type 2 diabetes > non-insulin-treated type 2 diabetes. In each type of diabetes, as expected, the ranking frequency
26 of hypoglycaemic episodes was as follows: study days with mean glucose values less than 150 mg/dL (8.3
27 mmol/L) > study days with mean glucose values between 150 and 180 mg/dL (8.3-10.0 mmol/L) > study days
28 with mean glucose values > 180mg/dL (10 mmol/L). More importantly in each subgroup of patients, the
29 frequency of hypoglycaemic episodes increased substantially when the glycaemic variability was increased
30 above the mean SD value. Glycaemic variability seems therefore to be strongly associated with hypoglycaemia,
31 although it is not possible to know whether glycaemic variability is the chicken or the egg. Glycaemic variability
32 being a causative risk factor for hypoglycaemia is a possibility as the majority of hypoglycaemic episodes
33 recorded in the present study were asymptomatic and thus were not subject to a subsequent glucose rebound due
34 to excess self-treatment with carbohydrate-containing drinks or snacks.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **6. Concomitant reduction of ambient hyperglycaemia and glycaemic variability: Is this possible?**

51
52 The benefits of attaining normal or near normal levels of HbA1c with its ensuing reduction in the risk of diabetic
53 complications should not be counterbalanced by an increased risk of hypoglycaemic episodes [8-10, 66].
54 Therefore, during the stage of “intermediate” dysglycaemia where basal hyperglycaemia is absent and glucose
55 values during nocturnal periods are within the normal range [34], therapeutic intervention should avoid the use
56 of anti-diabetic agents or strategies that increase the risk of hypoglycaemia. Such agents include the
57 sulfonylureas or glinides [70], basal insulin [71]. Instead insulin sensitizers [72] incretin-based therapies such as
58
59
60
61
62
63
64
65

1 the DPP-4 inhibitors or the GLP-1 receptor agonists should be employed at this stage of the natural history of
2 type 2 diabetes [73-78]. Unfortunately glycaemic variability and chronic ambient hyperglycaemia were rarely
3 studied in parallel in large interventional trials. For instance, in the 4-T study [79], HbA1c levels were improved
4 when patients were switched from a basal insulin therapy to a basal-bolus insulin regimen although this resulted
5 in an increased frequency of hypoglycaemic episodes. As the glycaemic variability was not assessed it is not
6 possible to know whether this effect was due or not to an increase in the magnitude of glucose excursions.
7 Another example is the 4B study [80] that was designed to compare the efficacy and safety of exenatide twice
8 daily or thrice daily mealtime insulin lispro in those inadequately controlled by optimised insulin glargine and
9 metformin therapy. Whilst similar improvements in ambient hyperglycaemia (HbA1c) were observed in both
10 groups, exenatide resulted in fewer episodes of hypoglycaemia considered as a whole (30% vs 41% respectively)
11 and more particularly during the day (15% vs 34% respectively). However, given the fact that exenatide is a
12 GLP-1 receptor agonist mainly aimed at reducing postprandial glucose excursions, it is highly likely that this
13 category of antidiabetic agents can be of value in reducing both ambient hyperglycaemia and glucose variability.
14 In a post-hoc analysis of the OPTIMA study [30] using CGM it has been demonstrated that the DPP-4 inhibitors
15 sitagliptin and vildagliptin, when prescribed as add-on therapy in persons with type 2 diabetes not sufficiently
16 controlled on metformin alone, resulted in a significant reduction in ambient hyperglycaemia (HbA1c, AUC
17 under glycaemic profiles). This reduction was concomitantly associated with significant decreases in
18 postprandial glucose exposure and glycaemic variability (MAGE) with a positive relationship between these 2
19 parameters [30]. These results demonstrate that it is possible to reduce both the ambient hyperglycaemia and
20 glucose variability in a concomitant manner. Therefore, the assessment of glycaemic variability should be more
21 extensively and thoroughly evaluated in all forthcoming interventional trials. Such assessments will be crucial to
22 define which drugs have the potential to reduce glucose fluctuations and to finally resolve the debate as to
23 whether reduction in glycaemic variability has a positive impact or not on cardiovascular outcomes.
24
25
26
27
28
29
30
31
32
33
34

35 **7. Conclusion**

36
37 The current situation can be summarized by 3 terminologies: “consensus”, “disensus”, and “non sensus”.
38 “Consensus” because, despite a small number of discordant opinions, it is now well accepted that chronic
39 exposure to ambient hyperglycaemia is a key player in the pathogenesis of diabetic complications. The
40 “disensus” is based on the fact that there exists a large spectrum of opinions on the role of both postprandial
41 glucose excursions and glycaemic variability as risk factors for adverse cardiovascular outcomes. The “non
42 sensus” would be to deny any impact of short-term excess postprandial glucose fluctuations on the development
43 or progression of vascular complications.
44
45
46
47

48 Although at present, we lack the evidence, we hope that the ongoing or future randomised control interventional
49 trials will be able to provide a clear answer to the question as to whether more consideration should be given to
50 postprandial hyperglycaemia and glucose variability which are regarded as minor or ancillary components of
51 dysglycaemia relative to overall hyperglycaemia.
52
53
54

55 **Disclosure of interest**

56 All authors declare that they have no conflict of interest with the content of this review
57
58
59
60
61
62
63
64
65

References

- 1
2 [1] Kirkman MS, Mc Carren M, Shah J, Duckworth W, Abraira C. The association between metabolic
3 control and prevalent macrovascular disease in type 2 diabetes: the VA Cooperative Study in diabetes. *J*
4 *Diabetes Complications* 2006;20:75-80.
5
6
7 [2] Selvin E, Marinopoulos S, Berkenblit G, Rami T, Brancati FL, Powe NR et al. Meta-analysis
8 glycosylated hemoglobin and cardiovascular disease in diabetes mellitus. *Ann Intern Med* 2004;141:421-
9 31.
10
11
12 [3] The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of
13 diabetes on the development and progression of long-term complications in insulin-dependent diabetes
14 mellitus. *N Engl J Med* 1993 ;329 :977-86.
15
16
17 [4] Stratton IM, Adler AI, Neil HAW, Matthews DR, Manley SE, Cull CA et al. Association of glycaemia
18 with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective
19 observational study. *BMJ* 2000; 321: 405-12.
20
21
22 [5] Sacks DB, Bruns DE, Goldstein DE, Maclaren NK, Mc Donald JM, Parrott M. Guidelines and
23 recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus (Position
24 statement). *Diabetes Care* 2002; 25:750-86.
25
26
27 [6] Sacks DB. Measurement of HbA1c. A new twist on the path to harmony. *Diabetes Care* 2012 ;35 :2674-
28 80.
29
30
31 [7] Selvin E, Steffes MW, Zhu H, Matsushita K, Wagenknecht L, Pankow J et al. Glycated hemoglobin,
32 diabetes and cardiovascular risk in non diabetic adults. *N Engl J Med* 2010 ;362 :800-11.
33
34
35 [8] The Action to Control Cardiovascular Risk in Diabetes Study Group. Effect of intensive glucose lowering
36 in type 2 diabetes. *N Engl J Med* 2008 ;358 :2545-59.
37
38
39 [9] The ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients
40 with type 2 diabetes. *N Engl J Med* 2008 ;358 :2560-72.
41
42
43 [10] Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD et al, for the VADT Investigators.
44 Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;
45 360:129-39.
46
47
48 [11] Turnbull FM, Abraira C, Anderson RJ, Byinton RP, Chalmers JP, Duckworth WC et al, for the Control
49 Group. Intensive glucose control and macrovascular outcomes in type 2 diabetes. *Diabetologia* 2009 ;52
50 :2288-98.
51
52
53 [12] Bejan-Angoulvant T, Cornu C, Archambault P, Tudrej B, Audier P, Brabant Y et al. Is HbA1c a valid
54 surrogate for macrovascular and microvascular complications in type 2 diabetes? *Diabetes Metab*
55 2015;41:195-201.
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [13] Inzucchi SE, Bergenstal RM, Buse JB, Diamant M, Ferrannini E, Nauck M et al. Management of hyperglycemia in type 2 diabetes, 2015 : A patient-centered approach. Update to a position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 2015 ;38 :140-9.
 - [14] The Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med* 2005 ;353 :2643-53.
 - [15] Lachin JM, Orchard TJ, Nathan DM, for the DCCT/EDIC Research Group. Update on cardiovascular outcomes at 30 years of the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study. *Diabetes Care* 2014 ;37 :39-43.
 - [16] Holman RR, Paul SK, Bethel MA, Matthews DR, Neil HAW. 10-years follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008 ;359 :1577-89.
 - [17] Hayward RA, Reaven PD, Wiitala WL, Bahn GD, Reda DJ, Ge L et al. Follow-up of glycemic control and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015 ;372 :2197-206.
 - [18] Gerstein HC, Miller ME, Ismail-Beigi F, Largay J, McDonald C, Lochnan HA et al, for the ACCORD study Group. Effects of intensive glycemic control on ischemic heart disease analysis of data from the randomised, controlled ACCORD trial. *Lancet* 2014;384 :1936-41.
 - [19] Gaede P, Lund-Andersen H, Parving HH, Pedersen O. Effect of multifactorial intervention on mortality in type 2 diabetes. *New Engl J Med* 2008 ;358 :580-91.
 - [20] Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S et al, for the EMPA-REG OUTCOME Investigators. Empagliflozin, cardiovascular outcomes and mortality in type 2 diabetes. *N Engl J Med* 2015.DOI :10.1056/NEJMoa 1504720.
 - [21] American Diabetes Association. Standards of Medical Care in Diabetes-2015. 6. Glycemic targets. *Diabetes Care* 2015 ;38 (Suppl 1) :S33-40.
 - [22] IDF 2012 Clinical Guidelines Task Force. Global guideline for type 2 diabetes. *Diabetes Res Clin Pract*.<http://dx.doi.org/10.1016/j.diabres.2012.10.001>.
 - [23] American Diabetes Association. Standards of Medical Care in Diabetes-2015. 2. Classification and diagnosis of diabetes. *Diabetes Care* 2015;38 (Suppl 1): S8-16.
 - [24] Bolli GB, Gerich JE. The "dawn phenomenon" - a common occurrence in both non-insulin-dependent and insulin-dependent diabetes mellitus. *N Engl J Med* 1984; 310: 746-50.
 - [25] Carroll MF, Schade DS. The dawn phenomenon revisited: implications for diabetes therapy. *Endocr Pract* 2005; 11: 55-64.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [26] Monnier L, Colette C, Dejager S, Owens D. Magnitude of the dawn phenomenon and its impact on the overall glucose exposure in type 2 diabetes. Is this of concern? *Diabetes Care* 2013; 36: 4057-62.
- [27] Monnier L, Colette C, Dunseath GJ, Owens DR. The loss of postprandial glycemic control precedes stepwise degradation of fasting with worsening diabetes. *Diabetes Care* 2007; 30: 263-9.
- [28] Monnier L, Lapinski H, Colette C. Contribution of fasting and postprandial plasma glucose increments to the overall diurnal hyperglycemia of type 2 diabetic patients: variations with increasing levels of HbA1c. *Diabetes Care*. 2003 ;26 :881-5.
- [29] Monnier L, Colette C, Owens D. Post-prandial and basal glucose in type 2 diabetes : assessment and respective impacts. *Diabetes Technol Ther* 2011 ;13 (Suppl 1) :S25-32.
- [30] Monnier L, Colette C, Dejager S, Owens DR. « Mild dysglycemia » in type 2 diabetes : to be neglected or not ? *J Diabetes Complications* 2015 ;29 :451-8.
- [31] Boden G, Chen X, Urbain JL. Evidence for a circadian rhythm of insulin sensitivity in patients with NIDDM caused by cyclic changes in hepatic glucose production. *Diabetes* 1996; 45:1044-50.
- [32] Porcellati F, Lucidi P, Bolli GB, Fanelli CG. Thirty years of research on the dawn phenomenon: lessons to optimize blood glucose control in diabetes. *Diabetes Care* 2013; 36: 3860-2.
- [33] De Feo P, Perriello G, Ventura MM, Calcinaro F, Basta G, Lolli C et al. Studies on overnight insulin requirements and metabolic clearance rate of insulin in normal and diabetic man: relevance to the pathogenesis of the dawn phenomenon. *Diabetologia* 1986; 29: 475-80.
- [34] Monnier L, Colette C, Dejager S, Owens D. Residual dysglycaemia when at target HbA1c of 7% (53 mmol/mol) in persons with type 2 diabetes. *Diabetes Res Clin Pract* 2014;104:371-5.
- [35] Zoungas S, Chalmers J, Ninomiya T, Li Q, Cooper ME, Colagiuri S et al. Association of HbA1c levels with vascular complications and death in patients with type 2 diabetes : evidence of glycemic thresholds. *Diabetologia* 2012 ;55 :636-43.
- [36] Peter R, Dunseath G, Luzio SD, Chudleigh R, Choudhury SR, Owens D. Relative and absolute contributions of postprandial and fasting plasma glucose to daytime hyperglycaemia and HbA(1c) in subjects with Type 2 diabetes. *Diabet Med* 2009 ;26 :974-80.
- [37] Monnier L, Colette C. Glycemic variability. Should we and can we prevent it? *Diabetes Care* 2008;31 (Suppl 2): S150-4.
- [38] Monnier L, Colette C. Glycemic variability : can we bridge the divide between controversies ? *Diabetes Care* 2011 ;34 :1058-9.
- [39] DeVries JH. Glucose variability : Where is it important and how to measure it. *Diabetes* 2013 ;62 :1405-8.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [40] Service FJ. Glucose variability. *Diabetes* 2013 ;62 :1398-404.
- [41] Rodbard D. New and improved methods to characterize glycemic variability using continuous glucose monitoring. *Diabetes Technol Ther* 2009 ;11 :551-65.
- [42] Weber C, Schnell O. The assessment of glycemic variability and its impact on diabetes-related complications : An overview. *Diabetes Technol Ther* 2009 ;11 :623-33.
- [43] Schlichtkrull J, Munck O, Jersild M. The M-value, an index of blood sugar control in diabetes. *Acta Med Scand* 1965;177:95-102.
- [44] Paty BW, Senior PA, Lakey JR, Shapiro AM, Ryan EA. Assessment of glycemic control after islet transplantation using the continuous glucose monitor in insulin-independent versus insulin-requiring type 1 diabetes subjects. *Diabetes Technol Ther* 2006;8:165-73.
- [45] Suh S, Joung JY, Jin S-M, Kim MY, Bae JC, Park H-D et al. Strong correlations between glycaemic variability and total glucose exposure in type 2 diabetes are limited to subjects with a satisfactory glycaemic control. *Diabetes Metab* 2014; 40: 272-7.
- [46] Guerci B, Monnier L, Serusclat P, Petit C, Valensi P, Huet D et al. Continuous glucose profiles with vildagliptin versus sitagliptin in add-on to metformin: Results from the randomized Optima study. *Diabetes Metab* 2012;38: 359-66.
- [47] Hirsch IB. Glycemic variability and diabetes complications : Does it matter ? Of course it does ! *Diabetes Care* 2015 ;38 :1610-14.
- [48] Bergenstal RM. Glycemic variability and diabetes complications : Does it matter ? Simply put, there are better glycemic markers ! *Diabetes Care* 2015 ;38 :1615-21.
- [49] Ceriello A, Quagliaro L, Catone B, Pascon R, Piazzola M, Bais B et al. Role of hyperglycemia in nitrotyrosine postprandial generation. *Diabetes Care* 2002 ;25 :1439-43.
- [50] Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001 ;414 :813-20.
- [51] Brownlee M. The pathobiology of diabetic complications : a unifying mechanism. *Diabetes* 2005 ;54 :1615-25.
- [52] Monnier L, Mas E, Ginet C, Michel F, Villon L, Cristol JP et al. Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA* 2006; 295:1681-7.
- [53] Ceriello A, Esposito K, Piconi L, Ihnat MA, Thorpe JE, Testa R et al. Oscillating glucose is more deleterious to endothelial function and oxidative stress than mean glucose in normal and type 2 diabetic patients. *Diabetes* 2008; 57: 1349-54.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [54] Monnier L, Colette C, Mas E, Michel F, Cristol JP, Boegner C et al. Regulation of oxidative stress by glycaemic control : evidence for an independent inhibitory effect of insulin therapy. *Diabetologia* 2010 ;53 :562-71.
- [55] Kilpatrick ES, Rigby AS, Atkin SL. The effect of glucose variability on the risk of microvascular complications in type 1 diabetes. *Diabetes Care* 2006;29:1486-90.
- [56] Lachin JM, Genuth S, Nathan DM, Zinman B, Rutledge BN for the DCCT/EDIC Research Group. Effect of glycemic exposure on the risk of microvascular complications in the Diabetes Control and Complications Trial-revisited. *Diabetes* 2008; 57: 995-1001.
- [57] Raz I, Wilson PWF, Strojek K, Kowalska I, Bozikov V, Gitt AK et al. Effects of prandial versus fasting hyperglycemia on cardiovascular outcomes in type 2 diabetes : the HEART2D trial. *Diabetes Care* 2009 ;32 :381-6.
- [58] Monnier L, Hanefeld M, Schnell O, Colette C, Owens D. Insulin and atherosclerosis: how are they related? *Diabetes Metab* 2013;39:111-7.
- [59] Hanefeld M. Use of insulin in type 2 diabetes: What we learned from recent clinical trials on the benefits of early insulin initiation. *Diabetes Metab* 2014;40:391-9.
- [60] The FLAT-SUGAR Trial Investigators. Design of FLAT-SUGAR randomized trial of prandial insulin versus prandial GLP-1 receptor agonist together with basal insulin and metformin for high-risk Type 2 diabetes. *Diabetes Care* 2015 ;38 :1558-66.
- [61] Amiel SA, Dixon T, Mann R, Jameson K. Hypoglycaemia in type 2 diabetes. *Diabet Med* 2008 ;25 :245-54.
- [62] Cryer PE. The barrier of hypoglycemia in diabetes. *Diabetes* 2008 ;57 :3169-76.
- [63] Zoungas S, Patel A, Chalmers J, de Galan BE, Li Q, Biostat M. Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 2010; 363: 1410-8.
- [64] Bonds DE, Miller ME, Bergenstal RM, Buse JB, Byington RP, Cutler JA et al. The association between symptomatic, severe hypoglycemia and mortality in type 2 diabetes: retrospective epidemiological analysis of the ACCORD Study. *BMJ* 2010; 340:b4909.doi:1136/bmj.b4909.
- [65] Frier BM, Schernthaner G, Heller SR. Hypoglycemia and cardiovascular risks. *Diabetes Care* 2011 ;34 (Suppl 2) : S132-7.
- [66] Cryer PE. Glycemic goals in diabetes: trade-off between glycemic control and iatrogenic hypoglycemia. *Diabetes* 2014; 63: 2188-95.
- [67] Murata GH, Hoffman RM, Shah JH, Wendel CS, Duckworth WC. A probabilistic model for predicting hypoglycemia in type 2 diabetes mellitus. The Diabetes Outcomes in Veterans Study (DOVES). *Arch Intern Med* 2004 ;164 :1445-50.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [68] Kilpatrick ES, Rigby AS, Goode K, Atkin SL. Relating mean blood glucose and glucose variability to the risk of multiple episodes of hypoglycemia in type 1 diabetes. *Diabetologia* 2007;50 :2553-61.
- [69] Monnier L, Wojtusciszyn A, Colette C, Owens D. The contribution of glucose variability to asymptomatic hypoglycemia in persons with type 2 diabetes. *Diabetes Technol Ther* 2011;13:813-8.
- [70] Lebovitz HE. Insulin secretagogues : old and new. *Diabetes Rev* 1999 ;7 :139-53.
- [71] The ORIGIN Trial Investigators. Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med* 2012; 367: 319-28.
- [72] Scheen A, Paquot N. Metformin revisited: a critical review of the benefit-risk balance in at-risk patients with type 2 diabetes. *Diabetes Metab* 2013; 39: 179-90.
- [73] Ahrén B, Lequizamo Dimas A, Miossec P, Saubadu S, Aronson R. Efficacy and safety of lixisenatide once-daily morning or evening injections in type 2 diabetes inadequately controlled on metformin (GetGoal M). *Diabetes Care* 2013; 36: 2543-50.
- [74] Bergenstal RM, Wysham C, Macconell L, Malloy J, Walsh B, Yan P et al. Duration-2 Study Group. Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomized trial. *Lancet* 2010; 376: 431-9.
- [75] Bolli GB, Owens DR. Lixisenatide, a novel GLP-1 receptor agonist: efficacy, safety and clinical implications for type 2 diabetes. *Diabetes Obes Metab* 2014; 15: 588-601.
- [76] DeFronzo RA, Hissa MN, Garber AJ, Luiz Gross J, Yuyan Duan R, Ravichandran S et al, for the saxagliptin 014 Study Group. The efficacy and safety of saxagliptin when added to metformin therapy in patients with inadequately controlled type 2 diabetes with metformin alone. *Diabetes Care* 2009;32:1649-55.
- [77] Drucker DJ, Nauck MA. The incretin system : glucagon-like peptide-1 receptor agonists and dipeptidyl-peptidase inhibitors in type 2 diabetes . *Lancet* 2006 ;368 :1696-705.
- [78] Owens DR, Monnier L, Bolli GB. Differential effects of GLP-1 receptor agonists on components of dysglycaemia in individuals with type 2 diabetes mellitus. *Diabetes Metab* 2013; 39:485-96.
- [79] Holman RR, Farmer AJ, Davies, Levy JC, Darbyshire JL, Keenan JF et al, 4-T Study Group. Three-year efficacy of complex insulin regimens in type 2 diabetes. *N Engl J Med* 2009;361:1736-47.
- [80] Diamant M, Nauck MA, Shaginan R, Malone JK, Cleall S, Reaney M et al, for the 4B Study Group. Glucagon-like peptide receptor agonist or bolus insulin with optimized basal insulin in type 2 diabetes. *Diabetes Care* 2014;37:2763-73.

Legends of figures

1
2 **Fig. 1** Influence of duration of follow-up upon the relative risk of major cardiovascular outcomes in patients with
3 type 2 diabetes submitted to intensive glucose lowering as compared with standard therapy. A significant
4 reduction in the relative risk was only observed when the duration of follow-up was longer than 10 years:
5 extensions of the UKPDS [16] and VADT [17] studies. In contrast the use of intensive therapy did not
6 significantly reduce major cardiovascular events when the follow-up remained shorter than [8-10] or nearly 10
7 years [4]
8
9

10
11 **Fig. 2** The respective contributions of the different components of glucose exposure in persons with type 2
12 diabetes are depicted on the left part of the figure (i) normal glucose exposure (grey shaded rectangle); (ii)
13 exposure to basal hyperglycaemia (black area) and (iii) exposure to postprandial hyperglycaemia (white area).
14 The dawn phenomenon corresponds to the spontaneous rise in basal hyperglycaemia observed before breakfast
15 time. The aims of treatment are indicated on the right part of the figure
16
17
18
19

20 **Fig. 3** Respective contributions of the 3 main glycaemic disorders (dawn phenomenon, basal and postprandial
21 hyperglycaemia) to the dysglycaemia of type 2 diabetes mellitus across the HbA1c spectrum. The dawn
22 phenomenon is a permanent feature of glycaemic disorders with worsening HbA1c
23
24
25

26 **Fig. 4** Mean 24-hour glycaemic profile in persons with type 2 diabetes, who were divided into 2 groups. Group
27 1: HbA1c < 6.5% group 2: HbA1c between 6.5 and 6.9% (n = 50). Exposures to basal and postprandial
28 hyperglycaemia are illustrated by black and white areas, respectively
29
30

31 **Fig. 5** Absolute impact of postprandial hyperglycaemia on HbA1c (expressed as percent points) in persons with
32 type 2 diabetes across the HbA1c continuum/spectrum [29]. Reprinted with the permission of Mary Ann Liebert,
33 Inc., publishers
34
35
36

37 **Fig. 6** Relationship between postprandial glucose excursions (x = increment from the breakfast glucose value to
38 the 90-min postbreakfast glucose value (mg/dL) and the glycaemic variability (y = SD around the 24-h mean
39 glucose value, mg/dL).
40
41

42 **Fig. 7** Total hypoglycaemic episodes (interstitial glucose concentration < 56 mg/dL during continuous glucose
43 monitoring) expressed as number/patient-day in 3 groups of persons with diabetes: type 1 (n = 313, figure 7a);
44 type 2 treated with insulin (n = 216, figure 7b) and type 2 treated only with diet or oral antidiabetic agents (n =
45 222, figure 7c).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1




Figure 2




Figure 3

Basal hyperglycaemia largely > Postprandial hyperglycaemia

Basal hyperglycaemia approximately equivalent to postprandial hyperglycaemia

Postprandial hyperglycaemia > Basal hyperglycaemia

Isolated postprandial hyperglycaemia

Dawn phenomenon

Figure 4





Figure 5

Absolute impact of postprandial hyper



Groups

1

2

3

Mean HbA1c (%)

5.9

6.8

7.4

Figure 6




Figure 7

