



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in :  
*Quaternary Geochronology*

Cronfa URL for this paper:  
<http://cronfa.swan.ac.uk/Record/cronfa26996>

---

### **Paper:**

Chen, X., Blockley, S., Tarasov, P., Xu, Y., McLean, D., Tomlinson, E., Albert, P., Liu, J., Müller, S., Wagner, M. & Menzies, M. (2016). Clarifying the distal to proximal tephrochronology of the Millennium (B–Tm) eruption, Changbaishan Volcano, northeast China. *Quaternary Geochronology*, 33, 61-75.

<http://dx.doi.org/10.1016/j.quageo.2016.02.003>

---

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

# Accepted Manuscript

Clarifying the distal to proximal tephrochronology of the Millennium (B-Tm) eruption, Changbaishan Volcano, northeast China

Xuan-Yu Chen, Simon P.E. Blockley, Pavel E. Tarasov, Yi-Gang Xu, Danielle McLean, Emma L. Tomlinson, Paul G. Albert, Jia-Qi Liu, Stefanie Müller, Mayke Wagner, Martin A. Menzies

PII: S1871-1014(16)30009-7

DOI: [10.1016/j.quageo.2016.02.003](https://doi.org/10.1016/j.quageo.2016.02.003)

Reference: QUAGEO 754

To appear in: *Quaternary Geochronology*


Received Date: 22 October 2015

Revised Date: 12 February 2016

Accepted Date: 16 February 2016

Please cite this article as: Chen, X.-Y., Blockley, S.P.E., Tarasov, P.E., Xu, Y.-G., McLean, D., Tomlinson, E.L., Albert, P.G., Liu, J.-Q., Müller, S., Wagner, M., Menzies, M.A., Clarifying the distal to proximal tephrochronology of the Millennium (B-Tm) eruption, Changbaishan Volcano, northeast China, *Quaternary Geochronology* (2016), doi: 10.1016/j.quageo.2016.02.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Xuan-Yu Chen**<sup>a,b,c,d,\*</sup>, **Simon P.E. Blockley**<sup>b</sup>, **Pavel E. Tarasov**<sup>e</sup>, **Yi-Gang Xu**<sup>c</sup>, **Danielle McLean**<sup>b,f</sup>, **Emma L. Tomlinson**<sup>g</sup>, **Paul G. Albert**<sup>f,h</sup>, **Jia-Qi Liu**<sup>i</sup>, **Stefanie Müller**<sup>e,j</sup>, **Mayke Wagner**<sup>k</sup> and **Martin A. Menzies**<sup>a,c</sup>

<sup>a</sup> Department of Earth Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK

<sup>b</sup> Department of Geography, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK

<sup>c</sup> State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

<sup>d</sup> University of Chinese Academy of Sciences, Beijing 100049, China

<sup>e</sup> Institute of Geological Sciences, Palaeontology, Free University Berlin, Malteserstr. 74–100, Building D, 12249 Berlin, Germany

<sup>f</sup> Research Laboratory for Archaeology and the History of Art, University of Oxford, Oxford OX1 3QY, UK

<sup>g</sup> Department of Geology, Trinity College Dublin, Dublin 2, Ireland

<sup>h</sup> Department of Geography, Swansea University, Swansea SA2 8PP, UK

<sup>i</sup> Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China

<sup>j</sup> Center for Ainu and Indigenous Studies, Hokkaido University, Kita 8, Nishi 6, Kita-ku, Sapporo 060-0808, Japan

<sup>k</sup> Eurasia Department and Beijing Branch Office, German Archaeological Institute, Im Dol 2–6, 14195 Berlin, Germany

## **Abstract**

Tephra dispersed during the Millennium eruption (ME), Changbaishan Volcano, NE China provides one of the key stratigraphic links between Asia and Greenland for the synchronization of palaeoenvironmental records. However, controversy surrounds proximal-distal tephra correlations because (a) the proposed proximal correlatives of the distal ME tephra (i.e. B-Tm) lack an unequivocal chronostratigraphic context, and (b) the ME tephra deposits have not been chemically characterized for a full spectrum of element using grain-specific techniques. Here we present grain-specific glass chemistry, including for the first time, single grain trace element data, for a composite proximal sequence and a distal tephra from Lake Kushu, northern Japan (ca. 1100 km away from Changbaishan). We demonstrate a robust proximal-distal correlation and that the Kushu tephra is chemically associated with the ME/B-Tm. We propose that three of the proximal pyroclastic fall units were erupted as part of the ME. The

supports our geochemical correlation. Two further Bayesian age-depth models were independently constructed each incorporating one of two ice-core derived ages for the B-Tm tephra, providing Bayesian modelled ages of 933-949 and 944-947 cal AD (95.4%) for the Kushu tephra. The high resolution ice-core tephra ages imported into the deposition models help test and ultimately constrain the radiocarbon chronology in this interval of the Lake Kushu sedimentary record. The observed geochemical diversity between proximal and distal ME tephra deposits clearly evidences the interaction of two compositionally distinct magma batches during this caldera forming eruption.

**Key words** : Changbaishan; Millennium eruption; B-Tm tephra; Lake Kushu; Cryptotephra; Tephrochronology; Radiocarbon; Bayesian age modelling

## 1. Introduction

The Millennium eruption (ME) from the Changbaishan Volcano was a very large (VEI $\approx$ 7) eruption with the potential for ash dispersal across the Northern Hemisphere. Whilst the eruption was estimated to have had substantial volatile and sulfate aerosol emissions (Horn and Schmincke, 2000; Guo et al., 2002), it lacked a global impact on climate (Xu et al., 2013; Sun et al., 2014a). Distal products of the ME (i.e. B-Tm) have been reported in many localities including the Sea of Japan, the Japanese Archipelago (Machida and Arai, 1983; Furuta et al., 1986; Machida et al., 1990; Fukusawa et al., 1998; Nanayama et al., 2003; Kamite et al., 2010; Okuno et al., 2011; Hughes et al., 2013), northeast China (Sun et al., 2015) and the Greenland ice cores (Sun et al., 2014a) (Fig. 1a). As such it provides one of the potential key stratigraphic links between Asia and Greenland for the synchronization of palaeoenvironmental records. The proposed correlations between the distal ME tephra deposits and the Changbaishan Volcano (Machida et al., 1990; Sun et al., 2014a; 2015) require detailed chronostratigraphic studies and geochemical analyses of the proximal deposits and distal records. However, the proximal correlatives reported therein lack chronostratigraphic context and the ME tephra units have not been chemically characterized for a full spectrum of element using grain-specific techniques. Here we present the results of geochemical investigations of tephra units from proximal deposits with chronostratigraphic context, and a distal archive from Japan. This is then used to re-evaluate the proximal deposits, distal correlatives and the overall

## 2.1. Proximal volcanic deposits

The Changbaishan Volcano, situated on the border between China and North Korea (42°00' N, 128°03' E, Fig. 1a), is a polygenetic central stratovolcano with three main eruptive stages: early shield building, middle cone construction and a late explosive stage (Wei et al., 2007; 2013). During the latest explosive stage (<20ka: Wei et al., 2013), a major caldera-forming eruption occurring at ~AD 1000 (i.e. the “Millennium” eruption) produced a ca. 4.5 km wide caldera containing the crater lake called Tianchi (Machida et al., 1990).

The latest explosive stage recorded at Tianwen summit, on the Chinese flank of the crater contains four sequentially deposited pyroclastic fall units, C-4 [base], C-3, C-2 and C-1 [top], which are colored yellow, light grey, dark grey and black, respectively (Fig. 1b). Stratigraphic and chronological studies of the Tianwen summit profile are summarized in Figure 2. The lowermost C-4 yellow unit was dated by different methods and yielded ages around 4~5 ka (Liu et al., 1998; Wang et al., 2001; Yang et al., 2014). Hence, it was believed to be the product of an older pre-caldera forming eruption (Fig. 2). The overlying C-3 light grey unit is a very widespread pyroclastic fall which covers the crater rim and extends eastwards for more than 15 km from the crater (Horn and Schmincke, 2000; Sun et al., 2014b). At the Tianwen summit profile, this light grey unit has been dated at ~1 ka by U-series TIMS (Wang et al., 2001) and  $^{40}\text{Ar}/^{39}\text{Ar}$  (Yang et al., 2014) methods. The same light grey-colored tephra was visually identified in the environs of the volcano where extensive radiocarbon dating studies of charcoal samples confirmed that it has an age consistent with the onset of the last millennium (e.g. Horn and Schmincke, 2000; Nakamura et al., 2007; Yatsuzuka et al., 2010; Yin et al., 2012; Xu et al., 2013; Table 1). Therefore, the C-3 unit was regarded as the product of the ME. However, there is considerable controversy regarding the stratigraphic and chronological significance of the overlying C-2 and C-1 units (Fig. 2). None of the previous studies reported a dark grey pyroclastic fall (here labelled C-2) overlying the C-3 light grey unit. Fan et al. (1999) described a light purplish red welded tuff unit with a post-ME age overlying the light grey unit. This cannot be correlated to our C-2 unit on the basis of their color and compositional discrepancies. Wang et al. (2001) mentioned two stratigraphically upper units without color description, whose age and compositional information did not help to clarify the stratigraphic correlation. The two more recent stratigraphic and

reviewing the historical records of “abnormal natural phenomenon”, Cui et al. (1995) proposed three possible dates (i.e. AD 1668, 1702 and 1903) for the post-caldera eruptions of the Changbaishan Volcano. Ji et al. (1999) used thermo-luminescence (TL) to date the K-feldspar from the black pumice (C-1) and reported an age of  $0.34 \pm 0.03$  ka. Based on these two pieces of evidence, Liu et al. (1999) concluded that the black unit (C-1) can be correlated to the AD 1668-1702 eruptive record. However, the “abnormal natural phenomenon” recorded by ancient people including “thunder” or “white ash rain” cannot be confidently and accurately attributed to a particular volcanic eruption. Moreover, the TL age from the black pumice cannot completely rule out the possibility that this C-1 unit could be an older eruptive, reset by a younger eruption. As such, the eruption age of the C-1 unit remains ambiguous and it could be either contemporaneous with the ME or post date the ME.

Controversy also surrounds geochemical studies of the proximal ME stratigraphy on the crater rim. Fan et al. (1999) and Yu et al. (2012) suggested that, at Tianwen summit, only the light grey unit (i.e. C-3) is the ME product and that it can be classified as pantelleritic rhyolite based on bulk rock chemistry. However, Horn and Schmincke (2000) argued that the ME deposits should have included two phases: a major rhyolitic phase producing white to grey colored pumice, and a minor later phase forming trachytic agglutinates mantling the inner crater walls. Meanwhile, their electron microprobe glass data indicated that the widespread light grey pumice has a peralkalic composition akin to comendite rather than pantellerite. More recently, Sun et al. (2014a) reported major element glass chemistry for tephra sampled from the Korean side, indicating that the proximal ME deposits have bimodal compositions (i.e. trachyte and rhyolite). Although they described the outcrop containing white-yellow, grey and black pyroclastic fall, it is not easy to correlate this sequence to the Tianwen summit profile because no detailed stratigraphic context was provided.

## **2.2. Distal B-Tm records and the proximal correlatives**

The B-Tm tephra, which was named after its source volcano Baegdusan (i.e. Changbaishan Volcano) and its type locality in Tomakomai, Hokkaido, was first reported to be found in the Sea of Japan and Hokkaido and the northern part of Honshu (Machida and Arai, 1983). It was suggested to be the distal equivalent of the ME on the basis of its composition and stratigraphic position (i.e. relative age), as well as petrographic features (Machida et al., 1990). Later on, the B-Tm tephra was

above-mentioned studies have focused on the distal-proximal tephra correlation and the proposed proximal correlatives lack either independent dating to corroborate their “millennium” age (e.g. Machida et al., 1990) or unequivocal stratigraphic context (e.g. Sun et al., 2014a; 2015). As such hardly any clearly described outcrops have been reliably confirmed as the proximal equivalent, which leaves the B-Tm tephra almost distally described only. Moreover, the identification, characterization and correlation of the B-Tm is typically based on major element glass chemistry. Whilst there are some effective major element discriminants that can be used to separate contemporaneous marker tephtras (Sun et al., 2014a), one has to be careful because volcanoes can produce eruptions with similar compositions over a considerable time window (e.g. Smith et al., 2011a; 2011c; Lane et al., 2012; MacLeod et al., 2015). Hence we highly recommend that a full spectrum of major, minor and trace elements should be applied when correlating tephtras as this greatly increases the reliability of the resultant correlations. Since the abundances of trace elements are more sensitive to magmatic processes, they are widely used to secure tephtra correlations (e.g. Allan et al., 2008; Smith et al., 2011a; Tomlinson et al., 2012; Lane et al., 2012; Albert et al., 2012; 2013; Tomlinson et al., 2014; Albert et al., 2015; Tomlinson et al., 2015; Lane et al., 2015).

### **3. Sampling**

#### **3.1. Changbaishan tephtra**

Proximal fall lapilli samples from the C-1, C-2 and C-3 units were collected at the Tianwen summit (42°01'33" N, 128°04'00" E) under tight stratigraphic control. Sub-samples of the light grey pumice from Yang et al. (2014) are included in our C-3 sample set as they were collected from the same profile and the same unit, and were dated by  $^{40}\text{Ar}/^{39}\text{Ar}$  to corroborate the millennium age. Lapilli sized samples were crushed, cleaned, dried and clean fragments were examined and picked under the microscope then mounted in epoxy resin for geochemical micro-beam analysis. The samples from these three fall units contain alkali feldspar as the dominant phenocryst phase, with some minor contributions of olivine, pyroxene and quartz. The C-1 unit contains the highest percentage of phenocryst among all the three units. SEM images were taken to reveal the vesicularity and the location of phenocrysts prior to micro-beam analysis (Fig. 3). SEM images show that the C-3 pumice samples are

highly vesiculated glasses (Fig. 3c) and have a similar level of vesicularity with the C-3 samples. The C-1 pumice samples mainly contain poorly vesiculated glasses (Fig. 3d), with the amount of vesicles decreasing significantly relative to the C-2 and C-3 samples. The vesicularity of C-1 sample was estimated to be less than 50% (Yu et al., 2012), which is consistent with the observation that the C-1 samples are slightly welded whereas the C-2 and C-3 samples are not.

### **3.2. Lake Kushu tephra**

Lake Kushu (45°25'55" N, 141°02'13" E), ca. 1100 km away from the Changbaishan Volcano, is the only fresh-water lake of Rebun Island in the Sea of Japan, northwest off Hokkaido (Fig. 1a). Located in the northern part of the island, about 230-400 m from the coast, the lake has a kidney bean shape and a maximum length of ca. 1100 m. The maximum water depth reaches ca. 6 m in the eastern part of the lake with average depths of about 3 to 5 m. The bottom sediment of Lake Kushu has been considered as a valuable natural archive which stores detailed, high-resolution information about post-glacial environmental changes (Kumano et al., 1990). Therefore, Lake Kushu has been selected as one of the key study sites in the Baikal-Hokkaido Archaeology Project (BHAP), aiming to fill the existing gap in the current knowledge and to address the role of climate and environmental change in the life of northern hunter-gatherers (Weber et al., 2013). In February 2012, two parallel sediment cores (RK12-01 and RK12-02) were collected in the central part of the lake from the thick ice cover. The recovered composite core (RK12) revealed a continuous, partly laminated, organic-rich ca. 19.5 m long sediment column. The RK12 core age model built upon the 57 calibrated AMS radiocarbon dates suggests that the sedimentation continued over the past ca. 17,000 years (Müller et al., in press).

This lake was also selected for visible tephra and crypto-tephra analyses due to the potential for tephra to be located from Japanese, Korean and Chinese volcanoes, and coupled with the high precision radiocarbon chronology and palaeoenvironmental record. The tephra study covered the entire core archive, although here we focus only on the tephra of approximate millennium age, based on the radiocarbon chronology. A detailed tephrochronological study of the whole core is currently in preparation (Chen et al., in progress). Distal tephra were separated using procedures outlined by Blockley et al. (2005). The extracted glass shards were mounted in Canada Balsam on glass slides and examined using an Olympus CX-41 microscope fitted with



revealed the quantitative distribution of shards per gram of dry sediment for the complete composite core. A detailed analysis phase termed point-sample count, with 1 or 2 cm sampling interval, was undertaken later to locate the precise depth of the tephra where range-finders showed high concentration of glass shards. The point-sample counts relative to this research clearly show that the initial onset of the tephra input appears at the composite depth of 152-151 cm whilst the peak of it sits at 151-150 cm (Fig. 4). No visible ash layer was identified at these depths during core opening and sediment description though the tephra counts show quite large numbers. Individual shards (Fig. 5) were picked in the interval of 150-152 cm and embedded in resin, sectioned and polished ready for geochemical analysis.

## **4. Analytical methods**

### **4.1. Electron micro-probe analysis (EMPA)**

Major and minor element concentrations of glasses of proximal pumices and distal tephra shards were determined using Jeol JXA-8600 wavelength-dispersive electron microprobe (WDS-EMPA), equipped with 4 spectrometers and SamX software, at the Research Laboratory for Archaeology and the History of Art, University of Oxford. An accelerating voltage of 15 kV, low beam current (6 nA), and defocused (10  $\mu\text{m}$ ) beam were used to minimize Na migration. Count times were 30 s for Si, Al, Mg, Ti, Ca, K and Fe on each peak. Na was analysed for 12 s, Cl and Mn for 50 s and P for 60 s on each peak. The instrument was calibrated for each set of beam conditions using a suite of appropriate mineral standards. The calibration was verified using a range of secondary glass standards (MPI-DING suite) including the ATHO-G (rhyolite), StHs6/80-G (andesite) and GOR132-G (komatiite) glasses from the Max Planck Institute (Jochum et al., 2006). Samples with analytical totals <94% were discarded. All the analyses presented in the text, plots and tables are normalised to 100 wt.% for comparative purposes. Analytical precision is typically <0.8% relative standard deviation (RSD) for Si; <5% RSD for analytes with concentrations >0.8 wt.% with the exception of Na (7-10%). All standard data is presented in the supplementary material.

Trace element analyses of glasses of proximal and distal tephtras were performed using a Thermo Scientific iCAP Q ICP-MS coupled to an Analyte Excite excimer laser-ablation system at the Department of Geology, Trinity College Dublin. Spot sizes of 18, 24, 30 and 36  $\mu\text{m}$  were used, depending on the size of the area available for analysis in different samples. The repetition rate was 5Hz and the count time was 40 s (200 pulses) on the sample and 40 s on the gas blank (background). The ablated sample was transported in He gas flow ( $0.65\text{Lmin}^{-1}$ ) with additional  $\text{N}_2$  ( $5\text{mlmin}^{-1}$ ) via a signal smoothing device. Concentrations were calibrated using NIST612 with  $^{29}\text{Si}$  as the internal standard. Data reduction was performed manually using Microsoft Excel that allowed removal of portions of the signal compromised by the microcryst, void or resin component. Full details of the analytical and data reduction methods are described in Tomlinson et al. (2010). Accuracies of ATHO-G and StHs6/80-G MPI-DING glass analyses are typically  $<5\%$  for most elements,  $<10\%$  for Y, Zr, Nb, Gd and  $<15\%$  for Ta. Reproducibility of ATHO-G analyses is  $<5\%$  RSD for all trace elements with the exception of U ( $<7\%$ ).

## 5. Results

Glasses from juvenile clasts found in both proximal and distal deposits have compositions ranging from trachyte to rhyolite, with the rhyolitic population straddling the alkaline-subalkaline boundary (Fig. 6a). On a molecular basis, all the rhyolitic glasses and the majority of trachytic glasses reported herein have  $(\text{Na}_2\text{O}+\text{K}_2\text{O}) > \text{Al}_2\text{O}_3$ , whereas only few trachytic glasses show subtle  $(\text{Na}_2\text{O}+\text{K}_2\text{O}) \leq \text{Al}_2\text{O}_3$ . By definition, they are peralkaline trachyte and rhyolite. According to the classificatory scheme for peralkaline extrusive rocks (MacDonald, 1974), they can be further classified as comenditic trachyte and comendite, respectively (Fig. 6b). Representative major, minor and trace element glass compositions of proximal pumices and distal shards are given in Table 2. The full geochemical data sets are presented in the supplementary material.

The uppermost C-1 fall unit at Tianwen summit has a comenditic trachyte glass composition (Fig. 6b). The glasses are characterized by low CaO (1.1-1.5 wt.%) and MgO (0.1-0.3 wt.%), high K<sub>2</sub>O (4.8-6.2 wt.%) and significantly high FeO<sub>t</sub> (4.3-5.1 wt.%) with homogeneous SiO<sub>2</sub> (64.7-66.6 wt.%; Fig. 7a-d). They show an overall trend of decreasing total alkalis with increasing SiO<sub>2</sub> (Fig. 6a). Trace element data reveal that these glasses have limited compositional variations with 9-16 ppm Th, 14-40 ppm Ba, 3-5 ppm Ta, 49-73 ppm Nd and 30-45 ppm Y. They form compositional clusters in most of the trace element bi-plots (Fig. 7f-h). C-1 glasses have light rare earth element (LREE) enrichment relative to the heavy rare earth element (HREE) with La/Yb ratios ranging from 21 to 25. Mantle-normalised spider diagram reveals that C-1 glasses are up to more than 200 times more enriched than the primitive mantle (Fig. 8). They are characterized by pronounced negative anomalies in Ba, Sr and Eu in response to the fractionation of K-feldspar, which is a dominant phenocryst phase in the juvenile clasts.

### **5.1.2. C-2 fall deposits**

There is significant overlap in glass compositions of the C-2 fall unit and the overlying C-1 unit. The C-2 glasses are also classified as comenditic trachyte (Fig. 6b). They extend to slightly more evolved SiO<sub>2</sub> (65.4-67.7 wt.%), have less variable K<sub>2</sub>O (5.5-6.1 wt.%), comparable CaO (1.0-1.4 wt.%), FeO<sub>t</sub> (4.4-5.2 wt.%) and MgO (0.1-0.3 wt.%) relative to those of the C-1 glasses (Fig. 7a-d). The compositional similarity between the two units can also be seen in the trace element data. C-2 glasses show trace element concentrations of 12-19 ppm Th, 14-39 ppm Ba, 4-6 ppm Ta, 61-78 ppm Nd, and 36-51 ppm Y, which form almost identical compositional clusters overlapping the C-1 glasses (Fig. 7e-h). Mantle-normalised spider diagram reveals that the multi-element profiles of C-2 glasses have similar distribution pattern and incompatible element enrichment level with those of the C-1 glasses (Fig. 8). Nevertheless, some subtle differences in composition can be observed: (1) C-1 glasses have some least evolved components that are not seen in C-2 deposits (e.g. lower SiO<sub>2</sub> content and lower incompatible element enrichment level; Figs. 6a, 8); (2) C-2 glasses extend to more evolved compositions than C-1 glasses (e.g. higher SiO<sub>2</sub> content and higher incompatible element enrichment level; Figs. 6a, 8).

between alkaline and subalkaline rhyolite (Fig. 6a) and can be specifically classified as comendite (Fig. 6b). C-3 glasses are highly homogeneous with restricted major element compositional ranges. They have concentrations of highly evolved SiO<sub>2</sub> (74.4-75.3 wt.%) with lower CaO (0.1-0.3 wt.%), MgO (<0.1 wt.%), K<sub>2</sub>O (4.3-4.6 wt.%) and FeO<sub>t</sub> (3.8-4.3 wt.%) than those of the overlying trachytic units (i.e. C-1 and C-2; Fig. 7a-d). Trace element concentration data for C-3 glasses form linear clusters in element bi-plots (Fig. 7e-h). They have higher contents of Th (44-57 ppm), Ta (14-18 ppm), Nd (111-149 ppm), Y (117-156 ppm) and lower Ba (7-10 ppm) relative to the upper trachyte units (Fig. 7e-h). C-3 glasses show LREE enrichment relative to the HREE with a lower LREE/HREE fractionation level (La/Yb=11-13) than those of the trachytic glasses. Mantle-normalised spider diagram shows that the C-3 glasses are up to more than 600 times more enriched than the primitive mantle, with more pronounced depletions in Ba, Sr and Eu relative to the trachytic units (Fig. 8). It is worth noting that, in their multi-element profiles, the trachytic glasses and the C-3 rhyolitic glasses show different trends in anomalies of Nb-Ta (Fig. 8).

## 5.2. Glass chemistry of distal tephra

The distal tephra found in Lake Kushu shows significant heterogeneity with compositions ranging from trachyte to rhyolite (Fig. 6a). This can be further classified as comenditic trachyte and comendite respectively (Fig. 6b). The less evolved trachytic components are characterized by relatively homogeneous SiO<sub>2</sub> (typically between 65.3-67.0 wt.%), low CaO (typically between 1.1-1.4 wt.%) and MgO (typically between 0.1-0.3 wt.%), high K<sub>2</sub>O (5.3-6.0 wt.%) and FeO<sub>t</sub> (4.4-5.5 wt.%), while the more evolved rhyolitic shards have heterogeneous SiO<sub>2</sub> (70.3-76.1 wt.%), lower CaO (typically <0.8 wt.%), MgO (<0.1 wt.%), K<sub>2</sub>O (typically <4.7 wt.%) and FeO<sub>t</sub> (typically < 4.2wt.%; Fig. 7a-d). The two populations collectively show an overall trend of decreasing TiO<sub>2</sub>, CaO, MgO, FeO<sub>t</sub> and K<sub>2</sub>O (Fig. 7a-d) with increasing SiO<sub>2</sub>, while the Na<sub>2</sub>O contents between the trachyte and rhyolite remain fairly constant (4.3-6.0 wt.% and 3.3-6.0 wt.% respectively).

Mantle-normalized spider diagrams shows that the distal trachytic and rhyolitic glasses are comparable in incompatible element distribution patterns and enrichment levels to the proximal trachytic and rhyolitic glasses respectively (Fig. 8). All the distal glasses show LREE enrichment relative to HREE and pronounced negative anomalies in Ba, Sr and Eu (Fig. 8). Nevertheless, the less evolved trachytic glasses

trends in most of the trace element bi-plots (Fig. 7e-h). The trachytic glasses have trace element variations with 11-18 ppm Th, 10-76 ppm Ba, 4-6 ppm Ta, 54-88 ppm Nd and 37-50 ppm Y, while the rhyolitic glasses show significantly higher contents of Th (30-56 ppm), Ta (10-17 ppm), Nd (78-144 ppm), Y (66-144 ppm) and lower Ba (6-12 ppm; Fig. 7e-h). The rhyolitic glasses contain an analysis that is characterized by noticeably lower concentrations in most of the measured trace elements (Th=30 ppm, see Table 2: RH554-37) and higher La/Yb ratio (La/Yb=22) than any other rhyolite. Nevertheless, this analysis lies on the evolutionary trend of the rhyolitic members (Fig. 7g).

## 6. Interpretation

### 6.1. Proximal-distal glass correlation

Proximal-distal relationships are assessed in this section based on the glass geochemistry presented above. Overall, glass shards found in Lake Kushu are highly heterogeneous and broadly overlap all three proximal fall units at a major, minor and trace element level (Figs. 7, 8).

The two proximal trachytic units, C-1 and C-2, share broadly overlapping major, minor and trace element glass chemistries, which makes it problematic to distinguish between these two units based on glass chemistry alone. The trachytic glasses of the distal Kushu tephra overlap with the two proximal trachytic units (Figs. 7, 8). Subtle differences can be recognized between the proximal and distal trachytic glasses: a) Kushu trachytic glasses have a least evolved composition that is not seen in proximal trachytic glasses (Fig. 7a); b) Kushu trachytic glasses have greater compositional variations of highly depleted trace element (e.g. Ba and Sr) than proximal trachytic glasses (Fig. 7e), which might be affected by analysis of microlite of distal shards in LA-ICP-MS.

The proximal rhyolitic unit C-3 geochemically overlaps the most evolved compositions of the distal rhyolitic glasses. These most evolved distal compositions ( $\text{SiO}_2 \geq 74.0$  wt.%, n=41) make up the majority of the Kushu rhyolitic glasses (n=46) and could not be separated from the C-3 glasses in the major or trace element bi-plots (Fig. 7). The mantle-normalised spider diagram corroborates the geochemical overlaps at a full spectrum of element level (Fig. 8). Nevertheless, Kushu rhyolitic glasses have some less fractionated compositions that are not seen in the C-3 glasses

units collectively comprise compositionally identical trachyte and rhyolite but without the intervening compositional continuum. Given the similarity between the trachyte and rhyolite end-members we propose a correlation between the distal Kushu tephra and all three of the investigated proximal units.

## 6.2. Comparison with the published data

In order to determine the provenance of the tephra reported herein and to investigate the geochemical diversity between the well-correlated proximal and distal tephra, we compare our glass chemistries with those of the published proximal ME tephra and B-Tm, along with other widespread late Quaternary marker tephra in and around the Northeast Asia area. Since all the available data are major element glass chemistry, comparison can only be made at a major element level.

Major element glass chemistry provides some basic discriminants for the identification of the B-Tm tephra. The B-Tm tephra has distinctive compositional heterogeneity ranging from trachyte to alkaline/subalkaline rhyolite with a fairly high total alkali content (8-12 wt.%; Fig. 9a). However, the late Quaternary Kyushu tephra layers (e.g. K-Ah, AT, Aso-4 and Ata) and Holocene Kamchatka tephra deposits typically plot into the subalkaline area, and the early Holocene Ulleungdo tephra (i.e. U-Oki) straddles the boundary between phonolite and trachyte (Fig. 9a). The B-Tm tephra has lower CaO than that of the Kyushu and Kamchatka tephra in a given SiO<sub>2</sub>, and higher SiO<sub>2</sub> than that of the Ulleungdo tephra (Fig. 9b). The FeO<sub>t</sub> vs. CaO bi-plot is an excellent discriminant (Fig. 9c) since the B-Tm tephra has a fairly high FeO<sub>t</sub> content (typically between 3.8-5.6 wt.%) coupled with a relatively low CaO content (typically <1.5 wt.%), given its highly evolved composition (SiO<sub>2</sub> >63 wt.%). All the proximal and distal tephra reported herein show major element compositions that correspond precisely to the published B-Tm chemistries, and can be separated from tephra from other volcanic settings in the same region (Fig. 9). On this basis we conclude that all our proximal and distal tephra deposits are the product of the ME.

Sun et al. (2014a) reported the major element glass chemistry for a proximal ME sequence in North Korea which included the white-yellow, grey and black fall pumice. The major element chemistry for glasses from this ME sequence reveal an overlap with the proximal units described herein (Fig. 10a-b), suggesting both sequences relate to the ME. Most importantly, the proximal ME tephra units at Chinese and Korean sequences on the crater rim are both characterized by significant geochemical

Lake (NE China; Sun et al., 2015), Sea of Japan (Machida et al., 1990), Utasai Bog (Hokkaido; Hughes et al., 2013) and Lake Kushu (Hokkaido, this study) reveal continuous compositions with data plotting in the compositional gap between the end-member compositions defined by proximal samples (Fig. 10c-d). Interestingly, although the ultra-distal B-Tm found in the Greenland ice cores contains very few shards (Coulter et al., 2012; Sun et al., 2014a), their compositions plot within the proximal compositional gap (Fig. 10c-d). Overall the proximal fall units deposited on the crater rim only preserve bimodal compositions for the caldera forming eruption, whereas the mid-range to distal tephra preserve a complete compositional range including the bimodal end-member compositions and intermediate compositions.

## **7. Discussion**

### **7.1. Proximal-distal expression of the ME**

As in the case of B-Tm tephra deposits reported from other localities, the single distal tephra layer found in Lake Kushu has compositional variation ranging from trachyte to rhyolite. This indicates the magmatic system that produced this distal ME tephra must have had the same compositional heterogeneity. In contrast the proximal C-3 unit is characterized by homogeneous rhyolite without any trachytic component. As such the C-3 unit alone cannot be the proximal equivalent of the B-Tm, though it is the most widespread unit in the Changbaishan volcanic region (Horn and Schmincke, 2000). Our glass chemistry shows that the geochemical bimodality observed between the proximal units C-3 to C-1 is comparable to the end-member compositions of the distal Kushu tephra. Importantly there is major, minor and trace element geochemical overlap between the proximal and distal end-member compositions which significantly increases the reliability of the tephra correlation. Hence, we propose that the proximal record of the ME at Tianwen summit is a composite sequence that includes the light grey, dark grey and black (i.e. C-3 to C-1) sub-units. This is also consistent with the field observation that there is no time break/erosional unconformity or paleosol between the three investigated fall units.

Proximal fall deposits preserved on the crater rim are dominated by bimodal compositions whereas distal tephra record a full range of geochemical heterogeneity. The presence of intermediate glass compositions in the distal tephra, not currently observed in the proximal records, clearly evidences the interaction of two

In fact, there is no such evidence in our pumice samples. Overall, the observed compositional diversity indicates that the trachytic and rhyolitic magmas were not thoroughly mixed so only a minor portion of intermediate compositions were generated during magmatic interaction, which were then transferred into the stratosphere during the syn-eruptive process and were consequently recorded in the distal realms.

The preservation of bimodal proximal deposits (comprising the least and most fractionated components) has been reported from calderas elsewhere including in Afro-Arabia (Ukstins Peate et al., 2003; Ukstins Peate et al., 2008). As reported herein, the proximal units in Arabia are markedly bimodal and contrast with the contemporaneous distal tephra layers from 3000 km away in the Indian Ocean. These distal tephras display a continuum of compositions with end-members identical to the bimodal end-members found proximally. The bias between proximal and distal magma sequences is controversial and may relate to high level sub-volcanic processes that include fractionation processes involving silica rich magmas (Grove and Donnelly-Nolan, 1986), large scale liquid immiscibility (Charlier et al., 2011) and/or melt-crystal dynamics (Dufek and Bachmann, 2010).

## **7.2. Bayesian age modelling for Kushu tephra**

The ME tephra deposits have been dated either directly or indirectly by multiple methods in many localities over the past two decades (Table 1). The proximal ages are mainly derived from the indirect dating method of  $^{14}\text{C}$  dating of charcoals preserved in the pyroclastic fall and flow deposits (e.g. Horn and Schmincke, 2000; Nakamura et al., 2007; Yatsuzuka et al., 2010; Yin et al., 2012; Xu et al., 2013). Besides, direct dating of primary minerals using U-series TIMS (Wang et al., 2001) and  $^{40}\text{Ar}/^{39}\text{Ar}$  (Yang et al., 2014) methods has also provided additional age constraints on the eruption. Distally, B-Tm tephra deposits were found in varve lakes in Japan and northeast China where the eruption was dated by varve chronology (Fukusawa et al., 1998; Kamite et al., 2010; Sun et al., 2015). Ultra-distal B-Tm tephra deposits were reported from the Greenland ice cores and as such the eruption has ages derived from the GICC05 and GISP2 ice-core chronologies (Sun et al., 2014a). Most recently, Sigl et al. (2015) has proposed a slight revision to the GICC05 chronology for the past 2,500 years. This new NS1-2011 chronology, based on  $^{10}\text{Be}$  and  $^{14}\text{C}$  synchronization of ice-core and tree-ring timescales, has shifted the age of the ME from AD 941 $\pm$ 1



Holocene radiocarbon ages in the Kushu core (Fig. 11), for comparison with the highest resolution distal age reported for this tephra. In this age model, we have modelled the AMS  $^{14}\text{C}$  dates reported in Müller et al. (in press), for Kushu along with the depth information (Table 3) in a Bayesian *P\_Sequence* model with the addition of sediment deposition modelling to calculate the highest likelihood age ranges for every 1.6 cm, incorporating formal outlier detection (Bronk Ramsey, 2009a; 2009b; 2013). As there are no radiocarbon dates in the same centimeter of the core as the proposed B-Tm tephra, it was necessary to determine the age range for its depth at 151.5 cm, which is the average position in the core for the initial onset of Kushu tephra. The age range for the depth of 151.5 cm is 913-1395 cal AD (Fig. 11; supplementary material), and while at fairly low resolution this comfortably overlaps with the ice-core ages of the ME/B-Tm.

Due to the low resolution generated by the requirement to model the ages and associated uncertainty of the depths between the radiocarbon ages in this approach we also attempted to test the chronological relationship between the Kushu tephra and the high resolution age for the B-Tm in more detail. The logic of the test is that, while the radiocarbon ages for the Kushu site (Fig. 11) show a broadly last millennium age for the tephra sitting at 151 cm, if our correlation is correct then the tephra and radiocarbon ages should all be compatible with a high precision age model. We have thus constructed two more constrained age models (Fig. 12). We again use the above radiocarbon ages and their depths (Table 3), without interpolation and have also imported the GICC05 (Sun et al., 2014a; Fig. 12a) and NS1-2011 (Sigl et al., 2015; Fig. 12b) ages for the ME/B-Tm at the depth of our proposed correlative tephra in Kushu. We have applied a *P\_Sequence* depositional model in Oxcal (Bronk Ramsey, 2013), with a variable K factor and automatic outlier detection, following Bronk Ramsey (2008; 2009a; 2009b) with boundaries at the top and bottom of the selected 2 meters sequence. The outlier model selection used the general outlier model and outlier probabilities were set to 0.05%. The results of the modelling exercise are reported as 95% confidence highest probability density function and depositional model plots. In these higher resolution, more constrained models, we are testing if the B-Tm ages are fully consistent with the Kushu radiocarbon ages and the depths in the core in which they sit. If our data is incompatible with the B-Tm on chronological grounds we would expect that the B-Tm ages we have incorporated, or the radiocarbon ages above and below the tephra would be reported as being anomalous by the outlier detection software. In these cases, however only the uppermost date Poz-51689 reported significant potential to be an outlier and thus the weighting of this

the depth and all chronological information are 933-949 and 944-947 cal AD (95.4%), respectively (Fig. 12; Table 3). We, thus, suggest on this rigorous basis, using formal deposition models the Kushu tephra is both chemically and chronologically well correlated to the ME/B-Tm.

### 7.3. Implication for future studies

High-resolution (annual to decadal-scale) reconstructions of past environments, and better understanding which role environmental and climatic changes played in the cultural dynamics remains extremely important, though an empirically challenging question for the Hokkaido region (Weber et al., 2013). Despite, the northern part of Japan, including Hokkaido and adjacent islands are rich in archaeological and environmental archives, the main challenge remains the scarcity of published records with high temporal resolution and adequate dating control (Nakagawa et al., 2012; Müller et al., in press and references therein). These two common problems hinder direct correlation between individual archives, thus preventing inter-regional comparison and identification of leads and lags in reconstructed climate variability. A multidisciplinary research on the RK12 core from Lake Kushu started in 2012 has a significant potential to fill the existing gap in the current knowledge and to build up the high-resolution environmental archive spanning the past ca. 17,000 years (Müller et al., in press). However, in order to serve as a link between the high-resolution and accurately-dated records from central Japan (e.g. Lake Suigetsu: Nakagawa et al., 2012), China (e.g. Sihailongwan Maar Lake: Stebich et al., 2009; 2015), and North Atlantic region, the Kushu  $^{14}\text{C}$  chronology must be checked, and when necessary, improved. The current study demonstrates the potential of cryptotephra analysis for improving the age constraints on the Kushu sedimentary record. The ice-core ages imported into the deposition models significantly help constrain the age of the sediments around the tephra layer and improve the radiocarbon chronology in this interval (Fig. 12). This provides encouragement for the identification of additional cryptotephra layers in the Lake Kushu sediment. Though it still could be improved, we believe that the existing RK12 core chronology based on the 57 AMS dates (Müller et al., in press) is robust enough to provide reliable age estimations for the tephra layers preserved in the sediment. This will facilitate search of the source volcano and identification of the eruption, which produced each given tephra.

the crater rim of Changbaishan Volcano, NE China display bimodal glass compositions comprising trachyte and rhyolite end-member compositions.

- b) A distal tephra layer in Lake Kushu of northern Japan contains glass shards that define a complete compositional continuum ranging from trachyte to rhyolite that in themselves are chemically identical to the investigated proximal glasses.
- c) The chemical constraint and radiocarbon-based Bayesian age-depth model indicate that the analysed Kushu tephra is the distal expression of the Millennium eruption.
- d) Proximal-distal geochemical correlation at a full spectrum of element level (i.e. major, minor and trace elements) requires that the three proximal pyroclastic fall units were erupted as part of the Millennium eruption as alluded to by previous studies (Horn and Schmincke, 2000; Sun et al., 2014a).
- e) Bayesian age-depth modelling of Kushu sediments involving the AMS  $^{14}\text{C}$  dates and two ice-core derived ages for the B-Tm provides 95% confidence interval ranges of 933-949 and 944-947 cal AD for the Kushu tephra. The high resolution ice-core tephra ages imported into the deposition models help test and ultimately constrain the radiocarbon chronology in this interval of the Lake Kushu sedimentary record.
- f) The presence of intermediate glass compositions in the distal tephra layers, not currently observed in the proximal records, clearly evidences the interaction of two compositionally distinct magma batches (with end-member compositions) during the caldera forming eruption.

This study was supported by the External Cooperation Program of BIC, Chinese Academy of Sciences, Grant No.132744KYSB20130005. XYC's stay at RHUL was supported by the collaboration scheme between GIGCAS and RHUL. Coring of Lake Kushu and core transportation costs were covered by the Japanese MEXT-Japan Kakenhi research grant No. 21101002 held by Dr. H. Yonenobu, whose generous help we greatly acknowledge. The Baikal–Hokkaido Archaeology Project and BHAP-related research activities on Rebun Island are supported by the Major Collaborative Research Initiative (MCRI) program of the Social Sciences and Humanities Research Council of Canada, and collaborating institutions, including University of Alberta, German Archaeological Institute (DAI), Free University Berlin and German Science Foundation (DFG TA 540/5), Hokkaido University, and Japan Society for the Promotion of Science. We would like to thank Prof. F. Wang for providing proximal tephra sub-samples; Dr. C. J. Manning for the assistance in sample preparation; Dr. S. Armitage for beneficial discussions; Dr. V. C. Smith and an anonymous reviewer for their detailed and constructive feedback. We thank N. Holloway for preparing samples in epoxy resin stubs; K. Flowers and S. Gibbons for their assistance in the lab.

## References

- Albert, P.G., Hardiman, M., Keller, J., Tomlinson, E.L., Smith, V.C., Bourne, A.J., Wulf, S., Zanchetta, G., Sulpizio, R., Müller, U.C., Pross, J., Ottolini, L., Matthews, I.P., Blockley, S.P.E. and Menzies, M.A., 2015. Revisiting the Y-3 tephrostratigraphic marker: a new diagnostic glass geochemistry, age estimate, and details on its climatostratigraphical context. *Quaternary Science Reviews*, 118: 105-121.
- Albert, P.G., Tomlinson, E.L., Lane, C.S., Wulf, S., Smith, V.C., Coltelli, M., Keller, J., Lo Castro, D., Manning, C.J., Müller, W. and Menzies, M.A., 2013. Late glacial explosive activity on Mount Etna: Implications for proximal–distal tephra correlations and the synchronisation of Mediterranean archives. *Journal of Volcanology and Geothermal Research*, 265(0): 9-26.
- Albert, P.G., Tomlinson, E.L., Smith, V.C., Di Roberto, A., Todman, A., Rosi, M., Marani, M., Muller, W. and Menzies, M.A., 2012. Marine-continental tephra correlations: Volcanic glass geochemistry from the Marsili Basin and the Aeolian Islands, Southern Tyrrhenian Sea, Italy. *Journal of Volcanology and Geothermal Research*, 229–230: 74-94.
- Allan, A.S.R., Baker, J.A., Carter, L. and Wysoczanski, R.J., 2008. Reconstructing the Quaternary evolution of the world's most active silicic volcanic system: insights from an ~1.65 Ma deep ocean tephra record sourced from Taupo Volcanic Zone, New Zealand. *Quaternary Science Reviews*, 27(25–26): 2341-2360.

- Bronk Ramsey, C., 2008. Deposition models for chronological records. *Quaternary Science Reviews*, 27(1–2): 42-60.
- Bronk Ramsey, C., 2009a. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51(1): 337-360.
- Bronk Ramsey, C., 2009b. Dealing with outliers and offsets in radiocarbon dating. *Radiocarbon*, 51(3): 1023-1045.
- Bronk Ramsey, C., 2013. OxCal 4.2. Available from: <http://c14.arch.ox.ac.uk/oxcal>.
- Charlier, B., Namur, O., Toplis, M.J., Schiano, P., Cluzel, N., Higgins, M.D. and Auwera, J.V., 2011. Large-scale silicate liquid immiscibility during differentiation of tholeiitic basalt to granite and the origin of the Daly gap. *Geology*, 39(10): 907-910.
- Chen, X.Y., Xu, Y.G. and Menzies, M., 2014. Tephrochronology: Principles and applications. *Acta Petrologica Sinica*, 30(12): 3491-3500 (in Chinese with English abstract).
- Coulter, S.E., Pilcher, J.R., Plunkett, G., Baillie, M., Hall, V.A., Steffensen, J.P., Vinther, B.M., Clausen, H.B. and Johnsen, S.J., 2012. Holocene tephra highlight complexity of volcanic signals in Greenland ice cores. *Journal of Geophysical Research: Atmospheres*, 117(D21303).
- Cui, Z., Wei, H. and Liu, R., 1995. Historical documentation research of eruptions of the Tianchi Volcano, Changbaishan. In: Liu, R. (ed). *Volcanism and Human Environment*. Beijing: Seismology Press: 36-39 (in Chinese).
- Dufek, J. and Bachmann, O., 2010. Quantum magmatism: Magmatic compositional gaps generated by melt-crystal dynamics. *Geology*, 38(8): 687-690.
- Fan, Q., Liu, R., Wei, H., Sui, J. and Li, N., 1999. Petrogeochemical Characteristics of Holocene Eruption of the Tianchi Volcano, Changbai Mountains. *Geological Review*, 45(Sup.): 263-271 (in Chinese with English abstract).
- Fukusawa, H., Tsukamoto, S., Tsukamoto, H., Ikeda, M., Okamura, M. and MATsuoka, H., 1998. Falling age of Baegdusan-Tomakomai tephra (B-Tm) estimated by using non-glacial varves. *Laguna*, 5: 55-62 (in Japanese with English abstract).
- Furuta, T., Fujioka, K. and Arai, F., 1986. Widespread submarine tephra around Japan — Petrographic and chemical properties. *Marine Geology*, 72(1): 125-142.
- Grove, T.L. and Donnelly-Nolan, J.M., 1986. The evolution of young silicic lavas at Medicine Lake Volcano, California: Implications for the origin of compositional gaps in calc-alkaline series lavas. *Contrib Mineral Petrol*, 92(3): 281-302.
- Guo, Z., Liu, J., Sui, S., Liu, Q., He, H. and Ni, Y., 2002. The mass estimation of volatile emission during 1199–1200 AD eruption of Baitoushan volcano and its significance. *Science in China Series D: Earth Sciences*, 45(6): 530-539.
- Horn, S. and Schmincke, H.-U., 2000. Volatile emission during the eruption of Baitoushan Volcano (China/North Korea) ca. 969 AD. *Bull Volcanol*, 61(8): 537-555.
- Hughes, P.D.M., Mallon, G., Brown, A., Essex, H.J., Stanford, J.D. and Hotes, S., 2013. The impact of high tephra loading on late-Holocene carbon accumulation and vegetation succession in peatland communities. *Quaternary Science Reviews*, 67(0): 160-175.
- Irvine, T.N. and Baragar, W.R.A., 1971. A Guide to the Chemical Classification of the Common Volcanic Rocks. *Canadian Journal of Earth Sciences*, 8(5): 523-548.

- Hellebrand, E., Mocek, B., Raczek, I., Stracke, A., Alard, O., Bouman, C., Becker, S., Dücking, M., Brätz, H., Klemm, R., de Bruin, D., Canil, D., Cornell, D., de Hoog, C.-J., Dalpé, C., Danyushevsky, L., Eisenhauer, A., Gao, Y., Snow, J.E., Groschopf, N., Günther, D., Latkoczy, C., Guillong, M., Hauri, E.H., Höfer, H.E., Lahaye, Y., Horz, K., Jacob, D.E., Kasemann, S.A., Kent, A.J.R., Ludwig, T., Zack, T., Mason, P.R.D., Meixner, A., Rosner, M., Misawa, K., Nash, B.P., Pfänder, J., Premo, W.R., Sun, W.D., Tiepolo, M., Vannucci, R., Vennemann, T., Wayne, D. and Woodhead, J.D., 2006. MPI-DING reference glasses for in situ microanalysis: New reference values for element concentrations and isotope ratios. *Geochemistry, Geophysics, Geosystems*, 7(2): Q02008.
- Kamite, M., Yamada, K., Saito-Kato, M., Okuno, M. and Yasuda, Y., 2010. Microscopic observations of varve sediments from Lake Ni-no-Megata and Lake San-no-Megata, Oga Peninsula, NE Japan, with reference to the fallout age of the B-Tm Tephra. *Jour. Geol. Soc. Japan*, 116(7): 349-359 (in Japanese with English abstract).
- Kumano, S., Ihira, M., Kuromi, M., Maeda, Y., Matsumoto, E., Nakamura, T., Matsushima, Y., Sato, H. and Matsuda, I., 1990. Holocene sedimentary history of some coastal plains in Hokkaido, Japan V. Sedimentary history of Kushu Lake and Akkeshi. *Ecol. Res.*, 5(3): 277-289.
- Kyle, P.R., Ponomareva, V.V. and Rourke Schlupe, R., 2011. Geochemical characterization of marker tephra layers from major Holocene eruptions, Kamchatka Peninsula, Russia. *International Geology Review*, 53(9): 1059-1097.
- Lane, C.S., Blockley, S.P.E., Mangerud, J., Smith, V.C., Lohne, Ø.S., Tomlinson, E.L., Matthews, I.P. and Lotter, A.F., 2012. Was the 12.1 ka Icelandic Vedde Ash one of a kind? *Quaternary Science Reviews*, 33(0): 87-99.
- Lane, C.S., Brauer, A., Martín-Puertas, C., Blockley, S.P.E., Smith, V.C. and Tomlinson, E.L., 2015. The Late Quaternary tephrostratigraphy of annually laminated sediments from Meerfelder Maar, Germany. *Quaternary Science Reviews*, 122: 192-206.
- Le Bas, M.J., Le Maitre, R.W., Streckeisen, A. and Zanettin, B., 1986. A Chemical Classification of Volcanic Rocks Based on the Total Alkali-Silica Diagram. *Journal of Petrology*, 27(3): 745-750.
- Liu, R., Fan, Q., Wei, H. and Li, N., 1999. Study on Active Volcanoes of China. *Geological Review*, 45(Sup.): 3-15 (in Chinese with English abstract).
- Liu, R., Wei, H. and Li, J., 1998. The latest eruptions from Tianchi volcano, Changbaishan. SciencePress, Beijing (in Chinese).
- MacDonald, R., 1974. Nomenclature and petrochemistry of the peralkaline oversaturated extrusive rocks. *Bulletin Volcanologique*, 38(2): 498-516.
- Machida, H. and Arai, F., 1983. Extensive ash falls in and around the sea of Japan from large late quaternary eruptions. *Journal of Volcanology and Geothermal Research*, 18(1-4): 151-164.
- Machida, H., Moriwaki, H. and Zhao, D.-C., 1990. The recent major eruption of Changbai Volcano and its environmental effects. *Geographical Reports of Tokyo Metropolitan University*(25): 1-20.
- MacLeod, A., Matthews, I.P., Lowe, J.J., Palmer, A.P. and Albert, P.G., 2015. A second tephra isochron for the Younger Dryas period in northern Europe: The Abernethy Tephra. *Quat Geochronol*, 28: 1-11.
- Müller, S., Schmidt, M., Kossler, A., Leipe, C., Irino, T., Yamamoto, M., Yonenobu, H., Goslar, T., Kato, H.,

- Takemura, K., Staff, R.A., Payne, R., Bronk Ramsey, C., Bryant, C., Brock, F., Schlolaut, G., Marshall, M., Tarasov, P. and Lamb, H., 2012. SG06, a fully continuous and varved sediment core from Lake Suigetsu, Japan: stratigraphy and potential for improving the radiocarbon calibration model and understanding of late Quaternary climate changes. *Quaternary Science Reviews*, 36: 164-176.
- Nakamura, T., Okuno, M., Kimura, K., Mitsutani, T., Moriwaki, H., Ishizuka, Y., Kim, K.H., Jing, B.L., Oda, H., Minami, M. and Takada, H., 2007. Application of <sup>14</sup>C Wiggle-Matching to Support Dendrochronological Analysis in Japan. *Tree-Ring Research*, 63(1): 37-46.
- Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, K. and Yamaki, S., 2003. Unusually large earthquakes inferred from tsunami deposits along the Kuril trench. *Nature*, 424(6949): 660-663.
- Okuno, M., Torii, M., Yamada, K., Shinozuka, Y., Danhara, T., Gotanda, K., Yonenobu, H. and Yasuda, Y., 2011. Widespread tephras in sediments from lake Ichi-no-Megata in northern Japan: Their description, correlation and significance. *Quaternary International*, 246(1–2): 270-277.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M. and van der Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon*, 55(4): 1869-1887.
- Sigl, M., Winstrup, M., McConnell, J.R., Welten, K.C., Plunkett, G., Ludlow, F., Buntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O.J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D.R., Pilcher, J.R., Salzer, M., Schupbach, S., Steffensen, J.P., Vinther, B.M. and Woodruff, T.E., 2015. Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature*, 523(7562): 543-549.
- Smith, V.C., Isaia, R. and Pearce, N.J.G., 2011a. Tephrostratigraphy and glass compositions of post-15 kyr Campi Flegrei eruptions: implications for eruption history and chronostratigraphic markers. *Quaternary Science Reviews*, 30(25–26): 3638-3660.
- Smith, V.C., Mark, D.F., Staff, R.A., Blockley, S.P.E., Ramsey, C.B., Bryant, C.L., Nakagawa, T., Han, K.K., Weh, A., Takemura, K. and Danhara, T., 2011b. Toward establishing precise <sup>40</sup>Ar/<sup>39</sup>Ar chronologies for Late Pleistocene palaeoclimate archives: an example from the Lake Suigetsu (Japan) sedimentary record. *Quaternary Science Reviews*, 30(21–22): 2845-2850.
- Smith, V.C., Pearce, N.J.G., Matthews, N.E., Westgate, J.A., Petraglia, M.D., Haslam, M., Lane, C.S., Korisettar, R. and Pal, J.N., 2011c. Geochemical fingerprinting of the widespread Toba tephra using biotite compositions. *Quaternary International*, 246(1–2): 97-104.
- Smith, V.C., Staff, R.A., Blockley, S.P.E., Bronk Ramsey, C., Nakagawa, T., Mark, D.F., Takemura, K. and Danhara, T., 2013. Identification and correlation of visible tephras in the Lake Suigetsu SG06 sedimentary archive, Japan: chronostratigraphic markers for synchronising of east Asian/west Pacific palaeoclimatic records across the last 150 ka. *Quaternary Science Reviews*, 67(0): 121-137.

- climate dynamics based on the pollen record from Sihailongwan Maar Lake, NE-China. *Quaternary Science Reviews*, in press.
- Sun, C., Plunkett, G., Liu, J., Zhao, H., Sigl, M., McConnell, J.R., Pilcher, J.R., Vinther, B., Steffensen, J.P. and Hall, V., 2014a. Ash from Changbaishan Millennium eruption recorded in Greenland ice: Implications for determining the eruption's timing and impact. *Geophysical Research Letters*: 2013GL058642.
- Sun, C., You, H., He, H., Zhang, L., Gao, J., Guo, W., Chen, S., Mao, Q., Liu, Q., Chu, G. and Liu, J., 2015. New evidence for the presence of Changbaishan Millennium eruption ash in the Longgang volcanic field, Northeast China. *Gondwana Research*, In Press.
- Sun, C.Q., You, H.T., Liu, J.Q., Li, X., Gao, J.L. and Chen, S.S., 2014b. Distribution, geochemistry and age of the Millennium eruptives of Changbaishan volcano, Northeast China - A review. *Front. Earth Sci.*, 8(2): 216-230.
- Sun, S.-s. and McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. Geological Society, London, Special Publications, 42(1): 313-345.
- Tomlinson, E.L., Albert, P.G., Wulf, S., Brown, R.J., Smith, V.C., Keller, J., Orsi, G., Bourne, A.J. and Menzies, M.A., 2014. Age and geochemistry of tephra layers from Ischia, Italy: constraints from proximal-distal correlations with Lago Grande di Monticchio. *Journal of Volcanology and Geothermal Research*, 287: 22-39.
- Tomlinson, E.L., Arienzo, I., Civetta, L., Wulf, S., Smith, V.C., Hardiman, M., Lane, C.S., Carandente, A., Orsi, G., Rosi, M., Müller, W. and Menzies, M.A., 2012. Geochemistry of the Phlegraean Fields (Italy) proximal sources for major Mediterranean tephra: Implications for the dispersal of Plinian and co-ignimbritic components of explosive eruptions. *Geochimica et Cosmochimica Acta*, 93(0): 102-128.
- Tomlinson, E.L., Smith, V.C., Albert, P.G., Aydar, E., Civetta, L., Cioni, R., Cubukcu, E., Gertisser, R., Isaia, R., Menzies, M.A., Orsi, G., Rosi, M. and Zanchetta, G., 2015. The major and trace element glass compositions of the productive Mediterranean volcanic sources: tools for correlating distal tephra layers in and around Europe. *Quaternary Science Reviews*, 118: 48-66.
- Tomlinson, E.L., Thordarson, T., Müller, W., Thirlwall, M. and Menzies, M.A., 2010. Microanalysis of tephra by LA-ICP-MS — Strategies, advantages and limitations assessed using the Thorsmörk ignimbrite (Southern Iceland). *Chemical Geology*, 279(3–4): 73-89.
- Ukstins Peate, I., Baker, J.A., Kent, A.J.R., Al-Kadasi, M., Al-Subbary, A., Ayalew, D. and Menzies, M., 2003. Correlation of Indian Ocean tephra to individual Oligocene silicic eruptions from Afro-Arabian flood volcanism. *Earth and Planetary Science Letters*, 211(3–4): 311-327.
- Ukstins Peate, I., Kent, A.J.R., Baker, J.A. and Menzies, M.A., 2008. Extreme geochemical heterogeneity in Afro-Arabian Oligocene tephra: Preserving fractional crystallization and mafic recharge processes in silicic magma chambers. *Lithos*, 102(1–2): 260-278.
- Wang, F., Chen, W.-j., Peng, Z.-c. and Li, Q., 2001. Activity of Cangbaishan Tianchi Volcano since Late Pleistocene: The constrain from geochronology of high precision U-series TIMS method. *Geochimica*, 30(1): 88-94 (in Chinese with English abstract).



- northeast China: implications for possible future eruptions. *Bull Volcanol*, 75(4): 1-14.
- Wei, H., Wang, Y., Jin, J., Gao, L., Yun, S.-H. and Jin, B., 2007. Timescale and evolution of the intracontinental Tianchi volcanic shield and ignimbrite-forming eruption, Changbaishan, Northeast China. *Lithos*, 96(1–2): 315-324.
- Xu, J., Pan, B., Liu, T., Hajdas, I., Zhao, B., Yu, H., Liu, R. and Zhao, P., 2013. Climatic impact of the Millennium eruption of Changbaishan volcano in China: New insights from high-precision radiocarbon wiggle-match dating. *Geophysical Research Letters*, 40(1): 54-59.
- Yang, L., Wang, F., Feng, H., Wu, L. and Shi, W., 2014.  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology of Holocene volcanic activity at Changbaishan Tianchi volcano, Northeast China. *Quat Geochronol*, 21: 106-114.
- Yatsuzuka, S., Okuno, M., Nakamura, T., Kimura, K., Setoma, Y., Miyamoto, T., Kim, K.H., Moriwaki, H., Nagase, T. and Jin, X., 2010.  $^{14}\text{C}$  wiggle-matching of the B-Tm tephra, Baitoushan volcano, China/North Korea. *Radiocarbon*, 52(3): 933-940.
- Yin, J., Jull, A.J.T., Burr, G.S. and Zheng, Y., 2012. A wiggle-match age for the Millennium eruption of Tianchi Volcano at Changbaishan, Northeastern China. *Quaternary Science Reviews*, 47(0): 150-159.
- Yu, H., Wu, J., Xu, J., Lin, C., Shi, L. and Chen, X., 2012. Microstructural Characteristics of the Holocene Pumice Erupted from Changbaishan Tianchi Volcano and Their Volcanological Implications. *Journal of Jilin University (Earth Science Edition)*, 42(Sup. 3): 132-144 (in Chinese with English abstract).

tephra where the geochemical data used for comparison come from are marked on the map as well. (b) Sampled stratigraphy at the Tianwen summit on the Chinese flank of the Tianchi crater, Changbaishan Volcano, modified from Chen et al. (2014).

**Fig. 2.** Schematic illustration of previous stratigraphic and chronological studies of the Tianwen summit profile, Changbaishan Volcano. The solid arrows (black, grey, yellow) indicate correlations based on stratigraphy and the dashed arrows plus question mark indicate where correlations cannot be confirmed due to conflicting or inadequate stratigraphic information. Asterisks denote no color description of tephra units and a question mark denotes uncertainty about the nature of an eruptive deposit. This paper focuses on the three upper pyroclastic fall units above the yellow pumice (i.e. light grey, dark grey and black units).

**Fig. 3.** SEM images of the proximal Changbaishan tephras. (a-b) the C-3 light grey pumice, both highly vesiculated and less vesiculated glasses with some highly vesiculated glasses being strongly stretched; (c) the C-2 dark grey pumice, vesicular and stretched glasses; (d) the C-1 black pumice samples that have the highest content of phenocryst and the least vesicularity among all the three proximal fall units.

**Fig. 4.** Tephra shard concentrations measured in sediments in the top 2 meters of samples from Lake Kushu. Shard counts are given as numbers of shards per gram of dry sediment. The star marks the peak of the tephra input.

**Fig. 5.** Light microscope photographs of tephra shards extracted from Lake Kushu sediments at 150-152 cm composite depth.

**Fig. 6.** (a) TAS classification (Le Bas et al., 1986) diagram showing the glass compositions of proximal fall deposits (i.e. C-1, C-2 and C-3) and distal Kushu tephra. The boundary separating the alkaline and subalkaline series is from Irvine and Baragar (1971). (b) Classificatory diagram for peralkaline extrusive rocks (MacDonald, 1974) showing the glasses compositions of the peralkaline proximal and distal tephras.

**Fig. 7.** Major and trace element variation diagrams showing the glass compositions of proximal fall deposits and distal Kushu tephra.

**Fig. 8.** Primitive mantle normalized trace element compositions of glasses from proximal fall deposits and distal Kushu tephra. Primitive mantle values are from Sun and McDonough (1989).

**Fig. 9.** (a) TAS classification (Le Bas et al., 1986) diagram and (b-c) major element variation diagrams showing the glass compositions of tephras reported herein along with the published data for comparison. The published data are glass compositions of the B-Tm and the widespread late Quaternary tephras from other volcanic settings in

represents late Quaternary tephra (e.g. K-Ah, AI, Aso-4 and Ata) from volcanoes in Kyushu Island, Japan (Smith et al., 2013). Ulleungdo represents the early Holocene U-Oki tephra erupted from Ulleungdo, South Korea (Smith et al., 2011b). Kamchatka represents the tephra layers from major Holocene eruptions in Kamchatka Peninsula, Russia (Kyle et al., 2011).

**Fig. 10.** Major element variation diagrams showing the glass compositions of (a-b) proximal tephra deposits in both Chinese (this study) and N. Korean ME sequences; (c-d) distal ME tephra deposits found in northeast China, Sea of Japan, Hokkaido and Greenland. ME-K represents the proximal ME sequence in N. Korean side of the crater (Sun et al., 2014a). Kushu represents the Lake Kushu in Rebun Island, Hokkaido, Japan (this study). Utsai represents the Utsai Bog in Hokkaido, Japan (Hughes et al., 2013). SHL represents the Sihailongwan Lake in northeast China (Sun et al., 2015). Japan Sea represents the distal ME tephra deposits found in Sea of Japan marine cores (Machida et al., 1990). NGRIP and NEEM represent the ultra-distal ME tephra deposits found in Greenland ice cores (Coulter et al., 2012; Sun et al., 2014a).

**Fig. 11.** 95% confidence Highest Probability Density output for radiocarbon-based Bayesian age-depth model for the uppermost 2 meters sequence of Lake Kushu (run using a Poisson process model). The model was constructed purely using AMS  $^{14}\text{C}$  dates of Kushu sediments from Müller et al. (in press), for providing independent age constraint on the Kushu tephra. The deposition model age range for the depth of 151.5 cm, which is the average position for the initial onset of the Kushu tephra, is 913-1395 cal AD.

**Fig. 12.** (a) 95% confidence Highest Probability Density output for Bayesian age-depth model for the uppermost 2 meters sequence of Lake Kushu (run using a Poisson process model). The model was constructed using the AMS  $^{14}\text{C}$  dates reported in Müller et al. (in press) and the GICC05 ice-core age for B-Tm tephra (Sun et al., 2014a) imported in the appropriate position where our proposed correlative tephra was identified; (b) As above but with the B-Tm age from the revised NS1-2011 timescale after Sigl et al. (2015).

**Table 1.**

Age results for the Millennium eruption from the dating of proximal and distal (B-Tm) tephra deposits.

**Table 2.**

Representative major, minor and trace element data for glasses from proximal volcanic deposits at Tianwen summit and distal Kushu tephra. Full geochemical data sets are available in the Supplementary material file.

shown in 95% confidence, and the relative depth information. AMS  $^{14}\text{C}$  uncal. dates are from Müller et al. (in press). Date shown in *italic* font may be an age inversion although after calibration this still contributes significant probability to the deposition model. GICC05 ice-core tephra age for the B-Tm is from Sun et al. (2014a), the revised B-Tm age based on NS1-2011 timescale is from Sigl et al. (2015). The Bayesian age-depth models are produced using OxCal v4.2.4 (Bronk Ramsey, 2013). The modelled date results for the set of 6 AMS-dated samples are from the model with GICC05 tephra age imported as shown in Fig. 12a.

ACCEPTED MANUSCRIPT

environs (pyroclastic fall and flow deposits)	dating	AD 925-959 (2 $\sigma$ ) AD 945-960 (2 $\sigma$ ) AD 930-943 (2 $\sigma$ ) AD 945-984 (2 $\sigma$ )	Yin et al. (2012) Yatsuzuka et al. (2010) Nakamura et al. (2007) Horn and Schmincke (2000)
Tianwen summit (light grey fall)	$^{40}\text{Ar}/^{39}\text{Ar}$ U-series TIMS	1.24 $\pm$ 0.51 ka 1.0 $\pm$ 0.6 ka	Yang et al. (2014) Wang et al. (2001)
<i>Distal</i>			
Lake Kushu (Rebun, northern Japan)	Bayesian Modelling	AD 933-949 (2 $\sigma$ ) AD 944-947 (2 $\sigma$ )	This study
Greenland ice cores	Ice-core chronology	AD 946-947 (NS1-2011 timescale)  AD 941 $\pm$ 1 (GICC05 timescale) AD 945 $\pm$ 4 (GISP2 timescale)	Sigl et al. (2015)  Sun et al. (2014a)
Sihailongwan Maar Lake (NE China)	Varve chronology Conventional $^{14}\text{C}$	AD 953 $\pm$ 37 AD 940-1020 (2 $\sigma$ )	Sun et al. (2015)
Lake Ni-no-Megata/ San-no-Megata (NE Japan)	Varve chronology	AD 929	Kamite et al. (2010)
Lake Ogawara (NE Japan)	Varve chronology	AD 937-938	Fukusawa et al. (1998)

---


**Table 2**

Locality	Tianwen summit							Lake		
Unit/group	C-3			C-2		C-1		Rhyolite		
Sample	10CB-2-1	14-1-1	14-1-4	14-5-1	14-5-9	14-4-1	14-4-5	RH490-20	RH555-33	RH554-3
Material	Pumice							Teph		
Major elements wt. %										
SiO <sub>2</sub>	74.98	74.73	74.93	66.06	66.42	66.08	66.24	74.90	74.97	70.34
TiO <sub>2</sub>	0.21	0.23	0.19	0.41	0.52	0.45	0.44	0.24	0.20	0.30
Al <sub>2</sub> O <sub>3</sub>	10.06	10.15	10.23	15.37	15.26	15.69	15.09	10.58	10.36	14.60
FeO <sub>t</sub>	4.15	4.23	4.08	4.96	4.91	4.37	4.92	3.92	4.13	2.98
MnO	0.13	0.10	0.02	0.13	0.12	0.08	0.20	0.10	0.08	0.09
MgO	0.00	0.04	0.00	0.19	0.17	0.22	0.17	0.00	0.00	0.00
CaO	0.20	0.16	0.21	1.25	1.30	1.25	1.23	0.23	0.23	0.19
Na <sub>2</sub> O	5.12	5.45	5.26	5.68	5.54	6.06	5.90	5.27	5.25	6.05
K <sub>2</sub> O	4.51	4.27	4.50	5.76	5.57	5.58	5.65	4.26	4.30	5.16
P <sub>2</sub> O <sub>5</sub>	0.01	0.02	0.00	0.04	0.06	0.09	0.04	0.00	0.00	0.03
Cl	0.63	0.62	0.57	0.15	0.13	0.14	0.12	0.51	0.48	0.25
Analytical total	98.02	99.61	98.08	99.29	98.47	98.13	99.89	97.02	97.97	98.60
K <sub>2</sub> O+Na <sub>2</sub> O	9.63	9.72	9.76	11.44	11.11	11.64	11.55	9.53	9.55	11.21
Trace elements ppm										
Rb	419	426	428	153	157	143	143	417	403	308
Sr	2.7	3.0	6.1	6.3	5.9	8.4	6.8	2.8	2.7	2.7
Y	125	143	156	43	43	37	42	144	130	66
Zr	2152	2377	2305	655	682	564	653	2395	2252	1260
Nb	280	302	299	93	95	79	88	303	286	195


Ba	7.7	9.1	9.9	19.2	17.6	28.6	19.9	8.5	9.7	10.8
La	135	152	173	89	92	77	87	163	148	127
Ce	288	316	336	175	175	154	170	328	307	246
Pr	31.8	35.4	39.6	19.0	19.1	16.5	18.3	34.7	33.2	23.6
Nd	118	130	149	71	71	62	69	136	123	78
Sm	26.0	29.3	32.2	13.4	14.3	11.7	13.0	31.6	26.6	16.5
Eu	<LOD	0.4	<LOD	<LOD	<LOD	0.4	0.4	0.4	<LOD	<LOD
Gd	23.9	27.2	29.3	10.3	10.5	9.6	10.1	27.4	23.0	13.3
Dy	23.3	28.2	29.7	9.1	9.3	7.8	8.7	25.9	24.0	12.6
Er	12.9	14.4	15.0	4.6	4.5	3.8	4.2	13.7	11.9	7.3
Yb	10.7	12.7	12.9	4.0	4.1	3.5	4.0	11.9	10.7	5.9
Hf	52.2	60.9	56.7	16.2	16.2	13.2	15.1	58.1	51.3	30.3
Ta	15.9	17.1	17.0	4.7	4.8	4.0	4.5	15.4	14.3	9.7
Th	48.9	56.8	52.4	15.1	15.2	12.4	14.5	52.9	46.1	30.1
U	11.4	12.7	12.8	3.4	3.4	2.8	3.0	11.5	10.7	7.0

<i>Poz-51689</i>	64.5	470 ± 25	528	325	1422	1626
<i>Poz-51700</i>	96.5	415 ± 30	569	449	1381	1502
<i>Poz-51713</i>	126.5	510 ± 35	653	500	1297	1450
<i>Poz-51721</i>	164.5	1065 ± 25	1055	1004	895	947
<i>Poz-51731</i>	196.5	1290 ± 30	1285	1180	666	771
<i>Poz-51735</i>	226.5	1445 ± 30	1386	1296	565	655
Imported tephra	Imported position	Imported ice-core ages				
B-Tm	151	AD 941±1 (GICC05 timescale)	1017	1001	933	949
		AD 946-947 (NS1-2011 timescale)	1006	1003	944	947







ACCEPTED MANUSCRIPT




ACCEPTED MANUSCRIPT




ACCEPTED MANUSCRIPT





ACCEPTED MANUSCRIPT











ACCEPTED MANUSCRIPT




ACCEPTED MANUSCRIPT






ACCEPTED MANUSCRIPT












ACCEPTED MANUSCRIPT





ACCEPTED



ACCEPTED





ACCEPTED MANUSCRIPT

A robust proximal-distal tephra correlation demonstrated chemically and chronologically.

Clarification on the problematic proximal stratigraphy via proximal-distal correlation.

The use of ice-core tephra ages to test and constrain Lake Koshu radiocarbon chronology.

The observed compositional diversity helps understand the nature of the eruption.

ACCEPTED MANUSCRIPT