



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Journal of Sport Psychology in Action

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa31635>

Paper:

Gross, M., Hall, R., Bringer, J., Cook, C., Kilduff, L. & Shearer, D. (2017). Resonant frequency training in elite sport: A case study example. *Journal of Sport Psychology in Action*, 1-11.

<http://dx.doi.org/10.1080/21520704.2017.1287797>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

1 Resonant Frequency Training in elite sport: A case study example

2 Mike J. Gross, and Ross Hall

3 Faculty of Life Sciences & Education, University of South Wales, UK. Welsh Institute of
4 Performance Science, Wales

5 Joy D. Bringer

6 Department of Sports Science, Sports Council for Wales, UK. Welsh Institute of Performance
7 Science, Wales

8 Christian J. Cook

9 School of Sport, Health and Exercise Science Bangor University Wales, UK

10 Liam P. Kilduff

11 A-STEM, Swansea University, UK. Welsh Institute of Performance Science, Wales

12 David A. Shearer

13 Faculty of Life Sciences & Education, University of South Wales, UK. Welsh Institute of
14 Performance Science, Wales

15
16 Mike J. Gross, Faculty of Life Sciences & Education, University of South Wales. Welsh Institute of
17 Performance Science, Wales; Ross Hall, Faculty of Life Sciences & Education, University of South
18 Wales; Joy D. Bringer, Department of Sports Science, Sports Council for Wales. Welsh Institute of
19 Performance Science, Wales; Christian J. Cook, School of Sport, Health and Exercise Science Bangor
20 University Wales, UK; Liam P. Kilduff, A-STEM, Swansea University, Welsh Institute of
21 Performance Science, Wales; David A. Shearer, Faculty of Life Sciences & Education, University of
22 South Wales. Welsh Institute of Performance Science, Wales

23 *Author Note*

24 This research was supported by European Regional Development Fund (ERDF) and Welsh
25 Government Academic Expertise for Business (A4B) funding.

26 The authors would like to thank the Welsh Target Shooting Federation for their continued co-
27 operation and support. The authors would also like to thank the Welsh Elite Performance Sport
28 Innovation Network (WEPSIN) for their time and expertise.

Resonant Frequency Training in Elite Sport: A case study example

Natural breaks during self-paced sports, such as moments between shots in shooting, make athletes susceptible to positive and negative emotions arising from corresponding facilitative and debilitating appraisals (e.g., Neil, Fletcher, Hanton, & Mellalieu, 2007; Uphill & Jones, 2007). Heart rate variability (HRV) biofeedback interventions (e.g., Lagos et al., 2008) teach athletes to perform breathing techniques which acutely increase HRV in order to manage emotional states on-demand and facilitate performance (Lehrer & Gevirtz, 2014). HRV is the change in intervals between heart beats and provides an indicator of regulatory influences on the heart (Shaffer, McCraty, & Zerr, 2014). Broadly, increased HRV indicates increased physiological and psychological adaptability to internal and external demands (Bertsch, Hagemann, Naumann, Schachunger, & Schulz, 2012).

Research demonstrates that HRV biofeedback interventions have benefits for a range of clinical and performance related conditions (see Gevirtz, 2013 for a review). Lehrer & Gevirtz (2014) propose a number of mechanisms for these benefits including; restoring autonomic homeostatic balance to reduce sympathetic overflows; stimulating the regulatory baroreflex to provide context-appropriate blood pressure; stimulating the cholinergic anti-inflammatory system; enhancing the vagus nerve mediated heart-brain link to provide emotional centres of the brain with a relaxed visceral state; increased gaseous exchange in the alveoli via phased relations of breathing and heart rate; mechanical stretching of airwaves by slow rhythmic breathing; and meditation or mindfulness related effects of focused attention on one's breathing. However, reducing autonomic stress by stimulating the regulatory baroreflex, and enhancing vagal afferent signals (heart-brain communications regarding the visceral state of the body) are the most widely evidenced therapeutic mechanisms (e.g., Thayer, Åhs, Fredrikson, Sollers III, & Wager, 2012). In sport, these two related mechanisms may combine with distraction-related effects of focused breathing to reduce cognitive and somatic competitive anxiety.

HRV biofeedback interventions have reduced anxiety-related deficits in dance technique (Gruzelier, Thompson, Redding, Brandt, Steffert, 2014) and basketball skills (Paul & Garg, 2012). Similar benefits for self-paced sport skills, such as enhanced positive emotions, decreases in cognitive and somatic anxiety, and objective improvements in performance have also been observed (Lagos et

1 al. 2008; Paul, Garg, & Sandhu, 2012). Improvements in major depression have been achieved after
2 four sessions of HRV biofeedback (e.g., Karavidas et al., 2007), and five-session interventions with 5-
3 minute daily home practice sessions improved the on-demand emotional regulation abilities of elite
4 sport support and management staff (Gross et al., 2016). This evidence indicates that HRV
5 biofeedback may be a time-efficient method to achieve affective and performance-related benefits.
6 Resonant Frequency Training (RFT) is a specific HRV biofeedback method that consists of rhythmic,
7 abdominal breathing at approximately six breaths per minute (0.1Hz; Lehrer, Vaschillo, & Vaschillo,
8 2000). The method identifies and teaches participants to perform their Resonant Frequency (RF;
9 bespoke breathing rate at which maximal HRV occurs; Lehrer et al.). Users become more aware of
10 psychophysiological interactions by observing live changes in their data on a display, and can use
11 their RF to preemptively or reactively manage the effects of anxiety and stress (Gross et al.). This
12 article presents an applied case study of a RFT intervention with an elite athlete to learn an on-
13 demand emotional regulation technique.

14 **Case Study**

15 Louise (pseudonym) is a 45 year old Olympic Trap shooter with 16 years competitive
16 experience. Louise was experiencing negative thoughts and emotions between shots (especially after
17 missed targets), and difficulties dealing with unexpected interruptions to the flow of a competitive
18 round (e.g., target malfunctions), both which were unhelpful for her preparation and performance of
19 the next shot. During an Olympic Trap round there is approximately 1-minute between shots in which
20 athletes can prepare for the next, and one lost target can mean the difference between silver or gold
21 medals. Louise had found reciting songs useful to substitute anxious thoughts, however this technique
22 was not consistently effective in major competitions where anxiety also manifested as somatic
23 symptoms (i.e., perceptible increases in breathing & heart rates). The cognitive and somatic anxiety
24 relieving effects of HRV biofeedback (Lehrer & Gevirtz, 2014) meant that RFT was a suitable
25 intervention to provide Louise with an on-demand emotional regulation tool for use between shots.
26 Additionally, logistical constraints and the potential for benefits from five-sessions (e.g., Gross et al.,
27 2016; Karavidas et al., 2007) meant that RFT was an efficient and convenient intervention for Louise.

28 **Measures**

1 **Lifestyle & Adherence Measures.** Factors that may influence HRV indices (e.g., caffeine;
2 Notarius & Floras, 2012) were recorded at the beginning of each RFT session (cf. Gross et al., 2016).
3 At the beginning of sessions three, five, and Simulated Competition, Louise self-reported the number
4 of advised home practice sessions that she had not performed. These numbers were used to calculate
5 the percentage of advised home practice sessions performed between practitioner-led sessions. Louise
6 was not contacted between practitioner-led sessions, nor asked to keep a diary of her practice sessions.

7 **Physiological Measures.** All RFT physiological measures were captured by the Nexus-10
8 encoder (Mind Media, Roermond-Herten, the Netherlands) with dedicated photoplethmograph (PPG;
9 placed on non-dominant middle finger; sampled at 128Hz) and respiration sensors (pneumony
10 impedance; Velcro attached 1-inch above the navel; sampled at 128Hz) via bluetooth to the Biotrace+
11 software. HRV indices included total spectral power (TP; indicative of total HRV), low frequency
12 band percentage of TP (LF%; representative of baroreflexive activity and vagal afferents; Goldstein et
13 al., 2011), low frequency peak percentage of TP (LFPK%; indicative of HRV % influenced by RF
14 breathing), and the route-mean square of successive differences (RMSSD; representative of
15 parasympathetic HRV). Respiratory indices included mean respiratory rate (mResp; breaths per
16 minute) and mean distance from the target resonant frequency (mDist; breaths per minute). During a
17 simulated competition, respiratory data was captured (via pneumony impedance) using a Hexoskin
18 undergarment & encoder (Hexoskin, Montréal, Canada). The Hexoskin mobile application was used
19 to view live computed statistics (i.e., 1Hz respiratory rates) and a 10-second window of live raw
20 respiratory data. Tri-axis accelerometer (sampled at 64Hz with 0.004g resolution) and
21 electrocardiogram (ECG; 1-channel sampled at 256Hz) data were also collected via the Hexoskin.
22 Hexoskin data were analysed using the EDFbrowser software (Version 1.55; Van Beelen, 2014). For
23 home RF practice, Louise's mobile device (iOS) was equipped with the Breathe2Relax (National
24 Centre for Telehealth & Technology; United States of America) breath pacing application.

25 **Client Feedback.** Two client feedback interviews were used to reflect on the intervention
26 experience, use of RF, and benefits to performance. At the final RFT session, Louise was asked
27 "Please tell me about your experiences of the intervention" and "When have you used your resonant
28 frequency?". Following the simulated competition, Louise and her coach were asked "What are the

1 benefits of using RF breathing during a shooting round?”. Responses were recorded verbatim in hand-
2 written notes during both interviews. Twelve months after the simulated competition, Louise
3 described the performance impact via email.

4 **Intervention**

5 Ethical approval was granted via the corresponding author’s institution and Louise provided informed
6 consent before the intervention began. Figure 1 illustrates Louise’s progression through the RFT
7 intervention and temporal gaps between sessions. The RFT sessions consisted of RF paced breathing
8 (PB) and accuracy test (AT) activities, each now described in succession.

9 **Session 1.** To record resting physiology data, Louise was connected to the PPG and
10 respiration sensors and asked to relax for 5-minutes while sitting up-right and alone (see Kleck et al.,
11 1976). Louise then completed an RF estimation procedure which guided her breathing via a bar graph
12 at seven consecutively reducing frequencies (i.e., 7 breaths/minute to 4 breaths/minute) for 2-minutes
13 each with an inhale to exhale ratio of 2:3 (cf. Lehrer et al., 2000). Louise’s RF was estimated using
14 TP, LF%, and LFPK% for each frequency derived from the Fast Fourier Transformation (FFT)
15 function in Biotrace+.

16 **Session 2.** To confirm her RF, Louise’s breathing was paced at three consecutively reducing
17 frequencies for 3-minutes each (the estimated RF from Session 1 was placed as the middle
18 frequency). LFPK% derived from the FFT was used to immediately confirm Louise’s RF, which she
19 then performed for 5-minutes using the breath pacer (PB1). Louise’s Breathe2Relax application was
20 programmed with her RF timings, she was advised to practice for 5-minutes each day, and told that
21 her RF accuracy would be tested in the future sessions.

22 **Session 3.** Louise was assessed on her ability to perform her RF for 3-minutes without a
23 breath pacer (AT1). As per Gross et al., (2016), verbal feedback on RF accuracy was provided at the
24 half-way point (e.g., ‘too slow, shorten your exhale’), and a discussion of TP, LFPK%, mResp and
25 mDist data followed the end of the accuracy test. Following AT1, Louise completed a 5-minute RF
26 paced breathing exercise (PB2; as per PB1) and then performed an identical 3-minute RF accuracy
27 test (AT2), again with data discussions.

1 **Session 4.** This session was identical to Session 3 except the accuracy tests either side of
2 paced RF breathing (PB3) were 4-minutes in length (AT3 & AT4). In a discussion around RF use
3 during performance, Louise suggested that the approximate 1-minute pause between shots would be
4 an appropriate time to incorporate her RF into her performance.

5 **Session 5.** The accuracy tests in this session (AT5 & AT6) were altered to simulate how
6 Louise would incorporate her RF into performance. Conducted either side of the final paced RF
7 breathing activity (PB4), both accuracy tests comprised five separate, 1-minute between-shot periods.
8 Louise was instructed to stand in her firing position, imagine that she had performed a shot and
9 reproduce the same movements (e.g., throwing away the empty cartridges), then perform her RF for
10 1-minute. No feedback was given during these accuracy tests, however data were discussed after each
11 activity (as per Session 4). As this was the final RFT session, Louise was debriefed and instructed to
12 use her RF on-demand for pre-competition and every-day emotional regulation, to use her RF
13 between shots, and to periodically practice her RF using Breathe2Relax.

14 **Simulated Competition.** Louise's RF accuracy between shots was assessed during a
15 simulated Olympic Trap round with five other shooters, following International Sport Shooting
16 Federation protocols. Louise wore the Hexoskin and computed statistics were visually monitored on
17 the mobile application, Louise suggested that wearing the Hexoskin did not impact her shooting. A
18 debrief including Louise's coach followed the round, quantitatively focused on RF accuracy between
19 shots (using mResp), and qualitatively focused on accuracy difficulties. Louise and her coach then
20 commented on the benefits of using her RF between shots and were provided with a written RF
21 accuracy report using mDist values.

22 **Data Analysis**

23 Using guidance for HRV data cleaning and analysis (Nunan, Sandercock, & Brodie, 2010;
24 Task Force, 1996), six recordings required some inter-beat intervals (IBI) to be removed due to
25 artifacts in the PPG signal (See Gross et al. 2016 for full details of these procedures). All PPG heart
26 beat waveforms were visually compared to three previous and three preceding beats to assess the
27 morphology for Biotrace+ to accurately determine IBI. Recordings with IBIs removed were
28 interpolated at 4 Hz. Mean IBIs and mResp for all recordings are available upon request to the

1 corresponding author. Four resting recordings required between 1.6% and 14.2% of IBI removal, two
2 were accuracy tests that required 0.7% (AT3) and 1.4% (AT4) IBI removal. For AT5 and AT6, the
3 combined mDist across all five 1-minute accuracy tests was used. Raw Hexoskin x-axis accelerometer
4 and abdominal respiration data were time-synchronised to identify shots during the simulated
5 competition, and thus highlight periods of RF breathing between shots. mResp and mDist were then
6 calculated as per the RFT session data, Hexoskin ECG data was not of sufficient quality to derive
7 HRV statistics from.

8 **Results**

9 **Lifestyle and Intervention Adherence**

10 Louise reported similar sleep length and quality, and similar water, food and caffeine intake before all
11 sessions. Louise's adherence to home practice increased across the intervention, from 54% (RFT 2 to
12 RFT 3), to 82% (RFT 4 to RFT 5) and 98% between RFT 5 and the simulated competition.

13 **RFT & Simulated Competition**

14 Compared to first resting measures, Louise's percentage of LF activity increased during the final
15 resting measures (from 38.95% to 71.22%), however a lower respiratory rate in the latter recording
16 (mResp = 14.05 vs mResp = 7.85) will have overemphasised this effect. Across the 5-minute paced
17 breathing activities, Louise increased LFPK% (PB1 = 50.23% to PB4 = 56.71%), LF% (PB1 =
18 87.66% to PB4 = 94.94%), and RMSSD (PB1 = 70.88ms to PB4 = 90.58ms). She achieved similar
19 effects in 4-minute RF accuracy tests (i.e., AT4; LFPK% = 50.65%, LF% = 81.45%, RMSSD =
20 82.24ms), and similar LF% effects in 1-minute accuracy tests (AT5; 70.15% \pm 10.07, & AT6; 62.31%
21 \pm 15.71) both of which increased in comparison to average resting LF% values (39.68% \pm 8.64). These
22 increases in HRV were accompanied by increased RF accuracy (decreased mDist) across the six
23 accuracy tests, see Figure 2. These results demonstrate that during standing RF breathing for 1-minute
24 Louise was able to increase HRV above resting levels and to a similar extent as seated paced
25 breathing. Hexoskin respiratory data indicated that Louise was able to accurately reproduce her RF
26 between shots during a simulated competition, however, there was some variance in her accuracy as
27 indicated by mDist in Figure 3.

28 **Client Feedback**

1 Louise indicated that RF tests enhanced her confidence to implement RF and efficacy beliefs in
2 generating the psychophysiological benefits:

3 I feel confident that I can use my breathing and still create those heart rate changes we see
4 after I've followed the pacer on the computer.

5 Louise also described an instance during a competition when she used her RF:

6 I used it almost as soon as I had missed the second target and I instantly felt calmer and more
7 controlled...I felt back to normal and cleared up the rest with my first barrel [using one shot
8 to hit each target].

9 Louise described the benefits she saw in RF breathing during performance:

10 ...I can keep calm during a round and handle unexpected interruptions, it's like a safety net.

11 Louise's coach suggested that he saw a kinematic benefit to Louise's technique when using her RF
12 between shots:

13 She's smoother to the shoulder [raising the gun into the shoulder before a shot] and just looks
14 more controlled.

15 After one year, Louise summarised improvements in her performance which RFT contributed to:

16 My average score has lifted approximately one target a round ... which doesn't sound much
17 but actually is a massive difference because 3 targets per comp is the difference from being an
18 "also ran" to making the final.

19 Discussion

20 Louise's case highlights how RFT interventions can raise an athlete's awareness of their
21 psychophysiological response to stress, swiftly teach a technique to combat cognitive and somatic
22 anxiety, and successfully apply the technique in elite sport competitions. Louise found that influencing
23 her HRV via paced breathing activities (e.g., RF estimation procedure in Session 1), and reviewing
24 the mechanisms in action raised her awareness of the psychophysiological control she could exert via
25 breathing. She suggested that the RF accuracy tests helped her learn her RF, along with varied (times
26 & situations) home practice and self-testing RF accuracy by periodically looking away from the
27 breath pacing mobile application. High adherence rates reflect Louise's engagement with the
28 intervention, which combined with the simulated competition RF accuracy discussions to reinforce

1 the application to performance. Louise suggested that her focus on RF breathing replaced negative
2 thoughts with a neutral image (i.e., a breath pacing bar-graph), and that the psychophysiological
3 control enabled her to consistently manage cognitive and somatic anxiety during competitions to
4 support optimal performance.

5 Similar to other HRV biofeedback interventions in sport, during paced RF breathing Louise
6 was able to increase HRV indicative of baroreflex stimulation and increased vagal afferents (cf. Lagos
7 et al., 2008; Paul et al., 2012). By the end of the intervention, she recreated these HRV effects via RF
8 breathing in seated accuracy tests and standing imagined simulations, then reproduced accurate RF
9 breathing patterns between shots during a simulated competition. Partially supporting other case
10 studies such as Lagos et al., Louise may have increased her resting HRV, however a slow respiratory
11 rate in the final resting measures and the removal of some resting data may have overemphasised this
12 effect. Supporting the coaches' kinematic observations, previous research has shown reductions in
13 choice-reaction and movement times (Paul et al.), and improved dance technique (Gruzelier et al.,
14 2014) following HRV biofeedback. Additionally, preparatory heart rate deceleration has been
15 implicated in expertise and success during self-paced closed skills (Cooke et al., 2010), and shooters
16 often exhale during their shot process which decreases heart rate, an effect strengthened by HRV
17 biofeedback (Lehrer et al., 2000).

18 Outside of RFT sessions, objective evidence for intervention effectiveness is limited to
19 simulated competition respiratory data and anecdotal evidence for both emotional regulation and
20 improved performance. Insufficient simulated competition HRV data quality may have been due to
21 inappropriate Hexoskin fit, and sufficient HRV data may have provided further evidence for
22 psychophysiological benefits. As this was a true case study of applied practice and not a research study,
23 it is important to note the lack of pre-intervention baseline measures of emotional regulation, and the
24 inherent limitations of client feedback. Much of the client feedback has been omitted due to limited
25 space, however Louise described multiple instances where she used RF breathing to manage
26 competitive and general life stressors. It is difficult to determine the precise contribution that Louise's
27 RFT made to performance improvements, and other interventions have similarly reported concomitant
28 long-term improvements (e.g., Lagos et al.). However, the intervention aim was to teach Louise an

1 on-demand emotional regulation technique for use during competition. The combination of
2 physiological and anecdotal evidence suggests that this aim was achieved.

3 More research is needed to explore potential psychomotor benefits of HRV biofeedback for
4 athletes, however growing evidence for psychological benefits alone suggests that these interventions
5 warrant wider adoption in sport psychology practice. Louise's RFT was data-driven using specialist
6 equipment, however the psychophysiological benefits from RFT can be achieved without such
7 investment. As a cost-free solution to teach the techniques without HRV data collection, practitioners
8 can use mobile applications to pace an individual's breathing at a human average RF (four seconds
9 inhale, six seconds exhale; Lehrer et al., 2000). Alternatively, there is growing support for the efficacy
10 and accuracy of low-cost (approximately \$120), data-based interventions using smart phone devices
11 (Heathers, 2013). These solutions are often supplied with finger or ear clip PPG sensors to measure
12 HRV during paced breathing activities, and may enable users to program bespoke breathing patterns
13 for differing purposes. Some newer generations of wearable, smart fitness trackers include paced
14 breathing activities to help users manage daily stress, and could provide another way to begin and
15 practice HRV biofeedback techniques. Regardless of the solution, we recommend practice in differing
16 situations and accuracy testing to reinforce confidence and efficacy in achieving the benefits in
17 performance situations.

18 **References**

- 19 Bertsch, K., Hagemann, D., Naumann, E., Schachinger, H., & Schulz, A. (2012). Stability of heart
20 rate variability indices reflecting parasympathetic activity. *Psychophysiology*, *49*, 672-682.
21 doi:10.1111/j.1469-8986.2011.01341.x
- 22 Cooke, A., Kavussanu, M., Gallicchio, G., Willoughby, A., McIntyre, D., & Ring, C. (2010).
23 Preparation for action: Psychophysiological activity preceding a motor skill as a function of
24 expertise, performance outcome, and psychological pressure. *Psychophysiology*, *51*, 374-384.
25 doi:10.1111/psyp.12182
- 26 Gevirtz, R. (2013). The promise of heart rate variability biofeedback: Evidence-based applications.
27 *Biofeedback*, *41*, 110-120. doi:10.5298/1081-5937-41. 3.01

- 1 Goldstein, D. S., Benthon, O., Park, M.-Y., & Sharabi, Y. (2011). Low-frequency power of heart rate
2 variability is not a measure of cardiac sympathetic tone but may be a measure of modulation
3 of cardiac autonomic outflows by baroreflexes. *Experimental Physiology*, *96*, 1255-1261.
4 doi:10.1113/expphysiol.2010.056259
- 5 Gross, M. J., Shearer, D. A., Bringer, J. D., Hall, R., Cook, C. J., & Kilduff, L. P. (2016). Abbreviated
6 resonant frequency training to augment heart rate variability and enhance on-demand
7 emotional regulation in elite sport support staff. *Applied Psychophysiology and Biofeedback*,
8 263-274. doi:10.1007/s10484-015-9330-9
- 9 Gruzelier, J. H., Thompson, T., Redding, E., Brandt, R., Steffert, T. (2014). Application of alpha/theta
10 neurofeedback and heart rate variability training to young contemporary dancers: State
11 anxiety and creativity. *International Journal of Psychophysiology*, *93*, 105-111.
12 doi:10.1016/j.ijpsycho.2013.05.004
- 13 Heathers, J. A. J. (2013). Smartphone-enabled pulse rate variability: An alternative methodology for
14 the collection of heart rate variability in psychophysiological research. *International Journal*
15 *of Psychophysiology*, *89*, 297-304. doi:10.1016/j.ijpsycho.2013.05.017
- 16 Karavidas, M. K., Lehrer, P. M., Vaschillo, E., Vaschillo, B., Marin, H., Buyske, S., & Hasset, A.
17 (2007). Preliminary results of an open label study of heart rate variability biofeedback for the
18 treatment of major depression. *Applied Psychophysiology and Biofeedback*, *32*, 19-30.
19 doi:10.1007/s10484-006-9029-z
- 20 Kleck, R. E., Vaughan, R. C., Cartwright-Smith, J., Vaughan, K. B., Colby, C. Z., & Lanzetta, J. T.
21 (1976). Effects of being observed on expressive, subjective and physiological responses to
22 painful stimuli. *Journal of Personality and Social Psychology*, *34*, 1211-1218.
23 doi:10.1037/0022-3514.34.6.1211
- 24 Lagos, L., Vaschillo, E., Vaschillo, B., Lehrer, P., Bates, M., & Pandina, R. (2008). Heart rate
25 variability as a strategy for dealing with competitive anxiety: A case study. *Biofeedback*, *36*,
26 109-115.

- 1 Lehrer, P. M., & Eddie, D. (2013). Dynamic processes in regulation and some implications for
2 biofeedback and biobehavioral interventions. *Applied Psychophysiology and Biofeedback, 38*,
3 143-155. doi:10.1007/s10484-013-9217-6
- 4 Lehrer, P. M., & Gevirtz, R. (2014). Heart rate variability biofeedback: How and why does it work?
5 *Frontiers in Psychology, 5*, 756. doi:10.3389/fpsyg.2014.00756
- 6 Lehrer, P. M., Vaschillo, E., & Vaschillo, B. (2000). Resonant frequency biofeedback training to
7 increase cardiac variability: Rationale and manual for training. *Applied Psychophysiology and*
8 *Biofeedback, 25*, 177-189.
- 9 Neil, R., Fletcher, D., Hanton, S., & Mellalieu, S. D. (2007). (Re)conceptualizing competition stress
10 in sport performers. *Sport and Exercise Psychology Review, 3*, 23-29.
- 11 Notarius, C. F., & Floras, J. S. (2012). Caffeine enhances heart rate variability in middle-aged healthy,
12 but not heart failure subjects. *Journal of Caffeine Research, 2*, 77-82.
13 doi:10.1089/jcr.2012.0010
- 14 Nunan, D., Sandercock, G. R., & Brodie, D. A. (2010). A quantitative systematic review of normal
15 values for short-term heart rate variability in healthy adults. *Pacing and Cardiac*
16 *Electrophysiology, 33*, 1407-1417. doi:10.1111/j.1540-8159.2010.02841.x
- 17 Paul, M., & Garg, K. (2012). The effect of heart rate variability biofeedback on performance
18 psychology of basketball players. *Applied Psychophysiology and Biofeedback, 37*, 131-144.
19 doi:10.1007/s10484-012-9185-2
- 20 Paul, M., Garg, K., & Sandhu, J. S. (2012). Role of biofeedback in optimizing psychomotor
21 performance in sports. *Asian Journal of Sports Medicine, 3*, 29-40.
- 22 Shaffer, F., McCraty, R., & Zerr, C. L. (2014). A healthy heart is not a metronome: An integrative
23 review of the heart's anatomy and heart rate variability. *Frontiers in Psychology, 5*, 1040.
24 doi:10.3389/fpsyg.2014.01040
- 25 Task Force. (1996). Heart Rate Variability: Standards of measurement, physiological interpretation
26 and clinical use. *Circulation, 93*, 1043-1065.
- 27 Thayer, J. F., Ahs, F., Fredrikson, M., Sollers III, J. J., & Wager, T. D. (2012). A meta-analysis of
28 heart rate variability and neuroimaging studies: Implications for heart rate variability as a

- 1 marker of stress and health. *Neuroscience and Biobehavioral Reviews*, 36, 747-756. Doi:
2 10.1016/j.neubiorev.2011.11.009
- 3 Uphill, M. A., & Jones, M. V. (2007). The antecedents of emotions in elite athletes. *Research*
4 *Quarterly for Sport and Exercise*, 78, 79-89.
- 5 Van Beelen, T. (2014). EDFbrowser [Computer software]. Retrieved from
6 <http://www.teuniz.net/edfbrowser/index.html>
7

1 *Figure 1. Case study timeline.*

2


3

4

5


6

7



8 *Figure 2. Resonant Frequency (RF) accuracy across six accuracy tests (AT). mDist = mean distance*
 9 *from RF in breaths per minute (bpm).*

10



11

12

13

14

15

1 *Figure 3.* Olympic Trap layout with timing and respiratory data. Targets appear from an underground
 2 Clay Trap, traveling away from the shooting positions, shooters fire outwards into the Shooting
 3 Range. Six shooters fire from positions 1 to 5, position 6 is a holding position. The average time
 4 Louise spent Resonant Frequency (RF) breathing between positions is presented in seconds with
 5 standard deviations. RF accuracy is presented as mDist values (mean distance from target RF in
 6 breaths per minute), where 0 represents perfect accuracy. Negative values indicate a breathing pace
 7 slower than the target RF, whereas positive values indicate a breathing pace faster than the target RF.

8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20

