



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa32149>

Paper:

Perkins, W., Dunbar, D., Godwin, J. & Jehu, G. (in press). Diagrammar in an Extended Theory of Gravity.

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Diagrammar in an Extended Theory of Gravity

David C. Dunbar, John H. Godwin, Guy R. Jehu and Warren B. Perkins

*College of Science,
Swansea University,
Swansea, SA2 8PP, UK*

Abstract

We show how the S -matrix of an extended theory of gravity defined by its three-point amplitudes can be constructed by demanding factorisation. The resultant S -matrix has tree amplitudes obeying the same soft singularity theorems as Einstein gravity including the sub-sub-leading terms.

Keywords:

1. Introduction

Scattering amplitudes are traditionally defined from a quantum field theory and the resulting Feynman vertices and Feynman diagrams. Alternatively, the amplitudes can be regarded as the fundamental objects which define the theory perturbatively. It is not very useful to define a theory by specifying the entire S -matrix explicitly but it is an important question whether the S -matrix can be defined from a minimal set of data and rules i.e. a “*diagrammar*” [1]. Once a minimal set of amplitudes is specified we aim to construct all other amplitudes by demanding they have the correct symmetries and singularities. Defining the S -matrix using its singularities is a long-standing programme which is still active and fruitful [2, 3, 4, 5, 6, 7].

In this letter we build an S -matrix from a set of three-point amplitudes using their singularity structure. The S -matrix corresponds to a theory of Einstein gravity extended by the addition of R^3 terms. We are working with massless theories and view the amplitude as a function of the twistor variables λ_i^a and $\bar{\lambda}_i^{\dot{a}}$, $M(\lambda_i, \bar{\lambda}_i)$. The spinor products $\langle ij \rangle, [ij]$ are $\langle ij \rangle = \epsilon_{ab} \lambda_i^a \lambda_j^b$, $[ij] = \epsilon_{\dot{a}\dot{b}} \bar{\lambda}_i^{\dot{a}} \bar{\lambda}_j^{\dot{b}}$. In this formalism amplitudes have a well-defined “spinor weight”. Counting λ_i as weight +1 and $\bar{\lambda}_i$ as -1, then the amplitude has

weight +4 for a negative helicity graviton and -4 for a positive helicity graviton.

We define the theory starting with the the usual three-point amplitudes of Einstein gravity:¹

$$\begin{aligned} V_3(1^-, 2^-, 3^+) &= \frac{\langle 1 2 \rangle^6}{\langle 1 3 \rangle^2 \langle 3 2 \rangle^2} , \\ V_3(1^+, 2^+, 3^-) &= \frac{[1 2]^6}{[1 3]^2 [3 2]^2} , \\ V_3(1^+, 2^+, 3^+) &= V_3(1^-, 2^-, 3^-) = 0 . \end{aligned} \tag{1}$$

These amplitudes have the correct spinor weight and are quadratic in the momenta. These amplitudes are only defined for complex momenta. For an on-shell three-point amplitude the condition $k_1 + k_2 + k_3 = 0$ demands $k_1 \cdot k_2 = 0$ etc. For real momenta this implies $\langle ij \rangle = [ij] = 0$ and the vertices are all zero. However if we consider complex momenta then we can have $\lambda_1 \sim \lambda_2 \sim \lambda_3$ but $[ij] \neq 0$.

The tree amplitudes for Einstein gravity can be computed recursively starting from these [8, 9, 10]. We show that a similar construction can be used for an extended theory.

We extend this theory by adding additional three-point amplitudes which are of higher power in momenta. To be non-trivial, these three-point amplitudes must either be functions of $\langle ij \rangle$ or $[ij]$ exclusively. The simplest polynomial amplitudes arise with six powers of momenta and are

$$\begin{aligned} V_3^\alpha(1^-, 2^-, 3^-) &= \alpha \langle 1 2 \rangle^2 \langle 2 3 \rangle^2 \langle 3 1 \rangle^2 , \\ V_3^\alpha(1^+, 2^+, 3^+) &= \alpha [1 2]^2 [2 3]^2 [3 1]^2 \end{aligned} \tag{2}$$

where α is an arbitrary constant. We also have

$$V_3^\alpha(1^-, 2^-, 3^+) = V_3^\alpha(1^+, 2^+, 3^-) = 0 , \tag{3}$$

there being no polynomial function with the correct spinor and momentum weight. These are essentially the unique choice for a three-point amplitude [11].

¹We remove a factor of $i(\kappa/2)^{n-2}$ from the n -point amplitude.




Figure 1: The non-zero three-point amplitudes

The amplitudes in this theory can be expanded as a power series in α ,

$$M_n(1, \dots, n) = \sum_{r=0} \alpha^r M_n^{(r)}(1, \dots, n) \quad (4)$$

where $M_n^{(0)}$ is the Einstein gravity amplitude. Here we focus on the $r = 1$ part of the extended theory. This being the leading deformation of the theory from Einstein gravity.

The theory we are considering would arise using field theory methods from the Lagrangian

$$L = \int d^D x \sqrt{-g} (R + C_\alpha R_{abcd} R^{cdef} R_{ef}{}^{ab}) \quad (5)$$

where $C_\alpha = \alpha/60$. However we note that to do so would involve determining increasingly complicated n -point vertices as the Lagrangian is expanded in the graviton field. As we will see the three-point amplitudes are sufficient to completely determine the S -matrix.

The key element is that the entire S -matrix is determined from these vertices if we demand that the amplitudes factorise on simple poles. Specifically, for any partition of the external legs into two sets, $\{k_{L_1}, k_{L_2}, \dots, k_{L_l}\}$ and $\{k_{R_1}, k_{R_2}, \dots, k_{R_m}\}$ with $l + m = n$ and $l, m \geq 2$, if $K = \sum_{j=1}^l k_{L_j}$, then when $K^2 \rightarrow 0$ the amplitude is singular with the simple pole being

$$M_n^{\text{tree}} \xrightarrow{K^2 \rightarrow 0} \sum_{\lambda=\pm} \left[M_{l+1}^{\text{tree}}(k_{L_1}, \dots, k_{L_l}, -K^\lambda) \frac{i}{K^2} M_{m+1}^{\text{tree}}(K^{-\lambda}, k_{R_1}, \dots, k_{R_m}) \right]. \quad (6)$$

We can excise the pole in K^2 by shifting to complex momenta and applying methods of complex analysis. There are two shifts which we use to

generate the S -matrix. Firstly there is the original Britto-Cachazo-Feng-Witten (BCFW) shift [5],

$$\lambda_i \longrightarrow \lambda_i + z\lambda_j, \quad \bar{\lambda}_j \longrightarrow \bar{\lambda}_j - z\bar{\lambda}_i. \quad (7)$$

For Einstein gravity this shift is sufficient to generate the tree level S -matrix [12]. Additionally we can use the Risager shift [13],

$$\begin{aligned} \lambda_i &\longrightarrow \lambda_i + z[jk]\lambda_\eta, \\ \lambda_j &\longrightarrow \lambda_j + z[kl]\lambda_\eta, \\ \lambda_k &\longrightarrow \lambda_k + z[lj]\lambda_\eta, \end{aligned} \quad (8)$$

where λ_η is an arbitrary spinor. Both shifts change the momenta to be functions of z whilst leaving all momenta null and preserving overall momentum conservation. We need both shifts to construct the S -matrix for the extended theory. By considering the integral

$$\int_\gamma \frac{M(z)}{z} \quad (9)$$

where γ is a closed contour, *provided* $M(z)$ vanishes at infinity the unshifted amplitude, $M(0)$, can be obtained from the singularities in the amplitude. These occur at points z_i where $K_i^2(z) = 0$. At these points,

$$K_i^2(z) = -\frac{(z - z_i)}{z_i} \times K_i^2(0) \quad (10)$$

and we obtain,

$$M_n^{\text{tree}}(0) = \sum_{i,\lambda} M_{l_i+1}^{\text{tree},\lambda}(z_i) \frac{i}{K_i^2(0)} M_{m_i+1}^{\text{tree},-\lambda}(z_i), \quad (11)$$

where the summation over i is only over factorisations where there are shifted legs on both sides of the pole. This is the on-shell recursive expression of [5]. Note that if $M(z)$ does not vanish at infinity this does not imply factorisation is insufficient to determine the amplitude but only that that particular shift can not be used to engineer the amplitude.

Expressions obtained from (11) are not manifestly symmetric as the choice of shift legs breaks crossing symmetry, however symmetry is restored in the sum. This is a highly non-trivial check that the amplitude has been computed successfully.

2. Generating the amplitudes

In this section we give some of the details of the process of generating the leading α contribution to the S -matrix.

Four-Point Amplitudes: The three-point amplitudes are our inputs so the first outputs are the four-point amplitudes. There are three independent helicity configurations,

$$M_4(1^+, 2^+, 3^+, 4^+), \quad M_4(1^-, 2^+, 3^+, 4^+), \quad M_4(1^-, 2^-, 3^+, 4^+). \quad (12)$$

Of these the first two are vanishing in Einstein gravity with only the last being non-zero: which is consequently termed the ‘‘Maximally-Helicity-Violating’’ (MHV) amplitude. For $M_4^{(1)}$ the reverse is true: $M_4^{(1)}(1^-, 2^-, 3^+, 4^+) = 0$ since there are no possible factorisations, while $M_4^{(1)}(1^+, 2^+, 3^+, 4^+)$ and $M_4^{(1)}(1^-, 2^+, 3^+, 4^+)$ are non-zero.

The factorisations of the n -point all-plus amplitude are shown in fig. 2, and the factorisations of the four-point single minus amplitude are shown on fig. 3.




Figure 2: Factorisations of the n -point all-plus




Figure 3: Factorisations of the four-point single minus amplitude

These factorisations can be excited using either of the shifts in (7) and (8). In the all-plus case only the second results in an amplitude with the correct symmetries. This in indication that (7) yields a shifted all-plus amplitude

that does not vanish at infinity. Conversely, for the single minus amplitude we must use the BCFW shift. Performing the shifts and evaluating the amplitudes we obtain

$$\begin{aligned} M_4^{(1)}(1^+, 2^+, 3^+, 4^+) &= 10 \left(\frac{st}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 41 \rangle} \right)^2 stu, \\ M_4^{(1)}(1^-, 2^+, 3^+, 4^+) &= \left(\frac{[24]^2}{[12] \langle 23 \rangle \langle 34 \rangle [41]} \right)^2 \frac{s^3 t^3}{u}. \end{aligned} \quad (13)$$

The other non-zero amplitudes are available by conjugation. For the all-plus amplitude the recursion generates terms that contain the arbitrary spinor λ_η , however the sum of terms is independent of λ_η and simplifies to the above. These four-point amplitudes due to a R^3 term have been computed using field theory methods long ago [14]. These amplitudes vanish to all orders in a supersymmetric theory: a fact used to show supergravity was two-loop ultraviolet finite [15, 16]. The above expressions are in a spinor helicity basis but agree once this is accounted for. In [17] these four-point amplitudes were also obtained using a ‘‘all-line recursion’’ technique where all legs have shifted momenta. These expressions also appear as the UV infinite pieces of both two-loop gravity in four dimensions [18, 19] and one-loop gravity in six dimensions [20].

Five-Point Amplitudes: As before the shift (8) yields an all-plus amplitude that is independent of λ_η and has full crossing symmetry:

$$M_5^{(1)}(1^+, 2^+, 3^+, 4^+, 5^+) = \left(\sum_{P_6} T^A_{(1,2,3),(4,5)} + \sum_{P_3} T^B_{(1,2,3),4,5} \right) \quad (14)$$

where

$$T^A_{(1,2,3),(4,5)} = 10 \frac{[14] [53] [52] [23]^2}{\langle 14 \rangle \langle 1\eta \rangle^2 \langle 4\eta \rangle \langle 45 \rangle} \times [5|K_{14}|\eta\rangle [2|K_{14}|\eta\rangle [3|K_{14}|\eta\rangle], \quad (15)$$

$$T^B_{(1,2,3),4,5} = -10 \frac{[14] [15] [23] [1|K_{23}|\eta\rangle^2 [5|K_{23}|\eta\rangle [4|K_{23}|\eta\rangle [45]}{\langle 23 \rangle \langle 2\eta \rangle^2 \langle 3\eta \rangle^2 \langle 45 \rangle} \quad (16)$$

and P_3 denotes summation over the three cyclic permutations of legs 1,2 and 3. P_6 denotes the three permutations of P_3 together with interchange of legs 4 and 5. The λ_η independence of $M_5^{(1)}(1^+, 2^+, 3^+, 4^+, 5^+)$ is not manifest.




Figure 4: Factorisations of the five-point single minus amplitude

The factorisations of the five-point single minus amplitudes are more varied as shown on fig. 4. Using the BCFW shift on $(\bar{\lambda}_1, \lambda_2)$ we obtain the amplitude

$$\begin{aligned}
M_5^{(1)}(1^-, 2^+, 3^+, 4^+, 5^+) &= \frac{10}{[12]^2} \left(\prod_{i,j=2,3,4,5, i<j} [ij] \right) \left(\frac{\langle 15 \rangle [25]^3}{[15] \langle 34 \rangle} + \frac{\langle 13 \rangle [23]^3}{[13] \langle 45 \rangle} + \frac{\langle 14 \rangle [24]^3}{[14] \langle 53 \rangle} \right) \\
&+ \frac{\langle 12 \rangle^2}{\langle 34 \rangle \langle 35 \rangle \langle 45 \rangle \prod_{i=3,4,5} \langle 1i \rangle} \left(\frac{[23]^5 [45] \langle 13 \rangle^5}{\langle 23 \rangle} + \frac{[24]^5 [53] \langle 14 \rangle^5}{\langle 24 \rangle} + \frac{[25]^5 [35] \langle 15 \rangle^5}{\langle 25 \rangle} \right) \\
&+ \frac{1}{\langle 12 \rangle^2 \langle 34 \rangle \langle 35 \rangle \langle 45 \rangle} \left(\frac{[23] [45]^5 \langle 15 \rangle^3 \langle 14 \rangle^3}{\langle 23 \rangle} + \frac{[24] [53]^5 \langle 13 \rangle^3 \langle 15 \rangle^3}{\langle 24 \rangle} \right. \\
&+ \left. \frac{[25] [34]^5 \langle 14 \rangle^3 \langle 13 \rangle^3}{\langle 25 \rangle} \right). \tag{17}
\end{aligned}$$

The five-point MHV amplitude is non-zero. The non-zero factorisations of the amplitude are shown in fig. 5.




Figure 5: Factorisations of the five-point MHV amplitude

This amplitude can be obtained using a BCFW shift of either the two negative helicity legs or of a negative-positive pair. Shifting the two negative legs generates the expression (using only the second factorisation of fig. 5),

$$M_5^{(1)}(1^-, 2^-, 3^+, 4^+, 5^+) = -s_{34} \frac{\langle 15 \rangle [34]^2 [35]^3 [45]^3}{[15] [12]^2 [23] [24]}$$

$$-s_{45} \frac{\langle 13 \rangle [45]^2 [43]^3 [53]^3}{[13] [12]^2 [24] [25]} - s_{53} \frac{\langle 14 \rangle [53]^2 [54]^3 [34]^3}{[14] [12]^2 [25] [23]} . \quad (18)$$

This completes the set of five-point amplitudes. We can continue in this way generating the tree-level S -matrix. We have made available $M_n^{(1)}$ for $n \leq 7$ in `Mathematica` format at <http://pyweb.swan.ac.uk/~dunbar/Smatrix.html>. The amplitudes have been generated up to $n = 8$ and have the correct symmetries, are η -independent and have the correct leading soft-limits.

We have evaluated amplitudes in a $R + \alpha R^3$ theory. In ref. [21] amplitudes in Yang-Mills theory extended by F^3 terms were studied. Then using double copy techniques and the KLT relations [22] graviton scattering amplitudes were derived upto $n = 6$. As noted in [21] these correspond to amplitudes in a $R + \alpha R^3 + \sqrt{\alpha} R^2 \phi$ theory. The four-point amplitudes in the two theories are proportional [17, 21] but beyond four-point the two sets of amplitudes are functionally different. The all-plus amplitude in the two theories remain proportional for $n > 4$ with

$$M_n^{(1), R^3 + R^2 \phi}(1^+, 2^+, \dots, n^+) = \frac{5}{2} M_n^{(1), R^3}(1^+, 2^+, \dots, n^+) \quad (19)$$

and we confirm this for $n \leq 7$.

3. Soft Limits

Graviton scattering amplitudes are singular as a leg becomes soft. Weinberg [23] many years ago presented the leading soft limit. If we parametrise the momentum of the n -th leg as $k_n^\mu = t \times k_s^\mu$ then in the limit $t \rightarrow 0$ the singularity in the n -point amplitude is

$$M_n \rightarrow \frac{1}{t} \times S^{(0)} \times M_{n-1} + O(t^0) \quad (20)$$

where M_{n-1} is the $n-1$ -point amplitude. The soft-factor $S^{(0)}$ is universal and Weinberg showed that (20) does not receive corrections in loop amplitudes.

Recently it has also been proposed [24, 25, 26] that the sub-leading and sub-sub-leading terms are also universal. This can be best exposed, when a positive helicity leg becomes soft, by setting

$$\lambda_n = t \times \lambda_s , \quad \bar{\lambda}_n = \bar{\lambda}_s . \quad (21)$$

In the $t \rightarrow 0$ limit the amplitude has t^{-3} singularities. At tree level the amplitudes satisfy soft-theorems [25] whereby their behaviour as $t \rightarrow 0$ is

$$M_n^{\text{tree}} = S_t M_{n-1}^{\text{tree}} + O(t^0) = \left(\frac{1}{t^3} S^{(0)} + \frac{1}{t^2} S^{(1)} + \frac{1}{t} S^{(2)} \right) M_{n-1}^{\text{tree}} + O(t^0) \quad (22)$$

where, for a positive helicity-leg becoming soft [25, 27, 28]

$$S^{(0)} = - \sum_{i=1}^{n-1} \frac{[s i] \langle i \alpha \rangle \langle i \beta \rangle}{\langle s i \rangle \langle s \alpha \rangle \langle s \beta \rangle}, \quad (23)$$

$$S^{(1)} = - \frac{1}{2} \sum_{i=1}^{n-1} \frac{[s i]}{\langle s i \rangle} \left(\frac{\langle i \alpha \rangle}{\langle s \alpha \rangle} + \frac{\langle i \beta \rangle}{\langle s \beta \rangle} \right) \bar{\lambda}_s^{\dot{a}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{a}}}, \quad (24)$$

$$S^{(2)} = \frac{1}{2} \sum_{i=1}^{n-1} \frac{[i s]}{\langle i s \rangle} \bar{\lambda}_s^{\dot{a}} \bar{\lambda}_s^{\dot{b}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{a}}} \frac{\partial}{\partial \bar{\lambda}_i^{\dot{b}}}. \quad (25)$$

The proof of the soft theorems follows from Ward identities of extended Bondi, van der Burg, Metzner and Sachs (BMS) symmetry [29]. Although exact for tree level amplitudes these receive loop corrections [27, 30, 31].

Whether the soft theorems extend beyond Einstein gravity has been examined before. In particular the leading soft behaviour can often be used as a check upon amplitudes such, e.g. in [21]. The leading and sub-leading limits were shown to hold for a R^3 insertion in [32]. Here we examine the amplitudes and, in particular, test the sub-sub-leading soft behaviour.

We can summarise the behaviour of the leading amplitudes, $M_n^{(1)}$, simply by stating:

**All the amplitudes calculated satisfy the soft limits of
(22) up to and including the sub-sub-leading term.**

We have verified this for all helicity amplitudes up to $n = 8$. Note: to check (22) one must implement momentum conservation consistently between the n -point amplitudes and the $n - 1$ -point amplitudes which in essence specifies how the point $t = 0$ is approached. These are several ways to do this. We have followed the prescription of [25] but alternative implementations are possible [27, 28].

In principle we could have found a behaviour of the form

$$M_n^{(1)} \rightarrow S_t M_{n-1}^{(1)} + S_t^\alpha M_{n-1}^{(0)} + R_n \quad (26)$$

where S_t^α would be an α correction to the soft functions and R_n is a non-factorising term. In terms of this we find $S_t^\alpha = R_n = 0$. Since the theory we are considering is higher derivative it is not surprising that the leading and sub-leading parts of S_t^α vanish however it is interesting that the vanishing continues for the sub-sub-leading - unlike the loop corrections to Einstein gravity.

Incidentally as a consequence of eq.(19) the amplitude $M_n^{(1),R^3+R^2\phi}(1^+, 2^+, \dots n^+)$ also satisfies the soft theorems to sub-sub leading level.

4. Other Theories

We have chosen to extend gravity using a three-point vertex and use a diagrammatic approach whereby we only consider the on-shell amplitudes. There is, of course, complementarity between this approach and that of Lagrangian based field theory. The single choice of three-point amplitude corresponds to the single R^3 field density that affects on-shell amplitudes. This makes the extended S -matrix simply depend upon the single parameter α .

If we were to deform Einstein gravity by an additional four-point amplitude then there are more choices consistent with symmetry and spinor weight, e.g. we could have

$$M_4(1^+, 2^+, 3^+, 4^+) = \alpha_1(\langle 1 2 \rangle^4 \langle 3 4 \rangle^4 + \langle 1 3 \rangle^4 \langle 2 4 \rangle^4 + \langle 1 4 \rangle^4 \langle 2 3 \rangle^4) + \alpha_2(\langle 1 2 \rangle \langle 2 3 \rangle \langle 3 4 \rangle \langle 4 1 \rangle + \text{permutations})^2 + \dots \quad (27)$$

From a field theory perspective this freedom corresponds to the observation that there are multiple R^4 tensors that contribute to on-shell amplitudes [33].

The same issue arises when we consider the further expansion in α . If we consider $M_4^{(2)}(1^-, 2^-, 3^+, 4^+)$ there is a single factorisation as shown in fig. 6. The amplitude

$$M_4^{(2)}(1^-, 2^-, 3^+, 4^+) = \langle 1 2 \rangle^4 [3 4]^4 \left(\frac{tu + \beta s^2}{s} \right) \quad (28)$$

has the correct factorisation for any choice of β . This ambiguity means we also have to specify the four-point amplitude to determine the S -matrix. In the diagrammatic approach this ambiguity arises due to the existence of a polynomial function with the correct symmetries and spinor and momentum weight. From a field theory perspective, additional counterterms can contribute to this amplitude. Specifically, we could deform the theory via

$$R \longrightarrow R + C_\alpha R^3 + C_\beta D^2 R^4 \quad (29)$$

and the four-point amplitude is only specified once C_α and C_β are determined.




Figure 6: Factorisations of the four-point MHV amplitude at α^2 .

5. Conclusion

We have constructed the (leading part) of the S -matrix of an extended theory of gravity starting from three-point amplitudes and only demanding factorisation. The theory is extended by the addition of amplitudes which are polynomial in momentum, thus implicitly imposing locality and unitarity on the S -matrix. We also require the amplitudes to have the correct spinor helicity as appropriate for massless particles. The S -matrix is then generated entirely from on-shell amplitudes by demanding factorisation. Specifically, we have extended the theory by the addition of three-point amplitudes which, from a field theory perspective, corresponds to introducing R^3 terms. This S -matrix differs from that obtained by applying double copy or KLT techniques to a F^3 extension of Yang-Mills.

Beyond the leading part, polynomial amplitudes exist at higher point and these must be specified to fully determine the S -matrix. Consistency of this approach and a field theoretic approach beyond leading order requires a correspondence between these polynomial amplitudes and the counter terms contributing to on-shell amplitudes.

We find that these amplitudes satisfy the same soft theorems as the tree amplitudes of Einstein gravity up to and including the sub-sub leading terms. It is interesting that these theorems are robust to deformations of Einstein gravity even at the sub-sub-leading level particularly given the link to BMS symmetry which plays an important role in the recent understanding of black hole soft hair [34].

6. Acknowledgements

This work was supported by STFC grant ST/L000369/1. GRJ was supported by STFC grant ST/M503848/1. JHG was supported by the College of Science (CoS) Doctoral Training Centre (DTC) at Swansea University.

- [1] G. 't Hooft and M. J. G. Veltman, NATO Sci. Ser. B **4** (1974) 177.
- [2] R.J. Eden, P.V. Landshoff, D.I. Olive, J.C. Polkinghorne, *The Analytic S Matrix*, (Cambridge University Press, 1966).
- [3] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **425** (1994) 217 doi:10.1016/0550-3213(94)90179-1 [hep-ph/9403226].
- [4] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **435** (1995) 59 doi:10.1016/0550-3213(94)00488-Z [hep-ph/9409265].
- [5] R. Britto, F. Cachazo, B. Feng and E. Witten, Phys. Rev. Lett. **94** (2005) 181602 [hep-th/0501052].
- [6] N. Arkani-Hamed and J. Trnka, JHEP **1410** (2014) 030 doi:10.1007/JHEP10(2014)030 [arXiv:1312.2007 [hep-th]].
- [7] N. Arkani-Hamed, L. Rodina and J. Trnka, arXiv:1612.02797 [hep-th].
- [8] J. Bedford, A. Brandhuber, B. J. Spence and G. Travaglini, Nucl. Phys. B **721** (2005) 98 doi:10.1016/j.nuclphysb.2005.05.016, 10.1016/j.nuclphysb.2005.016 [hep-th/0502146].
- [9] F. Cachazo and P. Svrcek, hep-th/0502160.
- [10] N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins and K. Risager, JHEP **0601** (2006) 009 doi:10.1088/1126-6708/2006/01/009 [hep-th/0509016].
- [11] P. Benincasa and F. Cachazo, arXiv:0705.4305 [hep-th].
- [12] P. Benincasa, C. Boucher-Veronneau and F. Cachazo, JHEP **0711** (2007) 057 doi:10.1088/1126-6708/2007/11/057 [hep-th/0702032 [HEP-TH]].

- [13] K. Risager, JHEP **0512** (2005) 003 doi:10.1088/1126-6708/2005/12/003 [hep-th/0508206].
evanNieuwenhuizen:1976vb
- [14] P. van Nieuwenhuizen and C. C. Wu, J. Math. Phys. **18** (1977) 182. doi:10.1063/1.523128
- [15] M. T. Grisaru, Phys. Lett. **66B** (1977) 75. doi:10.1016/0370-2693(77)90617-7
- [16] E. Tomboulis, Phys. Lett. **67B** (1977) 417. doi:10.1016/0370-2693(77)90434-8
- [17] T. Cohen, H. Elvang and M. Kiermaier, JHEP **1104** (2011) 053 doi:10.1007/JHEP04(2011)053 [arXiv:1010.0257 [hep-th]].
- [18] D. C. Dunbar, G. R. Jehu and W. B. Perkins, arXiv:1701.02934 [hep-th].
- [19] Z. Bern, H. H. Chi, L. Dixon and A. Edison, arXiv:1701.02422 [hep-th].
- [20] D. C. Dunbar and N. W. P. Turner, Class. Quant. Grav. **20** (2003) 2293 doi:10.1088/0264-9381/20/11/323 [hep-th/0212160].
- [21] J. Broedel and L. J. Dixon, JHEP **1210** (2012) 091 doi:10.1007/JHEP10(2012)091 [arXiv:1208.0876 [hep-th]].
- [22] H. Kawai, D. C. Lewellen and S. H. H. Tye, Nucl. Phys. B **269** (1986) 1. doi:10.1016/0550-3213(86)90362-7
- [23] Steven Weinberg, Phys.Rev. **140** (1965) B516-B524.
- [24] C. D. White, JHEP **1105** (2011) 060 [arXiv:1103.2981 [hep-th]].
- [25] F. Cachazo and A. Strominger, arXiv:1404.4091 [hep-th].
- [26] T. He, V. Lysov, P. Mitra and A. Strominger, JHEP **1505** (2015) 151 doi:10.1007/JHEP05(2015)151 [arXiv:1401.7026 [hep-th]].
- [27] Z. Bern, S. Davies and J. Nohle, Phys. Rev. D **90** (2014) 8, 085015 [arXiv:1405.1015 [hep-th]].
- [28] J. Broedel, M. de Leeuw, J. Plefka and M. Rosso, Phys. Rev. D **90** (2014) 6, 065024 [arXiv:1406.6574 [hep-th]].

- [29] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, Proc. Roy. Soc. Lond. A **269** (1962) 21;
R. K. Sachs, Proc. Roy. Soc. Lond. A **270** (1962) 103.
- [30] S. He, Y. t. Huang and C. Wen, JHEP **1412** (2014) 115 [arXiv:1405.1410 [hep-th]].
- [31] S. D. Alston, D. C. Dunbar and W. B. Perkins, Phys. Rev. D **86** (2012) 085022 [arXiv:1208.0190 [hep-th]].
- [32] M. Bianchi, S. He, Y. t. Huang and C. Wen, Phys. Rev. D **92** (2015) no.6, 065022 doi:10.1103/PhysRevD.92.065022 [arXiv:1406.5155 [hep-th]].
- [33] S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins, Class. Quant. Grav. **9** (1992) 1151. doi:10.1088/0264-9381/9/5/003
- [34] S. W. Hawking, M. J. Perry and A. Strominger, Phys. Rev. Lett. **116** (2016) no.23, 231301 doi:10.1103/PhysRevLett.116.231301 [arXiv:1601.00921 [hep-th]].