

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Effects of golf courses on local biodiversity

R. A. Tanner, A.C. Gange

*School of Biological Sciences, Royal Holloway, University of London,
Egham, Surrey UK TW20 0EX*

Author for correspondence:

A.C. Gange, address above

Tel: + 44 (0)1784 443188

Fax: + 44 (0)1784 470756

e-mail: a.gange@rhul.ac.uk

29 **Abstract**

30

31 There are approximately 2,600 golf courses in the UK, occupying 0.7% of the total
32 land cover. However, it is unknown whether these represent a significant resource, in
33 terms of biodiversity conservation, or if they are significantly less diverse than the
34 surrounding habitats.

35 The diversity of vegetation (tree and herbaceous species) and three indicator taxa
36 (birds, ground beetles (Coleoptera, Carabidae) and bumblebees (Hymenoptera,
37 Apidae)) was studied on nine golf courses and nine adjacent habitats (from which the
38 golf course had been created) in Surrey, UK. Two main objectives were addressed:
39 (1) To determine if golf courses support a higher diversity of organisms than the
40 farmland they frequently replace and (2) to examine whether biodiversity increases
41 with the age of the golf course.

42 Birds and both insect taxa showed higher species richness and higher abundance on
43 the golf course habitat than in nearby farmland. While there was no difference in the
44 diversity of herbaceous plant species, courses supported a greater diversity of tree
45 species. Furthermore, bird diversity showed a positive relation with tree diversity for
46 each habitat type. It was found that introduced tree species were more abundant on
47 the older golf courses, showing that attitudes to nature conservation on courses have
48 changed over time. Although the courses studied differed in age by up to 90 years,
49 the age of the course had no effect on diversity, abundance or species richness for any
50 of the animal taxa sampled. We conclude that golf courses of any age can enhance
51 the local biodiversity of an area by providing a greater variety of habitats than
52 intensively managed agricultural areas.

53

54 *Keywords:* Golf courses: Conservation: Carabid beetles: Birds: Bumblebees:
55 Biodiversity.

56

57 **1. Introduction**

58

59 Over the last 30 years, household expenditure on recreation has increased
60 substantially in the UK, for example in 1996 an estimated £9 billion was spent on day
61 trips to the countryside. Recreation and leisure activities do not always pose a
62 significant problem to the environment (Coppock and Duffield, 1975), though impacts

63 on wildlife (Chettri et al., 2001) and habitats (Boyle and Samson, 1985) have been
64 reported. Furthermore, the effect of transport (Cincotta et al., 2000), noise (Mikola et
65 al., 1994) and pollution (Sun and Walsh, 1998) are all concerns expressed by
66 governmental bodies. Activities including hill walking (Riffell et al., 1996), power
67 boating (Bell, 2000), wildlife-photography and skiing (Burger, 2000) have all been
68 shown to disturb wildlife and habitats.

69 Few of the aforementioned activities have such an intimate interaction with the
70 environment as golf. The game has seen a tremendous increase in popularity over the
71 last 100 years and there are now over 2,600 golf courses in the UK and over 31,500
72 worldwide. Golf course establishment has increased by over 42% in the last 30 years
73 (Daniels, 1972) and currently, the UK holds over half of all golf courses found in
74 Europe (Anon., 1996). Annual participation in the game increased by 18% between
75 1987 and 1996 and at that time an estimated 12% of the population played golf over a
76 12-month period, (Anon., 1996).

77 The demand on Britain's 22 million hectares from the growing human population
78 and the demand for golf courses has led to changes in course design and management.
79 Following the traditional style of links courses, established in Scotland during the 14th
80 century, the beginning of the 16th century saw the introduction of open inland courses
81 dominated by heathland habitat (Anon., 1989). It was not until the 19th century that a
82 new style of golf course was seen to evolve from the landscape garden designs of
83 Lancelot 'Capability' Brown. These parkland golf courses were different from
84 previous, in that shelter in the form of trees and bushes surrounded the course and
85 patches of woodland were found throughout. By 1972, over 54% of golf courses in
86 the UK were parkland courses (Dair and Schofield, 1990).

87 There has been increasing concern about the magnitude of global biodiversity loss
88 (Gaston, 1996). In the UK, biodiversity is highly concentrated in the south east of the
89 country, as is much of the human population and the majority of golf courses (Gaston,
90 1996; Lennon et al., 2000; Beebee, 2001). Habitat modification (Terman, 1997),
91 chemical contamination (Murphy and Aucott, 1998), water management (Cohen et al.,
92 1993) and urbanisation around golf courses (Markwick, 2000) are all concerns that
93 have been expressed by those who claim that courses are a poor use, ecologically
94 speaking, of land (Platt, 1994). However, until recently, there was little evidence to
95 support the view that golf courses are good or bad for the environment at a landscape
96 scale. What little information there is suggests that golf courses are not significant

97 sources of water pollution (Cohen et al., 1999) and may be the equal of many natural
98 habitats in terms of animal and plant diversity (Terman, 1997; Gange and Lindsay,
99 2002).

100 To date, there is only a handful of research studies that have employed a strict
101 scientific method to the study of wildlife on golf courses with most focusing on links
102 courses (Green and Marshall, 1986; Blair, 1996; Terman, 1997, 2000). All studies
103 have shown that golf courses compare well in terms of wildlife abundance and
104 diversity to that of adjacent areas of land. A feature of these studies (e.g. Blair, 1996,
105 Terman, 1997) is that the diversity of taxa on golf courses has been compared with
106 areas of pristine natural habitat. As shown by Gange and Lindsay (2002), a more
107 realistic question to ask, in terms of landscape ecology, is how the biological diversity
108 of a golf course compares with that of the habitat from which the course was
109 constructed. Gange and Lindsay (2002) present four simple case studies, where in
110 each instance it was found that the diversity of insects and birds on a golf course was
111 higher than that of the surrounding agricultural land. However, this was a short term
112 study of about 11 weeks and only two courses studied were in the UK.
113 Approximately 60% of the UK is arable and pasture farmland (equally divided
114 between the two), forms of land use known to be of low ecological value and shown
115 to degrade biodiversity (Altieri, 1999; Chamberlain et al., 2000). As land targeted for
116 golf development in the last 20 years has been almost exclusively farmland, it is the
117 aim of this paper to extend the studies of Gange and Lindsay (2002), in terms of
118 duration and replicate number. We sought to determine whether golf courses harbour
119 different levels of biodiversity than the habitats they replace and whether abundance
120 and species richness of certain animal taxa differ between old and young courses.
121 These studies are important, because it is well known that effective course
122 management lies in the understanding of the natural processes, which operate within
123 the course (Brennan, 1992).

124

125 **2. Materials and Methods**

126

127 *2.1. Sites and taxa studied*

128

129 All sites used in this study are located in the county of Surrey, UK and all golf
130 courses were of the parkland design. It has been suggested that the age of a golf

131 course is an important factor in its wildlife value (Dair and Schofield, 1990), and so
132 courses were selected that fell into one of three age groups, with three replicates in
133 each group. These groups were 1-10-yrs, 20-30-yrs and 90-yrs plus. Nine golf
134 courses and nine adjacent farmland areas were sampled in total. All adjacent areas of
135 land were within 0.5 km of the golf course and all consisted of pasture grassland used
136 for cattle or sheep grazing. The adjacent areas were demarcated by the farm boundary
137 and chosen to reflect the land use that was in existence before a course was created, or
138 what the land would support if it were not a golf course. Hereafter, each course or
139 adjacent area is termed a 'site', thus there were 18 sites in total. Aerial photographs
140 of each site were obtained and the total area of each was calculated.

141 When measuring diversity, a complete inventory for all species is impossible due
142 to time and effort. We chose well-known indicator species (Kremen, 1992; Pearson,
143 1996; Simberloff, 1998) that were relatively easy to observe and identify. Vegetation
144 was sampled, as this is the habitat template and dominance and diversity in plant
145 communities dictate the composition and diversity of animal species (Southwood et
146 al., 1983). Birds (Furness and Greenwood, 1993; Gregory and Baillie, 1998), ground
147 beetles (Coleoptera, Carabidae (Butterfield et al., 1995)) and bumblebees
148 (Hymenoptera, Apidae (Saville et al., 1997; Carvell, 2002)) have repeatedly been used
149 as indicator species and were the taxa chosen for this study.

150

151 *2.2. Recording techniques*

152

153 Eleven bird censuses were conducted, approximately one every three weeks at each
154 site, between 12th November 2001 and 30th June 2002. The Variable Circular Plot
155 (VCP) method was used (Reynolds et al., 1980), which is a form of distance sampling
156 developed from line transects (Buckland et al., 1993). The method does not assume
157 all individuals present are recorded and the observer can miss up to 50% of
158 individuals and still obtain reliable density (Bibby et al., 2000). The VCP method is
159 ideal for bird sampling when habitats within areas are patchy and has proved to be a
160 powerful reliable estimator of bird density for a range of different species (Buckland
161 et al., 1993; Fancy, 1997; Nelson and Fancy, 1999). Birds were only recorded if they
162 were seen utilising the site, i.e. perching, feeding or nesting. We did not record birds
163 by their song alone; because we were not confident of our ability to use sound
164 reliably. This was a deliberate decision, because we wanted to eliminate any possible

165 incidental use of a site (e.g. flying over) and to only record direct utilization (defined
166 above). Our bird estimates are therefore conservative, but as unbiased as possible.

167 A square grid drawn to scale and consisting of squares totalling 100 m x 100 m
168 was placed on each aerial photograph. Using this grid, 16 points were randomly
169 selected in each site, each being greater than 200 m apart. The observer stood at each
170 of the 16 points for five minutes and counted all the birds visible. Approaching each
171 point carefully and moving vigilantly between points avoided disturbing any birds.
172 Using reference points (trees, shrubs, fences, etc.) from the aerial photographs, the
173 distance each individual bird was from the point was recorded to the nearest metre.
174 One golf course and one adjacent site were sampled each day with no censuses
175 conducted in high winds or heavy rain. Sampling took place between the hours of
176 0600 and 1030 when bird activity is greatest (Bibby et al., 2000).

177 The program DISTANCE 3.5 (Thomas et al., 1998) was used to calculate bird
178 density. The program fits field data to a selection of different models (key functions)
179 using series expansions to fine-tune the fit. The data were ungrouped and in cases
180 where the model fit was weak, the data were truncated at varying lengths and
181 percentages, as recommended by Buckland et al. (1993). Means of the 16 data points
182 for density, species richness and diversity in each site on each date were calculated to
183 provide overall site values for analysis.

184 All invertebrate sampling was conducted for two months from May 1st to June 30th
185 2002. Pitfall traps, consisting of plastic containers 10 cm deep and 5 cm in diameter
186 filled with 30 ml of ethylene glycol as a preservative, were sunk into the ground.
187 These are the most commonly used and highly effective traps for catching ground
188 beetles (Greenslade, 1964; Southwood and Henderson, 2000). Using aerial
189 photographs, 20 traps were randomly placed throughout each site. Samples were
190 collected and stored every ten days, and identification of species was performed with
191 reference to Lindroth (1974) and Forsythe (2000). Means of the 20 data points for
192 density, species richness and diversity in each site on each date were calculated to
193 provide overall site values for analysis.

194 Line walking is the most frequently cited method for bumblebee censusing and was
195 the method adopted in this study (Saville et al., 1997; Walther-Hellwig and Frankl,
196 2000). Surveys were conducted between midday and 1500 and consisted of four x
197 100 m line transects, randomly located within each site using aerial photographs.
198 Each site was surveyed 15 times between May 1st and June 30th 2002. Every

199 bumblebee seen whilst walking was either identified on the wing or captured with a
200 net, identified, recorded and released. Recording was only conducted on clear bright
201 days, of low winds. Means of the four transects points for density, species richness
202 and diversity in each site on each date were calculated to provide overall site values
203 for analysis.

204 Vegetation sampling was divided into two categories (1) trees (sampled in
205 November 2001) and (2) herbaceous species (sampled from June 1st to July 18th
206 2002). Using aerial photographs, six 50 m x 50 m quadrats for tree sampling and
207 twenty 5 m x 5 m quadrats (hereafter termed 'plots') for herbaceous plants were
208 selected in each site. Tree quadrats were randomly placed, while herbaceous quadrats
209 conformed to stratified random samples, by the avoidance of heavily wooded areas or
210 the actual pasture, or greens, tees and fairways. Within each quadrat, total tree
211 abundance for each species present was recorded. Herbaceous species were sampled
212 using a 38 cm linear steel frame, containing ten 3 mm diameter point quadrat pins.
213 The frame was placed randomly 20 times in each plot, giving a total of 200 pins
214 sampled per plot. The number of touches of all living plant material was recorded in
215 2 cm (below 10 cm) or 5 cm (10 cm and above) height intervals on each pin. Data for
216 the 200 pins were summed and means calculated of diversity, height of vegetation and
217 species richness (Brown and Gange 1989). Values for each plot were then averaged
218 to provide site means for analysis.

219

220

221 2.3. *Statistical analysis*

222

223 All data on species richness, abundance, and diversity were analysed using site
224 means as replicates. Bird species were categorised by feeding type (1) insect feeders,
225 (2) other carnivores, (3) seedeaters and (4) omnivorous species. The Shannon-Wiener
226 diversity index (H) (Magurran, 1988) was used to estimate diversity for all taxa,
227 except for herbaceous species where Williams Alpha diversity (Southwood and
228 Henderson, 2000) was used. Data was tested for normality and homogeneity of
229 variance and where appropriate square root transformations were made. Zero values
230 were rare and did not compromise any of the analyses. A repeated measures Analysis
231 of Variance, using date and site as the main effects was performed on diversity,
232 abundance and species richness for each organism group. Meanwhile, single factor

233 Analysis of Variance was used to examine whether course age had an effect on
234 density and diversity of each group.

235

236 **3. Results**

237

238 *3.1. Vegetation*

239

240 Tree diversity was higher on the golf course habitats than the adjacent land sites
241 ($F_{1,16}=6.42$, $P < 0.05$), with a mean of 10.4 species per 2,500 m² being found in golf
242 course habitats compared to 7.4 species per 2,500 m² on the adjacent lands. The
243 proportion of native trees in the landscape differed between course types ($\chi^2 = 0.75$, P
244 < 0.01), with oldest courses having significantly fewer natives (74.1%) than middle
245 aged (81.8%) or young courses. The proportion of native trees on youngest courses
246 (84.7%) was lower, but not significantly so, compared with that of the surrounding
247 farmland (91.9%). No differences were found in the herbaceous vegetation (diversity,
248 species richness or height) between the two habitats.

249

250

251 *3.2. Bird species*

252

253 Bird diversity was significantly higher on the golf courses than the adjacent areas
254 of land ($F_{1,16} = 7.67$, $P < 0.05$; Fig. 1a). A significant interaction term between site
255 and date was found in the analysis ($F_{10,160} = 1.94$, $P < 0.05$), because the two habitats
256 did not show a similar pattern of change through the season. The golf course habitats
257 had higher species richness than the adjacent sites ($F_{1,16} = 13.92$, $P < 0.05$), with an
258 average of 13 bird species seen on each sample date, compared to 11 species on each
259 date in the adjacent sites.

260 There was no difference in the density of birds between the habitat types, and
261 neither was there any significant change in bird abundance over time (Fig. 1b).
262 However, there was a highly significant association between bird species diet and
263 habitat type ($\chi^2 = 19.36$, $P < 0.01$). Higher proportions of insect feeding birds (28%)
264 were found on the golf course habitats compared to the adjacent land types (19%).
265 Meanwhile, omnivorous species (e.g. the Rook (*Corvus frugilegus*) and Magpie (*Pica*

266 *pica*) were found in higher proportions within the adjacent sites (56%) than the golf
267 course habitat (46%). The age of the golf course had no effect on bird density,
268 diversity or species richness.

269 A significant relationship was found between bird diversity and tree diversity in
270 each habitat type (Fig. 2). Of most interest was the fact that the slopes of the
271 regression lines for each habitat type were significantly different ($t = 2.29$; $df = 14$; P
272 $= < 0.05$), indicating that for any given value of tree diversity, bird diversity was
273 higher on the golf courses than the adjacent land sites. However, the lines appeared to
274 converge, such that at high tree diversity, one might predict no difference between the
275 courses and adjacent areas.

276

277 3.3. *Carabid species*

278

279 There was some evidence that beetle diversity differed between the two habitat
280 types ($F_{1,16} = 4.21$, $P = 0.057$, Fig. 3a). However, numbers of beetle individuals
281 captured were much higher on the golf courses than the adjacent sites ($F_{1,16}=20.40$,
282 $P<0.001$, Fig. 3b) and an average of 8.4 different species were found on each date on
283 the golf courses, compared with 6.5 species on the adjacent sites ($F_{1,16}=6.59$, $P<0.05$).
284 There was no significant interaction term between site and date for any of the beetle
285 data, indicating that beetles followed similar temporal patterns in the different areas.
286 The age of the golf course had no effect on beetle abundance, diversity or species
287 richness.

288

289 3.4. *Bumblebee species*

290

291 There was no difference in diversity (Shannon Weiner H), of bumblebees between
292 golf course habitats and adjacent sites (Fig 4a). However, bumblebees showed a
293 highly significant difference in abundance and species richness per 100 m when
294 comparing the two habitat types. The golf courses had higher abundance ($F_{1,16} =$
295 19.41 , $P < 0.001$) and higher species richness ($F_{1,16} = 24.41$, $P < 0.001$) than the
296 adjacent farmland. An average of 6 species per transect were found on the courses,
297 compared with 3 species per transect at the adjacent sites. Both bumblebee diversity
298 ($F_{15,240} = 1.94$, $P < 0.05$) and abundance ($F_{15,240} = 2.12$, $P < 0.05$; Fig. 4b) showed a

299 significant interaction term between site and date. This was because for both
300 variables, values on the adjacent land stayed relatively constant through time, whereas
301 the course showed fluctuating values. In the case of diversity, values for the course
302 were higher early in the season, but lower in late season, thereby contributing to the
303 fact that there was no overall effect of site in the ANOVA (above). The age of the
304 golf course had no effect on diversity, abundance or species richness.

305

306 **4. Discussion**

307

308 These results show that, for the taxa studied, golf courses can contain levels of
309 biodiversity equal to or above that of the habitats they replace. Gange and Lindsay
310 (2002) discuss how enhancing biodiversity is about conserving species local to an
311 area, not just increasing numbers. Every species has specific habitat preferences and
312 green keepers can contribute greatly to conservation by providing such habitats for
313 endangered local species. We suggest that the variety of habitats that a golf course
314 provides is potentially greater than that of farmland, thus enabling a greater diversity
315 of species to exist. By increasing habitat heterogeneity within a landscape, golf
316 courses can enhance the diversity of a local area.

317 The age of the golf course had no effect on diversity for any of the taxa studied.
318 This was surprising, because one might think that over time a greater variety of
319 habitats on a golf course would become established, thereby enhancing biodiversity.
320 One possible explanation lies in the identity of the vegetation in the different sites.
321 Older courses were found to harbour a greater amount of introduced tree species,
322 many of which were planted for their aesthetic, rather than ecological value.
323 Introduced tree species provide poorer habitats for birds than native trees (Fuller,
324 1997) and they can affect biodiversity by changing the composition, structure and
325 community pattern of an ecosystem (Peterken, 2001). Although the diversity of
326 native trees was often lower on the golf courses, we found that for any given value of
327 tree diversity, bird diversity was higher on the golf courses than the adjacent land
328 sites, with each habitat displaying a different temporal change through time. These
329 results are consistent with other studies (Blair, 1996; Terman, 1997; Gange and
330 Lindsay, 2002). It is known that mass planting of introduced species in plantations,
331 like conifer forests, does reduce bird diversity (Fuller, 1997), but in the case of golf

332 courses, bird diversity could be reacting to the stand diversification produced by the
333 array of exotic and native species rather than individual introduced species.

334 It should be noted that the regression for golf courses is clearly dependent on one
335 datum, that of the lowest value for tree diversity, suggesting that further work needs to
336 be done to assess the validity of the relation. However, an important point is that if
337 the slope of the regression for golf courses was close to zero, this would imply that
338 bird diversity was high, irrespective of tree diversity. Such a result suggests that other
339 habitats on the golf courses are very important in affecting the diversity of birds that
340 inhabit the area. Furthermore, we found that the regression lines tended to converge,
341 suggesting that at high tree diversity, farmland would be the equal of the golf course
342 in terms of bird diversity and could even exceed it. Extrapolation of regression lines
343 is dangerous and only further research can confirm or refute this hypothesis.

344 A second explanation for the lack of course age effects is the mobility of the
345 groups we studied. Birds, ground beetles and bumblebees are all highly mobile
346 creatures and all of them would have no difficulty colonizing new golf course
347 developments. For taxa that are less mobile or slow to disperse, course age may well
348 affect their occurrence, and again this highlights the need for further research in this
349 area.

350 A final point regarding the lack of course age effects concerns the dietary
351 requirements of the organisms studied. Although different bird feeding types were
352 found between the sites (below), none of the birds, beetles or bumblebees found could
353 be considered as extreme dietary specialists. Even species common on the courses,
354 (but not recorded on the farmland) such as the green woodpecker (*Picus viridis*)
355 (which feeds on ants) or the song thrush (*Turdus philomelos*) (which prefers snails)
356 are just as likely to find food on a 5 y as they are on a 100 y old course. However, the
357 abundance of this food may change with time (e.g. one would expect ant colonies to
358 increase with course age) as will the structure of the habitat in which it is found.
359 Future research should take into account the degree of specialism of the taxa studied,
360 in order to determine whether older courses harbour greater numbers of specialist
361 species and whether this is related to food or habitat availability. Certainly, one
362 would expect more specialists in older sites (Southwood et al, 1983) and these are
363 often the rare species in a community (Gaston, 1996).

364 It is most likely that combinations of environmental factors are shaping bird
365 diversity on the golf courses including topography, nest sites, the 'health' of the site

366 (Furness and Greenwood, 1993), and food source. The two habitats attracted different
367 types of bird species (insect feeding birds were more common on the golf course
368 habitats compared to the adjacent sites, while omnivorous species were rarer) due to
369 the vegetation composition of each habitat, invertebrate abundance and the land-use
370 of each habitat. The adjacent sites were pasture farmland which has repeatedly been
371 shown to contain homogenous habitats and low levels of biodiversity (Gregory and
372 Baillie, 1998; Chamberlain et al 2000; Stoate et al, 2001). Birds do not abide by man-
373 made boundaries and confusion can arise as to which birds are using the site and
374 which are just using the course as a stepping stone to other habitats. To overcome this
375 problem, individuals were only recorded if they were seen utilising the site. Given
376 that we also did not use song as a measure of presence, we believe that our estimates
377 of bird diversity on courses are very conservative and show an encouraging diversity
378 of birds on courses. Many bird species are becoming increasingly rare due to
379 intensive agricultural farming, loss of preferred habitat, pollution and land-use
380 changes (Gill, 1990; Gregory and Baillie, 1998). It is possible that the presence of
381 golfers could disturb birds and impact on breeding patterns but evidence suggests bird
382 communities can withstand intermediate levels of human activity like golfers (Riffell
383 et al., 1996; Chettri et al., 2001).

384 It has been suggested that golf courses could act as 'sink' habitats, into which
385 species are attracted, only to be killed by exposure to pesticides (Terman, 1997).
386 While, in theory at least, this is quite possible, there appears to be no scientific
387 evidence to support or refute this suggestion. While not being specifically tested for
388 in our study, we found no evidence to support this idea. In all the bird surveys we
389 conducted, not a single dead bird was seen whose death could be attributed to
390 anything other than predation. Furthermore, certain bird species, whose decline in
391 numbers have been attributed to agricultural pesticides (e.g. *T. philomelos*, Bullfinch,
392 (*Pyrrhula pyrrhula*) and Kestrel, (*Falco tinnunculus*)) were all found feeding on golf
393 courses, but not on the adjacent areas.

394 Species richness and abundance of carabid ground beetles were higher on the golf
395 course habitat than the adjacent sites. The difference in beetle numbers can be
396 attributed to courses having heterogeneous habitats, which provide varying
397 microclimates (Gange and Lindsay, 2002). Carabid species are vital omnivorous
398 predators in arable fields, providing farmers with a natural self-regulating pest control,
399 but numbers and species in intensive agricultural cultivation have repeatedly been

400 shown to be low (Kromp, 1999). An interesting finding made by Lindsay (2003) is
401 that these beetles were never recorded crossing fairways on golf courses, indicating
402 that these are major barriers for some invertebrate species. Incorporating natural
403 buffer zones within the golf course and between adjoining sites, as suggested by
404 Terman (2000), could provide wildlife with natural corridors. It is a fact that between
405 40 and 70% of a golf course is non-play areas of varying habitats (Anon., 1989;
406 Terman, 1997) which has the potential to act as corridors within the course. More
407 studies such as that of Gange et al. (2003) are required so that management of golf
408 course habitats can be better informed by ecological research.

409 There is growing concern about the decline in the natural populations of several
410 species of bumblebee in Europe, and only six of Britain's 19 species are now
411 regularly found in the countryside (Carvell, 2002). Declines in populations have been
412 attributed to habitat loss and agriculture intensification (Saville et al., 1997). We
413 found that the species richness and abundance of bumblebees was higher on the golf
414 course habitats than the adjacent habitats. Nest site availability, abundant flowering
415 herbaceous species and low management intensity (in the rough) are possible
416 explanations for the higher numbers and species of bumblebees found on the courses.
417 Often golf courses have a varied ground surface with exposed banks, which are ideal
418 nesting sites for some bee species (Gange and Lindsay, 2002). Such heterogeneous
419 habitats with uneven, exposed ground are much less common on farmland and
420 pasture. Our data suggest that the presence of a golf course in a landscape could have
421 a positive effect on bumblebee populations, though as yet we do not know if courses
422 can act as reservoirs of these insects. If golf courses can act as source habitats for
423 bees, then they could greatly enhance crop pollination and production in nearby areas.

424 We are aware that our data only cover one season. Future studies in golf course
425 ecology should include multi-species sampling and large sample sizes, performed
426 over longer periods of time. Recording species movements within golf courses (and
427 between golf course and adjacent sites) is vital, so that green keepers and ecologists
428 can formulate biological action plans, which target specific endangered species and
429 promote their existence with the course. These problems are the subject of our current
430 research.

431

432 **5. Conclusion**

433

434 In the current age of golf expansion, the most meaningful question to address is
435 whether construction of a golf course can enhance local biodiversity, compared with
436 the farmland from which it is invariably formed. This study has shown that golf
437 courses can enhance the diversity of three indicator groups (birds, ground beetles and
438 bumblebees), relative to adjacent pasture farmland. More studies are needed to
439 determine if golf courses act as source or sink habitats for beneficial insects and rare
440 species, or conversely, whether they can act as refuges for pest species too. Different
441 forms of farmland, involving varying intensities of agriculture also need to be
442 considered.

443

444 **Acknowledgments**

445 We are most grateful to all the site owners and golf courses for allowing access to
446 their land.

447

448 **References**

449

- 450 Altieri, M.A., 1999. The ecological role of biodiversity in agroecosystems. *Agric.*
451 *Ecosyst. Environ.*, 74: 19-31.
- 452 Anonymous, 1989. On course conservation: Managing golf's natural heritage. English
453 Nature, Peterborough, UK.
- 454 Anonymous, 1996. An environmental management program for golf courses. European
455 Golf Association. Pisces Publications, Newbury, UK.
- 456 Beebee, T.J.C., 2001. British wildlife and human numbers: the ultimate conservation
457 issue? *British Wildlife*, 1: 56-59.
- 458 Bell, J.P., 2000. Contesting rural recreation: The battle over access to Windermere.
459 *Landscape Use Policy*, 17: 295-303.
- 460 Bibby, C.J., Burgess, N.D., Hill, D.A., Mustoe, S.H., 2000. Bird census techniques.
461 Academic Press, London.
- 462 Blair, R.B., 1996. Land use and avian species diversity along an urban gradient. *Ecol.*
463 *Appl.*, 6: 506-519.
- 464 Boyle, S.A., Samson, F.B., 1985. Effects of non-consumptive recreation on wildlife: A
465 review. *Wildlife Soc. Bull.*, 13: 110-116.
- 466 Brennan, A-M., 1992. The management of golf courses as potential nature reserves. *Asp.*
467 *Appl. Biol.*, 29: 241-248.

- 468 Brown, V.K. Gange, A.C., 1989. Differential effects of above- and below-ground insect
469 herbivory during early plant succession. *Oikos*, 54: 67-76.
- 470 Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., 1993. Distance sampling:
471 Estimating abundance of biological populations. Chapman and Hall, London.
- 472 Burger, J., 2000. Landscapes, tourism and conservation. *Sci. Total Environ.*, 249: 39-49.
- 473 Butterfield, J., Luff, M.L., Baines, M., Eyre, M.D., 1995. Carabid beetle communities as
474 indicators of conservation potential in upland forests. *Forest Ecol. Manage.*, 79: 63-
475 77.
- 476 Carvell, C., 2002. Habitat use and conservation of bumblebees (*Bombus* spp.) under
477 different grassland management regimes. *Biol. Cons.*, 103: 33-49.
- 478 Chamberlain, D.E., Fuller, R.T., Bunce, R.G.H., Duckworth, J.C., Shrubbs, M., 2000.
479 Changes in the abundance of farmland birds in relation to the timing of agricultural
480 intensification in England and Wales. *J. Appl. Ecol.*, 37: 771-788.
- 481 Chettri, N., Sharma, E., Deb, D. C., 2001. Bird community structure along a trekking
482 corridor of Sikkim Himalaya: a conservation perspective. *Biol. Cons.*, 102: 1-16.
- 483 Cincotta, R.P., Wisnewski, J., Engelman, R. 2000. Human population in the biodiversity
484 hotspots. *Nature*, 404: 990-992.
- 485 Cohen, S., Durborow, T., Barnes, N.L., 1993. Groundwater and surface-water risk
486 assessments for proposed golf courses. *ACS Symp. Ser.*, 522: 214-227.
- 487 Cohen S., Svrjcek A., Durborow T., Barnes, N.L. 1999. Water quality impacts on golf
488 courses. *J. Environ. Qual.*, 28: 798-809.
- 489 Coppock, J.T., Duffield, B.S., 1975. Recreation in the countryside, a spatial analysis.
490 Macmillan Press Ltd, London.
- 491 Dair, I., Schofield, J.M., 1990. Nature conservation, legislation and environmental
492 aspects of golf course management in England. In: A.J. Cochran (Editor). *Science*
493 *and Golf*. E. and F.N. Spon, London, pp. 330-335.
- 494 Daniels, R.E., 1972. Golf and wildlife conservation. *J. Devon Trust Nat. Cons.*, 3: 39-46.
- 495 Fancy, S.G., 1997. A new approach for analysing bird densities from variable circular
496 plot counts. *Pacific Sci.*, 51: 107-114.
- 497 Forsythe, T.G., 2000. Common Ground beetles. Richmond Publishing Co. Ltd,
498 Richmond.
- 499 Fuller, R.J., 1997. Native and non-native trees as factors in habitat selection by woodland
500 birds in Britain. In: P.R. Ratcliffe (Editor). *Native and non-native in British Forestry*.
501 Institute of Chartered Foresters, Edinburgh, pp. 132-141.

- 502 Furness, R.W., Greenwood, J.J.D., 1993. Birds as monitors of environmental change.
503 Chapman and Hall, London.
- 504 Gange, A.C., Lindsay, D.E., 2002. Can golf courses enhance local biodiversity? In: E.
505 Thain (Editor). Science and Golf IV. Routledge, pp. 721-736.
- 506 Gange, A.C., Lindsay, D.E., Schofield, J.M., 2003. The ecology of golf courses.
507 Biologist, 50: 63-68.
- 508 Gaston, K.J., 1996. Biodiversity. A biology of numbers and differences. Blackwell
509 Science, London.
- 510 Gill, F.B., 1990. Ornithology. W.H.Freeman & Company, London.
- 511 Green, B.H., Marshall, I.C., 1987. An assessment of the role of golf courses in Kent,
512 England, in protecting wildlife and landscapes. Landscape Urban Plan., 14: 143-154.
- 513 Greenslade, P.J.M., 1964. Pitfall trapping as a method for studying populations of
514 Carabidae (Coleoptera). J. Anim. Ecol., 33: 301-310.
- 515 Gregory, R.D., Baillie, S.R., 1998. Large-scale habitat use of some declining British
516 birds. J. Appl. Ecol., 35: 785-799.
- 517 Kremen, C., 1992. Assessing the indicator properties of species assemblages for natural
518 areas monitoring. Ecol. Appl., 2: 203-217.
- 519 Kromp, B. 1999. Carabid beetles in sustainable agriculture: a review on pest control
520 efficacy, cultivation impacts and enhancement. Agric. Ecosyst. Environ., 74: 187-228.
- 521 Lennon, J.J., Greenwood, J.J.D., Turner, J.R.G., 2000. Bird diversity and environmental
522 gradients in Britain: a test for the species-energy hypothesis. J. Anim. Ecol., 69: 581-
523 598.
- 524 Lindroth, C.H., 1974. Coleoptera, Carabidae. Handbooks for the identification of British
525 insects. Royal Entomological Society of London.
- 526 Lindsay, D.E., 2003. Conservation potential and patch dynamics of lowland heath on
527 golf courses. PhD Thesis, University of London.
- 528 Magurran, A.E., 1988. Ecological diversity and its measurements. Chapman and Hall,
529 London.
- 530 Markwick, M.C., 2000. Golf tourism development, stakeholders, differing discourses
531 and alternative agendas: the case of Malta. Tourism Manage., 21: 515-524.
- 532 Mikola, J., Miettinen, M., Lehikoinen, E., Lehtila, K. 1994. The effects of disturbance
533 caused by boating on survival and behaviour of velvet scoter *Melanitta fusca*
534 ducklings. Biol. Cons., 67: 119-124.

- 535 Murphy, E.A., Aucott, M., 1998. An assessment of the amounts of arsenical pesticides
536 used historically in a geographical area. *Sci. Total Environ.*, 218: 89-101.
- 537 Nelson, J.T., Fancy, S.G., 1999. A test of the variable circular-plot method where exact
538 density of a bird population was known. *Pacific Cons. Biol.*, 5: 139-143.
- 539 Pearson, D.L., 1996. Selecting indicator taxa for the quantitative assessment of
540 biodiversity. In: D.L. Hawksworth (Editor). *Biodiversity: Measurements and*
541 *Estimation*. Chapman and Hall, London, pp. 75-79.
- 542 Peterken, G.F., 2001. Ecological effects of introduced tree species in Britain. *Forest*
543 *Ecol. Manage.*, 141: 31-42.
- 544 Platt A.E. 1994. *Toxic green: the trouble with golf*. Worldwatch Institute, Washington
545 DC.
- 546 Reynolds, R.T., Scott, J.M., Nussbaum, R.A., 1980. A variable circular-plot method for
547 estimating bird numbers. *Condor*, 82: 309-313.
- 548 Riffell, S.K., Gutzwiller, K.J., Anderson, S.H., 1996. Does repeated human intrusion
549 cause cumulative declines in avian richness and abundance? *Ecol. Appl.*, 6: 492-505.
- 550 Saville, N.M., Dramstad, W.E., Fry, G.L.A., Corbet, S.A., 1997. Bumblebee movement
551 in a fragmented agricultural landscape. *Agric. Ecosyst. Environ.*, 61: 145-154.
- 552 Simberloff, D., 1998. Flagship, umbrella, and keystones: is single species management
553 passé in the landscape era? *Biol. Cons.*, 83: 247-257.
- 554 Southwood, T.R.E. and Henderson, P.A. 2000. *Ecological Methods with Particular*
555 *Reference to the Study of Insect Populations*. Kluwer, Dordrecht.
- 556 Southwood, T.R.E., Brown, U.K., Reader, P.M., 1983. Continuity of vegetation in space
557 and time: a comparison of insects' habitat template in different successional stages.
558 *Res. Popul. Ecol.*, 3: 61-74.
- 559 Stoate, C., Boatman, N.D., Borralho, R.T., Carvalho, C.R., Snoo, G.R., Eden, P., 2001.
560 Ecological impacts of arable intensification in Europe. *J. Environ. Manage.*, 63: 337-
561 365.
- 562 Sun, D., Walsh, D., 1998. Review of studies on environmental impacts of recreation and
563 tourism in Australia. *J. Environ. Manage.*, 53: 323-338.
- 564 Terman, M.R., 1997. Natural links: naturalistic golf courses as wildlife habitat.
565 *Landscape Urban Plan.*, 38: 183-197.
- 566 Terman, M.R., 2000. Ecology and golf: saving wildlife habitats on human landscapes.
567 *Golf Course Manage.*, 68: 183-197.

568 Thomas, L., Laake, J.L., Derry, J.F., Buckland, S.T., Borchers, D.L., Anderson, D.R.,
 569 Burnham, K.P., Strindberg, S., Hedley, S.L., Burt, M.L., Marques, F., Pollard, J.H.,
 570 Fewster, R.M., 1998. Distance 3.5. Research Unit for Wildlife Population
 571 Assessment, University of St. Andrews, UK.

572 Walther-Hellwig, K., Frankl, R., 2000. Foraging habitats and foraging distances of
 573 bumblebees, *Bombus* spp. (Hym., Apidae), in an agricultural landscape. J. Appl. Ent.,
 574 127: 299-306.

575

576

577

578 **Figure legends**

579

580 Fig. 1. (a) Mean bird diversity (H) and mean bird density (numbers per ha) (b) on golf
 581 course habitat (●) and adjacent land sites (▲). Vertical bars represent one standard
 582 error.

583

584 Fig. 2. The relationships between bird diversity and tree diversity, for nine golf
 585 courses ($r^2 = 0.706$; $F_{1,7} = 16.796$, $P < 0.05$) (●) ($y = 0.265x + 1.902$) and nine
 586 adjacent sites ($r^2 = 0.705$; $F_{1,7} = 16.75$, $P < 0.05$) (▲) ($y = 0.366x + 1.663$).

587

588 Fig. 3. (a) Mean Carabid diversity and mean total numbers caught (b) on the nine golf
 589 courses (●) and nine adjacent sites (▲). Vertical bars represent one standard error.

590

591 Fig. 4. (a) Mean bumblebee diversity and mean total numbers caught (b) on the nine
 592 golf courses (●) and nine adjacent sites (▲). Vertical bars represent one standard
 593 error.