



ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect)

## Redox Biology

journal homepage: [www.elsevier.com/locate/redox](http://www.elsevier.com/locate/redox)

## Review Article

## Thiol redox homeostasis in neurodegenerative disease

Gethin J. McBean<sup>a,\*</sup>, Mutay Aslan<sup>b</sup>, Helen R. Griffiths<sup>c</sup>, Rita C. Torrão<sup>c</sup><sup>a</sup> UCD School of Biomolecular and Biomedical Science, University College, Dublin, Ireland<sup>b</sup> Department of Medical Biochemistry, Akdeniz University School of Medicine, Antalya, Turkey<sup>c</sup> School of Life and Health Sciences, Aston University, Birmingham, UK

## ARTICLE INFO

## Article history:

Received 27 March 2015

Received in revised form

8 April 2015

Accepted 11 April 2015

Available online 22 April 2015

## Keywords:

Cysteine

Glutathione

Thioredoxin

Glutaredoxin

Peroxiredoxin

Neurodegenerative disease

## ABSTRACT

This review provides an overview of the biochemistry of thiol redox couples and the significance of thiol redox homeostasis in neurodegenerative disease. The discussion is centred on cysteine/cystine redox balance, the significance of the  $x_c^-$  cystine–glutamate exchanger and the association between protein thiol redox balance and neurodegeneration, with particular reference to Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis and glaucoma. The role of thiol disulphide oxidoreductases in providing neuroprotection is also discussed.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Contents

Introduction	186
Cellular redox couples	187
Cysteine	188
The $x_c^-$ exchanger in neurodegenerative disease	188
Thiol redox homeostasis in glaucoma	189
Protein redox state in neurodegeneration	189
Glutaredoxin as a neuronal survival factor	190
Thioredoxin and neuronal survival	190
Peroxiredoxin and neuronal survival	191
Evidence for redox disturbance in neurodegeneration	191
Conclusion	192
Acknowledgement	192
References	192

**Abbreviations:** ALS, amyotrophic lateral sclerosis; AD, Alzheimer's disease; AP1, activator protein 1; BOAA,  $\beta$ -N-oxalylamino-L-alanine; DAXX, death-associated receptor; GCL, ganglion cell layer; Grx, glutaredoxin; GSH, glutathione (reduced); GSR, glutathione reductase; GSSG, glutathione disulphide (oxidised glutathione); GST, glutathione-S-transferase; L-DOPA, L-3,4-dihydroxyphenylalanine; mGluR, metabotropic glutamate receptor; MMP, mitochondrial membrane potential; MPP<sup>+</sup>, 1-methyl-4-phenylpyridinium; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; NF- $\kappa$ B, nuclear factor- $\kappa$ B; NGF, nerve growth factor; NMDA, N-methyl-D-aspartate; NRF2, nuclear factor (erythroid-derived 2)-like 2; NTG, normal tension glaucoma; POAG, primary open angle glaucoma; PD, Parkinson's disease; Prx, peroxiredoxin; SOD1, superoxide dismutase 1; Trx, thioredoxin; TrxR, thioredoxin reductase; TxNIP, thioredoxin inhibitory protein; xCT, functional subunit of the  $x_c^-$  cystine–glutamate exchanger


\* Correspondence to: UCD School of Biomolecular and Biomedical Science, Conway Institute, University College Dublin, Belfield, Dublin 4, Ireland.

E-mail address: [gethin.mcbean@ucd.ie](mailto:gethin.mcbean@ucd.ie) (G.J. McBean).

<http://dx.doi.org/10.1016/j.redox.2015.04.004>

2213-2317/© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).





**Fig. 2.** Pathways of cysteine metabolism; cysteine is derived from extracellular cystine via the  $x_c^-$  cystine glutamate exchanger and from methionine, via the transsulphuration pathway. Cysteine is the precursor for GSH, hydrogen sulphide ( $H_2S$ ) and taurine.  $C\beta S$ , cystathionine- $\beta$ -synthase; CDO, cysteine dioxygenase;  $C\gamma L$ , cystathionine- $\gamma$ -lyase; CSA, cysteine sulphinic acid; GCS,  $\gamma$ -glutamylcysteine synthase; GS, glutathione synthase; GSH, glutathione; SAH, S-adenosylhomocysteine; SAM, S-adenosylmethionine; THF, tetrahydrofolate.

## Cysteine

In the brain, cysteine must be supplied either from the extracellular medium, or in situ synthesis from methionine (see Fig. 2). Astrocytes and microglial cells are the main sites of GSH production in the brain and import cystine, which is immediately reduced to cysteine on entering the cytosol. In astrocytes, uptake of cystine provides approximately two thirds of the cysteine required for GSH synthesis. The remaining one third is derived from methionine via the transsulphuration pathway, but this contribution increases during oxidative stress [13,14]. Uptake of cystine is mediated by the plasma-membrane  $x_c^-$  cystine-glutamate exchanger (SLC7A11 carrier) that releases glutamate in a 1:1 ratio with cystine uptake (Fig. 2). Immature neurons readily take up cystine, but this facility is lost in mature cells. Instead, neurons take up cysteine, which originates from astrocytic GSH that is exported and degraded extracellularly [15,16]. Therefore, de novo synthesis of GSH in astrocytes must meet the antioxidant requirements of both astrocytes and neurons.

It is increasingly recognised that the  $x_c^-$  exchanger participates in an intracellular-extracellular cysteine/cystine redox cycle that serves to regulate the extracellular redox potential and is not directly associated with GSH synthesis [17–19]. Key features of the cycle are cystine uptake, intracellular reduction to cysteine and secretion of cysteine that is surplus to intracellular requirements to the extracellular space. The net effect is a reduction in the extracellular cysteine/cystine redox potential [19]. Additional support for the cycle comes from the observation that  $x_c^-$ -deficient mice do not display depletion of GSH [20].

The  $x_c^-$  exchanger has a significant role in regulating the extracellular concentration of glutamate and, increasingly, attention is drawn to the physiological and pathological implications of exchanger-mediated glutamate release [21]. For example,  $x_c^-$ -derived glutamate stimulates group 2/3 metabotropic glutamate receptors (mGluR2/3) in tissue slices of the nucleus accumbens or

prefrontal cortex, leading to a reduction in excitatory tone [22]. Moreover, withdrawal from cocaine reduces extracellular glutamate levels in the nucleus accumbens by decreasing cystine-glutamate exchange [22].

### The $x_c^-$ exchanger in neurodegenerative disease

Increased expression of the  $x_c^-$  exchanger and consequent elevation of glutamate release has been recorded in a number of neurodegenerative disorders, including brain tumours [18,23], HIV infection [24], multiple sclerosis [25] and hereditary haemochromatosis, a neurodegenerative disorder of the visual system [26,27]. Direct sampling of peritumoural extracellular fluid in Grade IV glioblastoma multiforme (GBM) patients by microdialysis has revealed concentrations of glutamate in excess of 100  $\mu M$  at the tumour margin [28], leading to widespread glutamate-mediated excitotoxicity and neurodegeneration. siRNA-mediated silencing of the xCT subunit of the exchanger protects against neurodegeneration and reduces glioma-associated oedema [29]. Equally, blockade of  $x_c^-$  exchanger with sulphasalazine or the cyclic glutamate analogue 4-carboxyphenylglycine (4-CPG) has an anti-proliferative and cytotoxic effect on glioma cell lines [30–32]. While additional research is needed, the results cited here illustrate the therapeutic potential of treatments targeting the  $x_c^-$  exchanger in glioma therapy.


Aberrant activity of the cystine-glutamate exchanger is associated with neurodegenerative changes in AD and ALS. In AD, activated microglia display increased expression and functional activity of the  $x_c^-$  exchanger and elevated glutamate release [33]. The toxicity of amyloid beta peptide 1–40 to cultured neurons was increased on exposure of the cells to activated microglia due to glutamate release via the  $x_c^-$  exchanger. Inhibition of N-methyl-D-aspartate (NMDA) receptors or system  $x_c^-$  prevented the microglia-enhanced toxicity of the peptide [33]. More recently, it has been hypothesised that excessive glutamate release from activated

microglia may be a significant contributor to motor neuron degeneration in the symptomatic stages of ALS. In support of this theory, expression of xCT is elevated in spinal cord and isolated microglia from mutant superoxide dismutase 1 (SOD1) ALS mice and in the spinal cord of ALS patients [34]. Moreover, xCT expression correlates with increased inflammation and enhanced release of glutamate. xCT is absent from motor neurons [34]. Deletion of xCT in ALS mice leads to earlier symptom onset, but ultimately, increases the number of surviving motor neurons. Taken together, these observations raise the possibility that targeting the x<sub>c</sub><sup>-</sup> exchanger could provide a mechanism to slow disease progression after onset of clinical symptoms.

### Thiol redox homeostasis in glaucoma

Glaucoma is the leading cause of irreversible blindness worldwide [35] and is characterised by progressive damage or degeneration to the optic nerve [36]. Since the retina and optic nerve are part of the central nervous system, glaucoma shares mechanistic and epidemiological similarities with other CNS neurodegenerations, including PD, ALS, AD and Huntington's disease [35].

Several hypotheses exist on the mechanism of ganglion cell death in glaucoma, including high intraocular pressure (IOP) and hypoxia [37], oxidative or nitrate stress [38,39], glutamate toxicity [40], loss of neurotrophic factors [41] and autoimmune reactions assisting degeneration of retinal ganglion cells [42]. The retina is one of the most vascularised tissues in the body and is particularly vulnerable to oxidative stress due to its high consumption of oxygen and exposure to light [43]. It has one of the highest oxidative metabolic rates per tissue weight [44]. Retinal cells require the maintenance of normal GSH levels, turnover rates and oxidation state [45]. GSH deficiency is manifested largely through an increased susceptibility to oxidative stress, and the resulting damage is thought to be a key step in the onset and progression of glaucoma [46]. Enzymes involved in GSH synthesis and metabolism in different types of retinal cells are shown in Fig. 3.



**Fig. 3.** Scheme of enzymes involved in glutathione synthesis and metabolism in the retina. AC, amacrine cell; BPC, bipolar cell; GC, ganglion cell; GCL, ganglion cell layer;  $\gamma$ GCL,  $\gamma$ glutamate cysteine ligase; G6PD, glucose-6-phosphate dehydrogenase; 6PG, 6-phosphogluconate; GPx, glutathione peroxidase; GR, glutathione reductase; GS, glutathione synthase; GSH, glutathione; HC, horizontal cell; INL, inner nuclear layer; IPL, inner plexiform layer; IOSP, inner and outer segments of photoreceptors; NF, nerve fibre; ONL, Outer nuclear layer; OPL, Outer plexiform layer; protein-S<sub>2</sub>, protein-S-S; protein (SH)<sub>2</sub>, protein-SH. RPE, retinal pigment epithelium; TR, thioredoxin reductase; TRX (SH)<sub>2</sub>, thioredoxin; TRX-S<sub>2</sub>, oxidised thioredoxin.

GSH levels decline in human eyes with age [47] and decreased GSH levels have been associated with glaucoma [48]. Moreno et al. analysed retinal GSH levels in a rat glaucoma model induced by chronic injection of hyaluronic acid in the eye anterior chamber [48]. Reduced GSH levels were assessed in the retinas from hypertensive eyes at 6 weeks of ocular hypertension compared to controls, whereas, at 3 or 10 weeks of treatment, levels of GSH did not change compared to eyes injected with vehicle. In accordance with animal studies, plasma glutathione levels measured in 21 patients with newly diagnosed primary open-angle glaucoma and 34 age- and gender-matched control subjects revealed that glaucoma patients exhibited significantly lower levels of reduced and total glutathione than did control subjects [49].

In a recent study performed on humans, levels of circulating total GSH (GSH and GSSG) were analysed in 34 primary open angle glaucoma (POAG) patients, 30 normal tension glaucoma (NTG) patients and 53 controls [50]. Total blood GSH levels were also determined in this study by the glutathione reductase-5,5 dithio-bis-2-nitrobenzoic acid (DTNB) recycling procedure. Independent of age, POAG and NTG patients demonstrated significantly lower GSH and total GSH levels than age-matched controls. Additionally, a lower redox index, defined as the GSH/GSSG ratio, was found only in POAG patients, in comparison to both NTG and control groups. GSSG levels were similar between all study groups. This study demonstrated that both POAG and NTG patients exhibit lower GSH and total GSH levels than age-matched controls, indicating a compromise of the antioxidant defence systems in glaucoma.

Glutathione S-transferase (GST) catalyses the conjugation of GSH to electrophilic centres on a variety of substrates including toxic species [51]. In a recent study, the aqueous humour proteome was analysed by antibody microarray in samples obtained from 10 clinically uncontrolled POAG patients and 10 cataract patients immediately before trabeculectomy and cataract surgery, respectively. Aqueous humour GST levels were significantly lower in POAG patients than in controls reflecting a weakening of GST-mediated mechanism of protection against harmful electrophiles produced during oxidative damage [52]. Serum antibodies against GST have also been reported to be higher in patients with POAG than in controls [53].

Glutamate toxicity is an important mechanism of ganglion cell death in glaucoma [54]. Increased levels of glutamate have been reported in the vitreous of glaucomatous patients and a prolonged elevation of extracellular glutamate is possibly caused by chronic ocular hypertension [55]. The cystine–glutamate exchanger is upregulated by oxidative stress in retinal ganglion cells [56] and, given that glutamate is toxic to these cells via over-stimulation of the NMDA receptor [40], upregulation of the exchanger under oxidative stress conditions in glaucoma would have injurious consequences.

### Protein redox state in neurodegeneration

Alterations in protein redox state have been implicated in the pathology of neurodegenerative diseases [57–59]. Early important observations of cell death under redox stress have been reviewed and linked to more recent, elegant experiments describing mechanisms of redox-regulated cell death in AD, PD and ALS. Key pathways that are affected by redox state and predispose to death are the apoptosis signalling kinase 1 (ASK1)/JNK and ASK1/p38MAPkinase cascades [9,60]. Mitochondrial dysfunction is considered pivotal to cell death as a source of increases in thiol-oxidising reactive oxygen species and during intrinsic apoptotic signalling. Consequently, a number of studies have investigated Grx, Trx and Prx activity and mitochondrial function in neurodegeneration models. While the majority of articles focus on

mitochondria as a source of oxidants that contribute to cytotoxicity, mitochondria have also been reported to play an important role in removal of hydrogen peroxide, e.g. generated by redox cycling of chemicals such as paraquat [61,62]. Nicotinamide nucleotide transhydrogenase capitalises on the mitochondrial proton gradient to generate NADPH from NADH and NADP<sup>+</sup>, so providing a link between energetics and removal of hydrogen peroxide via the Trx2/TrxR and Prx systems [63].

#### *Glutaredoxin as a neuronal survival factor*

Glutathionylation/deglutathionylation reactions are regulated by the expression and activity of Grx and the availability of GSH substrate which is required for chemical reduction of Grx. Grx exists in at least 3 forms in humans, however, most studies focus on Grx1 found in the cytoplasm and Grx2 found in the nucleus and mitochondria. In 2004, Kenchappa et al. described that down-regulation of Grx using anti-sense oligonucleotides prevented recovery of mitochondrial complex I activity in the striatum after 1-methyl, 4-phenyl, 1,2,3,6-tetrahydropyridine (MPTP) treatment [64]. This study suggested a critical role for Grx in recovery of mitochondrial function in brain. A follow up study investigated the mechanism of toxicity of the excitotoxin  $\beta$ -N-oxalyl amino-L-alanine (BOAA) [65]. While male mice were susceptible to mitochondrial complex I loss and GSH depletion only ovariectomised female mice showed mitochondrial-dependent neuronal loss. Grx expression, central to neuroprotection via maintaining complex I activity, is regulated by oestrogen; treatment of SH-SY5Y cells with oestrogen upregulated Grx1 and protected from BOAA mediated mitochondrial membrane potential (MMP) loss. Later, Lee et al. demonstrated that Grx2 is involved in glutathionylation of protein cysteine sulphhydryl residues in the mitochondria and is required for iron sulphur cluster biogenesis and complex I activity. Following inhibition of Grx2 in vitro, complex I activity declines in a manner consistent with the pathological events associated with PD [66]. The concomitant rise in intracellular iron further predisposes neurones to oxidative stress. Consistent with these findings, Karunakaran elegantly demonstrated that downregulation of Grx2 using antisense oligonucleotides in the mouse brain in vivo resulted in partial loss of complex I activity supporting the concept that Grx2 may help maintain complex I function in the mitochondria. These findings were reinforced in vitro by over-expression of Grx2 in neuroblastoma cells which abolished 1-methyl-4-phenylpyridinium (MPP<sup>+</sup>)-mediated toxicity [67].

Despite its localisation to the cytoplasm, downregulation of Grx1 by shRNA resulted in loss of MMP that is associated with oxidation of thiols on voltage dependent anion channels [68]. Loss of MMP was preventable by the antioxidant,  $\alpha$ -lipoic acid, or by cyclosporine A, an inhibitor of mitochondrial permeability transition suggesting that either there is direct interaction between Grx1 and proteins in the mitochondrial membrane or that an intermediate redox active protein transduces the reducing signal.

Most commonly, familial ALS is associated with the aggregation of mutant SOD1, but whether SOD1 aggregates represent a cause, a correlate or a consequence of processes leading to cell death is unclear. As well as being localised in the cytoplasm, SOD1 is found in the intermembrane space and in ALS mitochondria, fission and dysfunction has been attributed to mutant SOD aggregation. Aggregation can be prevented by Grx2 overexpression, preserving mitochondrial function and preventing apoptosis [69]. Related to this, biochemical studies by Bouldin et al. have described the propensity for the intramolecular disulphide bond between two SOD monomers to be reduced by Grx1/GSH; they showed that ALS mutant SOD was more susceptible to reduction to monomers and inactivation than WT [70]. This is significant because it is the apo- and reduced forms of ALS mutant hSOD1 that are especially

vulnerable to destabilisation and readily aggregate under mild oxidative stress conditions. In ageing yeast, glutathionylation was shown to be essential for human SOD1 activation [71]. Further work is needed in this area to rationalise the role of Grx in mutant SOD aggregation.

Aberrant regulation of protein glutathionylation can disrupt both apoptotic and survival signalling pathways. More recent studies have focussed on potential signalling pathways by which Grx may support neuronal survival. DAXX, the death associated protein downstream from ASK1, has surfaced as an important transducer of death signals from the nucleus in PD. Based on studies of DJ-1, a putative gene recessively linked to early onset PD, reduced DAXX is normally retained in the nucleus. DJ-1 normally functions as an antioxidant, transcriptional co-activator, and molecular chaperone that is maintained active in a reduced state by Grx1. However, down-regulation of Grx 1 caused loss of DJ-1 protein, translocation of DAXX from nucleus, and cell death. This effect was eliminated in the DJ-1 cys mutant which was not susceptible to oxidation or degradation so maintained DAXX in the nucleus [72]. More recent studies using the brains from DJ-1 (–/–) mouse model of PD have suggested that the animals have adapted to loss of DJ-1, increasing consumption of hydrogen peroxide and through upregulation of the Trx/TrxR/Prx system [73] and in fact that the ability to adapt to stress may be the key factor in overt disease. In a model of AD, using SH-SY5Y exposed to amyloid beta, both Grx1 and Trx1 were oxidised, associated with nuclear export of DAXX. Moreover, amyloid beta toxicity was inhibited by insulin-like growth factor-I and by overexpressing Grx1 or Trx1 [74] highlighting the importance of maintaining redox state for neuronal survival.

Another hydrogen peroxide generating drug relevant to PD is L-3,4-dihydroxyphenylalanine (L-DOPA). While it is used as a therapeutic, L-DOPA causes oxidative inactivation of Trx and Grx, with concomitant activation of ASK1 [59,75]. A potential mechanism for this has been investigated by Sabens et al. [75]; in biochemical studies, Grx underwent irreversible adduction with dopaquinone to its nucleophilic active-site Cys-22. This caused enzymatic inactivation but did not cause protein degradation, consistent with observations after L-DOPA treatment. Whether this mechanism may also contribute to cognitive decline observed in PD patients remains to be explored.

#### *Thioredoxin and neuronal survival*

Trx exists in multiple forms and acts as a reducing agent within the cell and extracellularly. It restores the activity of oxidised Prx and is in turn recycled at the expense of NADPH by TrxR. In addition to regulation of expression, TrxR activity is also regulated post-translationally by thioredoxin inhibitory protein (TxNIP).

Trx has been recognised as a neurotrophic factor for over 10 years; the earliest studies showed it to support nerve growth factor (NGF) signalling and to play a critical role in NGF-mediated neurite outgrowth in PC12 cells [76]. It exists in at least two isoforms, Trx1 in the cytosol and Trx2 in the mitochondria. Trx1 is a small, 12-kDa, conserved and ubiquitous multifunctional protein with several redox-active cysteine residues. It reduces disulphide bonds and sulphenic acids and exhibits transnitrosylation activity [77–80]. Trx1 has many interaction partners depending on its cellular localisation. Through its reductase activity it may regulate apoptosis, cell growth, differentiation and development [81,82]. In the nucleus, Trx1 binds directly to a number of different transcription factors including p53, NF- $\kappa$ B, and activator protein-1 (AP1) and thereby modulates their DNA-binding activity [83,84]. With respect to apoptosis inhibition, at least three binding partners have been identified in the cytoplasm; ASK-1, TxNIP and actin, where actin protects Trx1 from degradation and preserves

its anti-apoptotic function [81,85]. The cytoprotective protein DJ-1 which is frequently mutated in PD, binds to ASK1 in a Cys-106 redox-sensitive manner and can be reduced by Trx1 [86,87]. It has been suggested that DJ-1 is an atypical peroxiredoxin-like peroxidase that scavenges hydrogen peroxide through oxidation of Cys-106. In WT mice, an increase of Cys-106 oxidised DJ-1 was observed after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) treatment [88]. DJ-1 has also been suggested to act as an upstream activator of the transcription factor, nuclear factor (erythroid-derived)2-like 2 (NRF2). NRF2 regulates Trx, and its overexpression inhibits ASK1/JNK and ASK1/p38 pathways that are frequently activated in neurodegenerative disease [89]. While overexpression of DJ-1 resulted in increased NRF2 protein levels, nuclear translocation and binding to the ARE site in the Trx1 promoter, NRF2 knockdown abolished DJ-1-mediated Trx1 induction and cytoprotection against hydrogen peroxide [11]. In PD patients with mutant DJ-1, alternative NRF2 activators may prove to be a useful strategy for upregulating Trx1 expression.

Neurotoxins associated with risk for PD frequently associate with oxidised Trx and activate unique pathways. For example, paraquat oxidised Trx1 impairing its ASK1 inhibitor activity and leading to JNK and caspase 3 activation whereas 1-methyl-4-phenylpyridinium (MPP<sup>+</sup>) and rotenone oxidised Trx2 without activating the JNK pathway [90,91]. Injection of MPP<sup>+</sup> into mouse brain decreased levels of TrxR 1 mRNA, protein and activity [92]. Others have shown that MPP<sup>+</sup> toxicity is due to ER stress which can be inhibited by Trx-1 overexpression; Trx-1 played a neuroprotective role in MPP<sup>+</sup>-mediated neurone loss by suppressing ER stress [93]. Interestingly, Trx acts not only as a reducing agent but also as a chaperone; when Trx mutants were studied in the Pael-R drosophila model of PD, it was the chaperone rather than redox activity of Trx that proved to be more significant for enhancing cell survival [94].

Trx1 also associates with the plasma membrane and is trafficked with a limited number of cytosolic proteins via the leaderless secretory pathway [95]. We have shown previously that under oxidative stress, Trx1 secretion is altered [96]. Trx1 and Grx1 are also released to the cerebrospinal fluid. In 120 patients the early stages of AD and with mild cognitive impairment Trx1 and Grx1 levels correlated with the established AD biomarkers tau and phospho-tau suggesting their potential involvement in the pathogenesis of disease [97].

The neuroprotective effect of Trx1 and Trx2 were investigated in a rat glaucoma model [98]. Expression of Trx1 and Trx2 was observed in the retinal ganglion cell layer (GCL), nerve fibre (NF) layer and inner nuclear layer (INL). Trx1 level decreased 2 weeks after glaucoma induction and more notably after 5 weeks. No change in Trx2 levels was reported. The effects of Trx1 and Trx2 overexpression on retinal ganglion cell survival were evaluated 5 weeks after glaucoma induction. Trx1 and Trx2 preserved 45 and 37% of cells, respectively that were destined to die in glaucomatous retinas.

#### *Peroxioredoxin and neuronal survival*

Basal expression of the six different Prx isozymes shows a distinctive distribution profile within brain regions and different cell types. On the one hand, Prx1 and 6 are expressed in glial cells but not in neurons; conversely, Prx2, 3, 4, and 5 are expressed in neurons [99]. Of these enzymes, it is Prx3 that is found in mitochondria. Prx 6 differs from the other Prx enzymes and is a 1-Cys Prx that lacks an internal resolving cysteine residue. The other mammalian Prx enzymes are 2-Cys Prx where the peroxidatic cysteine is first oxidised to a sulphenic acid and then it oxidises a second resolving cysteine to form a di-sulphide bond. The latter is reduced via Trx and TrxR.

Prx2 is highly expressed in many tissues and along with Prx 6 is the focus of most research in AD. In a transgenic Alzheimer's mouse diseased brain, Prx2 levels were significantly elevated. This was suggested to be an adaptive response that protected the mice against amyloid beta toxicity [100]. In a histochemical study of brains from AD patients, nitrated Prx2 was identified. Randall et al. investigated the functional consequences of Prx2 tyrosine nitration, and demonstrated that nitration was on a non-catalytic residue that resulted in increases in peroxidase activity and resistance to over-oxidation [101]. Prx2 is also S-nitrosylated forming SNO-Prx2 by reaction with nitric oxide at two critical cysteine residues (C51 and C172). In contrast to the effect of nitration, this prevents its reaction with peroxides [102].

The role of Prx6 is less clear; on the one hand, apoptosis was inhibited after amyloid beta treatment in PC12 cells overexpressing wild-type Prx6, but not in cells that overexpressed the C47S catalytic mutant. This indicates that the peroxidase activity of Prx6 protects PC12 cells from amyloid-induced neurotoxicity [103]. However, in a mouse model after amyloid beta infusion, memory impairment in Prx6 transgenic mice was worse than C57BL/6 mice. In addition, the astrocytes and microglia cells of amyloid-infused Prx6 transgenic mice were more activated, lipid peroxidation and protein carbonyl levels were increased and glutathione levels were lower, suggesting that Prx6 is promoting rather than preventing oxidative stress [104].

In the PD field, interest has been in Prx1, 2 and 3. In MN9D cells, overexpression of Prx1 protected against 6 hydroxydopamine toxicity, prevented p38 MAPK activation and subsequent activation of caspase-3. In contrast, apoptotic death signals were enhanced by RNA interference-targeted reduction of Prx1 [105]. Hu et al. explored the role of Prx2 and showed it inhibited 6 hydroxydopamine -induced ASK1 activation by modulating the redox state of Trx1 so preventing its dissociation from ASK1 [106]. In cells expressing a common p.G2019S LRRK2 gene mutation (rs34637584:A > G) that is responsible for up to 30–40% of PD cases in some ethnic populations, the phosphorylation of Prx3 is increased. LRRK2 interacts with Prx3 and mutations in the LRRK2 kinase domain significantly increased phosphorylation but decreased peroxidase activity and increased death in neuronal cells [107]. These findings point towards a common downstream pathway of redox imbalance in PD that may be mediated through different upstream effectors and by either environmental or genetic factors.

#### *Evidence for redox disturbance in neurodegeneration*

One of the more compelling arguments for thiol regulatory networks in neurodegenerative disease is evidence of altered expression in post-mortem tissue or in circulating cells and plasma. To this end, there is limited analysis of Grx1 in human neurodegenerative disease. However, of note, a recent study has examined post-mortem midbrain samples from PD patients, in which it was observed that Grx1 content is decreased in PD, specifically within the dopaminergic neurons [108]. This further supports a role for redox in PD. Although whether it is causal is unclear. Increased SNO-Prx2 has been described in human PD brains, and S-nitrosylation of Prx2 inhibits both its enzymatic activity and protective function from oxidative stress [102].

In post-mortem brain tissue of patients with AD, many blood vessels exhibited Prx6 staining that appeared to be due to the astrocytic foot processes [109]. Furthermore, an increase in Grx1 and a decrease in neuronal Trx1 have been described in AD brains by one group, while a second group reported from 120 patients that the Trx1, Trx2, Grx1, and Grx2 expression pattern was altered in hippocampal tissue sections from AD patients compared to

controls. In addition, oxidative modifications to Prx-2 and Prx-6 were increased in AD plasma [110].

## Conclusion

Much progress has been made in understanding the significance of thiol redox imbalance as a contributing factor to neurodegenerative disease. The biological reactivity of the sulphur atom in cysteine, whether as the free amino acid or incorporated into proteins and peptides, is a significant factor in determining susceptibility to oxidative damage, excitotoxicity and neurodegeneration. It is clear that restoration of redox balance may offer a useful approach to minimise neuronal loss during neurodegeneration.

## Acknowledgement

GMcB, MA and HRG were supported by the European Cooperation In Science and Technology (COST) Action BM1203, EU-ROS.

## References

- [1] F. Gu, V. Chauhan, A. Chauhan, Glutathione redox imbalance in brain disorders, *Current Opinion in Clinical Nutrition and Metabolic Care* 18 (2015) 89–95.
- [2] J. Ruzkiewicz, P. Albrecht, Changes in mitochondrial antioxidant systems in neurodegenerative disease and acute brain disorders, *Neurochemistry International* (2015).
- [3] K. Aoyama, T. Nakaki, Impaired glutathione synthesis in neurodegeneration, *International Journal of Molecular Science* 14 (10) (2013) 21021–21044. <http://dx.doi.org/10.3390/ijms141021021> 24145751.
- [4] K.G. Reddie, K.S. Carroll, Expanding the functional diversity of proteins through cysteine oxidation, *Current Opinion in Chemical Biology* 12 (6) (2008) 746–754. <http://dx.doi.org/10.1016/j.cbpa.2008.07.028> 18804173.
- [5] K.G. Reddie, Y.H. Seo, W.B. Mue III, S.E. Leonard, K.S. Carroll, A chemical approach for detecting sulfenic acid-modified proteins in living cells, *Molecular Biosystems* 4 (6) (2008) 521–531. <http://dx.doi.org/10.1039/b719986d18493649>.
- [6] S. Ratnayake, I.H. Dias, E. Lattman, H.R. Griffiths, Stabilising cysteinyl thiol oxidation and nitrosation for proteomic analysis, *Journal of Proteomics* 92 (2013) 160–170. <http://dx.doi.org/10.1016/j.jprot.2013.06.019> 23796488.
- [7] S. Fernandez-Fernandez, A. Almeida, J.P. Bolaños, Antioxidant and bioenergetic coupling between neurons and astrocytes, *Biochemical Journal* 443 (1) (2012) 3–11. <http://dx.doi.org/10.1042/BJ20111943> 22417747.
- [8] A. Herrero-Mendez, A. Almeida, E. Fernández, C. Maestre, S. Moncada, J.P. Bolaños, The bioenergetic and antioxidant status of neurons is controlled by continuous degradation of a key glycolytic enzyme by APC/C-Cdh1, *Nature Cell Biology* 11 (6) (2009) 747–752. <http://dx.doi.org/10.1038/ncb1881> 19448625.
- [9] J.J. Mielay, M.M. Gallogly, S. Qanungo, E.A. Sabens, M.D. Shelton, Molecular mechanisms and clinical implications of reversible protein S-glutathionylation, *Antioxidants and Redox Signaling* 10 (11) (2008) 1941–1988. <http://dx.doi.org/10.1089/ars.2008.2089> 18774901.
- [10] J.R. Roede, K. Uppal, Y. Liang, D.E. Promislow, L.M. Wachtman, D.P. Jones, Characterization of plasma thiol redox potential in a common marmoset model of aging, *Redox Biology* 18 (2013) 387–393.
- [11] J.W. Zhu, J.F. Yuan, H.M. Yang, S.T. Wang, C.G. Zhang, L.L. Sun, H. Yang, H. Zhang, Extracellular cysteine (cys)/cystine (CySS) redox regulates metabotropic glutamate receptor 5 activity, *Biochimie* 94 (3) (2012) 617–627. <http://dx.doi.org/10.1016/j.biochi.2011.09.013> 21964032.
- [12] D. Ghosh, G.J. Brewer, External cys/cySS redox state modification controls the intracellular redox state and neurodegeneration via Akt in aging and Alzheimer's disease mouse model neurons, *Journal of Alzheimer's Disease* 42 (1) (2014) 313–324. <http://dx.doi.org/10.3233/JAD-132756> 24844688.
- [13] L. Diwakar, V. Ravindranath, Inhibition of cystathionine-gamma-lyase leads to loss of glutathione and aggravation of mitochondrial dysfunction mediated by excitatory amino acid in the CNS, *Neurochemistry International* 50 (2) (2007) 418–426. <http://dx.doi.org/10.1016/j.neuint.2006.09.014> 17095121.
- [14] S. Kandil, L. Brennan, G.J. McBean, Glutathione depletion causes a JNK and p38MAPK-mediated increase in expression of cystathionine-gamma-lyase and upregulation of the transsulfuration pathway in C6 glioma cells, *Neurochemistry International* 56 (4) (2010) 611–619. <http://dx.doi.org/10.1016/j.neuint.2010.01.004> 20060865.
- [15] J.I. Sagara, K. Miura, S. Bannai, Maintenance of neuronal glutathione by glial cells, *Journal of Neurochemistry* 61 (5) (1993) 1672–1676. <http://dx.doi.org/10.1111/j.1471-4159.1993.tb09802.x> 8228986.
- [16] R. Dringen, B. Pfeiffer, B. Hamprecht, Synthesis of the antioxidant glutathione in neurons: supply by astrocytes of CysGly as precursor for neuronal glutathione, *Journal of Neuroscience* 19 (2) (1999) 562–569 9880576.
- [17] A. Banjac, T. Perisic, H. Sato, A. Seiler, S. Bannai, N. Weiss, P. Kölle, K. Tschopp, R.D. Issels, P.T. Daniel, M. Conrad, G.W. Bornkamm, The cystine/cysteine cycle: a redox cycle regulating susceptibility versus resistance to cell death, *Oncogene* 27 (11) (2008) 1618–1628. <http://dx.doi.org/10.1038/sj.onc.1210796> 17828297.
- [18] J. Lewerenz, P. Baxter, R. Kassubek, P. Albrecht, J. Van Liefvering, M. A. Westhoff, M.E. Halatsch, G. Karpel-Massler, P.J. Meakin, J.D. Hayes, E. Aronica, I. Smolders, A.C. Ludolph, A. Methner, M. Conrad, A. Massie, G. E. Hardingham, P. Maher, Phosphoinositide 3-kinases upregulate system xc<sup>-</sup> via eukaryotic initiation factor 2 $\alpha$  and activating transcription factor 4 – a pathway active in glioblastomas and epilepsy, *Antioxidants and Redox Signaling* 20 (2014) 2907–2922.
- [19] M. Conrad, H. Sato, The oxidative stress-inducible cystine/glutamate antiporter, system xc<sup>-</sup> (cystine supplier and beyond), *Amino Acids* 42 (1) (2012) 231–246. <http://dx.doi.org/10.1007/s00726-011-0867-5> 21409388.
- [20] D. De Blundel, A. Schallier, E. Loyens, R. Fernando, H. Miyashita, J. Van Liefvering, K. Vermoessen, S. Bannai, H. Sato, Y. Michotte, I. Smolders, A. Massie, Loss of system xc<sup>-</sup> formula does not induce oxidative stress but decreases extracellular glutamate in hippocampus and influences spatial working memory and limbic seizure susceptibility, *Journal of Neuroscience* 31 (2011) 3592–3603.
- [21] R. Bridges, V. Lutgen, D. Lobner, D.A. Baker, Thinking outside the cleft to understand synaptic activity: contribution of the cystine-glutamate antiporter (System xc<sup>-</sup>) to normal and pathological glutamatergic signaling, *Pharmacological Reviews* 64 (3) (2012) 780–802. <http://dx.doi.org/10.1124/pr.110.003889> 22759795.
- [22] M.M. Moran, K. McFarland, R.I. Melendez, P.W. Kalivas, J.K. Seamans, Cystine/glutamate exchange regulates metabotropic glutamate receptor presynaptic inhibition of excitatory transmission and vulnerability to cocaine seeking, *Journal of Neuroscience* 25 (27) (2005) 6389–6393. <http://dx.doi.org/10.1523/JNEUROSCI.1007-05.2005> 16000629.
- [23] J.F. De Groot, H. Sontheimer, Glutamate and the biology of gliomas, *Glia* 59 (8) (2011) 1181–1189. <http://dx.doi.org/10.1002/glia.21113> 21192095.
- [24] C.C. Bridges, H. Hu, S. Miyauchi, U.N. Siddaramappa, M.E. Ganapathy, L. Ignatowicz, D.M. Maddox, S.B. Smith, V. Ganapathy, Induction of cystine-glutamate transporter xc<sup>-</sup> by human immunodeficiency virus type 1 transactivator protein Tat in retinal pigment epithelium, *Investigative Ophthalmology and Visual Science* 45 (9) (2004) 2906–2914. <http://dx.doi.org/10.1167/iovs.03-1355> 15326101.
- [25] O. Pampliega, M. Domercq, F.N. Soria, P. Villoslada, A. Rodríguez-Antigüedad, C. Matute, Increased expression of cystine/glutamate antiporter in multiple sclerosis, *Journal of Neuroinflammation* 8 (2011) 63. <http://dx.doi.org/10.1186/1742-2094-8-63> 21639880.
- [26] J.P. Gnana-Prakasam, A. Tawfik, M. Romej, S. Ananth, P.M. Martin, S.B. Smith, V. Ganapathy, Iron-mediated retinal degeneration in haemojuvelin-knockout mice, *Biochemical Journal* 441 (2) (2012) 599–608. <http://dx.doi.org/10.1042/BJ20111148> 21943374.
- [27] J.P. Gnana-Prakasam, M. Thangaraju, K. Liu, Y. Ha, P.M. Martin, S.B. Smith, V. Ganapathy, Absence of iron-regulatory protein Hfe results in hyperproliferation of retinal pigment epithelium: role of cystine/glutamate exchanger, *Biochemical Journal* 424 (2) (2009) 243–252. <http://dx.doi.org/10.1042/BJ20090424> 19715555.
- [28] H.J. Marcus, K.L. Carpenter, S.J. Price, P.J. Hutchinson, In vivo assessment of high-grade glioma biochemistry using microdialysis: a study of energy-related molecules, growth factors and cytokines, *Journal of Neuro-Oncology* 97 (1) (2010) 11–23. <http://dx.doi.org/10.1007/s11060-009-9990-5> 19714445.
- [29] N.E. Savaskan, A. Heckel, E. Hahnen, T. Engelhorn, A. Doerfler, O. Ganslandt, C. Nimsky, M. Buchfelder, I.Y. Eypoglu, Small interfering RNA-mediated xCT silencing in gliomas inhibits neurodegeneration and alleviates brain edema, *Nature Medicine* 14 (6) (2008) 629–632. <http://dx.doi.org/10.1038/nm1772> 18469825.
- [30] W.J. Chung, S.A. Lyons, G.M. Nelson, H. Hamza, C.L. Gladson, G.Y. Gillespie, H. Sontheimer, Inhibition of cystine uptake disrupts the growth of primary brain tumors, *Journal of Neuroscience* 25 (31) (2005) 7101–7110. <http://dx.doi.org/10.1523/JNEUROSCI.5258-04.2005> 16079392.
- [31] P.W. Gout, A.R. Kang, C.R. Buckley, N. Bruchovsky, A.R. Buckley, Sulfasalazine, a potent suppressor of lymphoma growth by inhibition of the xc<sup>-</sup> cystine transporter: a new action for an old drug, *Leukemia* 15 (10) (2001) 1633–1640. <http://dx.doi.org/10.1038/sj.leu.2402238> 11587223.
- [32] L. Sleire, B.S. Skeie, I.A. Netland, H.E. Førde, E. Dodoo, F. Selheim, L. Leiss, J. I. Heggdal, P.H. Pedersen, J. Wang, P.Ø. Enger, Drug repurposing: sulfasalazine sensitizes gliomas to gamma knife radiosurgery by blocking cystine uptake through system xc<sup>-</sup>, leading to glutathione depletion, *Oncogene* (2015). <http://dx.doi.org/10.1038/onc.2015.60>, epub ahead of print].
- [33] S. Qin, C. Colin, I. Hinners, A. Gervais, C. Cheret, M. Mallat, System xc<sup>-</sup> and apolipoprotein E expressed by microglia have opposite effects on the neurotoxicity of amyloid-beta peptide 1–40, *Journal of Neuroscience* 26 (12) (2006) 3345–3356. <http://dx.doi.org/10.1523/JNEUROSCI.5186-05.2006> 16554485.

- [34] P. Mesci, S. Zaïdi, C.S. Lobsiger, S. Millemcamps, C. Escartin, D. Seilhean, H. Sato, M. Mallat, S. Boillée, System xc<sup>-</sup> is a mediator of microglial function and its deletion slows symptoms in amyotrophic lateral sclerosis mice, *Brain* 138 (1) (2015) 53–68. <http://dx.doi.org/10.1093/brain/awu312> 25384799.
- [35] S.D. Crish, D.J. Calkins, Neurodegeneration in glaucoma: progression and calcium-dependent intracellular mechanisms, *Neuroscience* 176 (2011) 1–11. <http://dx.doi.org/10.1016/j.neuroscience.2010.12.036> 21187126.
- [36] H.A. Quigley, *Glaucoma*, *Lancet* 377 (9774) (2011) 1367–1377. [http://dx.doi.org/10.1016/S0140-6736\(10\)61423-7](http://dx.doi.org/10.1016/S0140-6736(10)61423-7) 21453963.
- [37] Z. He, A.J. Vingrys, J.A. Armitage, C.T. Nguyen, B.V. Bui, Chronic hypertension increases susceptibility to acute IOP challenge in rats, *Investigative Ophthalmology and Visual Science* 55 (12) (2014) 7888–7895. <http://dx.doi.org/10.1167/jovs.14-15207> 25370512.
- [38] S.C. Saccá, A. Izzotti, P. Rossi, C. Traverso, Glaucomatous outflow pathway and oxidative stress, *Experimental Eye Research* 84 (3) (2007) 389–399. <http://dx.doi.org/10.1016/j.exer.2006.10.008> 17196589.
- [39] M. Aslan, S. Dogan, E. Kucuksayan, Oxidative stress and potential applications of free radical scavengers in glaucoma, *Redox Report* 18 (2) (2013) 76–87. <http://dx.doi.org/10.1179/1351000212Y.0000000033> 23485101.
- [40] N.J. Sucher, S.A. Lipton, E.B. Dreyer, Molecular basis of glutamate toxicity in retinal ganglion cells, *Vision Research* 37 (24) (1997) 3483–3493. [http://dx.doi.org/10.1016/S0042-6989\(97\)00047-3](http://dx.doi.org/10.1016/S0042-6989(97)00047-3) 9425525.
- [41] N. Gupta, Y.H. Yücel, Glaucoma as a neurodegenerative disease, *Current Opinion in Ophthalmology* 18 (2) (2007) 110–114. <http://dx.doi.org/10.1097/ICU.0b013e3280895aea> 17301611.
- [42] M.B. Wax, G. Tezel, Immunoregulation of retinal ganglion cell fate in glaucoma, *Experimental Eye Research* 88 (4) (2009) 825–830. <http://dx.doi.org/10.1016/j.exer.2009.02.005> 19233171.
- [43] D. Athanasiou, M. Aguilá, D. Bevilacqua, S.S. Novoselov, D.A. Parfitt, M. E. Cheetham, The cell stress machinery and retinal degeneration, *FEBS Letters* 587 (13) (2013) 2008–2017. <http://dx.doi.org/10.1016/j.febslet.2013.05.020> 23684651.
- [44] E.S. Shin, C.M. Sorenson, N. Sheibani, Diabetes and retinal vascular dysfunction, *Journal of Ophthalmic and Vision Research* 9 (3) (2014) 362–373. <http://dx.doi.org/10.4103/2008-322X.143378> 25667739.
- [45] E. Ganea, J.J. Harding, Glutathione-related enzymes and the eye, *Current Eye Research* 31 (1) (2006) 1–11. <http://dx.doi.org/10.1080/02713680500477347> 16421014.
- [46] M. Aslan, A. Cort, I. Yucel, Oxidative and nitrate stress markers in glaucoma, *Free Radical Biology and Medicine* 45 (4) (2008) 367–376. <http://dx.doi.org/10.1016/j.freeradbiomed.2008.04.026> 18489911.
- [47] L.M. Bova, M.H. Sweeney, Jamie F. Truscott, R.J. [!(%<nReflec:surname)!], Major changes in human ocular UV protection with age, *Investigative Ophthalmology and Vision Science* 42 (1) (2001) 200–205 11133868.
- [48] M.C. Moreno, J. Campanelli, P. Sande, D.A. Sáñez, M.I. Keller Sarmiento, R. E. Rosenstein, Retinal oxidative stress induced by high intraocular pressure, *Free Radical Biology and Medicine* 37 (6) (2004) 803–812. <http://dx.doi.org/10.1016/j.freeradbiomed.2004.06.001> 15384194.
- [49] D. Gherghel, H.R. Griffiths, E.J. Hilton, I.A. Cunliffe, S.L. Hosking, Systemic reduction in glutathione levels occurs in patients with primary open-angle glaucoma, *Investigative Ophthalmology and Vision Science* 46 (3) (2005) 877–883. <http://dx.doi.org/10.1167/jovs.04-0777> 15728543.
- [50] D. Gherghel, S. Mroczkowska, L. Qin, Reduction in blood glutathione levels occurs similarly in patients with primary-open angle or normal tension glaucoma, *Investigative Ophthalmology and Vision Science* 54 (5) (2013) 3333–3339. <http://dx.doi.org/10.1167/jovs.12-11256> 23599328.
- [51] A.P. Mazzetti, M.C. Fiorile, A. Primavera, M. Lo Bello, Glutathione transferases and neurodegenerative diseases, *Neurochemistry International* 82 (2015) 10–18. <http://dx.doi.org/10.1016/j.neuint.2015.01.008>.
- [52] A. Bagnis, A. Izzotti, M. Centofanti, S.C. Saccá, Aqueous humor oxidative stress proteomic levels in primary open angle glaucoma, *Experimental Eye Research* 103 (2012) 55–62. <http://dx.doi.org/10.1016/j.exer.2012.07.011> 22974818.
- [53] J. Yang, G. Tezel, R.V. Patil, C. Romano, M.B. Wax, Serum autoantibody against glutathione S-transferase in patients with glaucoma, *Investigative Ophthalmology and Vision Science* 42 (6) (2001) 1273–1276 11328739.
- [54] M. Almasieh, A.M. Wilson, B. Morquette, J.L. Cueva Vargas, A. Di Polo, The molecular basis of retinal ganglion cell death in glaucoma, *Progress in Retinal and Eye Research* 31 (2) (2012) 152–181. <http://dx.doi.org/10.1016/j.preteyeres.2011.11.002> 22155051.
- [55] E.B. Dreyer, D. Zurakowski, R.A. Schumer, S.M. Podos, S.A. Lipton, Elevated glutamate levels in the vitreous body of humans and monkeys with glaucoma, *Archives of Ophthalmology* 114 (3) (1996) 299–305. <http://dx.doi.org/10.1001/archophth.1996.01100130295012> 8600890.
- [56] Y. Dun, B. Mysona, T. Van Eils, L. Amarnath, M.S. Ola, V. Ganapathy, S. B. Smith, Expression of the cystine-glutamate exchanger (xc<sup>-</sup>) in retinal ganglion cells and regulation by nitric oxide and oxidative stress, *Cell Tissue Research* 324 (2) (2006) 189–202. <http://dx.doi.org/10.1007/s00441-005-0116-x> 16609915.
- [57] E.M. Allen, J.J. Mielay, Protein-thiol oxidation and cell death: regulatory role of glutaredoxins, *Antioxidants and Redox Signaling* 17 (12) (2012) 1748–1763. <http://dx.doi.org/10.1089/ars.2012.4644> 22530666.
- [58] A. García-García, L. Zavala-Flores, H. Rodríguez-Rocha, R. Franco, Thiol-redox signaling, dopaminergic cell death, and Parkinson's disease, *Antioxidants and Redox Signaling* 17 (12) (2012) 1764–1784. <http://dx.doi.org/10.1089/ars.2011.4501> 22369136.
- [59] E.A. Sabens Liedhegner, X.H. Gao, J.J. Mielay, Mechanisms of altered redox regulation in neurodegenerative diseases – focus on S-glutathionylation, *Antioxidants and Redox Signaling* 16 (6) (2012) 543–566. <http://dx.doi.org/10.1089/ars.2011.4119> 22066468.
- [60] K. Nomura, M. Lee, C. Banks, G. Lee, B.J. Morris, An ASK1-p38 signalling pathway mediates hydrogen peroxide-induced toxicity in NG108-15 neuronal cells, *Neuroscience Letters* 549 (2013) 163–167. <http://dx.doi.org/10.1016/j.neulet.2013.05.045> 23742763.
- [61] D.A. Drechsel, M. Patel, Respiration-dependent H<sub>2</sub>O<sub>2</sub> removal in brain mitochondria via the thioredoxin/peroxiredoxin system, *Journal of Biological Chemistry* 285 (36) (2010) 27850–27858. <http://dx.doi.org/10.1074/jbc.M110.101196> 20558743.
- [62] P. Lopert, B.J. Day, M. Patel, Thioredoxin reductase deficiency potentiates oxidative stress, mitochondrial dysfunction and cell death in dopaminergic cells, *PLoS One* 7 (11) (2012) e50683. <http://dx.doi.org/10.1371/journal.pone.0050683> 23226354.
- [63] P. Lopert, M. Patel, Nicotinamide nucleotide transhydrogenase (Nnt) links the substrate requirement in brain mitochondria for hydrogen peroxide removal to the thioredoxin/peroxiredoxin (Trx/Prx) system, *Journal of Biological Chemistry* 289 (22) (2014) 15611–15620. <http://dx.doi.org/10.1074/jbc.M113.533653> 24722990.
- [64] R.S. Kenchappa, V. Ravindranath, Glutaredoxin is essential for maintenance of brain mitochondrial complex I: Studies with MPTP, *FASEB Journal* 17 (6) (2003) 717–719. <http://dx.doi.org/10.1096/fj.02-0771> 12594173.
- [65] L. Diwakar, R.S. Kenchappa, J. Annepu, V. Ravindranath, Downregulation of glutaredoxin but not glutathione loss leads to mitochondrial dysfunction in female mice CNS: implications in excitotoxicity, *Neurochemistry International* 51 (1) (2007) 37–46. <http://dx.doi.org/10.1016/j.neuint.2007.03.008> 17512091.
- [66] D.W. Lee, D. Kaur, S.J. Chinta, S. Rajagopalan, J.K. Andersen, A disruption in iron-sulfur center biogenesis via inhibition of mitochondrial dithiol glutaredoxin 2 may contribute to mitochondrial and cellular iron dysregulation in mammalian glutathione-depleted dopaminergic cells: implications for Parkinson's disease, *Antioxidants and Redox Signaling* 11 (9) (2009) 2083–2094. <http://dx.doi.org/10.1089/ARS.2009.2489> 19290777.
- [67] S. Karunakaran, U. Saeed, S. Ramakrishnan, R.C. Koumar, V. Ravindranath, Constitutive expression and functional characterization of mitochondrial glutaredoxin (Grx2) in mouse and human brain, *Brain Research* 1185 (2007) 8–17. <http://dx.doi.org/10.1016/j.brainres.2007.09.019> 17961515.
- [68] U. Saeed, L. Durgados, R.K. Valli, D.C. Joshi, P.G. Joshi, V. Ravindranath, Knockdown of cytosolic glutaredoxin 1 leads to loss of mitochondrial membrane potential: implication in neurodegenerative diseases, *PLoS One* 3 (6) (2008) e2459. <http://dx.doi.org/10.1371/journal.pone.0002459> 18560520.
- [69] A. Ferri, P. Fiorenza, M. Nencini, M. Cozzolino, M.G. Pesaresi, C. Valle, S. Sepe, S. Moreno, M.T. Carrì, Glutaredoxin 2 prevents aggregation of mutant SOD1 in mitochondria and abolishes its toxicity, *Human Molecular Genetics* 19 (22) (2010) 4529–4542. <http://dx.doi.org/10.1093/hmg/ddq383> 20829229.
- [70] S.D. Bouldin, M.A. Darch, P.J. Hart, C.E. Outten, Redox properties of the disulfide bond of human Cu,Zn superoxide dismutase and the effects of human glutaredoxin 1, *Biochemical Journal* 446 (1) (2012) 59–67. <http://dx.doi.org/10.1042/BJ20120075> 22651090.
- [71] A.A. Brasil, A. Belati, S.C. Mannarino, A.D. Panek, E.C. Eleutherio, M.D. Pereira, The involvement of GSH in the activation of human Sod1 linked to FALS in chronologically aged yeast cells, *FEMS Yeast Research* 13 (5) (2013) 433–440. <http://dx.doi.org/10.1111/1567-1364.12045> 23496883.
- [72] U. Saeed, A. Ray, R.K. Valli, A.M. Kumar, V. Ravindranath, DJ-1 loss by glutaredoxin but not glutathione depletion triggers Daxx translocation and cell death, *Antioxidants and Redox Signaling* 13 (2) (2010) 127–144. <http://dx.doi.org/10.1089/ars.2009.2832> 20014998.
- [73] P. Lopert, M. Patel, Brain mitochondria from DJ-1 knockout mice show increased respiration-dependent hydrogen peroxide consumption, *Redox Biology* 2 (2014) 667–672. <http://dx.doi.org/10.1016/j.redox.2014.04.010> 24936441.
- [74] S. Akterin, R.F. Cowburn, A. Miranda-Vizueta, A. Jiménez, N. Bogdanovic, B. Winblad, A. Cedazo-Minguez, Involvement of glutaredoxin-1 and thioredoxin-1 in beta-amyloid toxicity and Alzheimer's disease, *Cell Death and Differentiation* 13 (9) (2006) 1454–1465. <http://dx.doi.org/10.1038/sj.cdd.4401818> 16311508.
- [75] E.A. Sabens, A.M. Distler, J.J. Mielay, Levodopa deactivates enzymes that regulate thiol-disulfide homeostasis and promotes neuronal cell death: implications for therapy of Parkinson's disease, *Biochemistry* 49 (12) (2010) 2715–2724. <http://dx.doi.org/10.1021/bi9018658> 20411169.
- [76] H. Masutani, J. Bai, Y.C. Kim, J. Yodoi, Thioredoxin as a neurotrophic cofactor and an important regulator of neuroprotection, *Molecular Neurobiology* 29 (3) (2004) 229–242. <http://dx.doi.org/10.1385/MN:29:3:229> 15181236.
- [77] J. Ungerstedt, Y. Du, H. Zhang, D. Nair, A. Holmgren, In vivo redox state of human thioredoxin and redox shift by the histone deacetylase inhibitor suberoylanilide hydroxamic acid (SAHA), *Free Radical Biology and Medicine* 53 (11) (2012) 2002–2007. <http://dx.doi.org/10.1016/j.freeradbiomed.2012.09.019> 23010496.
- [78] A. Holmgren, J. Lu, Thioredoxin and thioredoxin reductase: current research with special reference to human disease, *Biochemical and Biophysical Research Communications* 396 (1) (2010) 120–124. <http://dx.doi.org/10.1016/j.bbrc.2010.03.083> 20494123.
- [79] S.I. Hashemy, A. Holmgren, Regulation of the catalytic activity and structure of human thioredoxin 1 via oxidation and S-nitrosylation of cysteine



- residues, *Journal of Biological Chemistry* 283 (32) (2008) 21890–21898. <http://dx.doi.org/10.1074/jbc.M801047200.18544525>.
- [80] A. Holmgren, Thioredoxin, *Annual Review of Biochemistry* 54 (1985) 237–271. <http://dx.doi.org/10.1146/annurev.bi.54.070185.001321.3896121>.
- [81] C.H. Lillig, A. Holmgren, Thioredoxin and related molecules – from biology to health and disease, *Antioxidants and Redox Signaling* 9 (1) (2007) 25–47. <http://dx.doi.org/10.1089/ars.2007.9.25.17115886>.
- [82] G. Powis, W.R. Montfort, Properties and biological activities of thioredoxins, *Annual Review of Biophysics and Biomolecular Structure* 30 (2001) 421–455. <http://dx.doi.org/10.1146/annurev.biophys.30.1.421.11441809>.
- [83] K. Hirota, M. Matsui, S. Iwata, A. Nishiyama, K. Mori, J. Yodoi, AP-1 transcriptional activity is regulated by a direct association between thioredoxin and Ref-1, *Proceedings of the National Academy of Sciences of the United States of America* 94 (8) (1997) 3633–3638. <http://dx.doi.org/10.1073/pnas.94.8.3633.9108029>.
- [84] Y. Kabe, K. Ando, S. Hirao, M. Yoshida, H. Handa, Redox regulation of NF-kappaB activation: distinct redox regulation between the cytoplasm and the nucleus, *Antioxidants and Redox Signaling* 7 (3–4) (2005) 395–403. <http://dx.doi.org/10.1089/ars.2005.7.395.15706086>.
- [85] T.C. Zschauer, K. Kunze, S. Jakob, J. Haendeler, J. Altschmied, Oxidative stress-induced degradation of thioredoxin-1 and apoptosis is inhibited by thioredoxin-1-actin interaction in endothelial cells, *Arteriosclerosis, Thrombosis and Vascular Biology* 31 (3) (2011) 650–656. <http://dx.doi.org/10.1161/ATVBAHA.110.218982.21212402>.
- [86] J.Y. Im, K.W. Lee, E. Junn, M.M. Mouradian, DJ-1 protects against oxidative damage by regulating the thioredoxin/ASK1 complex, *Neuroscience Research* 67 (2010) 203–208. <http://dx.doi.org/10.1016/j.neures.2010.04.002.20385180>.
- [87] J. Waak, S.S. Weber, K. Görner, C. Schall, H. Ichijo, T. Stehle, P.J. Kahle, Oxidizable residues mediating protein stability and cytoprotective interaction of DJ-1 with apoptosis signal-regulating kinase 1, *Journal of Biological Chemistry* 284 (21) (2009) 14245–14257. <http://dx.doi.org/10.1074/jbc.M806902200.19293155>.
- [88] E. Andres-Mateos, C. Perier, L. Zhang, B. Blanchard-Fillion, T.M. Greco, B. Thomas, H.S. Ko, M. Sasaki, H. Ischiropoulos, S. Przedborski, T.M. Dawson, V.L. Dawson, DJ-1 gene deletion reveals that DJ-1 is an atypical peroxiredoxin-like peroxidase, *Proceedings of the National Academy of Sciences of the United States of America* 104 (37) (2007) 14807–14812. <http://dx.doi.org/10.1073/pnas.0703219104.17766438>.
- [89] M. Niso-Santano, R.A. González-Polo, J.M. Bravo-San Pedro, R. Gómez-Sánchez, I. Lastres-Becker, M.A. Ortiz-Ortiz, G. Soler, J.M. Morán, A. Cuadrado, J. M. Fuentes, Activation of apoptosis signal-regulating kinase 1 is a key factor in paraquat-induced cell death: modulation by the Nrf2/Trx axis, *Free Radical Biology and Medicine* 48 (10) (2010) 1370–1381. <http://dx.doi.org/10.1016/j.freeradbiomed.2010.02.024.20202476>.
- [90] S. Ramachandiran, J.M. Hansen, D.P. Jones, J.R. Richardson, G.W. Miller, Divergent mechanisms of paraquat, MPP<sup>+</sup>, and rotenone toxicity: oxidation of thioredoxin and caspase-3 activation, *Toxicological Sciences* 95 (1) (2007) 163–171. <http://dx.doi.org/10.1093/toxsci/kfl125.17018646>.
- [91] W. Yang, E. Tiffany-Castiglioni, H.C. Koh, I.H. Son, Paraquat activates the IRE1/ASK1/JNK cascade associated with apoptosis in human neuroblastoma SH-SY5Y cells, *Toxicology Letters* 191 (2–3) (2009) 203–210. <http://dx.doi.org/10.1016/j.toxlet.2009.08.024.19735704>.
- [92] Z. Liu, Y. Jing, J. Yin, J. Mu, T. Yao, L. Gao, Downregulation of thioredoxin reductase 1 expression in the substantia nigra pars compacta of Parkinson's disease mice, *Neural Regeneration Research* 8 (35) (2013) 3275–3283. <http://dx.doi.org/10.3969/j.issn.1673-5374.2013.35.002.25206649>.
- [93] X.S. Zeng, J.J. Jia, Y. Kwon, S.D. Wang, J. Bai, The role of thioredoxin-1 in suppression of endoplasmic reticulum stress in Parkinson disease, *Free Radical Biology and Medicine* 67 (2014) 10–18. <http://dx.doi.org/10.1016/j.freeradbiomed.2013.10.013.24140863>.
- [94] Y. Umeda-Kameyama, M. Tsuda, C. Ohkura, T. Matsuo, Y. Namba, Y. Ohuchi, T. Aigaki, Thioredoxin suppresses Parkin-associated endothelin receptor-like receptor-induced neurotoxicity and extends longevity in *Drosophila*, *Journal of Biological Chemistry* 282 (15) (2007) 11180–11187. <http://dx.doi.org/10.1074/jbc.M700937200.17301052>.
- [95] L. Meng, J.H. Wong, L.J. Feldman, P.G. Lemaux, B.B. Buchanan, A membrane-associated thioredoxin required for plant growth moves from cell to cell, suggestive of a role in intercellular communication, *Proceedings of the National Academy of Sciences of the United States of America* 107 (8) (2010) 3900–3905. <http://dx.doi.org/10.1073/pnas.0913759107.20133584>.
- [96] R.B. Carilho Torrao, I.H. Dias, S.J. Bennett, C.R. Dunston, H.R. Griffiths, Healthy ageing and depletion of intracellular glutathione influences T cell membrane thioredoxin-1 levels and cytokine secretion, *Chemistry Central Journal* 7 (1) (2013) 150. <http://dx.doi.org/10.1186/1752-153X-7-150.24007191>.
- [97] L. Arohin, H. Lamparter, H. Karlsson, I. Nennesmo, M. Björnstedt, J. Schröder, A.P. Fernandes, Alteration of thioredoxin and glutaredoxin in the progression of Alzheimer's disease, *Journal of Alzheimer's Disease* 39 (4) (2014) 787–797. <http://dx.doi.org/10.3233/JAD-131814.24270206>.
- [98] Y. Munemasa, J.H. Ahn, J.M. Kwong, J. Caprioli, N. Piri, Redox proteins thioredoxin 1 and thioredoxin 2 support retinal ganglion cell death in glaucoma, *Gene Therapy* 16 (2009) 17–25.
- [99] F. Hattori, S. Oikawa, Peroxiredoxins in the central nervous system, *Subcellular Biochemistry* 44 (2007) 357–374. [http://dx.doi.org/10.1007/978-1-4020-6051-9\\_17.18084903](http://dx.doi.org/10.1007/978-1-4020-6051-9_17.18084903).
- [100] J. Yao, M. Taylor, F. Davey, Y. Ren, J. Aiton, P. Coote, F. Fang, J.X. Chen, S.D. Yan, F.J. Gunn-Moore, Interaction of amyloid binding alcohol dehydrogenase/Abeta mediates up-regulation of peroxiredoxin II in the brains of Alzheimer's disease patients and a transgenic Alzheimer's disease mouse model, *Molecular and Cellular Neuroscience* 35 (2) (2007) 377–382. <http://dx.doi.org/10.1016/j.mcn.2007.03.013.17490890>.
- [101] L.M. Randall, B. Manta, M. Hugo, M. Gil, C. Batthyány, M. Trujillo, L.B. Poole, A. Denicola, Nitration transforms a sensitive peroxiredoxin 2 into a more active and robust peroxidase, *Journal of Biological Chemistry* 289 (22) (2014) 15536–15543. <http://dx.doi.org/10.1074/jbc.M113.539213.24719319>.
- [102] J. Fang, T. Nakamura, D.H. Cho, Z. Gu, S.A. Lipton, S-nitrosylation of peroxiredoxin 2 promotes oxidative stress-induced neuronal cell death in Parkinson's disease, *Proceedings of the National Academy of Sciences of the United States of America* 104 (47) (2007) 18742–18747. <http://dx.doi.org/10.1073/pnas.0705904104.18003920>.
- [103] I.K. Kim, K.J. Lee, S. Rhee, S.B. Seo, J.H. Pak, Protective effects of peroxiredoxin 6 overexpression on amyloid beta-induced apoptosis in PC12 cells, *Free Radical Research* 47 (10) (2013) 836–846. <http://dx.doi.org/10.3109/10715762.2013.833330.23937564>.
- [104] H.M. Yun, P. Jin, J.Y. Han, M.S. Lee, S.B. Han, K.W. Oh, S.H. Hong, E.Y. Jung, J. T. Hong, Acceleration of the development of Alzheimer's disease in amyloid beta-infused peroxiredoxin 6 overexpression transgenic mice, *Molecular Neurobiology* 48 (3) (2013) 941–951. <http://dx.doi.org/10.1007/s12035-013-8479-6.23771816>.
- [105] Y.M. Lee, S.H. Park, D.I. Shin, J.Y. Hwang, B. Park, Y.J. Park, T.H. Lee, H.Z. Chae, B.K. Jin, T.H. Oh, Y.J. Oh, Oxidative modification of peroxiredoxin is associated with drug-induced apoptotic signaling in experimental models of Parkinson disease, *Journal of Biological Chemistry* 283 (15) (2008) 9986–9998. <http://dx.doi.org/10.1074/jbc.M800426200.18250162>.
- [106] X. Hu, Z. Weng, C.T. Chu, L. Zhang, G. Cao, Y. Gao, A. Signore, J. Zhu, T. Hastings, J.T. Greenamyre, J. Chen, Peroxiredoxin-2 protects against 6-hydroxydopamine-induced dopaminergic neurodegeneration via attenuation of the apoptosis signal-regulating kinase (ASK1) signaling cascade, *Journal of Neuroscience* 31 (1) (2011) 247–261. <http://dx.doi.org/10.1523/JNEUROSCI.4589-10.2011.21209210>.
- [107] D.C. Angeles, B.H. Gan, L. Onstead, Y. Zhao, K.L. Lim, J. Dachselt, H. Melrose, M. Farrer, Z.K. Wszolek, D.W. Dickson, E.K. Tan, Mutations in LRRK2 increase phosphorylation of peroxiredoxin 3 exacerbating oxidative stress-induced neuronal death, *Human Mutation* 32 (12) (2011) 1390–1397. <http://dx.doi.org/10.1002/humu.21582.21850687>.
- [108] W.M. Johnson, C. Yao, S.L. Siedlak, W. Wang, X. Zhu, G.A. Caldwell, A. L. Wilson-Delfosse, J.J. Mielay, S.G. Chen, Glutaredoxin deficiency exacerbates neurodegeneration in *C. elegans* models of Parkinson's disease, *Human Molecular Genetics* 24 (5) (2015) 1322–1335. <http://dx.doi.org/10.1093/hmg/ddu542.25355420>.
- [109] J.H. Power, S. Asad, T.K. Chataway, F. Chegini, J. Manavis, J.A. Temlett, P. H. Jensen, P.C. Blumbergs, W.P. Gai, Peroxiredoxin 6 in human brain: molecular forms, cellular distribution and association with Alzheimer's disease pathology, *Acta Neuropathology* 115 (6) (2008) 611–622. <http://dx.doi.org/10.1007/s00401-008-0373-3.18386021>.
- [110] Y. Yoshida, A. Yoshikawa, T. Kinumi, Y. Ogawa, Y. Saito, K. OHara, H. Yamamoto, Y. Imai, E. Niki, Hydroxyoctadecadienoic acid and oxidatively modified peroxiredoxins in the blood of Alzheimer's disease patients and their potential as biomarkers, *Neurobiology of Aging* 30 (2) (2009) 174–185. <http://dx.doi.org/10.1016/j.neurobiolaging.2007.06.012.17688973>.