

1 WATER, AIR AND SOIL POLLUTION

2

3 **Effects of single, binary and quinary mixtures of phenanthrene and its N-PAHs on *Eisenia fetida* in soil**

4

5 **Ihuoma N. Anyanwu • Ojerime I. Clifford • Kirk T. Semple**

6

7 Ihuoma N. Anyanwu (✉) • Ojerime I. Clifford • Kirk T. Semple

8 Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom

9

10 I. N. Anyanwu (✉)

11 Department of Biological Sciences, Federal University Ndufu-Alike Ikwo, P.M.B 1010, Abakaliki, Ebonyi

12 State, Nigeria

13

14 (✉) Corresponding author:

15 [E-mail: ihuomal@yahoo.com](mailto:ihuomal@yahoo.com)

16

17

18

19

20

21

22

23 **Abstract** It is now acknowledged that aromatic hydrocarbons present in contaminated soils occur in mixtures.  
24 The effect of single, binary and quinary mixtures of phenanthrene and selected N-PAHs were investigated on  
25 the survival, growth and behavioural index of earthworms (*Eisenia fetida*) over a 21 d incubation in soil. The  
26 results showed that the LC<sub>50</sub> values ranged from (not detected) ND - 329.3 mg kg<sup>-1</sup> (single mixture), ND – 219.8  
27 mg kg<sup>-1</sup> (binary mixtures) and 148.4 mg kg<sup>-1</sup> (quinary mixture), while the EC<sub>50</sub> values (based on weight loss)  
28 ranged from 13.3 - 148.4 mg kg<sup>-1</sup> (single mixture), 63.8 - 148.4 mg kg<sup>-1</sup> (binary mixture) and 24.2 mg kg<sup>-1</sup>  
29 (quinary mixture). Greater impacts were recorded where N-PAHs are present with phenanthrene. Further,  
30 behavioural index of *E. fetida* was affected after 24 h exposure to N-PAH amended soils. Among the N-PAHs  
31 however, benzo[h]quinoline recorded the greatest impact on the survival, growth and behavioural index of *E.*  
32 *fetida* in soil. Findings from this study showed that 3 ring-N-PAHs are more toxic than phenanthrene as  
33 expected from their physico-chemical properties. Binary and quinary mixtures of phenanthrene and N-PAHs in  
34 soil intensified toxicity, suggesting that PAHs-N-PAHs mixtures represent greater risk to soil biota.

35

36 **Keywords** Phenanthrene • Nitrogen-containing PAHs • Toxicity • Behavioural index • Soil • Earthworm.

37

38

39

40

41

42

43

44

45

46

47

## 48 **1 Introduction**

49 Polycyclic aromatic hydrocarbons (PAHs) are a key group of contaminants present at many contaminated sites  
50 (e.g. ex-industrial sites, oil contaminated areas) (Brumley et al. 1991; Sverdrup et al. 2002; EC 2011; IARC  
51 2012; Anyanwu and Semple 2016b). What is less well described is that their nitrogen-containing polycyclic  
52 aromatic hydrocarbons (N-PAHs) can also be present, often at high concentrations (De Voogt and Laane 2009).  
53 Many of these compounds are known to be toxic, carcinogenic, mutagenic and genotoxic, thus, are of concern to  
54 both biota and human exposure (Bleeker et al. 2002; Sverdrup et al. 2002; Kobetičová et al. 2008; Brar et al.  
55 2010; IARC 2012; Anyanwu and Semple 2015a, b, c; 2016a, b). In addition, their toxicity cannot be determined  
56 by the number of aromatic rings alone (IARC 2012). N-PAHs contain one or more nitrogen atom(s) in place of  
57 carbon atom (Table 1). Due to the substitution of N-atom into one or more of the aromatic rings, and the  
58 resulting physico-chemical properties, they differ in terms of persistence, mobility, bioavailability and toxicity  
59 compared to analogue PAHs (Sverdrup et al. 2002; Kobetičová et al. 2008; Anyanwu et al. 2013; Anyanwu and  
60 Semple 2015a; 2016b). For example, the substitution with N-atom(s) makes N-PAHs more polar, mobile and  
61 soluble in the soil environment, and as such, putatively more toxic (Sverdrup et al. 2002; Kobetičová et al. 2008;  
62 Anyanwu and Semple 2015 c; 2016a, b).

63         Apart from their environmental occurrence, N-PAHs are discharged into soil from anthropogenic  
64 sources such as petroleum related activities, oil spills and combustion processes (Brumley et al. 1991; Webber  
65 1994; Švábenský et al. 2009). Previously, N-PAH levels were reported to be 1-10 orders of magnitude lower  
66 than those of the homocyclic analogues, however, De Voogt and Laane (2009) measured concentrations ranging  
67 from similar to equal to one order of magnitude higher than their homocyclic analogues in sediments.  
68 Furthermore, studies have reported that N-PAH chemicals are phyto-toxic, inhibitors to soil microflora and have  
69 cellular autolytic impact (Willumsen et al. 2001; 2005; Anyanwu and Semple 2015b; c; 2016a, b). In addition, it  
70 has been reported that losses of PAH degradation capacity in contaminated soil may be due to the existence of  
71 N-PAHs (Meyer and Steinhart 2000). Irrespective of this, only few studies have investigated N-PAH toxicity on  
72 soil invertebrates (Sverdrup et al. 2002; Kobetičová et al. 2008; 2011) and their observed effects ranged from  
73 (LC<sub>50</sub> (not detected) ND - <2000 mg kg<sup>-1</sup> and EC<sub>50</sub> 94 - 1033 mg k<sup>-1</sup>) (Sverdrup et al. 2002; Kobetičová et al.  
74 2008; 2011). Although there are some toxicity data for a few selected N-PAH compounds, all the investigations  
75 focused on the traditional Organization for Economic Cooperation and Development (OECD) methodologies of  
76 percentage survival / effects judgements (OECD, 1984) without considering other criteria, such as the

77 behavioural index (general physical condition) of the organisms during exposure. Monitoring of behavioural  
78 index of biota in contaminated soils allows assessment of possible impact of various contaminants on different  
79 invertebrate species.

80 Generally, contaminated soils contain complex mixtures of compounds that differ in their physico-  
81 chemical properties and toxicity to soil biota. Thus, effects of range of aromatic compounds can be intensified  
82 by the presence of other compounds. However, the effect of binary and quinary mixtures of phenanthrene and its  
83 N-PAH analogues on soil biota have not been reported in literature. Since information on toxicity is essential for  
84 effective assessment of poorly managed chemicals and/or chemical groups for environmental risk assessment,  
85 this study therefore investigated the effect of single, binary and quinary mixtures of phenanthrene and its  
86 structurally similar nitrogen-containing analogues on mature earthworms (*Eisenia fetida*) in soil using OECD  
87 (1984) guidelines (with little deviation) and behavioural index tool.

88

## 89 **2 Materials and methods**

### 90 2.1 Chemicals

91 Phenanthrene (Phen), 1,10-phenanthroline (1,10-Phen), 1,7-phenanthroline (1,7-Phen), 4,7-phenanthroline (4,7-  
92 Phen) and benzo[h]quinoline (B[h]Q) were purchased from Sigma Aldrich Company Ltd, UK (Table 1).

93

### 94 2.2 Test organisms

95 Mature earthworms (*Eisenia fetida*) were purchased from Blades Biological Ltd, UK.

96

### 97 2.3 Soil preparation

98 Soil (without contamination history) from Myerscough Agricultural College in Lancashire, UK was collected  
99 from the top layer of field under pasture (from a depth of approximately 5 - 20 cm). The soil was sandy-loam  
100 (19.5% clay, 60.4% sand, 20.0% silt) with an organic matter content of 2.7% and pH 6.5 (Doick et al. 2003).  
101 The soil was air dried at room temperature, sieved with 2 mm mesh size, and rehydrated with deionised water  
102 back to original water holding capacity (WHC). Spiking procedure followed those described in Doick et al.

103 (2003). One third of soil ( $\frac{1}{3}$ ; 100 g), placed in a bowl, were amended with dissolved chemical standards  
104 containing acetone (10 ml) to give concentrations of 10, 100 and 500 mg kg<sup>-1</sup>. The soils were left to evaporate  
105 for 4 hours in the fume hood; after which soils were mixed with the remaining  $\frac{2}{3}$  (200 g) soil and amended to  
106 80% WHC with deionized water. Control samples were prepared using soil amended with acetone only. After  
107 amendment, soils (50 g) were weighed into glass jars and the effects on *E. fetida* was measured for 21 d.  
108 Recoveries of tested compounds are at a range of 49.50 ± 0.55% and 104.72 ± 8.00%, except for 1,10-  
109 Phenanthroline which was not determined since it could not be detected by GC-MS

#### 110 2.4 Earthworm toxicity assay

111 Determination of the effects of the N-PAHs on soil biota was carried out using the earthworm toxicity assay  
112 (OECD, 1984) (with little deviation). Mature *E. fetida* (0.3 – 0.4 g) were selected for the bioassay. Prior to test,  
113 the earthworms were depurated for 24 h in petri dishes containing moist filter paper, after which the earthworms  
114 were weighed. Soils (50 g) were used for the assay. Three exposure concentrations (10, 100 and 500 mg kg<sup>-1</sup>)  
115 were used for each single, binary and quinary mixture, as well as control samples (0 mg kg<sup>-1</sup>). Five replicates  
116 were used for each concentration. The soils were mixed thoroughly and single earthworm was added to each  
117 replicate. Samples were covered with perforated lids and incubated at 12 ± 2°C. Mortality, biomass and  
118 behavioural index were the test parameters. Assessment of behavioural index (0 - 2) was carried out after 0, 1, 3,  
119 5, 7, 14 and 21 d, with 0 = dead, 1 = moribund and 2 = healthy (Langdon et al. 1999; Anyanwu and Semple  
120 2016b). Earthworms were presumed dead if no contractile response was observed when prodded lightly with a  
121 blunt probe; moribund when weak contractile was observed; and healthy when sharp and vigorous response was  
122 observed (Langdon et al. 1999; Anyanwu and Semple 2016b). The depurated earthworms were weighed before  
123 and after exposure to determine changes in weight (biomass). Other physical changes on different parts of the  
124 earthworms were also recorded.

125

#### 126 2.5 Statistical analysis

127 The mean weight losses of the earthworms for the replicates were used as a measure of biomass change. The  
128 concentration that caused weight loss as compared to control values were calculated on the basis of initial  
129 measurement and ANOVA was used to determine the significant impact on biomass. Differences are found to  
130 be statistically significant when p<0.05. Further, correlations were determined between the exposure  
131 concentrations and changes in biomass using Pearsons correlation. Estimation of the concentration of the

132 chemicals in amended soils that caused 50% mortality ( $LC_{50}$ ), and the concentration that provoked a response  
133 halfway between the baseline and maximum ( $EC_{50}$ ) (i.e. weight-reduction) during the test period were  
134 determined by probit analysis in SPSS 20 software package. Graphs were plotted with SigmaPlot 10.0 version.

135

### 136 3 Results

#### 137 3.1 Assessment of the aromatic hydrocarbons on the behavioural index of *E. fetida* in soil

138 Apart from the traditional OECD guidelines relating to survival / effect measurements, the study investigated  
139 the general health condition of *E. fetida* during exposure. Fig. 1 shows the behavioural indices of *E. fetida*  
140 exposed to single, binary and quinary mixtures of phenanthrene and its N-PAH analogues in soil. The 21 d  
141 assessment of behavioural index (0 - 2) showed that all the compounds negatively impacted on the health of *E.*  
142 *fetida* at the highest concentration of 500 mg kg<sup>-1</sup> ( $p < 0.05$ ;  $r^2 = 0.982$ ), with the exception of 1,10-Phen, 4,7-  
143 Phen (single mixture) and 1,10-Phen + Phen (binary mixture). Among the N-PAHs, benzo[h]quinoline (single  
144 mixture) exhibited greater impact on the general wellbeing of *E. fetida* in soil. For example, it was observed that  
145 *E. fetida* became moribund after few hours of exposure to B[h]Q soils and experienced mortality at 3 d in the  
146 500 mg kg<sup>-1</sup> amendments (Fig. 1).

147 However, 4,7-phenanthroline recorded significant effect on the behavioural index of *E. fetida* in the  
148 presence of phenanthrene (binary mixture) at the 500 mg kg<sup>-1</sup> amendments ( $p < 0.05$ ) (Fig. 1). In addition,  
149 benzo[h]quinoline toxicity was slightly reduced in the presence of phenanthrene because, *E. fetida* experienced  
150 moribund phase of 5 d in health prior to mortality. Furthermore, the data showed that quinary mixtures of  
151 phenanthrene and N-PAHs significantly affected the behavioural index of the exposed earthworms even at 100  
152 mg kg<sup>-1</sup> ( $p < 0.05$ ) (Fig. 1). Other observed health effect includes; breakage of clitellum, excretion of yellowish  
153 fluid and sores.

154

#### 155 3.2 Assessment of the aromatic hydrocarbons on the mortality ( $LC_{50}$ ) of *E. fetida* in soil

156 During the 21 d exposure, neither mortality nor significant weight losses were observed in the control  
157 incubations. The survival (%) of *E. fetida* exposed to single, binary and quinary mixtures of phenanthrene and  
158 its nitrogen-containing analogues are shown in Fig. 2. From the result, significant mortality rates were recorded

159 in the 500 mg kg<sup>-1</sup> amendments (p<0.05). For example, the concentration–response graph shows that survival  
160 (%) of the exposed earthworms ranged from 0% – 20% in soils amended with 500 mg kg<sup>-1</sup> chemicals, with the  
161 exception of 1,10-Phen, 4,7-Phen (single mixture) and 1,10-Phen + phenanthrene (binary mixture) (Fig. 2). In  
162 addition, *E. fetida* suffered high mortality (100%) in the 500 mg kg<sup>-1</sup> B[h]Q amendment (Fig. 2a).

163 Furthermore, 100% mortality were observed in the 500 mg kg<sup>-1</sup> phenanthrene + 4,7-phenantrolone  
164 amendment (binary mixture) (p<0.05). In quinary mixtures however, *E. fetida* recorded 20% – 100% mortality  
165 in the 100 mg kg<sup>-1</sup> and 500 mg kg<sup>-1</sup> amendments, respectively (Fig. 2b). From the result, a trend of increased  
166 mortality with increase in concentration was observed compared to the control soil (p<0.05). The toxicity data  
167 (LC<sub>50</sub>) of single, binary and quinary mixtures of phenanthrene and selected N-PAHs on *E. fetida* after 21 d  
168 incubation in soil are summarized in Table 2. From the data, the LC<sub>50</sub> ranged from ND – 329.3 mg kg<sup>-1</sup>, with  
169 quinary mixture recording the lowest value of 148.4 mg kg<sup>-1</sup> (Table 2).

170

### 171 3.3 Assessment of the aromatic hydrocarbons on the biomass (EC<sub>50</sub>) of *E. fetida* in soil

172 Impact of the chemicals on the biomass of the exposed earthworms was measured (Table 2). The biomass of  
173 dead earthworms was not determined and therefore were assigned zero (0). Figs 3 – 4 shows the weight-changes  
174 (biomass) of *E. fetida* exposed to single, binary and quinary mixtures of phenanthrene and its N-PAH analogues  
175 in soil. From the result, all the aromatics showed significant biomass-effect on *E. fetida*, and there was a trend of  
176 decrease in biomass with increase in chemical concentrations. The concentration–biomass–effect plots recorded  
177 notable decline in the weight of the earthworms after 21 d exposure to the aromatic hydrocarbons (Fig. 3).  
178 Biomass reductions were observed to be pronounced in the N-PAH amendments, especially 1,7-Phen and B[h]Q  
179 (single mixture), and in the binary and quinary mixtures of phenanthrene + N-PAHs (p<0.05) (Figs. 3 – 4).

180 The EC<sub>50</sub> measurement of single, binary and quinary mixtures of phenanthrene and its structurally  
181 similar N-PAHs on the biomass of *E. fetida* after 21 d incubation are summarized in Table 2. The calculated  
182 EC<sub>50</sub> values ranged from 13.3 – 148.4 mg kg<sup>-1</sup>. Among the chemical treatments, B[h]Q recorded the lowest EC<sub>50</sub>  
183 of 17.00 mg kg<sup>-1</sup> (Table 2). Analysis of data using ANOVA showed statistically significant difference in the  
184 earthworms biomass (weight before exposure and weight after exposure) in all the chemicals, at all the  
185 concentrations (p<0.05). Further analysis of data showed high negative correlation between the concentrations  
186 and biomass after exposure (p<0.05) in all the chemicals, with the exception of 1,10-Phen and 4,7-Phen which  
187 showed medium but non-statistically significant relationships (p>0.05) (Table 2).

188

#### 189 **4 Discussion**

190 Despite the majority of published studies having focused on PAHs and to a lesser extent their heterocyclic  
191 analogues, the toxicity data relating to *Eisenia fetida* are still fragmented, with most of the studies investigating  
192 single compounds and artificial soil (Sverdrup et al. 2002; Kobetičová et al. 2008; Kobetičová et al. 2011;  
193 Anyanwu and Semple 2016b). This study brings to focus information on the behavioural index, survival and  
194 biomass-effect of *E. fetida* following exposure to single, binary and quinary mixtures of phenanthrene and its  
195 structurally similar N-PAHs.

196 The endpoint of an earthworm toxicity test is normally a measure of mortality; however, mortality is  
197 unlikely to be the most sensitive parameter for risk assessment (Langdon et al. 1999; Anyanwu et al. 2013;  
198 Anyanwu and Semple 2016b). This current study showed that most of the health effects, such as healthy (2) –  
199 moribund (1) – mortality (0) and antagonistic interactions occurred within <24 h – 3 d of exposure in the N-  
200 PAHs amended soils with particular reference to benzo[h]quinoline. This pattern of mortality and/or interaction  
201 may be attributed more to uptake of the chemical(s) across the dermis rather than through ingestion. This is  
202 because, earthworms added to soils amended with 500 mg kg<sup>-1</sup> N-PAHs were initially healthy and very active,  
203 but became moribund after a few hours and died within 3 d. This phenomenon has also been reported for the  
204 highest concentration of metal and aromatic hydrocarbon amended soils (Spurgeon et al. 1994; Anyanwu and  
205 Semple 2016b). Presumably, as earthworm moves through the soil, they are exposed to N-PAHs, which may  
206 lead to absorption of the chemicals across the dermis. This further showed that earthworms were particularly  
207 sensitive to N-PAHs compared to homocyclic phenanthrene analogue. Furthermore, other behavioral response  
208 such as lack of burrowing into the amended soil and avoidance (moving away from the amended soil) was  
209 observed at the highest concentration (500 mg kg<sup>-1</sup>).

210 In this current study, benzo[h]quinoline was noted to be the most toxic chemical to *E. fetida*. It may be  
211 that N-PAHs such as benzo[h]quinoline, having log K<sub>ow</sub> of 3.43 and log K<sub>oc</sub> of 4.32, may be existing more in the  
212 aqueous phase (even though some may be associated with soil organic matter), hence, undergoing rapid  
213 absorption and thus, triggering physical injury to the dermis of the earthworms as observed by the formation of  
214 lesions, breakage of the clitellum, sores, loss of weight and eventually death (Broholm et al. 1999; Anyanwu and  
215 Semple 2015b).



216 Variations in aromatics toxicity were observed. The variations in toxicity could be ascribed to the  
217 differences in physico-chemical properties and N-position (Anyanwu et al. 2013; Anyanwu and Semple 2015a,  
218 b, c; 2016 a, b). Furthermore, variability was measured among 1,7-phen, 1,10-phen and 4,7-phen irrespective of  
219 their similarities in physico-chemical properties. Molecular structure and/or chemical bioavailability may be  
220 attributed in this study. From the results, the recorded LC<sub>50</sub> and EC<sub>50</sub> values for phenanthrene are in agreement  
221 with the values reported by Sverdrup et al. (2002) (Table 3). However, the values (LC<sub>50</sub> and EC<sub>50</sub>) were lower  
222 than those reported by Kobetičová et al. (2008; 2011); soil physico-chemical properties and/or species  
223 differences may be attributable. Further, the absence of mortality recorded in the 1,10-Phen amendment (in the  
224 study) is in agreement with the findings of Sochová et al. (2007), who reported 1,10-Phen as the least toxic  
225 chemical to nematode (*Caenorhabditis elegans*) in soil. However, the observation was different from the reports  
226 of Kobetičová et al. (2008; 2011), who recorded high toxicity for 1,10-Phen; chemical concentrations and/or  
227 media (type of matrix) may be ascribed. In support, Anderson et al. (1999) reported that during chemical-biota-  
228 interactions, the receptor, route of entry, test duration and the matrix containing the contaminant need to be  
229 considered. Therefore, with a soil organic matter content of 2.7% and pH 6.5, the toxicity and speciation of  
230 1,10-Phen may have been influenced (in this study). Similarly, (although not in the soil environment) the  
231 observation with 4,7-Phen (single mixture) is in agreement with Feldmannová et al. (2006), who recorded 4,7-  
232 Phen as the least toxic compound on the survival, fecundity and reproduction of *Daphnia magna*.

233 It has been reported that biomass of organisms (such as earthworms) can be impaired by: (i) direct  
234 toxic effects on the physiology of exposed organism, (ii) changes in the body function as the organism tries to  
235 prevent accumulation in the biological membranes and/or (iii) avoidance due to lack of feeding. Thus, the  
236 significant effect on biomass (weight) of *E. fetida* recorded in this current study may be ascribed to toxicity  
237 and/or changes in body function. However, it should be noted that the toxicological effects of N-PAHs could  
238 cause reductions in weight (Sverdrup et al. 2002; Anyanwu et al. 2013; Anyanwu and Semple 2016b). In  
239 addition, Widdows and Donkin (1989) reported changes in the individual energy budget as an organism (such as  
240 earthworm) spends energy resisting the contaminant by avoidance, elimination or reluctant to feed in a polluted  
241 environment. According to their report, the extra energy requirement will decrease the capacity for growth and  
242 development of the organism. This suggests that the effect on biomass of the exposed earthworms is as a result  
243 of changes in the energy budget as they strive to resist bioaccumulation in the site of biological response, or  
244 weight loss through lack of feeding and/or N-PAHs toxicity. Furthermore, the study showed that the biomass–

245 effect ratios increased with increase in N-PAHs burden on the earthworms as demonstrated by the correlation  
246 relationships; indicating greater toxicity of N-PAHs to soil biota.

247

## 248 **5 Conclusions**

249 In this study, the effects of single, binary and quinary mixtures of phenanthrene and its structurally similar N-  
250 PAHs were investigated since mixtures of contaminants are found in contaminated soils. The results showed  
251 that single N-PAHs possess high ecotoxicity risk as expected of their solubility and lower  $K_{ow}$  values, while  
252 mixtures of phenanthrene and N-PAHs in contaminated sites represent greater risks to soil biota. However,  
253 studies are required on the impact of N-PAHs in chronically contaminated / aged soil.

254

## 255 **Acknowledgment**

256 The authors are grateful to Petroleum Technology Development Fund (PTDF), Nigeria for providing financial  
257 support.

258

## 259 **References**

- 260 Anderson, W.C., Loehr, R.C., & Smith, B. P. (1999). Environmental availability of chlorinated organics,  
261 explosives and metals in soils. Annapolis, MD: *American Academy of Environmental Engineer.*
- 262 Anyanwu, I.N., & Semple, K.T. (2016a). Assessment of the effects of phenanthrene and its nitrogen  
263 heterocyclic analogues on microbial activity in soil. *SpringerPlus* 5, 279.
- 264 Anyanwu, I.N., & Semple, K.T. (2016b). Effects of phenanthrene and its nitrogen-heterocyclic analogues aged  
265 in soil on the earthworm *Eisenia fetida*. *Applied Soil Ecology* 105, 151-159.
- 266 Anyanwu, I.N., & Semple, K.T. (2015a). Fate and behaviour of nitrogen-containing polycyclic aromatic  
267 hydrocarbons in soil. *Environmental Technology and Innovation* 3, 108-120
- 268 Anyanwu, I.N., & Semple, K.T. (2015b). Biodegradation of phenanthrene-nitrogen-containing analogues in soil.  
269 *Water, Air and Soil Pollution* 226, 252.
- 270 Anyanwu, I.N., & Semple, K.T. (2015c). Phytotoxicity of phenanthrene and its nitrogen polycyclic aromatic  
271 hydrocarbon analogues in ageing soil. *Water, Air and Soil Pollution* 226, 347.

272 Anyanwu, I.N., Clifford, O.I., & Semple, K.T. (2013). Effects of three ringed PAH and N-PAHs on earthworm  
273 (*E. fetida*) in soil. In: *Proceedings of the International Conference on Environmental Pollution and*  
274 *Remediation*. International Academy of Science Engineering and Technology, Toronto (Ontario,  
275 Canada), pp. 57.

276 Bleeker, E.A.J., Wiegman, S., Kraak, M., de Voogt, P., Leslie, H.A., Haas, E., & Admiraal, W. (2002). Toxicity  
277 of azaarenes. *Reviews of Environmental Contamination and Toxicology* 173, 39-83.

278 Brar, N.K., Waggoner, C., Reyes, J.A., Fairey, R., Kelley, & K.M. (2010). Evidence for thyroid endocrine  
279 disruption in wild fish in San Francisco Bay, California, USA. Relationships to contaminant exposures.  
280 *Aquatic Toxicology* 93, 203-2015.

281 Broholm, M.M., Broholm, K., & Arvin, E. (1999). Sorption of heterocyclic compounds on natural clay till.  
282 *Journal of Contaminant Hydrology*, 39, 183-200.

283 Brumley, W.C., Brownrigg, C.M., & Brilis, G.M. (1991). Characterization of nitrogen-containing aromatic  
284 compounds in soil and sediment by capillary gas chromatography-mass spectrometry after  
285 fractionation. *Journal of Chromatography* 558, 223-233.

286 Environment Canada 2011. Risk management scope for quinoline. Chemical Abstract Service Registry Number  
287 (CAS RN), 91-22-5: <http://www.ec.gc.ca/ese-ees/default.asp?lang>.

288 De Voogt, P., & Laane, R.W.P.W. (2009). Assessment of azaarenes and azaarones (oxidized azaarene  
289 derivatives) in the Dutch coastal zone of the North Sea. *Chemosphere*, 76, 1067-1074

290 Doick, K.J., Lee, P.H., & Semple, K.T. (2003). Assessment of spiking procedures for the introduction of  
291 phenanthrene-LNAPL mixture into field-wet soil. *Environmental Pollution* 126, 399-406.

292 Feldmannová, M., Hilscherová, K., Marsálek, B., & Bláha, L. (2006). Effects of N-heterocyclic polyaromatic  
293 hydrocarbons on survival, reproduction and biochemical parameter in *Daphnia magna*.  
294 *Environmental Toxicology and Chemistry* 21, 425-431.

295 IARC 2012. In: IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Agents  
296 classified by IARC Monographs. *World Health Organization* 1-105.

297 Kobetičová, K., Bezchlebová, J., Lána, J., Sochová, I., & Hofman, J. (2008). Toxicity of four nitrogen  
298 heterocyclic polyaromatic hydrocarbons (NPAHs) to soil organisms. *Ecotoxicology and Environmental*  
299 *Safety* 71, 650-660.

300 Kobetičová, K., Šimek, Z., Brezovský, B., & Hofman, J. (2011). Toxic effects of nine polycyclic aromatic  
301 compounds on *Enchytraeus crypticus* in artificial soil in relation to their properties. *Ecotoxicology and*  
302 *Environmental Safety* 74, 1727-1733.

303 Langdon, C.J., Pearce, T.G., Black, S., & Semple, K.T. (1999). Resistance of arsenic toxicity in a population of  
304 earthworm *Lumbricus rubella*. *Soil Biology and Chemistry* 31, 1963-1967

305 Meyer, S., & Steinhart, H. (2000). Effects of heterocyclic PAHs (N, S, O) on the biodegradation of typical tar  
306 oil PAHs in a soil/compost mixture. *Chemosphere* 40, 357-367.

307 OECD, 1984. Guideline draft 207 for the testing of chemicals. Earthworm acute toxicity tests.

308 Sochová, I., Hofman, J., & Holoubek, I. (2007). Effects of seven organic pollutants soil nematode  
309 *Caenorhabditis elegans*. *Environment International* 33, 798-804.

310 Spurgeon, D.J., Hopkin, S.P., & Jones, D.T. (1994). Effects of cadmium, copper, lead and zinc on growth,  
311 reproduction and survival of the earthworm *Eisenia fetida* (savigny): assessing the environmental  
312 impact of the point-sources metal contamination in terrestrial ecosystems. *Environmental Pollution* 84,  
313 123-130.

314 Švábenský, R., Oravec, M., & Šimek, Z. (2009). Determination of polycyclic aromatic nitrogen heterocycles in  
315 soil using liquid chromatography/tandem mass spectrometry. *International Journal of Environment and*  
316 *Analytical Chemistry* 89, 167-181.

317 Sverdrup, L.E., Krogh, P.H., Nielsen, T., & Stenersen, J. (2002). Relative sensitivity of three terrestrial  
318 invertebrate tests to polycyclic aromatic compounds. *Environmental Toxicology and Chemistry* 21,  
319 1927-1933.

320 Webber, M.D. (1994). Industrial organic compounds in selected Canadian municipal sludges and agricultural  
321 soils. Final report for Land Resource Division, Centre for Land and Biological Resources Research  
322 Agriculture and Agri-Food Canada. Burlington: Environment Canada, Wastewater Technology Centre,  
323 pp. 100.

324 Widdows, J., & Donkin, P. (1989). The application of combined tissue residue chemistry and physiological  
325 measurements of mussels (*Mytilus edulis*) for the assessment of environmental pollution.  
326 *Hydrobiologia*, 188/189, 455-461.

327 Willumsen, P.A., Johansen, J.E., Karlson, U., & Hansen, B.M. (2005). Isolation and taxonomic affiliation of N-  
328 heterocyclic aromatic hydrocarbon-transforming bacteria. *Applied Microbiology and Biotechnology* 67,  
329 420-428.

330 Willumsen, P.A., Nielsen, J.K., & Karl, U. (2001). Degradation of phenanthrene-analogue azaarenes by  
331 *Mycobacterium gilvum* strain LB307T under aerobic conditions. *Applied Microbiology and*  
332 *Biotechnology* 56, 539-544.

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347






348

349

350

351

352 **Table 1** Test chemicals and their physico-chemical properties

Chemical	Chemical formula	Chemical structure	Molecular mass	Boiling point (°C)	Log $K_{ow}$	Solubility 25°C (mg L <sup>-1</sup> )	% purity
Phenanthrene	C <sub>14</sub> H <sub>10</sub>		178.20	340.00	4.46	1.15	96.00
1,10-Phenanthroline	C <sub>12</sub> H <sub>8</sub> N <sub>2</sub>		180.21	365.10	2.51	30.64	99.00
1,7-Phenanthroline	C <sub>12</sub> H <sub>8</sub> N <sub>2</sub>		180.21	365.10	2.51	30.64	99.00
4,7-Phenanthroline	C <sub>12</sub> H <sub>8</sub> N <sub>2</sub>		180.21	361.20	2.40	38.04	98.00
Benzo[h]-quinoline	C <sub>12</sub> H <sub>9</sub> N		179.20	339.00	3.43	78.70	97.00

353 Source: [www.chemspider.com/Chemical-Structure](http://www.chemspider.com/Chemical-Structure), Anyanwu and Semple (2015a, b, c ; 2016 a, b)

354

355

356

357

358

359

360

361

362

363

364

365

366

367 **Table 2** Summary of the effect of single, binary and quinary mixtures of phenanthrene and its N-PAH analogues  
 368 on the survival (LC<sub>50</sub>) and biomass (EC<sub>50</sub>) of *E. fetida* after 21 d incubation in soil

Chemical	LC <sub>50</sub> (mg kg <sup>-1</sup> )	EC <sub>50</sub> (mg kg <sup>-1</sup> )	R <sup>2</sup> (biomass)
Phen	329.30 (ND)	148.40 (ND)	-0.976
1,10-Phen	ND	102.30 (ND)	-0.686
1,7-Phen	219.80 (ND)	93.90 (ND)	-1.00
4,7-Phen	ND	13.30 (ND)	-0.690
B[h]Q	219.80 (ND)	17.00 (0-88.10)	-0.972
1,7-Phen + Phen	219.80 (ND)	63.80 (ND)	-0.979
4,7-Phen + Phen	219.80 (ND)	148.40 (ND)	-0.956
B[h]Q + Phen	219.80 (ND)	63.80 (5.5-526.70)	-0.979
Phen + N-PAHs	148.40 (ND)	24.20 (ND)	-0.961

369 Values show LC<sub>50</sub>, EC<sub>50</sub> and 95% confidence interval (in parenthesis), ND = (not determined), n = 5.

370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386

387 **Table 3** LC<sub>50</sub> and EC<sub>50</sub> ranges of phenanthrene and its nitrogen-containing analogues on survival and growth of  
 388 soil organisms

Chemical	Test organism	Test duration	LC <sub>50</sub> (mg kg <sup>-1</sup> )	EC <sub>50</sub> (mg kg <sup>-1</sup> )	Reference
	<i>E. veneta</i>	28 d	134 (ND)	94 (64-125)	Sverdrup <i>et al.</i> 2002
Phen	<i>E. cryptius</i>	28 d	1708 (1494-1920)	869 (627-1110)	Kobetičová <i>et al.</i> 2011
	<i>E. fetida</i>	21 d	329.3 (ND)	148.4 (ND)	This study
1,10-Phen	<i>E. fetida</i>	4 wks	1500<LC <sub>50</sub> <2000	1033 (986-1097)	Kobetičová <i>et al.</i> 2008
	<i>E. cryptius</i>	28 d	ND	798 (653-939)	Kobetičová <i>et al.</i> 2011
	<i>E. fetida</i>	21 d	ND	102.3 (ND)	This study
1,7-Phen	<i>E. fetida</i>	21 d	219.8 (ND)	93.9 (ND)	This study
4,7-Phen	<i>E. fetida</i>	21 d	ND	13.3 (ND)	This study
B[h]Q	<i>E. fetida</i>	21 d	219.8 (ND)	17.0 (0-88.1)	This study

389 Values show LC<sub>50</sub>, EC<sub>50</sub> and 95% confidence interval (in parenthesis), ND = (not determined).

390

391

392

393

394

395


396

397

398

399





400

401 **Fig. 1** Behavioral index of *E. fetida* exposed to 500 mg kg<sup>-1</sup> of single, binary and quinary mixtures of  
 402 phenanthrene and its N-PAHs analogues in soil during 21 d incubation. Data shows: Phen (●); B[h]Q (▼); 1,7-  
 403 Phen (■); 1,10 (◇) and 4,7-Phen (▲) (single mixture) (**Fig. 1a**). B[h]Q + Phen (●); 1,7-Phen + Phen (○); 1,10-  
 404 Phen + Phen (▼); 4,7-Phen + Phen (Δ) and Phen + NPAHs (■) (binary and quinary mixtures) (**Fig. 1b**).

405

406

407

408

409

410

411

412

413

414

415


416

417

418

419

420



421

422 **Fig. 2** Survival (%) of *E. fetida* exposed to single, binary and quinary mixtures of phenanthrene and its N-PAH  
 423 analogues in soil at 0 (control), 10, 100 and 500 mg kg<sup>-1</sup> after 21 d incubation. Data shows: 1,10-Phen(●); 1,7-  
 424 Phen (○); 4,7-Phen (▼); B[h]Q (Δ) and Phen (■) (single amendment) (**Fig. 2a**). 1,10-Phen + Phen (●); 1,7-Phen  
 425 + Phen (○); 4,7-Phen + Phen (▼); B[h]Q + Phen (Δ) and Phen + NPAHs (■) (binary and ternary mixtures) (**Fig.**  
 426 **2b**).

427

428

429

430

431

432

433


434

435


436

437


438



439





440





441 **Fig. 3** Mean biomass (g) of *E. fetida* exposed to single amendments of phenanthrene and its N-PAH analogues  
442 in soil at 0 (control), 10, 100 and 500 mg kg<sup>-1</sup> after 21 d incubation. Data shows: weight before exposure (●) and  
443 weight after exposure (▼).


444



445



446



447

448 **Fig. 4** Mean biomass (g) of *E. fetida* exposed to binary and quinary mixtures of phenanthrene and its N-PAH  
 449 analogues in soil at 0 (control), 10, 100 and 500 mg kg<sup>-1</sup> after 21 d incubation. Data shows: weight before  
 450 exposure (●) and weight after exposure (▼).